



SCUOLA INTERNAZIONALE SUPERIORE DI STUDI AVANZATI

SISSA Digital Library

Superconformal indices at large N and the entropy of AdS5 x SE5 black holes

Original

Superconformal indices at large N and the entropy of AdS5 x SE5 black holes / Benini, F., Colombo, E., Soltani, S., Zaffaroni, A., Zhang, Z.. - In: CLASSICAL AND QUANTUM GRAVITY. - ISSN 0264-9381. - 37:21(2020), pp. 1-62. [10.1088/1361-6382/abb39b]

Availability:

This version is available at: 20.500.11767/117576 since: 2021-01-15T19:52:27Z

Publisher:

Published

DOI:10.1088/1361-6382/abb39b

Terms of use:

Testo definito dall'ateneo relativo alle clausole di concessione d'uso

Publisher copyright

IOS Press

This version is available for education and non-commercial purposes.

note finali coverpage

(Article begins on next page)

Superconformal indices at large N and the entropy of $\text{AdS}_5 \times \text{SE}_5$ black holes

Francesco Benini^{1,2,3}, Edoardo Colombo⁴, Saman Soltani^{1,2},
Alberto Zaffaroni^{4,5}, and Ziruo Zhang^{1,2}

¹ *SISSA, Via Bonomea 265, 34136 Trieste, Italy*

² *INFN, Sezione di Trieste, Via Valerio 2, 34127 Trieste, Italy*

³ *ICTP, Strada Costiera 11, 34151 Trieste, Italy*

⁴ *Dipartimento di Fisica, Università di Milano-Bicocca, I-20126 Milano, Italy*

⁵ *INFN, Sezione di Milano-Bicocca, I-20126 Milano, Italy*

Abstract

The large N limit of the four-dimensional superconformal index was computed and successfully compared to the entropy of a class of AdS_5 black holes only in the particular case of equal angular momenta. Using the Bethe Ansatz formulation, we compute the index at large N with arbitrary chemical potentials for all charges and angular momenta, for general $\mathcal{N} = 1$ four-dimensional conformal theories with a holographic dual. We conjecture and bring some evidence that a particular universal contribution to the sum over Bethe vacua dominates the index at large N . For $\mathcal{N} = 4$ SYM, this contribution correctly leads to the entropy of BPS Kerr-Newman black holes in $\text{AdS}_5 \times S^5$ for arbitrary values of the conserved charges, thus completing the microscopic derivation of their microstates. We also consider theories dual to $\text{AdS}_5 \times \text{SE}_5$, where SE_5 is a Sasaki-Einstein manifold. We first check our results against the so-called universal black hole. We then explicitly construct the near-horizon geometry of BPS Kerr-Newman black holes in $\text{AdS}_5 \times T^{1,1}$, charged under the baryonic symmetry of the conifold theory and with equal angular momenta. We compute the entropy of these black holes using the attractor mechanism and find complete agreement with the field theory predictions.

Contents

1	Introduction	1
2	The index of $\mathcal{N} = 4$ SYM at large N	5
2.1	The building block	8
2.2	The index and the entropy function	11
3	The index of quiver theories with a holographic dual	12
3.1	Example: the conifold	16
3.2	Example: toric models	19
3.3	The entropy function	23
4	The universal rotating black hole	24
5	AdS₅ Kerr-Newman black holes in $T^{1,1}$	26
5.1	Reduction from 5d to 4d and the attractor mechanism	27
5.2	Example: the conifold	31
A	Subleading effect of simplifications	36
A.1	Simplifications of the building block	36
A.2	$SU(N)$ vs. $U(N)$ holonomies	39
A.3	Generic N	43
B	5d $\mathcal{N} = 2$ Abelian gauged supergravity	44
B.1	Conifold truncation in the general framework	51
C	4d $\mathcal{N} = 2$ Abelian gauged supergravity	52
D	Reduction with background gauge fields	56
D.1	Reduction of the conifold truncation	61
E	Black hole charges and their reduction	61
E.1	Baryonic charge quantization in the conifold theory	65

1 Introduction

There has been some progress in the microscopic explanation of the entropy of BPS asymptotically-AdS black holes, initiated with the counting of microstates of static magnetically-charged black holes in $\text{AdS}_4 \times S^7$ [1–3] and continued, more recently, with the counting for

Kerr-Newman black holes in $\text{AdS}_5 \times S^5$ [4–6].¹ The latter result, in particular, shed light on a long-standing puzzle. The holographic description of electrically-charged and rotating BPS black holes in $\text{AdS}_5 \times S^5$ is in terms of 1/16 BPS states of the dual four-dimensional $\mathcal{N} = 4$ super-Yang-Mills (SYM) boundary theory on S^3 . These states are counted (with sign) by the superconformal index [11–13], and one would expect that the contribution from black holes saturates it at large N . However, the large N computation of the superconformal index performed in [12] gave a result of order one, while the entropy for the black holes is of order N^2 , suggesting a large cancellation between bosonic and fermionic BPS states. On the other hand, it has been argued in [5, 6] that non-trivial complex phases of the fugacities for flavor symmetries can obstruct such a cancellation between bosonic and fermionic states — as already observed in [1, 3] — and that the entropy of Kerr-Newman black holes is indeed correctly captured by the index for complex values of the chemical potentials associated with electric charges and angular momenta.

The family of $\text{AdS}_5 \times S^5$ supersymmetric black holes found in [14–18] depends on three charges Q_a associated with the Cartan subgroup of the internal isometry $\text{SO}(6)$, and two angular momenta J_i in AdS_5 , subject to a non-linear constraint.² The entropy can be written as the value at the critical point (*i.e.*, as a Legendre transform) of the function [21]

$$\mathcal{S}(X_a, \tau, \sigma) = -i\pi N^2 \frac{X_1 X_2 X_3}{\tau \sigma} - 2\pi i \left(\sum_{a=1}^3 X_a Q_a + \tau J_1 + \sigma J_2 \right) \quad (1.1)$$

with the constraint $X_1 + X_2 + X_3 - \tau - \sigma = \pm 1$, where N is the number of colors of the dual 4d $\mathcal{N} = 4$ $\text{SU}(N)$ SYM theory. The same entropy function can also be obtained by computing the zero-temperature limit of the on-shell action of a class of supersymmetric but non-extremal complexified Euclidean black holes [4, 22]. The two constraints with \pm sign lead to the same value for the entropy, which is real precisely when the non-linear constraint on the black hole charges is imposed. The parameters X_a , τ and σ are chemical potentials for the conserved charges Q_a and J_i and can also be identified with the parameters the superconformal index depends on. With this identification, we expect that the entropy $S(Q_a, J_1, J_2)$ is just the constrained Legendre transform of $\log \mathcal{I}(X_a, \tau, \sigma)$, where $\mathcal{I}(X_a, \tau, \sigma)$ is the superconformal index.

Up to now, the entropy of $\text{AdS}_5 \times S^5$ Kerr-Newman black holes has been derived from the superconformal index and shown to be in agreement with (1.1) only in particular limits. In [5], the entropy was derived for large black holes (whose size is much larger than the AdS

¹The microstate counting for rotating black holes in AdS_4 was performed in [7–9]. Those results have been extended to other compactifications and other dimensions. See [10] for a more complete list of references.

²Supersymmetric hairy black holes depending on all charges have been recently found in [19, 20], but their entropy seems to be parametrically smaller in the range of parameters where our considerations apply.

radius) using a Cardy limit of the superconformal index where $\text{Im}(X_a), \tau, \sigma \ll 1$. In [6], the entropy was instead derived in the large N limit in the case of black holes with equal angular momenta, $J_1 = J_2$.³ The large N limit has been evaluated by writing the index as a sum over Bethe vacua [24], an approach that has been successful for AdS black holes in many other contexts.

It is one of the purposes of this paper to extend the derivation of [6] to the case of unequal angular momenta, thus providing a large N microscopic counting of the microstates of BPS Kerr-Newman black holes in $\text{AdS}_5 \times S^5$ for arbitrary values of the conserved charges. We will make use of the Bethe Ansatz formulation of the superconformal index derived for $\tau = \sigma$ in [25] and generalized to unequal angular chemical potentials in [24]. This formulation allows us to write the index as a sum over the solutions to a set of Bethe Ansatz Equations (BAEs) — whose explicit form and solutions have been studied in [6, 26–30] — and over some auxiliary integer parameters m_i . We expect that, in the large N limit, one particular solution dominates the sum.⁴ We will show that the “basic solution” to the BAEs, already used in [6], correctly reproduces the entropy of black holes in the form (1.1) for a choice of integers m_i . We stress that our result comes from a single contribution to the index, which is an infinite sum. Such a contribution might not be the dominant one — and so our estimate of the index might be incorrect — in some regions of the space of chemical potentials. It is known from the analysis in [6] that when the charges become smaller than a given threshold, new solutions take over and dominate the asymptotic behavior of the index. This suggests the existence of a rich structure where other black holes might also contribute. However, we conjecture and we will bring some evidence that the contribution of the basic solution is the dominant one in the region of the space of chemical potentials corresponding to sufficiently large charges.

We will also extend the large N computation of the index to a general class of superconformal theories dual to $\text{AdS}_5 \times \text{SE}_5$, where SE_5 is a five-dimensional Sasaki-Einstein manifold. The analysis for $J_1 = J_2$ was already performed in [28]. For toric holographic quiver gauge theories, we find a prediction for the entropy of black holes in $\text{AdS}_5 \times \text{SE}_5$ in the form of the entropy function

$$\mathcal{S}(X_a, \tau, \sigma) = -\frac{i\pi N^2}{6} \sum_{a,b,c}^D C_{abc} \frac{X_a X_b X_c}{\tau \sigma} - 2\pi i \left(\sum_{a=1}^D X_a Q_a + \tau J_1 + \sigma J_2 \right), \quad (1.2)$$

with the constraint $\sum_{a=1}^D X_a - \tau - \sigma = \pm 1$, in terms of chemical potentials X_a for a basis of independent R-symmetries R_a . The coefficients $C_{abc} N^2 = \frac{1}{4} \text{Tr} R_a R_b R_c$ are the 't Hooft

³The same result has been later reproduced with a different approach in [23].

⁴It is argued in [29] that there exist families of continuous solutions. This does not affect our argument provided the corresponding contribution to the index is subleading.

anomaly coefficients for this basis of R-symmetries. The form of the entropy function (1.2) was conjectured in [31] and reproduced for various toric models in the special case $\tau = \sigma$ in [28]. We will give a general derivation, valid for all toric quivers and even more. We will also show that both constraints in (1.2), which lead to the same value for the entropy, naturally arise from the index in different regions of the space of chemical potentials. The function (1.2) was also derived in the Cardy limit in [32].

In the last part of the paper we will provide some evidence that (1.2) correctly reproduces the entropy of black holes in $\text{AdS}_5 \times \text{SE}_5$. We first check that our formula correctly reproduce the entropy of the *universal* black hole that arises as a solution in five-dimensional minimal gauged supergravity, and, as such, can be embedded in any $\text{AdS}_5 \times \text{SE}_5$ compactification. It corresponds to a black hole with electric charges aligned with the exact R-symmetry of the dual superconformal field theory and with arbitrary angular momenta J_1 and J_2 . Since the solution is universal, the computation can be reduced to that of $\mathcal{N} = 4$ SYM and it is almost trivial. More interesting are black holes with general electric charges. Unfortunately, to the best of our knowledge, there are no available such black hole solutions in compactifications based on Sasaki-Einstein manifolds SE_5 other than S^5 . To overcome this obstacle, we will explicitly construct the near-horizon geometry of supersymmetric black holes in $\text{AdS}_5 \times T^{1,1}$ with equal angular momenta and charged under the baryonic symmetry of the dual Klebanov-Witten theory [33]. Luckily, the background $\text{AdS}_5 \times T^{1,1}$ admits a consistent truncation to a five-dimensional gauged supergravity containing the massless gauge field associated to the baryonic symmetry [34–36]. We then use the strategy suggested in [21]: a rotating black hole in five dimensions with $J_1 = J_2$ can be dimensionally reduced along the Hopf fiber of the horizon three-sphere to a static solution of four-dimensional $\mathcal{N} = 2$ gauged supergravity. We will explicitly solve the BPS equations [37–39] for the horizon of static black holes with the appropriate electric and magnetic charges in $\mathcal{N} = 2$ gauge supergravity in four dimensions. The main complication is the presence of hypermultiplets. By solving the hyperino equations at the horizon, we will be able to recast all other supersymmetric conditions as a set of attractor equations, and we will show that these are equivalent to the extremization of (1.2) for the Klebanov-Witten theory with $\tau = \sigma$. This provides a highly non-trivial check of our result, and the conjecture that the basic solution to the BAEs dominates the index.

The paper is organized as follows. In Section 2 we review the setting introduced in [6] and we evaluate the large N contribution of the “basic solution” to the BAEs to the index for generic angular fugacities. We show that it correctly captures the semiclassical Bekenstein-Hawking entropy of BPS black holes in $\text{AdS}_5 \times S^5$. In Section 3 we discuss the generalization of this result to general toric quiver theories and find agreement with the entropy function prediction (1.2) in certain corners of the space of chemical potentials. In Section 4 we discuss the particular case of the universal black hole, which can be embedded in all string

and M-theory supersymmetric compactifications with an AdS_5 factor. In Section 5 we match formula (1.2) with the entropy of a supersymmetric black hole in $\text{AdS}_5 \times T^{1,1}$, whose near-horizon geometry we explicitly construct. Technical computations as well as some review material can be found in several appendices.

Note added: while this work was ready to be posted on the arXiv, the preprint [40] appeared, which discusses the index in the particular case $\tau = \sigma$ using a different approach.

2 The index of $\mathcal{N} = 4$ SYM at large N

We are interested in evaluating the large N limit of the superconformal index of 4d $\mathcal{N} = 1$ holographic theories. We will consider in this section the simplest example, namely $\mathcal{N} = 4$ $\text{SU}(N)$ SYM. The superconformal index counts (with sign) the 1/16 BPS states of the theory on $\mathbb{R} \times S^3$ that preserve one complex supercharge \mathcal{Q} . These states are characterized by two angular momenta $J_{1,2}$ on S^3 and three R-charges for $\text{U}(1)^3 \subset \text{SO}(6)_R$. We write $\mathcal{N} = 4$ SYM in $\mathcal{N} = 1$ notation in terms of a vector multiplet and three chiral multiplets Φ_I and introduce a symmetric basis of R-symmetry generators $R_{1,2,3}$ such that R_I assigns R-charge 2 to Φ_I and zero to Φ_J with $J \neq I$. The index is defined by the trace [11, 12]

$$\mathcal{I}(p, q, v_1, v_2) = \text{Tr} (-1)^F e^{-\beta\{\mathcal{Q}, \mathcal{Q}^\dagger\}} p^{J_1 + \frac{\tau}{2}} q^{J_2 + \frac{\tau}{2}} v_1^{q_1} v_2^{q_2} , \quad (2.1)$$

in terms of two flavor generators $q_{1,2} = \frac{1}{2}(R_{1,2} - R_3)$ commuting with \mathcal{Q} , and the R-charge $r = \frac{1}{3}(R_1 + R_2 + R_3)$. Notice that $(-1)^F = e^{2\pi i J_{1,2}} = e^{i\pi R_{1,2,3}}$. Here p, q, v_I with $I = 1, 2$ are complex fugacities associated to the various quantum numbers, while the corresponding chemical potentials τ, σ, ξ_I are defined by

$$p = e^{2\pi i \tau} , \quad q = e^{2\pi i \sigma} , \quad v_I = e^{2\pi i \xi_I} . \quad (2.2)$$

The index is well-defined for $|p|, |q| < 1$.

It is convenient to redefine the flavor chemical potentials in terms of

$$\Delta_I = \xi_I + \frac{\tau + \sigma}{3} \quad \text{for} \quad I = 1, 2 . \quad (2.3)$$

It is also convenient to introduce an auxiliary chemical potential Δ_3 such that

$$\tau + \sigma - \Delta_1 - \Delta_2 - \Delta_3 \in 2\mathbb{Z} + 1 , \quad (2.4)$$

and use the corresponding fugacities

$$y_I = e^{2\pi i \Delta_I} . \quad (2.5)$$

The index then takes the more transparent form

$$\mathcal{I} = \text{Tr}_{\text{BPS}} p^{J_1} q^{J_2} y_1^{R_1/2} y_2^{R_2/2} y_3^{R_3/2} . \quad (2.6)$$

It shows that the constrained fugacities p, q, y_I with $I = 1, 2, 3$ are associated to the angular momenta $J_{1,2}$ and the charges $Q_I \equiv \frac{1}{2}R_I$.

Our starting point is the so-called Bethe Ansatz formulation of the superconformal index [25, 24]. The special case that the two angular chemical potentials are equal, $\tau = \sigma$, was already studied in [6] (see also [29]). Here we take them unequal. The formula of [24] can be applied when the ratio between the two angular chemical potentials is a rational number.⁵ We thus set

$$\tau = a\omega , \quad \sigma = b\omega \quad \text{with} \quad \text{Im } \omega > 0 \quad (2.7)$$

and with $a, b \in \mathbb{N}$ coprime positive integers. We call $\mathbb{H} = \{\omega \mid \text{Im } \omega > 0\}$ the upper half-plane. We then have the fugacities

$$h = e^{2\pi i\omega} , \quad p = h^a = e^{2\pi i\tau} , \quad q = h^b = e^{2\pi i\sigma} \quad \text{with} \quad |h|, |p|, |q| < 1 . \quad (2.8)$$

The formula in [24] allows us to write the superconformal index as a sum over the solutions to a set of Bethe Ansatz Equations (BAEs). Explicitly, the index reads

$$\mathcal{I} = \kappa_N \sum_{\hat{u} \in \text{BAE}} \mathcal{Z}_{\text{tot}} H^{-1} \Big|_{\hat{u}} . \quad (2.9)$$

The expressions of κ_N , H and \mathcal{Z}_{tot} for a generic $\mathcal{N} = 1$ theory are given in [24]. Here, we specialize them to $\mathcal{N} = 4$ $\text{SU}(N)$ SYM. The quantity

$$\kappa_N = \frac{1}{N!} \left(\frac{(p; p)_\infty (q; q)_\infty \tilde{\Gamma}(\Delta_1; \tau, \sigma) \tilde{\Gamma}(\Delta_2; \tau, \sigma)}{\tilde{\Gamma}(\Delta_1 + \Delta_2; \tau, \sigma)} \right)^{N-1} \quad (2.10)$$

is a prefactor written in terms of the elliptic gamma function $\tilde{\Gamma}$ and the Pochhammer symbol:

$$\tilde{\Gamma}(u; \tau, \sigma) \equiv \Gamma(z; p, q) = \prod_{m,n=0}^{\infty} \frac{1 - p^{m+1}q^{n+1}/z}{1 - p^m q^n z} , \quad (z; q)_\infty = \prod_{n=0}^{\infty} (1 - zq^n) , \quad (2.11)$$

where $z = e^{2\pi i u}$. The sum in (2.9) is over the solution set to the following BAEs:⁶

$$1 = Q_i(u; \Delta, \omega) \equiv e^{2\pi i(\lambda + 3\sum_j u_{ij})} \prod_{j=1}^N \frac{\theta_0(u_{ji} + \Delta_1; \omega) \theta_0(u_{ji} + \Delta_2; \omega) \theta_0(u_{ji} - \Delta_1 - \Delta_2; \omega)}{\theta_0(u_{ij} + \Delta_1; \omega) \theta_0(u_{ij} + \Delta_2; \omega) \theta_0(u_{ij} - \Delta_1 - \Delta_2; \omega)} \quad (2.12)$$

⁵This might sound like a strong limitation. However, the index (2.6) is invariant under integer shifts of τ and σ compatible with (2.4). As proven in [24], the set of complex number pairs $\{\tau, \sigma\} \in \mathbb{H}^2$ (two copies of the upper half-plane) whose ratio becomes a (real) rational number after some integer shifts of τ and σ , is dense in \mathbb{H}^2 . Thus, by continuity, the formula of [24] fixes the large N limit of the superconformal index for generic complex chemical potentials.

⁶The Bethe operators Q_i should not be confused with the charges Q_I introduced before.

written in terms of $u_{ij} = u_i - u_j$ with $i, j = 1, \dots, N$ and the theta function

$$\theta_0(u; \omega) = (z; h)_\infty (h/z; h)_\infty . \quad (2.13)$$

The unknowns are the ‘‘complexified $SU(N)$ holonomies’’, which are expressed here in terms of $U(N)$ holonomies u_i further constrained by

$$\sum_{i=1}^N u_i = 0 \pmod{\mathbb{Z}} , \quad (2.14)$$

as well as a ‘‘Lagrange multiplier’’ λ . The $SU(N)$ holonomies are to be identified with the first $N - 1$ variables $u_{i=1, \dots, N-1}$. As unknowns in the BAEs, they are subject to the identifications

$$u_i \sim u_i + 1 \sim u_i + \omega , \quad (2.15)$$

meaning that each one of them naturally lives on a torus of modular parameter ω . Instead, the last holonomy u_N is determined by the constraint (2.14). The relation between $SU(N)$ and $U(N)$ holonomies will be further clarified in Appendix A.2. The prescription in (2.9) is to sum over all the inequivalent solutions on the torus [24]. The function H is the Jacobian

$$H = \det \left[\frac{1}{2\pi i} \frac{\partial(Q_1, \dots, Q_N)}{\partial(u_1, \dots, u_{N-1}, \lambda)} \right] . \quad (2.16)$$

Finally, the function \mathcal{Z}_{tot} is the following sum over a set of integers $m_i = 1, \dots, ab$:

$$\mathcal{Z}_{\text{tot}} = \sum_{\{m_i\}=1}^{ab} \mathcal{Z}(u - m\omega; \tau, \sigma) , \quad (2.17)$$

where \mathcal{Z} , for $\mathcal{N} = 4$ $SU(N)$ SYM, reads

$$\mathcal{Z} = \prod_{\substack{i,j=1 \\ i \neq j}}^N \frac{\tilde{\Gamma}(u_{ij} + \Delta_1; \tau, \sigma) \tilde{\Gamma}(u_{ij} + \Delta_2; \tau, \sigma)}{\tilde{\Gamma}(u_{ij} + \Delta_1 + \Delta_2; \tau, \sigma) \tilde{\Gamma}(u_{ij}; \tau, \sigma)} . \quad (2.18)$$

The sum in (2.17) freely varies over the first $N - 1$ integers $m_{i=1, \dots, N-1}$ as indicated, while m_N is determined by the constraint

$$\sum_{i=1}^N m_i = 0 . \quad (2.19)$$

More details can be found in [24, 6]. In the following, when a double sum starts from 1 we will leave it implicit.

2.1 The building block

We will show that one particular contribution to the sums in (2.9) and (2.17) alone reproduces the entropy function of [21] and therefore it captures the Bekenstein-Hawking entropy of BPS black holes in $\text{AdS}_5 \times S^5$. To that aim, we are interested in the contribution from the so-called “basic solution” to the BAEs [26, 27, 6], namely

$$u_i = \frac{N-i}{N} \omega + \bar{u}, \quad u_{ij} \equiv u_i - u_j = \frac{j-i}{N} \omega, \quad \lambda = \frac{N-1}{2}. \quad (2.20)$$

Here \bar{u} is fixed by enforcing the constraint (2.14). We also consider the contribution from a particular choice for the integers $\{m_j\}$:

$$m_j \in \{1, \dots, ab\} \quad \text{such that} \quad m_j = j \pmod{ab}. \quad (2.21)$$

Note that this choice for $\{m_j\}$ does not satisfy the constraint (2.19). Nevertheless, we show in Appendix A.2 that this does not affect the contribution to leading order in N , in the sense that changing the single entry m_N has a subleading effect.

Now, the crucial technical point is to evaluate the following basic building block:

$$\Psi = \sum_{i \neq j}^N \log \tilde{\Gamma} \left(\Delta + \omega \frac{j-i}{N} + \omega(m_j - m_i); a\omega, b\omega \right) \quad (2.22)$$

for $N \rightarrow \infty$. Here Δ plays the role of an electric chemical potential. In order to simplify the discussion, we assume that N is a multiple of ab , *i.e.*, we take $N = ab\tilde{N}$. As we show in Appendix A.3 this assumption can be removed without affecting the leading behavior at large N .

We make use of the following identity [41]:

$$\tilde{\Gamma}(u; \tau, \sigma) = \prod_{r=0}^{a-1} \prod_{s=0}^{b-1} \tilde{\Gamma}(u + (r\tau + s\sigma); a\tau, b\sigma) \quad (2.23)$$

for any $\tau, \sigma \in \mathbb{H}$ and any $a, b \in \mathbb{N}$. This is immediate to prove exploiting the infinite product expression of $\tilde{\Gamma}$. Now, exchanging $a \leftrightarrow b$ and $r \leftrightarrow s$ in the formula, and then setting $\tau \rightarrow a\omega$, $\sigma \rightarrow b\omega$, we obtain the formula of [42]:

$$\tilde{\Gamma}(u; a\omega, b\omega) = \prod_{r=0}^{a-1} \prod_{s=0}^{b-1} \tilde{\Gamma}(u + (as + br)\omega; ab\omega, ab\omega). \quad (2.24)$$

Going back to Ψ , we can thus write

$$\Psi = \sum_{r=0}^{a-1} \sum_{s=0}^{b-1} \sum_{i \neq j}^N \log \tilde{\Gamma} \left(\Delta + \omega \frac{j-i}{N} + \omega(m_j - m_i + as + br); ab\omega, ab\omega \right). \quad (2.25)$$

Let us set $i = \gamma ab + c$, $j = \delta ab + d$ with $\gamma, \delta = 0, \dots, \tilde{N} - 1$ and $c, d = 1, \dots, ab$. Then

$$\Psi = \sum_{r=0}^{a-1} \sum_{s=0}^{b-1} \underbrace{\sum_{\gamma, \delta=0}^{\tilde{N}-1} \sum_{c, d=1}^{ab}}_{\text{s.t. } i \neq j} \log \tilde{\Gamma} \left(\Delta + \omega \frac{\delta - \gamma}{\tilde{N}} + \omega \frac{d - c}{N} + \omega(d - c + as + br); ab\omega, ab\omega \right). \quad (2.26)$$

We will now perform two simplifications, and prove in Appendix A.1 that their effect is of subleading order at large N . More precisely, Ψ is of order N^2 while the two simplifications modify it at most at order N if $\mathbb{Im}(\Delta/\omega) \notin \mathbb{Z} \times \mathbb{Im}(1/\omega)$, or at most at order $N \log N$ if $\Delta = 0$. First, we substitute the condition $i \neq j$ with the condition $\gamma \neq \delta$ in the summation. Second and more importantly, we drop the term $\omega(d - c)/N$ in the argument. We then redefine $c \rightarrow ab - c$, $d \rightarrow d + 1$, $\gamma \rightarrow \gamma - 1$, $\delta \rightarrow \delta - 1$ and obtain

$$\Psi \simeq \sum_{r=0}^{a-1} \sum_{s=0}^{b-1} \sum_{\gamma \neq \delta}^{\tilde{N}} \sum_{c, d=0}^{ab-1} \log \tilde{\Gamma} \left(\Delta + \omega \frac{\delta - \gamma}{\tilde{N}} + \omega(d + c + 1 - ab + as + br); ab\omega, ab\omega \right) \quad (2.27)$$

where \simeq means equality at leading order in N . At this point we can resum over c, d using (2.23) (with $\tau, \sigma \rightarrow \omega$ and $a, b \rightarrow ab$):

$$\Psi \simeq \sum_{r=0}^{a-1} \sum_{s=0}^{b-1} \sum_{\gamma \neq \delta}^{\tilde{N}} \log \tilde{\Gamma} \left(\Delta + \omega \frac{\delta - \gamma}{\tilde{N}} + \omega(1 - ab + as + br); \omega, \omega \right). \quad (2.28)$$

We recall the large N limit computed in [6]:

$$\sum_{i \neq j}^N \log \tilde{\Gamma} \left(\Delta + \omega \frac{j - i}{N}; \omega, \omega \right) = -\pi i N^2 \frac{B_3([\Delta]'_\omega - \omega)}{3\omega^2} + \mathcal{O}(N) \quad (2.29)$$

valid for $\mathbb{Im}(\Delta/\omega) \notin \mathbb{Z} \times \mathbb{Im}(1/\omega)$. Here $B_3(x)$ is a Bernoulli polynomial:

$$B_3(x) = x \left(x - \frac{1}{2} \right) (x - 1). \quad (2.30)$$

It has the property that $B_3(1 - x) = -B_3(x)$. The function $[\Delta]'_\omega$ was defined in [6] in the following way:

$$[\Delta]'_\omega = \left\{ z \mid z = \Delta \bmod 1, \ 0 > \mathbb{Im} \left(\frac{z}{\omega} \right) > \mathbb{Im} \left(\frac{1}{\omega} \right) \right\}. \quad (2.31)$$

This function is only defined for $\mathbb{Im}(\Delta/\omega) \notin \mathbb{Z} \times \mathbb{Im}(1/\omega)$, it is continuous in each open connected domain, and it is periodic by construction under $\Delta \rightarrow \Delta + 1$. In the following we will also use the function $[\Delta]_\omega = [\Delta]'_\omega - 1$,

$$[\Delta]_\omega = \left\{ z \mid z = \Delta \bmod 1, \ \mathbb{Im} \left(-\frac{1}{\omega} \right) > \mathbb{Im} \left(\frac{z}{\omega} \right) > 0 \right\}. \quad (2.32)$$




Figure 1: Fundamental strips for $[\Delta]_\omega$ and $[\Delta]'_\omega$. The function $[\Delta]_\omega$ is the restriction of $\Delta \bmod 1$ to the region $\text{Im}(-1/\omega) > \text{Im}(\Delta/\omega) > 0$ (in yellow, on the left), while $[\Delta]'_\omega$ is the restriction of $\Delta \bmod 1$ to the region $0 > \text{Im}(\Delta/\omega) > \text{Im}(1/\omega)$ (in blue, on the right).

The functions $[\Delta]_\omega$ and $[\Delta]'_\omega$ are the mod 1 reductions of Δ to the fundamental strips shown in Figure 1. Then we use the following formula:

$$\begin{aligned} \frac{1}{ab} \sum_{r=0}^{a-1} \sum_{s=0}^{b-1} B_3(x + \omega(as + br - ab)) &= \\ &= B_3\left(x - \frac{a+b}{2}\omega\right) + \frac{2a^2b^2 - a^2 - b^2}{4} \omega^2 B_1\left(x - \frac{a+b}{2}\omega\right), \end{aligned} \quad (2.33)$$

where $B_1(x) = x - \frac{1}{2}$ is another Bernoulli polynomial — and $B_1(1-x) = -B_1(x)$. Thus

$$\Psi = -\pi i N^2 \frac{B_3\left([\Delta]'_\omega - \frac{\tau+\sigma}{2}\right)}{3\tau\sigma} - \frac{\pi i N^2}{12} \left(2ab - \frac{a}{b} - \frac{b}{a}\right) B_1\left([\Delta]'_\omega - \frac{\tau+\sigma}{2}\right) + \mathcal{O}(N) \quad (2.34)$$

for $\text{Im}(\Delta/\omega) \notin \mathbb{Z} \times \text{Im}(1/\omega)$. As a check, notice that $[\tau + \sigma - \Delta]'_\omega = \tau + \sigma + 1 - [\Delta]'_\omega$. From the properties of $B_{1,3}(x)$ noticed above, it follows

$$\Psi(\tau + \sigma - \Delta) = -\Psi(\Delta) \quad (2.35)$$

at leading order in N . This is in accordance with the inversion formula of the elliptic gamma function:

$$\tilde{\Gamma}(u; \tau, \sigma) = 1/\tilde{\Gamma}(\tau + \sigma - u; \tau, \sigma). \quad (2.36)$$

The case $\Delta = 0$ requires some care, because $[0]_\omega$ is undefined. Taking the limit of Ψ as $\Delta \rightarrow 0$ from the left or the right, one obtains two values that differ by an imaginary quantity. The limit from the right corresponds to taking $[\Delta]'_\omega \rightarrow 0$ in (2.34), while the limit from the left corresponds to $[\Delta]_\omega \rightarrow 0$ (i.e., $[\Delta]'_\omega \rightarrow 1$). The difference is

$$\Psi\Big|_{[\Delta]'_\omega \rightarrow 0} - \Psi\Big|_{[\Delta]_\omega \rightarrow 0} = \frac{i\pi N^2}{6} \left(3 + ab + \frac{a}{b} + \frac{b}{a}\right). \quad (2.37)$$

Since Ψ is in any case ambiguous by shifts of $2\pi i$ because it is a logarithm, only the remainder modulo $2\pi i$ is meaningful but this is an order 1 quantity which can be neglected. In fact it

turns out that, with $N = ab\tilde{N}$, the quantity on the right-hand-side of (2.37) is always an integer multiple of $i\pi\tilde{N}$, and so its exponential is a sign. We should also notice that, for $\Delta = 0$, our approximation gets corrections at order $N \log N$.

2.2 The index and the entropy function

We are now ready to put all the ingredients together. Our working assumption is that, in the large N limit, the index (2.9) is dominated by the basic solution (2.20) and the choice of integers (2.21). Some evidence that the basic solution dominates the index for $\tau = \sigma$ has been given in [6] (see also [29]).

The leading contribution to (2.9) originates from \mathcal{Z}_{tot} that can be evaluated using (2.34). Indeed, the term κ_N is manifestly sub-leading. That the contribution of H is also subleading follows from the analysis in [6] for $\tau = \sigma$, since H only depends on the solutions to the BAEs and not explicitly on τ and σ . The large N limit of the index at leading order is then

$$\log \mathcal{I} = \Psi(\Delta_1) + \Psi(\Delta_2) - \Psi(\Delta_1 + \Delta_2) - \Psi(0) , \quad (2.38)$$

where the definition of the last term has an ambiguity of order 1.

Recall that in (2.4) we introduced the auxiliary chemical potential Δ_3 . Notice in particular that the chemical potentials are defined modulo 1. Using the basic properties

$$[\Delta + 1]_\omega = [\Delta]_\omega , \quad [\Delta + \omega]_\omega = [\Delta]_\omega + \omega , \quad [-\Delta]_\omega = -[\Delta]_\omega - 1 , \quad (2.39)$$

we find

$$[\Delta_3]_\omega = \tau + \sigma - 1 - [\Delta_1 + \Delta_2]_\omega . \quad (2.40)$$

It follows from the definition of the function $[\Delta]_\omega$ that $[\Delta_1 + \Delta_2]_\omega = [\Delta_1]_\omega + [\Delta_2]_\omega + n$ where $n = 0$ or $n = 1$. The result then breaks into two cases.

If $[\Delta_1 + \Delta_2]_\omega = [\Delta_1]_\omega + [\Delta_2]_\omega$ then

$$[\Delta_1]_\omega + [\Delta_2]_\omega + [\Delta_3]_\omega - \tau - \sigma = -1 , \quad (2.41)$$

and, using (2.38) and (2.34),

$$\begin{aligned} \log \mathcal{I} &= -\pi i N^2 \frac{[\Delta_1]_\omega [\Delta_2]_\omega (\tau + \sigma - 1 - [\Delta_1]_\omega - [\Delta_2]_\omega)}{\tau \sigma} \\ &= -i\pi N^2 \frac{[\Delta_1]_\omega [\Delta_2]_\omega [\Delta_3]_\omega}{\tau \sigma} . \end{aligned} \quad (2.42)$$

To obtain this formula we used $\Psi(0) = \Psi|_{[\Delta]_\omega \rightarrow 0}$. Notice that the contributions from B_1 cancel out. As we will see in Section 3, this is a consequence of the relation $a = c$ among the two four-dimensional central charges in the large N limit.

If $[\Delta_1 + \Delta_2]_\omega = [\Delta_1]_\omega + [\Delta_2]_\omega + 1$, namely $[\Delta_1 + \Delta_2]'_\omega = [\Delta_1]'_\omega + [\Delta_2]'_\omega$, then

$$[\Delta_1]'_\omega + [\Delta_2]'_\omega + [\Delta_3]'_\omega - \tau - \sigma = 1, \quad (2.43)$$

and

$$\begin{aligned} \log \mathcal{I} &= -\pi i N^2 \frac{[\Delta_1]'_\omega [\Delta_2]'_\omega (\tau + \sigma + 1 - [\Delta_1]'_\omega - [\Delta_2]'_\omega)}{\tau \sigma} \\ &= -i\pi N^2 \frac{[\Delta_1]'_\omega [\Delta_2]'_\omega [\Delta_3]'_\omega}{\tau \sigma}. \end{aligned} \quad (2.44)$$

This time we used $\Psi(0) = \Psi|_{[\Delta]'_\omega \rightarrow 0}$.

As in [6], we can extract the entropy of the dual black holes by taking the Legendre transform of the logarithm of the index. The precise identification of the charges associated with the chemical potentials follows from (2.6). The prediction for the entropy can then be combined into two constrained entropy functions

$$\begin{aligned} \mathcal{S}_\pm(X_I, \tau, \sigma, \Lambda) &= -i\pi N^2 \frac{X_1 X_2 X_3}{\tau \sigma} - 2\pi i \left(\sum_{I=1}^3 X_I Q_I + \tau J_1 + \sigma J_2 \right) \\ &\quad - 2\pi i \Lambda \left(X_1 + X_2 + X_3 - \tau - \sigma \pm 1 \right), \end{aligned} \quad (2.45)$$

where we used a neutral variable X_I to denote either $[\Delta_I]_\omega$ or $[\Delta_I]'_\omega$, we introduced a Lagrange multiplier Λ to enforce the constraint, and we recall that $Q_I = \frac{1}{2} R_I$. This completes our derivation of the entropy of supersymmetric black holes in $\text{AdS}_5 \times S^5$ for general angular momenta and electric charges. The expression (2.45) represents indeed the two entropy functions derived in [21], where it was shown that the (constrained) extremization of (2.45) reproduces the entropy of a black hole of angular momenta J_1 and J_2 and charges Q_I . The two results correspond to the two entropy functions that reproduce the same black hole entropy, and are associated to two Euclidean complex solutions that regularize the black hole horizon [4].

3 The index of quiver theories with a holographic dual

We want to generalize the large N computation of the superconformal index to theories dual to $\text{AdS}_5 \times \text{SE}_5$ compactifications, where SE_5 is a five-dimensional Sasaki-Einstein manifold. We can write general formulæ with very few assumptions. We consider 4d $\mathcal{N} = 1$ theories with $\text{SU}(N)$ gauge groups as well as adjoint and bi-fundamental chiral multiplet fields. To cancel gauge anomalies, the total number of fields transforming in the fundamental representation of a group must be the same as the number of anti-fundamentals. We also

require equality of the conformal central charges $c = a$ in the large N limit, as dictated by holography. Our analysis extends the results found in [28] for equal angular momenta.

We then assume that in the large N limit, as for $\mathcal{N} = 4$ SYM, the leading contribution to the superconformal index comes from the basic solution and the choice of integers $\{m_i\}$ discussed in (2.21). As already shown in [28, 30], the basic solution to the BAEs for $\mathcal{N} = 4$ SYM [26, 27, 6] can easily be extended to quiver gauge theories by setting

$$u_{ij}^{\alpha\beta} \equiv u_i^\alpha - u_j^\beta = \frac{j-i}{N} \omega \quad \alpha, \beta = 1, \dots, G, \quad (3.1)$$

where α, β run over the various gauge groups in the theory and G is the number of gauge groups. Similarly, we choose the integers

$$m_j^\alpha \in \{1, \dots, ab\} \quad \text{such that} \quad m_j^\alpha = j \pmod{ab}. \quad (3.2)$$

Notice in particular that neither $u_{ij}^{\alpha\beta}$ nor m_j^α depend on α, β . As for $\mathcal{N} = 4$ SYM, the contribution of the determinant H to the Bethe Ansatz expansion (2.9) is subleading [28].

Using the general expressions given in [24] and following the logic of Section 2, it is easy to write the large N limit of the leading contribution to the superconformal index of a holographic theory, with adjoint and bi-fundamental chiral fields. We find

$$\log \mathcal{I} = \sum_{i \neq j}^N \left[\sum_{I_{\alpha\beta}} \log \tilde{\Gamma} \left(u_{ij}^{\alpha\beta} - \omega(m_i^\alpha - m_j^\beta) + \Delta_{I_{\alpha\beta}}; \tau, \sigma \right) - \sum_{\alpha=1}^G \log \tilde{\Gamma} \left(u_{ij}^{\alpha\alpha} - \omega(m_i^\alpha - m_j^\beta); \tau, \sigma \right) \right] \quad (3.3)$$

where $z_i^\alpha = e^{2\pi i u_i^\alpha}$ are the gauge fugacities, u_i^α represent the basic solution (3.1) and m_i^α are given in (3.2). The sum over $I_{\alpha\beta}$ is over all adjoint (if $\alpha = \beta$) and bi-fundamental (if $\alpha \neq \beta$) chiral multiplets in the theory. The second sum is the contribution of vector multiplets. When no confusion is possible, we will keep the gauge group indices implicit and just write $\Delta_{I_{\alpha\beta}} \equiv \Delta_I$. In the previous formula,

$$\Delta_I = \xi_I + r_I \frac{\tau + \sigma}{2}, \quad (3.4)$$

where r_I is the exact R-charge of the field and ξ_I are the flavor chemical potentials. The R-charges satisfy

$$\sum_{I \in W} r_I = 2 \quad (3.5)$$

for each superpotential term W in the Lagrangian. In this notation, the index W runs over the monomials in the superpotential, while $I \in W$ indicates all chiral fields appearing in a given monomial. Using that each superpotential term must be invariant under the flavor symmetries, but chemical potentials are only defined up to integers, we also require

$$\sum_{I \in W} \xi_I = n_W \quad \text{for some} \quad n_W \in \mathbb{Z}. \quad (3.6)$$

The values $n_W \equiv n_0 = \pm 1$ have been used in [43, 44] to study the Cardy limit. As a consequence of the previous formulæ, for each superpotential term we have

$$\sum_{I \in W} \Delta_I = \tau + \sigma + n_W . \quad (3.7)$$

Hence, we stress that the chemical potentials Δ_I are *not* independent. Notice that the expression (3.3) correctly reduces to the one for $\mathcal{N} = 4$ SYM, Eqn. (2.18), once we use the definition (2.4) as well as the inversion formula for the elliptic gamma function (2.36). We also need to use the exact R-charges $r_I = 2/3$ of the chiral fields Φ_I .

Applying (2.34), we can evaluate the large N limit of (3.3) and obtain

$$\begin{aligned} \log \mathcal{I} \simeq & -\frac{\pi i N^2}{3\tau\sigma} \sum_I \left[B_3\left([\Delta_I]_\omega + 1 - \frac{\tau+\sigma}{2}\right) + \frac{\tau\sigma}{4} \left(2ab - \frac{a}{b} - \frac{b}{a}\right) B_1\left([\Delta_I]_\omega + 1 - \frac{\tau+\sigma}{2}\right) \right] \\ & + \frac{\pi i G N^2}{3\tau\sigma} \left[B_3\left(1 - \frac{\tau+\sigma}{2}\right) + \frac{\tau\sigma}{4} \left(2ab - \frac{a}{b} - \frac{b}{a}\right) B_1\left(1 - \frac{\tau+\sigma}{2}\right) \right] . \end{aligned} \quad (3.8)$$

The corrections are of order $N \log N$ or smaller. The formula is obtained by summing (2.34) for each chiral multiplet, as well as (2.34) with $[\Delta]_\omega \rightarrow 0$ (and opposite sign) for each vector multiplet. We stress that (3.8) comes from a *single* contribution — in the Bethe Ansatz expansion — to the index. Such a contribution might not be the dominant one, and so our estimate of the index might be incorrect, in some regions of the space of chemical potentials. However, we conjecture and we will bring some evidence that this contribution always captures the semiclassical Bekenstein-Hawking entropy of BPS black holes.

Due to the presence of the brackets $[\Delta_I]_\omega$, the expression (3.8) assumes different analytic forms in different regions of the space of chemical potentials Δ_I . There are two regions where the expression greatly simplifies. They correspond to the natural generalization of the two regions for $\mathcal{N} = 4$ SYM discussed in Section 2.2 and are expected to lead to the correct black hole entropy. In particular, they smoothly reduce to the results obtained in the Cardy limit [43, 44, 32] and match the previous analysis done for equal angular momenta [28]. The first region corresponds to chemical potentials Δ_I satisfying

$$\sum_{I \in W} [\Delta_I]_\omega = \tau + \sigma - 1 . \quad (3.9)$$

As we will discuss later, many models — in particular all toric ones — exhibit a corner in the space of chemical potentials where this constraint is satisfied. We can define the rescaled variables

$$\widehat{\Delta}_I = 2 \frac{[\Delta_I]_\omega}{\tau + \sigma - 1} \quad (3.10)$$

which, under the assumption (3.9), satisfy

$$\sum_{I \in W} \widehat{\Delta}_I = 2 \quad (3.11)$$

and can be interpreted as an assignment of R-charges to the chiral fields in the theory. In terms of $\widehat{\Delta}_I$ the contributions in (3.8) combine into

$$\begin{aligned} \log \mathcal{I} \simeq & -\frac{\pi i N^2}{24} \frac{(\tau + \sigma - 1)^3}{\tau \sigma} \left[\sum_I (\widehat{\Delta}_I - 1)^3 + G \right] \\ & + \frac{\pi i N^2}{24} \frac{(\tau + \sigma - 1)}{\tau \sigma} \left(1 - \tau \sigma \left(2ab - \frac{a}{b} - \frac{b}{a} \right) \right) \left[\sum_I (\widehat{\Delta}_I - 1) + G \right]. \end{aligned} \quad (3.12)$$

Introducing the charge operator $R(\widehat{\Delta})$ of R-charges parametrized by $\widehat{\Delta}_I$ and indicating with Tr the sum over all fermions in the theory, we can also write

$$\log \mathcal{I} \simeq -\frac{\pi i}{24} \left[\frac{(\tau + \sigma - 1)^3}{\tau \sigma} \text{Tr} R(\widehat{\Delta})^3 - \frac{(\tau + \sigma - 1)}{\tau \sigma} \left(1 - \tau \sigma \left(2ab - \frac{a}{b} - \frac{b}{a} \right) \right) \text{Tr} R(\widehat{\Delta}) \right], \quad (3.13)$$

valid at leading order in N .

In the large N limit, theories with a holographic dual satisfy $c = a$. Using standard formulæ for the central charges a and c in terms of the fermion R-charges [45], one finds $\text{Tr} R = \mathcal{O}(1)$ and $a = \frac{9}{32} \text{Tr} R^3 + \mathcal{O}(1)$ from which we obtain the final expression

$$\log \mathcal{I} \simeq -\frac{4\pi i}{27} \frac{(\tau + \sigma - 1)^3}{\tau \sigma} a(\widehat{\Delta}), \quad (3.14)$$

where

$$a = \frac{9}{32} N^2 \left(\sum_I (\widehat{\Delta}_I - 1)^3 + G \right) \quad (3.15)$$

at leading order in N . The result (3.14) was conjectured in [31] — see Eqn. (A.7). It is also compatible with the Cardy limit performed in [43, 44].

We can find an analogous result in a second region of chemical potentials where

$$\sum_{I \in W} [\Delta_I]'_{\omega} = \tau + \sigma + 1, \quad (3.16)$$

written in terms of the primed bracket $[\Delta]'_{\omega} = [\Delta]_{\omega} + 1$. As discussed at the end of Section 2, the contribution of vector multiplets can be written, up to subleading terms, as minus the contribution of a chiral multiplet with $[\Delta_I]'_{\omega} \rightarrow 0$. After defining another set of normalised R-charges,

$$\widehat{\Delta}'_I = 2 \frac{[\Delta_I]'_{\omega}}{\tau + \sigma + 1} \quad (3.17)$$

which satisfy

$$\sum_{I \in W} \widehat{\Delta}'_I = 2 \quad (3.18)$$

under the assumption (3.16), we can rewrite the index as

$$\log \mathcal{I} \simeq -\frac{\pi i}{24} \left[\frac{(\tau + \sigma + 1)^3}{\tau\sigma} \text{Tr} R(\widehat{\Delta}')^3 - \frac{(\tau + \sigma + 1)}{\tau\sigma} \left(1 - \tau\sigma \left(2ab - \frac{a}{b} - \frac{b}{a} \right) \right) \text{Tr} R(\widehat{\Delta}') \right] \quad (3.19)$$

at leading order in N . This reduces to the simple expression

$$\log \mathcal{I} \simeq -\frac{4\pi i}{27} \frac{(\tau + \sigma + 1)^3}{\tau\sigma} a(\widehat{\Delta}') \quad (3.20)$$

for holographic theories.

In the remainder of this section we will interpret the general results (3.14) and (3.20) and provide examples. In particular, we will show that both regions (3.9) and (3.16) in the space of chemical potentials always exist in toric quiver gauge theories. We will also see that the two expressions (3.14) and (3.20) lead to the very same result for the semiclassical entropy of dual black holes, generalizing what happens for $\mathcal{N} = 4$ SYM.

3.1 Example: the conifold

We start with the example of the Klebanov-Witten theory dual to $\text{AdS}_5 \times T^{1,1}$, the near-horizon limit of a set of N D3-branes sitting at a conifold singularity [33]. This example was already studied for equal angular momenta in [28] and our results are consistent with those found there when we set $\tau = \sigma$.

The theory has gauge group $\text{SU}(N) \times \text{SU}(N)$, bi-fundamental chiral multiplets A_1, A_2 transforming in the representation (N, \overline{N}) and B_1, B_2 transforming in the representation (\overline{N}, N) , and a superpotential

$$W = \text{Tr} (A_1 B_1 A_2 B_2 - A_1 B_2 A_2 B_1) . \quad (3.21)$$

The global symmetry of the theory is $U(1)_R \times SU(2)_{F_1} \times SU(2)_{F_2} \times U(1)_B$, where the first factor is the superconformal R-symmetry with charge r , while the other three factors are flavor symmetries. The charge assignments of chiral multiplets under the maximal torus are in Table 1. The index is defined as

$$\mathcal{I} = \text{Tr} (-1)^F e^{-\beta\{\mathcal{Q}, \mathcal{Q}^\dagger\}} p^{J_1+r/2} q^{J_2+r/2} v_{F_1}^{Q_{F_1}} v_{F_2}^{Q_{F_2}} v_B^{Q_B} . \quad (3.22)$$

It is convenient to introduce an alternative basis of R-charges R_I with $I = 1, 2, 3, 4$, such that each of them assigns R-charge 2 to one of the chiral multiplets and zero to the other ones. Correspondingly, we associate a variable Δ_I to each chiral multiplet. Notice that

Field	r	Q_{F_1}	Q_{F_2}	Q_B	R_1	R_2	R_3	R_4
A_1	$\frac{1}{2}$	1	0	1	2	0	0	0
A_2	$\frac{1}{2}$	-1	0	1	0	2	0	0
B_1	$\frac{1}{2}$	0	1	-1	0	0	2	0
B_2	$\frac{1}{2}$	0	-1	-1	0	0	0	2

Table 1: Charges of chiral multiplets in the Klebanov-Witten theory, under the maximal torus of the global symmetry $U(1)_R \times SU(2)_{F_1} \times SU(2)_{F_2} \times U(1)_B$. In the table we indicate two useful basis. Notice that r and R_I are R-charges, while $Q_{F_{1,2}}$ and Q_B are flavor charges.

$(-1)^F = e^{2\pi i J_{1,2}} = e^{\pi i R_{1,2,3,4}}$. According to (3.4) and up to integer ambiguities, the variables Δ_I are related to the chemical potentials for the charges in Table 1 by

$$\begin{aligned}
\Delta_1 &= \xi_{F_1} + \xi_B + \frac{\tau + \sigma}{4}, & \Delta_3 &= \xi_{F_2} - \xi_B + \frac{\tau + \sigma}{4}, \\
\Delta_2 &= -\xi_{F_1} + \xi_B + \frac{\tau + \sigma}{4}, & \Delta_4 &= -\xi_{F_2} - \xi_B + \frac{\tau + \sigma}{4} + (2\mathbb{Z} + 1).
\end{aligned} \tag{3.23}$$

Then, the constraint (3.7) reads

$$\Delta_1 + \Delta_2 + \Delta_3 + \Delta_4 = \tau + \sigma + n_W \tag{3.24}$$

and the index takes the more transparent form

$$\mathcal{I} = \text{Tr}_{\text{BPS}} p^{J_1} q^{J_2} y_1^{R_1/2} y_2^{R_2/2} y_3^{R_3/2} y_4^{R_4/2}. \tag{3.25}$$

This shows that Δ_I are the chemical potentials associated to the charges $Q_I \equiv R_I/2$.

We select three independent variables, say Δ_1, Δ_2 and Δ_3 . Then, using (2.39) we find that

$$[\Delta_4]_\omega = \tau + \sigma - 1 - [\Delta_1 + \Delta_2 + \Delta_3]_\omega. \tag{3.26}$$

In general there are three possible cases:

$$[\Delta_1 + \Delta_2 + \Delta_3]_\omega = [\Delta_1]_\omega + [\Delta_2]_\omega + [\Delta_3]_\omega + n \quad \text{with} \quad n = 0, 1, 2 \tag{3.27}$$

that we call Case I, II and III, respectively.⁷

Case I corresponds to the corner of moduli space (3.9) where

$$[\Delta_1]_\omega + [\Delta_2]_\omega + [\Delta_3]_\omega + [\Delta_4]_\omega = \tau + \sigma - 1. \tag{3.28}$$

⁷For the sake of comparison, the notation is the same as in [28].

In this corner, we can use (3.14). One can explicitly compute, at leading order in N ,

$$\mathrm{Tr} R(\widehat{\Delta})^3 = N^2 \left(2 + \sum_{I=1}^4 (\widehat{\Delta}_I - 1)^3 \right) = 3N^2 \left(\widehat{\Delta}_1 \widehat{\Delta}_2 \widehat{\Delta}_3 + \widehat{\Delta}_1 \widehat{\Delta}_2 \widehat{\Delta}_4 + \widehat{\Delta}_1 \widehat{\Delta}_3 \widehat{\Delta}_4 + \widehat{\Delta}_2 \widehat{\Delta}_3 \widehat{\Delta}_4 \right) \quad (3.29)$$

imposing $\sum_{I=1}^4 \widehat{\Delta}_I = 2$. Using (3.10), we can write the index (3.14) as

$$\log \mathcal{I} \simeq -\frac{\pi i N^2}{\tau \sigma} \left([\Delta_1]_\omega [\Delta_2]_\omega [\Delta_3]_\omega + [\Delta_1]_\omega [\Delta_2]_\omega [\Delta_4]_\omega + [\Delta_1]_\omega [\Delta_3]_\omega [\Delta_4]_\omega + [\Delta_2]_\omega [\Delta_3]_\omega [\Delta_4]_\omega \right) \quad (3.30)$$

with the constraint (3.28).⁸

Case III corresponds to the corner of moduli space (3.16). Indeed

$$[\Delta_1]'_\omega + [\Delta_2]'_\omega + [\Delta_3]'_\omega + [\Delta_4]'_\omega = \tau + \sigma + 1 . \quad (3.31)$$

In this corner, we can use (3.20) and (3.17) and find

$$\log \mathcal{I} \simeq -\frac{\pi i N^2}{\tau \sigma} \left([\Delta_1]'_\omega [\Delta_2]'_\omega [\Delta_3]'_\omega + [\Delta_1]'_\omega [\Delta_2]'_\omega [\Delta_4]'_\omega + [\Delta_1]'_\omega [\Delta_3]'_\omega [\Delta_4]'_\omega + [\Delta_2]'_\omega [\Delta_3]'_\omega [\Delta_4]'_\omega \right) \quad (3.32)$$

with the constraint (3.31).

The entropy, which is the logarithm of the number of states, is given by the Legendre transform of the index, *i.e.*, by the critical value of the entropy function

$$\begin{aligned} \mathcal{S} = & -\frac{\pi i N^2}{\tau \sigma} \left(X_1 X_2 X_3 + X_1 X_2 X_4 + X_1 X_3 X_4 + X_2 X_3 X_4 \right) \\ & - 2\pi i \left(\tau J_1 + \sigma J_2 + \sum_{I=1}^4 X_I Q_I \right) - 2\pi i \Lambda \left(\sum_{I=1}^4 X_I - \tau - \sigma \pm 1 \right) . \end{aligned} \quad (3.33)$$

Here the variables X_I stand for $[\Delta_I]_\omega$ or $[\Delta_I]'_\omega$ depending on whether we are in case I or III, respectively, and the \pm sign is chosen accordingly. One can check that the two signs lead to the same entropy. We will give a general argument in Section 3.3.

In Section 5 we will compare the field theory result (3.33) with the entropy of black holes in $\mathrm{AdS}_5 \times T^{1,1}$, in the special case that $J_1 = J_2 \equiv J$ and the $SU(2)_{F_1} \times SU(2)_{F_2}$ symmetry is unbroken. To that purpose, let us specialize the index to the case that $\tau = \sigma$ and $\xi_{F_1} = \xi_{F_2} = 0$, which corresponds to $X_1 = X_2$ and $X_3 = X_4$. It is then useful to define the new variables

$$X_R = X_1 + X_3 , \quad X_B = \frac{X_1 - X_3}{2} , \quad (3.34)$$

⁸For toric models, discussed in detail in Section 3.2, we can compute the index using formula (3.52). The 't Hooft coefficients are expressed in terms of toric data as $C_{abc} = |\det\{v_a, v_b, v_c\}|$, where v_a are the integer vectors defining the toric fan [46]. For the conifold: $v_1 = (1, 0, 0)$, $v_2 = (1, 1, 0)$, $v_3 = (1, 1, 1)$, $v_4 = (1, 0, 1)$ and thus $C_{123} = C_{124} = C_{134} = C_{234} = 1$ (and symmetrizations), recovering the expression above.

associated to R-symmetry and baryonic symmetry, respectively. The entropy function takes the simplified form

$$\mathcal{S} = -\frac{\pi i N^2}{2\tau^2} X_R (X_R^2 - 4X_B^2) - 2\pi i (2\tau J + X_R r + X_B Q_B) - 2\pi i \Lambda (2X_R - 2\tau \pm 1). \quad (3.35)$$

3.2 Example: toric models

In this section we consider the gauge theory dual to an $\text{AdS}_5 \times \text{SE}_5$ geometry, where SE_5 is a toric Sasaki-Einstein manifold. The theory lives on a stack of N D3-branes sitting at the toric Calabi-Yau singularity $C(\text{SE}_5)$ obtained by taking the cone over SE_5 [33, 47]. There is a general construction to extract gauge theory data from the geometry of the Calabi-Yau singularity [48–51]. The main complication compared to the \mathbb{C}^3 and the conifold cases is that there is no one-to-one correspondence between bi-fundamental fields Φ_I (and associated variables Δ_I) and R-symmetries R_a . However, we will argue in general that there always exist two corners of the space of chemical potentials where (3.9) and (3.16) are satisfied and the results (3.14) and (3.20) are valid. There are also other corners that should be analyzed separately for every specific model. Our findings are consistent with the case-by-case analysis performed in [28] for equal angular momenta.

We first need to understand how to write the trial central charges $a(\widehat{\Delta})$ and $a(\widehat{\Delta}')$ that enter in the expressions (3.14) and (3.20). Since the quantities $\widehat{\Delta}_I$ and $\widehat{\Delta}'_I$ satisfy the constraints (3.11), they can be interpreted as a set of trial R-charges for the chiral fields in the quiver. In the toric case, we can find an efficient parametrization of the trial R-charges of fields using the data of the toric diagram. Let us review how this is done.

A toric Calabi-Yau threefold singularity can be specified by a fan, *i.e.*, a convex cone in \mathbb{R}^3 defined by D integer vectors $v_a = (1, \vec{v}_a)$ lying on a plane. The restrictions \vec{v}_a of those vectors to the plane define a regular convex polygon with integer vertices called the toric diagram. In the list $\{v_a\}$ we should include all integer vectors such that \vec{v}_a is along the perimeter of the polygon, *i.e.*, we should include all integer points along the edges of the toric diagram. Moreover, we take the points \vec{v}_a to be ordered in a counterclockwise fashion. The number of vectors in the fan is associated with the total rank of the global symmetry of the dual field theory [50]: for a toric model with D vectors in the fan (including integer points along the edges of the toric diagram) there is a flavor symmetry of rank $D - 1$, besides the R-symmetry $U(1)_R$.⁹ This allows us to parametrize flavor and R-symmetries in terms of variables associated with the vertices of (and integer points along) the toric diagram. In particular, the possible R-charges of fields in a toric theory can be parametrized using D

⁹The distinction between R- and flavor symmetries changes in the case of extended supersymmetry.

variables δ_a satisfying the constraint

$$\sum_{a=1}^D \delta_a = 2, \quad (3.36)$$

and the corresponding R-charge can be written as

$$R(\delta) = \sum_{a=1}^D \frac{\delta_a}{2} R_a \quad (3.37)$$

in terms of a basis $\{R_a\}$. This is done as follows [52]. In a minimal toric phase,¹⁰ the theory contains a number G of gauge group factors $SU(N)$ equal to twice the area of the toric diagram. Moreover, defining the vectors $\vec{w}_a = \vec{v}_{a+1} - \vec{v}_a$ lying in the plane (we identify indices modulo D , so that, for example, $\vec{v}_{D+1} \equiv \vec{v}_1$), for each pair (a, b) such that \vec{w}_a can be rotated counterclockwise into \vec{w}_b in the plane with an angle smaller than π , there are precisely¹¹ $\det\{\vec{w}_a, \vec{w}_b\}$ bi-fundamental chiral fields Φ_{ab} with R-charge

$$R[\Phi_{ab}] = \delta_{a+1} + \delta_{a+2} + \dots + \delta_b. \quad (3.38)$$

Interestingly, for all toric models the trial central charge $a(\delta)$ is a homogeneous function of degree three at large N :

$$a(\delta) = \frac{9}{32} \text{Tr} R(\delta)^3 = \frac{9N^2}{64} \sum_{a,b,c=1}^D C_{abc} \delta_a \delta_b \delta_c. \quad (3.39)$$

Here $N^2 C_{abc} = \frac{1}{4} \text{Tr} R_a R_b R_c$ are the 't Hooft anomaly coefficients, which can be read from the toric data through $C_{abc} = |\det\{v_a, v_b, v_c\}|$ [46]. Another important property of toric models that we will use in the following is that the constraints

$$\sum_{I \in W} R[\Phi_I] = 2, \quad (3.40)$$

that must be satisfied for each monomial term W in the superpotential, always reduce to (3.36). Indeed, it follows from tiling techniques [48–52] that the R-charges $R[\Phi_I]$, $I \in W$, of the chiral fields entering in a superpotential monomial W correspond to a partition of the

¹⁰There are many different quiver theories that describe the same IR SCFT. They are called “phases”, and are related by Seiberg dualities. The toric phases are the quiver theories where all gauge groups are $SU(N)$ with the same rank N . It turns out that all toric phases have the same number G of gauge groups, but have different matter content. The “minimal” phases correspond to the quivers with the smallest number of chiral fields. There could be one or more minimal toric phases, for a given IR SCFT.

¹¹The condition on the angle guarantees that the formula for the number of fields gives a non-negative integer.

D elementary R-charges $\{\delta_1, \dots, \delta_D\}$ into sums of the form (3.38), with each δ_a entering in just one $R[\Phi_I]$.

We can similarly parametrize the chemical potentials $\Delta[\Phi]$ entering the superconformal index in terms of D basic quantities Δ_a , $a = 1, \dots, D$. For the chiral fields Φ_{ab} we have

$$\Delta[\Phi_{ab}] = \Delta_{a+1} + \Delta_{a+2} + \dots + \Delta_b . \quad (3.41)$$

The conditions

$$\sum_{I \in W} \Delta[\Phi_I] = \tau + \sigma + n_W , \quad (3.42)$$

to be imposed for each monomial term W in the superpotential (and where n_W is the same for all monomial terms), are then equivalent to

$$\sum_{a=1}^D \Delta_a = \tau + \sigma + n_W . \quad (3.43)$$

Independently of the value of n_W , we have

$$[\Delta_D]_\omega = \tau + \sigma - 1 - \left[\sum_{a=1}^{D-1} \Delta_a \right]_\omega . \quad (3.44)$$

In general

$$\left[\sum_{a=1}^{D-1} \Delta_a \right]_\omega = \sum_{a=1}^{D-1} [\Delta_a]_\omega + n \quad (3.45)$$

where $n = 0, \dots, D - 2$, thus dividing the space of parameters into $D - 1$ regions.

Two regions are particularly important for our analysis. The region $n = 0$ corresponds to

$$\sum_{a=1}^D [\Delta_a]_\omega = \tau + \sigma - 1 , \quad (3.46)$$

while $n = D - 2$ corresponds to

$$\sum_{a=1}^D [\Delta_a]'_\omega = \tau + \sigma + 1 . \quad (3.47)$$

We can argue that the two regions (3.46) and (3.47) are always realized somewhere in the space of parameters. For example, we can choose one elementary variable, say Δ_1 , to live in the fundamental strip $\text{Im}(-1/\omega) > \text{Im}(\Delta_1/\omega) > 0$ (see Fig. 1) and slightly on the right of the vertical line passing through $\tau + \sigma - 1$, while all the other Δ_a to live in the fundamental strip and slightly on the left of the vertical line passing through zero. One easily verifies that they can be arranged to satisfy (3.46). A similar construction gives parameters satisfying

(3.47). We now argue that (3.46) and (3.47) imply (3.9) and (3.16), respectively. We start noticing that

$$\sum_{a=1}^D [\Delta_a]_\omega = \tau + \sigma - 1 \quad \Rightarrow \quad \text{Im} \left(\frac{1}{\omega} \sum_{a=1}^D [\Delta_a]_\omega \right) = \text{Im} \left(-\frac{1}{\omega} \right). \quad (3.48)$$

Since each of the $[\Delta_a]_\omega$ lives in the fundamental strip $\text{Im}(-1/\omega) > \text{Im}([\Delta_a]_\omega/\omega) > 0$, the previous equation implies that $\text{Im}(-1/\omega) > \text{Im}(\sum_{a \in S} [\Delta_a]_\omega/\omega) > 0$ for any proper subset S of the indices $\{1, \dots, D\}$. Thus (3.46) implies that

$$\left[\sum_{a \in S} \Delta_a \right]_\omega = \sum_{a \in S} [\Delta_a]_\omega \quad (3.49)$$

for any proper subset $S \subsetneq \{1, \dots, D\}$. This implies that all charges in (3.41) *split*, in the sense that $[\Delta_{a+1} + \dots + \Delta_b]_\omega = [\Delta_{a+1}]_\omega + \dots + [\Delta_b]_\omega$. At this point, since all $[\Delta[\Phi_I]]_\omega$ split and each Δ_a enters precisely once in every superpotential constraint, the condition (3.9) is a consequence of (3.46).¹² A similar argument shows that (3.47) implies (3.16). Notice that the region specified by (3.9) can be larger than (3.46) and, similarly, the region specified by (3.16) can be larger than (3.47). This, in particular, happens for Calabi-Yau cones with codimension-one orbifold singularities. This is the case of the models SPP and dP₄ discussed in [28].¹³ For all the cones without orbifold singularities that we checked, the two regions (3.9) and (3.46) coincide. It would be interesting to see if this is a general result.

We are now ready to evaluate the index. Consider region (3.9) first. Since the chemical potentials $[\Delta_I]_\omega$ split, the rescaled quantities

$$\widehat{\Delta}_a = 2 \frac{[\Delta_a]_\omega}{\tau + \sigma - 1} \quad \text{with} \quad \sum_{a=1}^D \widehat{\Delta}_a = 2 \quad (3.50)$$

¹²There is an alternative algorithm that produces potentials Δ_I satisfying (3.9). Choose a perfect matching p_α of the dimer model of the theory [50]. It divides the chiral fields into two groups: those Φ_P appearing in the perfect matching, and those Φ_{NP} not doing so. Choose the potentials Δ_{NP} to be in the fundamental strip and slightly on the left of the origin. Each superpotential term W contains one and only one of the fields Φ_P (by definition of perfect matching): choose the corresponding Δ_P to be in the fundamental strip and slightly on the right of the point $\tau + \sigma - 1$, in such a way that (3.9) for that particular W is satisfied. The drawback of this construction is that it does not tell us what the independent variables Δ_a are.

¹³Models with codimension-one orbifold singularities are characterized by toric diagrams where at least one vector \vec{v}_a lies in the interior of an edge. The parameters δ_a associated with integer points lying in the interior of an edge of the polygon enter in the parametrization (3.38) of the R-charges of chiral fields, but no elementary field carries precisely charge δ_a . In order to recover the region (3.9), we can require the following. Construct a set M by grouping the points $\{1, \dots, D\}$ along the toric diagram in the following way: Break each edge in two pieces at a non-integer point, and then for each vertex form a group (that will be an element of M) that contains the vertex itself and all other integer points (if any) along the two pieces of edges on the two sides. (In the absence of orbifold singularity, M necessarily coincides with $\{1, \dots, D\}$.) Then require that the sums split over the groups in M for every proper subgroup $S' \subsetneq M$, and for every possible choice of M . This region is typically larger than (3.46).

provide a parametrization of the R-charges of chiral fields in the quiver in the sense discussed above. Using the general formula (3.39) we can then write

$$a(\widehat{\Delta}) = \frac{9N^2}{64} \sum_{a,b,c=1}^D C_{abc} \widehat{\Delta}_a \widehat{\Delta}_b \widehat{\Delta}_c . \quad (3.51)$$

Plugging it into (3.14) and re-expressing the result in terms of the chemical potentials $[\Delta_a]_\omega$, we find the large N limit of the superconformal index in region (3.9):

$$\log \mathcal{I} \simeq -\pi i N^2 \sum_{a,b,c=1}^D \frac{C_{abc}}{6} \frac{[\Delta_a]_\omega [\Delta_b]_\omega [\Delta_c]_\omega}{\tau \sigma} , \quad \sum_{a=1}^D [\Delta_a]_\omega = \tau + \sigma - 1 . \quad (3.52)$$

A similar argument shows that, in region (3.16),

$$\log \mathcal{I} \simeq -\pi i N^2 \sum_{a,b,c=1}^D \frac{C_{abc}}{6} \frac{[\Delta_a]'_\omega [\Delta_b]'_\omega [\Delta_c]'_\omega}{\tau \sigma} , \quad \sum_{a=1}^D [\Delta_a]'_\omega = \tau + \sigma + 1 . \quad (3.53)$$

We will show in the next section that both (3.52) and (3.53) lead to the same entropy.

3.3 The entropy function

For toric holographic quivers, we have found two different expressions, (3.52) and (3.53), for the large N limit of the superconformal index that are valid in two different regions in the space of chemical potentials. The two expressions differ only for the constraint and give rise to the very same entropy. This generalizes an observation made in [4] for $\mathcal{N} = 4$ SYM and holds for general quivers.

To show that, we define two entropy functions

$$S_\pm = -\pi i N^2 \sum_{a,b,c=1}^D \frac{C_{abc}}{6} \frac{X_a X_b X_c}{\tau \sigma} - 2\pi i \left(\tau J_1 + \sigma J_2 + \sum_{a=1}^D X_a Q_a \right) - 2\pi i \Lambda \left(\sum_{a=1}^D X_a - \tau - \sigma \pm 1 \right) , \quad (3.54)$$

where Λ is a Lagrange multiplier and we used neutral variables X_a to denote either $[\Delta_a]_\omega$ or $[\Delta_a]'_\omega$. Each of the electric charges $Q_a \equiv R_a/2$ is defined in terms of an R-charge R_a that assigns charge 2 to all chiral multiplets Φ_{ab} such that δ_a appears in the decomposition (3.38), and zero to all the other ones. The 't Hooft anomaly coefficients are defined by

$$C_{abc} N^2 = \frac{1}{4} \text{Tr} R_a R_b R_c . \quad (3.55)$$

Above, S_+ is the prediction for the entropy of the dual black hole based on the superconformal index in the region of parameters (3.9) while S_- in the region (3.16). The form of the entropy function (3.54) was first conjectured in [31].

Observe that, since $S_{\pm} \pm 2\pi i\Lambda$ are homogeneous functions of degree one in (X_a, τ, σ) , the values of the functions $S_{\pm}(X_a, \tau, \sigma, \Lambda)$ at the critical point are related to the Lagrange multiplier by

$$S_{\pm}|_{\text{crit}} = \mp 2\pi i\Lambda . \quad (3.56)$$

Observe also that, if Q_a, J_i are real (as charges should be), then the two functions are related by $\overline{S_+(X_a, \tau, \sigma, \Lambda)} = S_-(-\overline{X}_a, -\overline{\tau}, -\overline{\sigma}, \overline{\Lambda})$. Hence, if $(X_a, \tau, \sigma, \Lambda)$ is a critical point of S_+ , then $(-\overline{X}_a, -\overline{\tau}, -\overline{\sigma}, \overline{\Lambda})$ is a critical point of S_- with critical value

$$S_-|_{\text{crit}} = \overline{S_+}|_{\text{crit}} . \quad (3.57)$$

For arbitrary and general real charges Q_a and J_i , the critical value of S_+ is not real. For $\mathcal{N} = 4$ SYM, however, it becomes real and equal to the entropy when imposing the non-linear constraint on conserved charges that characterizes supersymmetric black holes [21, 4]. The same phenomenon was already observed in AdS_4 in [3]. We expect the same to be true for general black holes in Sasaki-Einstein compactifications. Even if this were wrong and S_+ were not real, it would still makes sense to identify the entropy with $\mathbb{R}e S_+$. In all cases, we see from (3.57) that both constraints in (3.54) lead to the very same result for the entropy.

The entropy functions (3.54) give our general result for the entropy of black holes in $\text{AdS}_5 \times \text{SE}_5$. We derived it for toric quiver gauge theories, but the very same argument can be extended to a class of more general non-toric quivers. In particular, the expression (3.54) only depends on the 't Hooft anomaly coefficients C_{abc} for a basis of R-symmetries and, as such, we expect that it is the correct result for generic holographic quiver theories.

4 The universal rotating black hole

In this section we discuss the case of the *universal rotating black hole* which has electric charge aligned with the exact R-symmetry of the theory. The black hole arises as a solution of minimal gauged supergravity in five dimensions and, as such, it can be embedded in any $\text{AdS}_5 \times \text{SE}_5$ compactification of type IIB and, more generally, in any AdS_5 solution of type II or M theory.¹⁴ Due to its universal character, most of the analysis is identical to the one for $\text{AdS}_5 \times S^5$. It is however interesting to see how the details work.

¹⁴It is believed and checked in many cases that the effective theory for all such compactifications can be consistently truncated to minimal gauged supergravity.

The universal black hole in AdS₅ was found in [17] in minimal gauged supergravity in five dimensions. It has charge Q under the graviphoton and angular momenta J_1 and J_2 in AdS₅.¹⁵ The entropy can be compactly written as [53]

$$S(Q, J) = 2\pi\sqrt{3Q^2 - 2a(J_1 + J_2)} \quad (4.1)$$

where we introduced the quantity

$$a = \frac{\pi\ell_5^3}{8G_N^{(5)}}, \quad (4.2)$$

where $G_N^{(5)}$ is the five-dimensional Newton constant and ℓ_5 is the radius of AdS₅. The conserved charges must satisfy the nonlinear constraint

$$8Q^3 + 6aQ^2 - 6a(J_1 + J_2)Q - 2aJ_1J_2 - 4a^2(J_1 + J_2) = 0 \quad (4.3)$$

for the BPS black hole to have a smooth horizon.

Consider now the uplift of the universal black hole to AdS₅ × SE₅, where SE₅ is a Sasaki-Einstein manifold. In such an embedding, the standard holographic dictionary identifies a with the central charge of the dual CFT₄. The black hole carries angular momenta J_1 and J_2 and an electric charge aligned with the exact R-symmetry of the dual CFT₄. We need to check that its entropy is reproduced by our result (3.14) (the same result can be similarly obtained using (3.20) instead). It is convenient to parametrize the chemical potentials as

$$\Delta_a = \frac{\tau + \sigma - 1}{2} \left(\widehat{\Delta}_a^{(0)} + \widehat{\delta}_a \right), \quad (4.4)$$

where $\widehat{\Delta}_a^{(0)}$ is the *exact* superconformal R-symmetry of the dual CFT₄ while $\widehat{\delta}_a$ parametrize a basis of flavor symmetries. These quantities satisfy

$$\sum_{a=1}^D \widehat{\Delta}_a^{(0)} = 2, \quad \sum_{a=1}^D \widehat{\delta}_a = 0. \quad (4.5)$$

The entropy of the universal black hole is given by the Legendre transform of (3.14). Using (3.46) we can write the entropy function as

$$\mathcal{S} = -\frac{4\pi i}{27} \frac{(\tau + \sigma - 1)^3}{\tau\sigma} a \left(\widehat{\Delta}^{(0)} + \widehat{\delta} \right) - 2\pi i \left((\tau + \sigma - 1)Q + \tau J_1 + \sigma J_2 \right), \quad (4.6)$$

where we introduced a charge $Q = \frac{1}{2} \sum_{a=1}^D \widehat{\Delta}_a^{(0)} Q_a$ in the direction of the exact R-symmetry, and set all other charges to zero. We need to extremize the function \mathcal{S} with respect to τ ,

¹⁵To compare with the notations of [17]: $Q_{\text{there}} = -\sqrt{3}gQ_{\text{here}}$ and $G_N^{(5)} = 1$, $\ell_5 = 1/g$.

σ and $\widehat{\delta}_a$ subject to the constraint (4.5). By a -maximization, since $\widehat{\Delta}_a^{(0)}$ is the exact R-symmetry, the function is extremized at $\widehat{\delta}_a = 0$. We can then restrict the entropy function to

$$\mathcal{S} = -\frac{4\pi i a}{27} \frac{(\tau + \sigma - 1)^3}{\tau\sigma} - 2\pi i \left((\tau + \sigma - 1)Q + \tau J_1 + \sigma J_2 \right), \quad (4.7)$$

where $a \equiv a(\widehat{\Delta}^{(0)})$ is the central charge of the CFT_4 , or, introducing a Lagrange multiplier Λ ,

$$\mathcal{S} = -4\pi i a \frac{\Delta^3}{\tau\sigma} - 2\pi i \left(3\Delta Q + \tau J_1 + \sigma J_2 \right) - 2\pi i \Lambda (3\Delta - \tau - \sigma + 1). \quad (4.8)$$

If we set $a = a_{\mathcal{N}=4} = \frac{1}{4}N^2$, the function (4.8) becomes identical to the entropy function of $\mathcal{N} = 4$ SYM for equal charges $Q_1 = Q_2 = Q_3 \equiv Q$, which is known to correctly reproduce (4.1) [21]. An analytic derivation of (4.1) and (4.3) for $\mathcal{N} = 4$ SYM is explicitly discussed in [4] and for equal angular momenta in [6]. The charge constraint (4.3) is obtained as the requirement that the extremum of \mathcal{S} be real.

At this point, the result for the universal black hole simply follows from the homogeneity properties of (4.8):

$$S(Q, J_1, J_2) = \frac{a}{a_{\mathcal{N}=4}} S_{\mathcal{N}=4} \left(\frac{a_{\mathcal{N}=4}}{a} Q, \frac{a_{\mathcal{N}=4}}{a} J_1, \frac{a_{\mathcal{N}=4}}{a} J_2 \right). \quad (4.9)$$

It is then immediate to derive the relations (4.1) and (4.3), thus completing our derivation.

5 AdS₅ Kerr-Newman black holes in $T^{1,1}$

We would like to compare the entropy function we obtained in Section 3 from the large N limit of the superconformal index of generic (toric) quiver gauge theories, with the Bekenstein-Hawking entropy of BPS black holes in the corresponding 5d gauged supergravities. In particular, the setup we would like to analyze is that of type IIB supergravity on asymptotically AdS₅ × SE₅ spacetimes, where SE₅ is a toric Sasaki-Einstein manifold,¹⁶ reduced and truncated to a 5d $\mathcal{N} = 2$ gauged supergravity on AdS₅. Unfortunately, with the exception of the case of S^5 truncated to the so-called 5d STU model, and the case of any SE₅ truncated to minimal $\mathcal{N} = 2$ gauged supergravity (that we analyzed in Section 4), all other known consistent truncations are to gauged supergravities with hypermultiplets (besides vector multiplets), and no supersymmetric black hole solutions have been constructed in such theories to date.

The strategy we propose to perform a test of our field theory results is as in [21]. We assume that a 5d BPS rotating black hole solution exists. Such a solution has the topology

¹⁶More precisely, the cone over SE₅ is a toric Calabi-Yau threefold.

of a fibration of AdS_2 over S^3 (the three-sphere being the topology of the event horizon), and thus we can reduce it along the Hopf fiber of S^3 . This gives a (putative) 4d BPS rotating black hole solution, with the same entropy.¹⁷ The reduction generates an extra vector field A^0 , corresponding to the isometry along the Hopf fiber. The 4d black hole has one unit of magnetic charge under A^0 , corresponding to the first Chern class of the Hopf fibration. Calling J_1 and J_2 the 5d angular momenta along two orthogonal planes, the quantity $J_1 + J_2$ appears in 4d as the electric charge under A^0 , while $J_1 - J_2$ becomes the angular momentum of the 4d black hole. Constructing such a 4d rotating black hole solution is still a difficult task, and an attractor mechanism is not known in general.¹⁸ However, if we restrict to 5d black holes with two equal angular momenta $J_1 = J_2$ (so that the isometry of the squashed S^3 is enhanced from $U(1)^2$ to $U(1) \times SU(2)$), then the 4d black hole is static: in this case we can determine its entropy by exploiting the attractor mechanism in the near-horizon geometry [37–39], without actually constructing the whole solution.

The simplest non-trivial example is when SE_5 is $T^{1,1}$, the base of the conifold Calabi-Yau threefold, whose holographic dual is the Klebanov-Witten gauge theory [33]. We already presented the field theory analysis in Section 3.1. On the other hand, starting from 10d type IIB supergravity on $T^{1,1}$, we can exploit a consistent truncation that preserves $SU(2)^2 \times U(1)$ isometry, down to a 5d $\mathcal{N} = 2$ gauged supergravity with the graviton multiplet, two vector multiplets and two hypermultiplets. This is the second truncation presented in Section 7 of [34] (see also [35, 36]). On the AdS_5 vacuum, one vector multiplet (sometimes called “Betti multiplet”) is massless and is associated to the baryonic symmetry, while the other vector multiplet is massive.

Hence, with the simplification that $J_1 = J_2$ and only the R-symmetry and baryonic symmetry charges are turned on (while the $SU(2)^2$ isometry of $T^{1,1}$ is unbroken), we will be able to match the Legendre transform of the superconformal index at large N with the extremization problem that comes from the attractor mechanism in supergravity. It follows that the bulk and boundary computations of the entropy exactly match.

5.1 Reduction from 5d to 4d and the attractor mechanism

A 5d $\mathcal{N} = 2$ Abelian gauged supergravity with n_V vector multiplets and n_H hypermultiplets — whose main building blocks we summarize in Appendix B — is specified by the following data [57–59]:

¹⁷The 4d solution has an exotic asymptotic behavior, that follows from the reduction of AdS_5 [54]. Nonetheless, it has a regular extremal horizon, whose area determines the entropy.

¹⁸There are however some general results for theories with vector multiplets [55, 56].

1. A very special real manifold \mathcal{SM} of real dimension n_V , specified by a symmetric tensor of Chern-Simons couplings C_{IJK} with $I, J, K = 1, \dots, n_V + 1$. The coordinates are Φ^I with the cubic constraint

$$\mathcal{V}(\Phi) \equiv \frac{1}{6} C_{IJK} \Phi^I \Phi^J \Phi^K = 1. \quad (5.1)$$

2. A quaternionic-Kähler manifold \mathcal{QM} of real dimension $4n_H$ with coordinates q^u .
3. A set of $n_V + 1$ Killing vectors k_I^u (that could be linearly dependent, or vanish) on \mathcal{QM} , compatible with the quaternionic-Kähler structure, representing the isometries to be gauged by the vector fields A^I . Each Killing vector comes equipped with a triplet of moment maps \vec{P}_I .¹⁹

On the other hand, a 4d $\mathcal{N} = 2$ Abelian gauged supergravity with $n_V + 1$ vector multiplets and n_H hypermultiplets — that we summarize in Appendix C — is specified by the following data (see for instance [60, 61]):

1. A special Kähler manifold \mathcal{KM} of complex dimension $n_V + 1$, with coordinates z^I and $I = 1, \dots, n_V + 1$. We will work in a duality frame in which the geometry is specified by holomorphic sections $X^\Lambda(z)$, with $\Lambda = 0, \dots, n_V + 1$, and a holomorphic prepotential $F(X)$, homogeneous of degree two.
2. A quaternionic-Kähler manifold \mathcal{QM} of real dimension $4n_H$ with coordinates q^u .
3. In duality frames in which all gaugings are purely electric, a set of $n_V + 2$ Killing vectors k_Λ^u (that could be linearly dependent, or vanish) on \mathcal{QM} , compatible with the quaternionic-Kähler structure, representing the isometries to be electrically gauged by the vector fields A^Λ . Each Killing vector comes equipped with a triplet of moment maps \vec{P}_Λ (see footnote 19).

We reduce the 5d theory on a circle, that will eventually be the Hopf fiber of S^3 . Following [62–66, 21] we use the ansatz

$$\begin{aligned} ds_{(5)}^2 &= e^{2\tilde{\phi}} ds_{(4)}^2 + e^{-4\tilde{\phi}} (dy - A_{(4)}^0)^2 \\ \Phi^I &= -e^{2\tilde{\phi}} \text{Im } z^I. \end{aligned} \quad (5.2)$$

Here y is the direction of the circular fiber, that we take with range $4\pi/g$ in terms of the coupling $g = \ell_5^{-1}$ inversely proportional to the AdS₅ radius ℓ_5 , therefore the size of the circle is $e^{-2\tilde{\phi}} 4\pi/g$. Because of the constraint $\mathcal{V}(\Phi) = 1$ in (5.1), the field $\tilde{\phi}$ is redundant and can

¹⁹If $n_H = 0$, instead, one has to specify n_V Fayet-Iliopoulos parameters ζ^I , not all vanishing.

be eliminated with $e^{-6\tilde{\phi}} = -\mathcal{V}(\text{Im } z^I)$. On the other hand, $A_{(4)}^0$ is the Kaluza-Klein vector. As noted in [67, 21], a Scherk-Schwarz twist for the gravitino as in [65] is necessary to satisfy the BPS conditions in 4d. We prefer to work in a gauge in which all bosonic fields are periodic around the circle, but there are flat gauge connections ξ^I turned on along y . This corresponds to the ansatz

$$A_{(5)}^I = A_{(4)}^I + \text{Re } z^I (dy - A_{(4)}^0) + \xi^I dy, \quad (5.3)$$

together with no y -dependence for any field. Notice that this ansatz is invariant under the redefinitions

$$z^I \rightarrow z^I + \delta\xi^I, \quad A_{(4)}^I \rightarrow A_{(4)}^I + \delta\xi^I A_{(4)}^0, \quad \xi^I \rightarrow \xi^I - \delta\xi^I \quad (5.4)$$

where $\delta\xi^I$ are real parameters. We will fix this redundancy below. The reduction of the 5d theory can be found in Appendix D. The resulting 4d data in terms of 5d ones are as follows.

1. The special Kähler manifold in 4d is described by the prepotential

$$F(X) = \frac{1}{6} C_{IJK} \frac{\check{X}^I \check{X}^J \check{X}^K}{X^0} \quad \text{with} \quad \check{X}^I = X^I + \xi^I X^0. \quad (5.5)$$

The holomorphic sections X^Λ can be used as homogeneous coordinates, and the physical scalars are identified with the special coordinates $z^I = X^I/X^0$.

2. The quaternionic-Kähler manifold in 4d is the same as in 5d.
3. The 4d Killing vectors k_I^u are inherited from 5d, while the additional Killing vector is

$$k_0^u = \xi^I k_I^u \quad \Rightarrow \quad \vec{P}_0 = \xi^I \vec{P}_I, \quad (5.6)$$

and is gauged by the Kaluza-Klein vector field $A_{(4)}^0$.

Next, we study the attractor equations for the near-horizon limit of 4d BPS static black hole solutions [37–39]. Our goal is to use the BPS equations to fix the VEVs in massive vector multiplets and hypermultiplets, and be left with an extremization principle for the scalars in massless vector multiplets, similarly to [68, 69]. We consider the near-horizon geometry $\text{AdS}_2 \times S^2$:

$$ds_{\text{near-horizon}}^2 = -\frac{r^2}{L_A^2} dt^2 + \frac{L_A^2}{r^2} dr^2 + L_S^2 ds_{S^2}^2, \quad (5.7)$$

where L_A and L_S are the radii of AdS_2 and S^2 , respectively. Electric and magnetic charges are defined as appropriate integrals over S^2 in the near-horizon region, respectively:

$$q_\Lambda = \frac{g}{4\pi} \int_{S^2} 16\pi G_N^{(4)} \frac{\delta S_{4d}}{\delta F^\Lambda}, \quad p^\Lambda = \frac{g}{4\pi} \int_{S^2} F^\Lambda. \quad (5.8)$$

Here $G_N^{(4)}$ is the 4d Newton constant, related to the 5d one by

$$\frac{4\pi}{G_N^{(5)}g} = \frac{1}{G_N^{(4)}} , \quad (5.9)$$

while S_{4d} is the 4d supergravity action. The 4d black holes we are interested in have both electric and magnetic charges. The magnetic charge $p^0 = 1$ is equal to the first Chern class of the Hopf fibration. On the other hand, we fix the redundancy (5.4) by setting the remaining magnetic charges to zero. In Appendix E we compute the relation of the 5d charges Q_I and angular momentum J measured at infinity, with the 4d charges measured at the horizon. We should be careful that only massless vector fields are associated to conserved charges. We indicate as \mathbb{B}^I_J the matrix of linear redefinitions such that $\mathbb{B}^I_J A_\mu^J$ are the 5d mass eigenstates in the AdS₅ vacuum, and we take the index \mathfrak{T} to run only over the massless vectors $\mathbb{B}^{\mathfrak{T}}_J A_\mu^J$. The corresponding conserved charges are $Q_{\mathfrak{T}} \equiv Q_J (\mathbb{B}^{-1})^J_{\mathfrak{T}}$. We find

$$\begin{aligned} p^0 &= 1 , & q_0 &= 4G_N^{(4)}g^2 J + \frac{1}{3}C_{IJK}\xi^I\xi^J\xi^K , \\ p^I &= 0 , & q_{\mathfrak{T}} &= 4G_N^{(4)}g^2 Q_{\mathfrak{T}} + \frac{1}{2}C_{\mathfrak{T}JK}\xi^J\xi^K , \end{aligned} \quad (5.10)$$

where $J_1 = J_2 \equiv J$, while the ‘‘non-conserved charges’’ $q_{J \neq \mathfrak{T}}$ will be fixed by the equations of motion. Notice that the conserved charges $Q_{\mathfrak{T}}$ are the same, but possibly in a different basis, as the charges Q_a introduced in Sections 3.2 and 3.3.²⁰

Using a symplectic covariant notation, electric and magnetic charges form a symplectic vector

$$\mathcal{Q} = (p^\Lambda, q_\Lambda) . \quad (5.11)$$

One also defines

$$\vec{\mathcal{P}} = (0, \vec{P}_\Lambda) , \quad \vec{\mathcal{Q}} = \langle \vec{\mathcal{P}}, \mathcal{Q} \rangle , \quad (5.12)$$

where vectors are triplets and $\langle V, W \rangle = V_\Lambda W^\Lambda - V^\Lambda W_\Lambda$ is the symplectic-invariant antisymmetric form.

To find covariantly-constant spinors, we impose the following twisting ansatz:

$$\epsilon_i = -\vec{\mathcal{Q}} \cdot \vec{\sigma}_i^j \Gamma^{\hat{t}\hat{r}} \epsilon_j , \quad (5.13)$$

whose square gives $\vec{\mathcal{Q}} \cdot \vec{\mathcal{Q}} = 1$. Here $\Gamma^{\hat{t}\hat{r}}$ is the antisymmetric product of two gamma matrices with flat indices \hat{t} and \hat{r} . We choose a gauge in which $\mathcal{Q}^1 = \mathcal{Q}^2 = 0$ and

$$\mathcal{Q}^3 = -1 \quad (5.14)$$

²⁰Similarly, the restriction of C_{IJK} to $C_{\mathfrak{T}\mathfrak{J}\mathfrak{K}}$ with curly indices is the same, but possibly in a different basis, as the 't Hooft anomaly coefficients C_{abc} previously defined.

at the horizon, as in [68].

The remaining BPS conditions are in general complicated, but they simplify at the horizon [37–39]. First, Maxwell’s equations give

$$\mathcal{K}^u h_{uv} \langle \mathcal{K}^v, \mathcal{Q} \rangle = 0, \quad (5.15)$$

where we defined

$$\mathcal{K}^u = (0, k_\Lambda^u) \quad (5.16)$$

because we work in a duality frame with purely electric gaugings. In fact, (5.15) in this case is equivalent to

$$p^\Lambda k_\Lambda^u = 0 \quad (5.17)$$

that must hold in the full solution simply because of spherical symmetry (see Appendix E). Second, vanishing of the hyperino variation implies

$$\langle \mathcal{K}^u, \mathcal{V} \rangle = 0, \quad (5.18)$$

where $\mathcal{V}(z, \bar{z}) = e^{\mathcal{K}/2}(X^\Lambda, F_\Lambda)$ is the covariantly-holomorphic section defined in (C.3) and $F_\Lambda = \partial_\Lambda F(X)$. Third, we have the attractor equations²¹

$$\frac{\partial}{\partial z^I} \left(\frac{\mathcal{Z}}{\mathcal{L}} \right) = 0, \quad \frac{\mathcal{Z}}{\mathcal{L}} = 2ig^2 L_S^2, \quad (5.19)$$

where the derivatives are with respect to the physical scalars z^I and we defined

$$\mathcal{Z} = \langle \mathcal{Q}, \mathcal{V} \rangle, \quad \mathcal{L} = \langle \mathcal{P}^3, \mathcal{V} \rangle. \quad (5.20)$$

The equation on the right in (5.19) determines L_S , and thus the horizon area.

5.2 Example: the conifold

We apply the general strategy to the case of the conifold. We start with the 5d $\mathcal{N} = 2$ gauged supergravity with $n_V = 2$ vector multiplets and $n_H = 2$ hypermultiplets constructed in Section 7 of [34] (called the “second model” in that paper), obtained from a consistent reduction of 10d type IIB supergravity on $T^{1,1}$ that preserves the $SU(2)^2 \times U(1)$ isometry. In Appendix B.1 we have recast its action as in the general formalism, and in Appendix D.1 we have reduced it down to 4d $\mathcal{N} = 2$ gauged supergravity. We are now ready to look for BPS near-horizon black hole solutions.

²¹There is an extra factor of 2 in front of L_S^2 compared to [60, 3, 68] due to the different normalization of kinetic terms in the Lagrangian (C.2): this is noticed footnote 4 of [65] and in footnote 10 of [21].

Using (B.46) and (B.47), the conditions (5.14) and (5.17) take the form:

$$\begin{cases} P_0^3 = -1 \\ k_0^u = 0 \end{cases} \Rightarrow \quad b_{1,2}^\Omega = c_{1,2}^\Omega = 0, \quad \xi^1 = -\xi^2 = -\frac{1}{3}, \quad (5.21)$$

where $b_{1,2}^\Omega, c_{1,2}^\Omega, a, \phi, C_0, u$ are the scalar fields in hypermultiplets. In fact, since (5.17) must hold in the whole solution, so (5.21) does. Using the form (B.47) of the moment maps, this is consistent with $\mathcal{Q}^1 = \mathcal{Q}^2 = 0$. The hyperino condition (5.18) then gives

$$X^1 + X^2 = 0 \quad (5.22)$$

at the horizon, where X^Λ are the holomorphic sections. The fields C_0 and ϕ are not fixed by the equations of motion. However, together they form the axiodilaton of type IIB supergravity and are thus fixed by the boundary conditions that set them in terms of the complexified gauge coupling of the boundary theory. As apparent from the expression of k_2^u in (B.46), a is a Stückelberg field that breaks an Abelian gauge symmetry and is eaten up as the corresponding gauge field becomes massive via Higgs mechanism.

The remaining BPS conditions are the attractor equations (5.19). Given C_{IJK} in (B.42), the prepotential is

$$F(X) = \frac{\check{X}^1((\check{X}^2)^2 - (\check{X}^3)^2)}{X^0} \quad \text{where} \quad \check{X}^I = X^I + \xi^I X^0. \quad (5.23)$$

Using special coordinates $z^I = X^I/X^0$ as well as homogeneity of the prepotential $F(X)$, one can easily show that the two equations in (5.19) are equivalent to

$$\partial_\Lambda \left[e^{-\kappa/2} \left(\mathcal{Z}(X) - 2ig^2 L_S^2 \mathcal{L}(X) \right) \right] = 0, \quad (5.24)$$

where the derivatives are with respect to *independent* sections X^Λ . In these equations L_S should be regarded as one of the unknowns. Notice that (5.19) or (5.24) give, in general, isolated solutions in terms of (z^I, L_S) , however the sections X^Λ are only fixed up to the “gauge” redundancy (related to Kähler transformations on \mathcal{KM}) $X^\Lambda \rightarrow e^f X^\Lambda$. In order to remove the redundancy, we choose to fix $\mathcal{L}(X)$ to a constant, which can elegantly be imposed by taking a derivative of the square bracket in (5.24) with respect to L_S^2 as well. More precisely, expanding \mathcal{Z} and \mathcal{L} using (B.47), we consider the following set of equations:

$$\begin{aligned} \partial_\Lambda \left[\frac{X^1((X^2)^2 - (X^3)^2)}{(X^0)^2} + \hat{q}_\Lambda X^\Lambda - 2ig^2 L_S^2 \left(3X^1 - X^0 - 2e^{-4u}(X^1 + X^2) - \alpha \right) \right] &= 0 \\ \frac{\partial}{\partial L_S^2} \left[\frac{X^1((X^2)^2 - (X^3)^2)}{(X^0)^2} + \hat{q}_\Lambda X^\Lambda - 2ig^2 L_S^2 \left(3X^1 - X^0 - 2e^{-4u}(X^1 + X^2) - \alpha \right) \right] &= 0 \end{aligned} \quad (5.25)$$

where

$$\widehat{q}_I = q_I - \frac{1}{2}C_{IJK}\xi^J\xi^K, \quad \widehat{q}_0 = q_0 - \frac{1}{3}C_{IJK}\xi^I\xi^J\xi^K. \quad (5.26)$$

The first line is the same as (5.24), except for the addition of the constant α that does not affect the equations. The second line fixes the gauge $\mathcal{L} = \alpha$. Notice that (5.22) should be imposed after solving (5.25).

From the point of view of AdS/CFT, only massless vector fields correspond to symmetries of the boundary theory and only their charges are conserved and fixed by the boundary conditions. On the contrary, the “charges” under massive vector fields are not conserved, and their radial profile should be determined by the equations of motion. The spectrum of the 5d supergravity under consideration around its supersymmetric AdS₅ vacuum was computed in [34] and we report it in our conventions in (B.49). In the basis

$$\begin{aligned} A^R &\equiv A^1 - 2A^2, & A^3, & & A^W &\equiv A^1 + A^2, \\ k_R &\equiv \frac{1}{3}(k_1 - k_2), & k_3, & & k_W &\equiv \frac{1}{3}(2k_1 + k_2), \end{aligned} \quad (5.27)$$

it turns out that A^R (corresponding to the R-symmetry) and A^3 are massless, while A^W is massive because of Higgs mechanism eating up the Stückelberg field a . In (5.27) we have indicated also the Killing vectors of the corresponding gauged isometries. On the black hole background the mass eigenstates may change (because the gauge kinetic functions have a non-trivial radial profile), however the fact that

$$k_R = k_3 = 0 \quad (5.28)$$

everywhere — which follows from (5.21) — guarantees that there is no hypermultiplet source in the 5d Maxwell equations (E.3) and thus the Page charges Q_R and Q_3 are conserved (while Q_W is not).

Indeed, the variation in (5.25) with respect to X^2 gives the complex equation

$$2\frac{X^1X^2}{(X^0)^2} + \widehat{q}_2 + 4ig^2L_S^2e^{-4u} = 0 \quad (5.29)$$

that fixes u and the “non-conserved charge” q_2 in terms of the sections and L_S . We can then use the hyperino condition (5.22) to eliminate X^2 as well. Notice that the second condition in (5.21) implies that in 5d we cannot turn on a “flat connection” for A^W along the circle.

We are left with the unknowns X^0, X^1, X^3, L_S^2 . One can check that, when (5.22) and (5.29) are in place, the remaining equations in (5.25) are equivalent to the conditions of extremization of the function

$$\mathcal{S} = \beta \left[\frac{X^1((X^1)^2 - (X^3)^2)}{(X^0)^2} + \widehat{q}_0X^0 + 3\widehat{q}_RX^1 + \widehat{q}_3X^3 - 2ig^2L_S^2(3X^1 - X^0 - \alpha) \right] \quad (5.30)$$

with respect to the variables X^0, X^1, X^3, L_S^2 . Here β is a constant included for later convenience, while \widehat{q}_R is the charge with respect to the massless vector A_R :

$$\widehat{q}_R = \frac{\widehat{q}_1 - \widehat{q}_2}{3} = \frac{g}{4\pi} \int_{S^2} 16\pi G_N^{(4)} \frac{\delta S_{4d}}{\delta FR} - \frac{1}{6} (C_{1JK} - C_{2JK}) \xi^J \xi^K = 4g^2 G_N^{(4)} Q_R. \quad (5.31)$$

It is encouraging that we find an extremization problem in which only conserved charges appear. Since \mathcal{S} is homogeneous in X^Λ of degree 1 except for the term involving α , it follows that $\mathcal{S}|_{\text{crit}} = 2i\alpha\beta g^2 L_S^2$ at the critical point. With the choice

$$\alpha\beta = \frac{\pi}{2iG_N^{(4)} g^2} \quad (5.32)$$

we obtain that $\mathcal{S}|_{\text{crit}}$ is the black hole entropy:

$$\mathcal{S}|_{\text{crit}} = \frac{4\pi L_S^2}{4G_N^{(4)}} = S_{\text{BH}}, \quad (5.33)$$

and therefore \mathcal{S} is the entropy function. Using (5.10) and (5.26) we can express the 4d charges $\widehat{q}_0, \widehat{q}_{\mathfrak{T}}$ computed at the horizon in terms of the 5d black hole charges $J, Q_{\mathfrak{T}}$ computed at infinity:

$$\mathcal{S} = \frac{1}{\alpha} \left[\frac{\pi}{2iG_N^{(4)} g^2} \frac{(X^1)^3 - X^1(X^3)^2}{(X^0)^2} - 2\pi i (JX^0 + 3Q_R X^1 + Q_3 X^3) - 2\pi i \Lambda (3X^1 - X^0 - \alpha) \right], \quad (5.34)$$

where we redefined the Lagrange multiplier $L_S^2 = 2iG_N^{(4)} \Lambda$ for convenience.

It remains to spell out the AdS/CFT dictionary between gravity and field theory charges. First, the gauge group ranks in field theory are determined by (see Appendix E.1)

$$N^2 = \frac{8\pi}{27 G_N^{(5)} g^3} = \frac{2}{27 G_N^{(4)} g^2}. \quad (5.35)$$

This is in agreement with (4.2) using $a = \frac{27}{64} N^2$ for the Klebanov-Witten theory. Second, the angular momentum J is the same in gravity and in field theory. Third, the electric charges are identified as

$$r = 2Q_R, \quad Q_B = \frac{4}{3} Q_3. \quad (5.36)$$

This is determined as follows. From (B.36) we infer that the gravitino components have charge $Q_R = \pm \frac{1}{2}$. In the boundary field theory, the corresponding operators are of the schematic form $\text{Tr}(F_{\mu\nu} \Gamma^\nu \lambda)$ (where F is a field strength and λ a gaugino) and have charge $r = \pm 1$ under $U(1)_R$. We deduce the first relation in (5.36). Obtaining the second relation is

more subtle because no supergravity field is charged under A^3 : what is charged are massive particles obtained from D3-branes wrapped on the 3-cycle of $T^{1,1}$, corresponding to dibaryon operators $A_{1,2}^N$ or $B_{1,2}^N$ in field theory. The 5d supergravity gauge field A^3 comes from the reduction of the Ramond-Ramond field strength F_5^{RR} of 10d type IIB supergravity on $T^{1,1}$. Therefore, from the 10d flux quantization condition we can deduce the 5d charge quantization condition $4Q_3/3N \in \mathbb{Z}$ (see the details in Appendix E.1). In field theory the dibaryon operators have charge $Q_B = \pm N$, implying the second relation in (5.36). Alternatively, we could compare the Chern-Simons terms restricted to massless vector fields in the 5d Lagrangian with the 't Hooft anomalies of the boundary theory. Taking into account the 't Hooft anomalies $\text{Tr}(r^3) = \frac{3}{2}N^2$ and $\text{Tr}(rQ_B^2) = -2N^2$ at leading order in N , the restriction of the 5d Chern-Simons action in (B.2) to $A^W \rightarrow 0$ matches the general expression

$$S_{\text{CS}} = \frac{g^3}{24\pi^2} \int \text{Tr}(Q_a Q_b Q_c) F^a \wedge F^b \wedge A^c \quad (5.37)$$

after setting $A^R \rightarrow 2A_r$ and $A^3 \rightarrow \frac{4}{3}A_B$. These correspond to (5.36).

Rewriting the entropy function (5.34) in terms of field theory charges, we find

$$\mathcal{S} = \frac{1}{\alpha} \left[-\frac{27\pi i N^2}{4} \frac{(X^1)^3 - X^1(X^3)^2}{(X^0)^2} - 2\pi i \left(JX^0 + \frac{3}{2}rX^1 + \frac{3}{4}Q_B X^3 \right) - 2\pi i \Lambda \left(3X^1 - X^0 - \alpha \right) \right]. \quad (5.38)$$

This exactly matches the entropy function (3.35) we found in field theory from the large N limit of the superconformal index of the Klebanov-Witten theory, after the change of coordinates $X^0 \rightarrow 2\alpha\tau$, $X^1 \rightarrow 2\alpha X_R/3$, $X^3 \rightarrow 4\alpha X_B/3$.

Acknowledgements

We thank Arash Arabi Ardehali, Seyed Morteza Hosseini, Paolo Milan, and Antoine Van Proeyen for useful conversations and correspondence. F.B., S.S., and Z.Z. are partially supported by the MIUR-PRIN contract 2015 MP2CX4, and by the INFN ‘‘Iniziativa Specifica ST&FI’’. F.B. is also supported by the MIUR-SIR grant RBSI1471GJ, and by the ERC-COG grant NP-QFT no. 864583. A.Z. is partially supported by the INFN, the ERC-STG grant 637844-HBQFTNCER, and the MIUR-PRIN contract 2017 CC72MK003.

A Subleading effect of simplifications

A.1 Simplifications of the building block

We want to show that the terms neglected in passing from (2.26) to (2.27) are subleading at large N . We will first analyze the effect of dropping the term $\omega(d-c)/N$ from the arguments of the gamma functions, in all those terms with $\gamma \neq \delta$. We will later estimate the contribution from the terms with $\gamma = \delta$ that were discarded from the sum.

Defining

$$f(z) = \sum_{\gamma \neq \delta}^{\tilde{N}} \log \tilde{\Gamma} \left(z + \omega \frac{\delta - \gamma}{\tilde{N}}; ab\omega, ab\omega \right), \quad (\text{A.1})$$

we want to show that

$$\left| f(z + C\omega/\tilde{N}) - f(z) \right| \leq \mathcal{O}(N \log N), \quad (\text{A.2})$$

where $C = (d-c)/ab$, $z = \Delta + \omega(d-c + as + br)$ and $c, d = 1, \dots, ab$, $r = 0, \dots, a-1$, $s = 0, \dots, b-1$. Without loss of generality we can assume $C > 0$, because the case $C < 0$ is analogous while $C = 0$ is trivial. As in [6], we discard the Stokes lines $\Delta \in \mathbb{Z} + \mathbb{R}\omega$ except the point $\Delta = 0$, because the limit we compute would be singular along those lines anyway. If Δ is not on a Stokes line, then the restriction of f to a straight line in the complex plane passing through the points z and $z + \omega$ is a C^∞ complex function. In the case $\Delta = 0$, instead, we consider the restriction of f to a straight closed segment from z to $z + C\omega/\tilde{N}$ and one can check that f is C^∞ along that segment, because for $\gamma \neq \delta$ the segment, suitably shifted, does not hit zeros nor poles of any of the gamma functions in (A.1) (in both cases, f is a holomorphic function in a neighbourhood of the restricted domain). A complex analogue of the Mean Value Theorem (MVT) then states that

$$\begin{aligned} \operatorname{Re} \frac{f(z + C\omega/\tilde{N}) - f(z)}{\omega} &= \frac{C}{\tilde{N}} \operatorname{Re} f'(z + \bar{c}_1\omega/\tilde{N}) \\ \operatorname{Im} \frac{f(z + C\omega/\tilde{N}) - f(z)}{\omega} &= \frac{C}{\tilde{N}} \operatorname{Im} f'(z + \bar{c}_2\omega/\tilde{N}) \end{aligned} \quad (\text{A.3})$$

with $\bar{c}_1, \bar{c}_2 \in (0, C)$. Summing the absolute values, it follows the bound

$$\left| \frac{f(z + C\omega/\tilde{N}) - f(z)}{\omega} \right| \leq \frac{1}{\tilde{N}} \left(\left| f'(z + \bar{c}_1\omega/\tilde{N}) \right| + \left| f'(z + \bar{c}_2\omega/\tilde{N}) \right| \right) \quad (\text{A.4})$$

where we used $|C| \leq 1 - \frac{1}{ab} < 1$. It is therefore sufficient to show that

$$\frac{1}{\tilde{N}} \left| f'(z + \bar{c}\omega/\tilde{N}) \right| \leq \mathcal{O}(N \log N) \quad (\text{A.5})$$

for any $\bar{c} \in (0, C)$. Notice that $0 < \bar{c} < 1 - \frac{1}{ab}$.

We reason as follows. For $\Delta \notin \mathbb{Z} + \mathbb{R}\omega$, the arguments of the elliptic gamma functions in (A.1) remain at an \tilde{N} -independent distance from the zeros and poles, that in our case are placed at the points

$$u_{0,i} = (1+i)ab\omega, \quad u_{\infty,j} = (1-j)ab\omega \quad \text{for} \quad i, j \in \mathbb{Z}_{\geq 1}, \quad (\text{A.6})$$

respectively. The orders of the zeros and poles are i and j respectively. The ratio $|\tilde{\Gamma}'/\tilde{\Gamma}|$ is bounded on the range of possible arguments, therefore

$$\frac{1}{\tilde{N}} \left| f'(z + \bar{c}\omega/\tilde{N}) \right| \leq \tilde{N} \max_{t \in [-ab, 3ab-a-b]} \left| \frac{\tilde{\Gamma}'(\Delta + t\omega; ab\omega, ab\omega)}{\tilde{\Gamma}(\Delta + t\omega; ab\omega, ab\omega)} \right| = \mathcal{O}(\tilde{N}). \quad (\text{A.7})$$

The case $\Delta = 0$ is more subtle since, as \tilde{N} grows, the arguments of some of the gamma functions can get increasingly close to zeros or poles instead of staying at an \tilde{N} -independent distance, and the \tilde{N} -independent bound above does not apply. This happens when

$$z = \bar{u}_{0,i} \in \{u_{0,i}, u_{0,i} \pm \omega\} \quad \text{or} \quad z = \bar{u}_{\infty,j} \in \{u_{\infty,j}, u_{\infty,j} \pm \omega\}. \quad (\text{A.8})$$

One can easily see that for $\Delta = 0$, z can range from $(1-ab)\omega$ to $(3ab-a-b-1)\omega$, so that the problematic points we may approach are the simple zero at $u_{0,1}$, the simple pole at $u_{\infty,1}$, and the double pole at $u_{\infty,2}$.

We now introduce a few results for later use. For a meromorphic function g whose zeros include $\{z_i\}$ of order $\{m_i\}$ and whose poles include $\{p_j\}$ of order $\{n_j\}$, one can write

$$g(z) = \frac{\prod_i (z - z_i)^{m_i}}{\prod_j (z - p_j)^{n_j}} s(z), \quad (\text{A.9})$$

where $s(z)$ is meromorphic with zeros and poles at the remaining zeros and poles of g that were not included in $\{z_i\}$ and $\{p_j\}$. Taking the derivative of this expression and computing g'/g , one finds

$$\frac{g'(z)}{g(z)} = \sum_i \frac{m_i}{z - z_i} - \sum_j \frac{n_j}{z - p_j} + h(z), \quad (\text{A.10})$$

where $h(z) = s'(z)/s(z)$ is meromorphic with simple poles at the remaining zeros and poles of g that were not included in $\{z_i\}$ and $\{p_j\}$. Therefore we can apply (A.10) to the meromorphic function $\tilde{\Gamma}$ and say that

$$\begin{aligned} \frac{1}{\tilde{N}} \left| f'(z + \bar{c}\omega/\tilde{N}) \right| &\leq \frac{1}{\tilde{N}} \sum_{\gamma \neq \delta}^{\tilde{N}} \left| \frac{\tilde{\Gamma}'(z + u_{\gamma,\delta}^{\bar{c}}; ab\omega, ab\omega)}{\tilde{\Gamma}(z + u_{\gamma,\delta}^{\bar{c}}; ab\omega, ab\omega)} \right| \\ &\leq \frac{1}{\tilde{N}} \sum_{\gamma \neq \delta}^{\tilde{N}} \left(\frac{1}{|z + u_{\gamma,\delta}^{\bar{c}} - 2ab\omega|} + \frac{1}{|z + u_{\gamma,\delta}^{\bar{c}}|} + \frac{2}{|z + u_{\gamma,\delta}^{\bar{c}} + ab\omega|} \right) + (\tilde{N} - 1)K \end{aligned} \quad (\text{A.11})$$

where we defined

$$u_{\gamma,\delta}^{\bar{c}} = \omega \frac{\delta - \gamma + \bar{c}}{\tilde{N}}, \quad K = \max_{t \in [-ab, 3ab-a-b]} |h_{\tilde{\Gamma}}(t\omega)| \quad (\text{A.12})$$

and $h_{\tilde{\Gamma}}$ is the meromorphic function associated to $\tilde{\Gamma}$ in (A.10). We can bound its value with an \tilde{N} -independent quantity because it is holomorphic on the range of possible arguments. If $z \neq \bar{u}_{0,1}, \bar{u}_{\infty,1}, \bar{u}_{\infty,2}$, the outlying sums in (A.11) will be of order $\mathcal{O}(\tilde{N})$ since $z + u_{\gamma,\delta}^{\bar{c}}$ will be at least at a distance $|\omega|$ away from the zeros and poles. To complete our proof when $z = \bar{u}_{0,1}, \bar{u}_{\infty,1}, \bar{u}_{\infty,2}$, we now need to bound the quantities

$$R_x = \frac{1}{N} \sum_{\gamma \neq \delta}^N \frac{1}{\left| x + \frac{\delta - \gamma + \bar{c}}{N} \right|} \quad \text{with } x = 0, \pm 1, \quad (\text{A.13})$$

where we wrote N in place of \tilde{N} in order not to clutter the formulae. We recall that $0 < \bar{c} < 1 - 1/ab$. Considering $x = 0$ first, we reparametrize the sum in terms of $\delta - \gamma$ so that, after some manipulations, it becomes

$$R_0 = \sum_{M=1}^{N-1} \left(\frac{N-M}{M+\bar{c}} + \frac{N-M}{M-\bar{c}} \right) < 2 \sum_{M=1}^{N-1} \frac{N-M}{M-\bar{c}}. \quad (\text{A.14})$$

The summand on the right is a positive decreasing function of M , therefore it can be bound by its integral:

$$R_0 < \frac{2(N-1)}{1-\bar{c}} + 2 \int_1^{N-1} \frac{N-x}{x-\bar{c}} dx = \mathcal{O}(N \log N). \quad (\text{A.15})$$

To ensure convergence of sums and integrals it is crucial to recall that $1 - \bar{c} > (ab)^{-1}$. In a similar way, for $x = +1$ we can write

$$R_1 = \sum_{M=1}^{N-1} \left(\frac{N-M}{N+M+\bar{c}} + \frac{N-M}{N-M+\bar{c}} \right) < \sum_{M=1}^{N-1} 2 = \mathcal{O}(N), \quad (\text{A.16})$$

while for $x = -1$ we can write

$$R_{-1} = \sum_{M=1}^{N-1} \left(\frac{N-M}{N-M-\bar{c}} + \frac{N-M}{N+M-\bar{c}} \right) < 2 \sum_{M=1}^{N-1} \frac{N-M}{N-M-\bar{c}} < \frac{2(N-1)}{1-\bar{c}} = \mathcal{O}(N). \quad (\text{A.17})$$

It remains to show that the terms we discarded from (2.26) when substituting the condition $i \neq j$ with the condition $\gamma \neq \delta$ give a subleading contribution. These are the terms in (2.26) with $\gamma = \delta$, whose total contribution is

$$\Phi = \tilde{N} \sum_{r=0}^{a-1} \sum_{s=0}^{b-1} \sum_{c \neq d}^{ab} \log \tilde{\Gamma} \left(\Delta + \omega \frac{d-c}{N} + \omega(d-c+as+br); ab\omega, ab\omega \right). \quad (\text{A.18})$$

We need to show that this is subleading in the large N limit. We will bound the absolute value of the summand for all possible $c \neq d, r, s$ and drop the sums since they give an overall order $\mathcal{O}(1)$ factor. After choosing a branch of the logarithm, the phases of $\tilde{\Gamma}$ can clearly only give an order \tilde{N} contribution to (the imaginary part of) Φ .

For what concerns the absolute value of $\tilde{\Gamma}$, reasoning in a very similar way to the $\gamma \neq \delta$ case discussed above, we see that if Δ is not on a Stokes line then $|\log |\tilde{\Gamma}|$ is bounded above by an N -independent quantity and thus Φ is of order $\mathcal{O}(N)$. When $\Delta = 0$, the argument of $\tilde{\Gamma}$ can only approach zeros or poles if $z = \omega(d - c + as + br) \in \{u_{0,1}, u_{\infty,1}, u_{\infty,2}\}$. Using (A.9), we can write

$$\log \left| \tilde{\Gamma} \left(z + \omega \frac{d-c}{N}; ab\omega, ab\omega \right) \right| = \log \left| \frac{(z + \omega \frac{d-c}{N} - u_{0,1}) s_{\tilde{\Gamma}} \left(z + \omega \frac{d-c}{N} \right)}{(z + \omega \frac{d-c}{N} - u_{\infty,1}) (z + \omega \frac{d-c}{N} - u_{\infty,2})^2} \right| \quad (\text{A.19})$$

where $s_{\tilde{\Gamma}}$ is a function which is regular at $u_{\infty,1}, u_{\infty,2}$ and non-zero at $u_{0,1}$. We can therefore bound $|\log |s_{\tilde{\Gamma}}||$ over its possible arguments with an N -independent constant, so that it contributes to Φ at order $\mathcal{O}(N)$. When $z = u_{0,1}, u_{\infty,1}, u_{\infty,2}$, only one of the factors multiplying $s_{\tilde{\Gamma}}$ is of order $\mathcal{O}(\log N)$ while the other two do not approach zero and can be bounded by an N -independent constant. Explicitly,

$$\tilde{N} \left| \log \left| \tilde{\Gamma} \left(z + \omega \frac{d-c}{N}; ab\omega, ab\omega \right) \right| \right| \leq 2\tilde{N} \left| \log \left| \omega \frac{d-c}{N} \right| \right| + \mathcal{O}(N) = \mathcal{O}(N \log N). \quad (\text{A.20})$$

A.2 $\text{SU}(N)$ vs. $\text{U}(N)$ holonomies

In what follows, as it is done in Section 2 and Section 3, in order to parametrize the $\text{SU}(N)$ holonomies u^{SU} we introduce $\text{U}(N)$ holonomies u^{U} , constrained by

$$\sum_{i=1}^N u_i^{\text{U}} = 0. \quad (\text{A.21})$$

With the choice of bases for the Cartan subalgebras of $\text{SU}(N)$ and $\text{U}(N)$ required to write the BA operators as in (2.12), the relation between the two sets of holonomies when expressing a generic element of the Cartan subalgebra of $\text{SU}(N)$ is

$$u_i^{\text{U}} = u_i^{\text{SU}} \quad \text{for } i \neq N, \quad u_N^{\text{U}} = - \sum_{j=1}^{N-1} u_j^{\text{SU}}. \quad (\text{A.22})$$

Note that the holonomies are only defined modulo \mathbb{Z} .

The $\text{SU}(N)$ superconformal index defined by (2.9) contains a sum over $\{m_i^{\text{SU}}\}$ that picks up (representatives of) solutions to the BAEs whose residue can contribute to the index, as

explained in [24] and made explicit in (2.17). Under a shift $\{m_i^{\text{SU}}\}$ of the $\text{SU}(N)$ holonomies, the $\text{U}(N)$ holonomies shift by corresponding amounts given by

$$m_i^{\text{U}} = m_i^{\text{SU}}, \quad m_N^{\text{U}} = - \sum_{j=1}^{N-1} m_j^{\text{SU}}. \quad (\text{A.23})$$

Given these identifications for the holonomies and shifts, the $\text{SU}(N)$ quantities are always equal to the first $N-1$ $\text{U}(N)$ quantities, so that in the following we will drop the superscripts SU and U , remembering that $u_{1,\dots,N-1}$ and $m_{1,\dots,N-1}$ are independent while u_N and m_N are determined by (A.22) and (A.23), respectively.

One might then worry that the choice of $\{m_j\}$ given in (2.21) is not allowed, since the last integer m_N there does not satisfy (A.23). Specifically, let us choose

$$m_j \in \{1, \dots, ab\} \quad \text{such that} \quad m_j = j \pmod{ab}, \quad \text{for } j = 1, \dots, N-1, \quad (\text{A.24})$$

so that m_N is fixed by (A.23) to be a negative integer of $\mathcal{O}(N)$. To match with the choice in (2.21), we want to replace this with $m_N = N \pmod{ab}$ and in $\{1, \dots, ab\}$. We will show that this replacement does not affect the value of \mathcal{Z} to leading order in N . This will be done in two steps. We will first show that the function \mathcal{Z} evaluated on a configuration $\{u_1, \dots, u_N\}$ which is obtained from the basic solution by shifting one or more variables u_i by multiples of $2ab\omega$ (or even of $ab\omega$, in many cases), is the same as \mathcal{Z} evaluated on the basic solution. Using this property, \mathcal{Z} is unaltered if evaluated on the following shifted value of m_N :

$$\tilde{m}_N \in \{1, \dots, 2ab\} \quad \text{such that} \quad \tilde{m}_N = \left(- \sum_{i=1}^{N-1} m_i \right) \pmod{2ab}. \quad (\text{A.25})$$

We will then show that the contribution to \mathcal{Z} of the single holonomy u_N is subleading, provided $\tilde{m}_N \in \{1, \dots, 2ab\}$. Therefore, choosing instead $m_N = N \pmod{ab}$ and in $\{1, \dots, ab\}$ as we did in (2.21) does not change \mathcal{Z} at leading order in N . This completes the proof.

As shown in [24], when evaluated on solutions to the BAEs, the function \mathcal{Z} for a general semi-simple gauge group is invariant under independent shifts of any gauge holonomy by $ab\omega$. This is proven assuming that gauge and global symmetries are non-anomalous. In our case, this result only allows us to shift the u_i 's while preserving the $\text{SU}(N)$ constraint. This property does not allow us to independently shift the last holonomy u_N , since it is always fixed by the $\text{SU}(N)$ constraint. We now show that an *independent* shift of u_N by a multiple of $ab\omega$ is also an invariance of \mathcal{Z} for $\mathcal{N} = 4$ $\text{SU}(N)$ SYM, when this function is evaluated on the basic solution. In order to prove this, one has to use the property

$$\begin{aligned} & \tilde{\Gamma}(u + mab\omega, a\omega, b\omega) \\ &= (-e^{2\pi i u})^{-\frac{ab}{2}m^2 + \frac{a+b-1}{2}m} (e^{2\pi i \omega})^{-\frac{ab}{6}m^3 + \frac{ab(a+b)}{4}m^2 - \frac{a^2+b^2+3ab-1}{12}m} \theta_0(u, \omega)^m \tilde{\Gamma}(u, a\omega, b\omega) \end{aligned} \quad (\text{A.26})$$

that was proven in [24], the fact that the $U(N)$ BA operators are periodic modulo ω in the u_i 's, and the explicit form of the basic solution (2.20). Applying (A.26), we first have that

$$\begin{aligned} & \prod_{i \neq j} \tilde{\Gamma}(u_{ij} + \Delta + mab\omega(\delta_{iN} - \delta_{jN}); a\omega, b\omega) \\ &= e^{-\pi i abm^2(1+2\Delta) + 2\pi i(a+b-1)m \sum_i u_{iN} + \pi i ab(a+b)m^2\omega} \prod_i \frac{\theta_0(u_{Ni} + \Delta, \omega)^m}{\theta_0(u_{iN} + \Delta; \omega)^m} \prod_{i \neq j} \tilde{\Gamma}(u_{ij} + \Delta; a\omega, b\omega), \end{aligned} \quad (\text{A.27})$$

and so from (2.12), (2.18) and (2.20) one obtains

$$\begin{aligned} & \mathcal{Z}(u_i + mab\omega\delta_{iN}; a\omega, b\omega, \Delta) \\ &= \prod_i \left(\frac{\theta_0(u_{Ni} + \Delta_1, \omega) \theta_0(u_{Ni} + \Delta_2, \omega) \theta_0(u_{iN}, \omega) \theta_0(u_{iN} + \Delta_1 + \Delta_2, \omega)}{\theta_0(u_{iN} + \Delta_1, \omega) \theta_0(u_{iN} + \Delta_2, \omega) \theta_0(u_{Ni}, \omega) \theta_0(u_{Ni} + \Delta_1 + \Delta_2, \omega)} \right)^m \mathcal{Z}(u_i; a\omega, b\omega, \Delta) \\ &= (-1)^{m(N-1)} e^{2\pi i m \lambda} Q_N^{-m}(u_i; \omega, \Delta) \mathcal{Z}(u_i; a\omega, b\omega, \Delta) \\ &= \mathcal{Z}(u_i; a\omega, b\omega, \Delta). \end{aligned} \quad (\text{A.28})$$

In the steps above we also used the theta function reflection property

$$\theta_0(u; \omega) = -e^{2\pi i u} \theta_0(-u; \omega). \quad (\text{A.29})$$

More generally, we can show that this shift invariance is true for quiver gauge theories, when \mathcal{Z} is evaluated on the basic solution and the chemical potentials u_N^α are shifted by a multiple of $2ab\omega$ (or even of $ab\omega$, in many cases) simultaneously for all gauge groups $SU(N)_\alpha$. The steps are the same as in (A.28). We should notice that the expression for any particular Lagrange multiplier λ_α is more complicated than for $\mathcal{N} = 4$ SYM, but the sum of all Lagrange multipliers is simple:

$$e^{2\pi i \sum_{\alpha=1}^G \lambda_\alpha} = (-1)^{n_\chi(N-1)}, \quad (\text{A.30})$$

where α runs over the G $SU(N)$ gauge group factors and n_χ is the number of chiral multiplets in the theory. Performing these steps one obtains

$$\begin{aligned} & \mathcal{Z}(u_i^\alpha + mab\omega\delta_{iN}; a\omega, b\omega, \Delta) \\ &= \frac{e^{2\pi i m \sum_{\alpha=1}^G \lambda_\alpha} (-1)^{-mG(N-1) + mab(N-1)(n_\chi - G)}}{\left(\prod_\alpha Q_N^\alpha(u_i^\alpha; \omega, \Delta) \right)^m} \mathcal{Z}(u_i^\alpha; a\omega, b\omega, \Delta) \\ &= (-1)^{m(G - n_\chi)(ab+1)(N-1)} \mathcal{Z}(u_i^\alpha; a\omega, b\omega, \Delta). \end{aligned} \quad (\text{A.31})$$

There are now different cases in which the sign in the last line disappears. First, in the case of toric quiver gauge theories one uses the relation [48, 49]

$$G - n_\chi + N_W = 0 \quad (\text{A.32})$$

between the number of gauge groups, of chiral multiplets, and of superpotential terms, as well as the fact that the number N_W of superpotential terms is even, to show that the sign disappears. Second, if N is odd then the sign disappears. Third, if the coprime integers a, b are both odd²² then the sign disappears. Fourth and most importantly, if we take m even then the sign disappears.

We now proceed to show that the contribution to \mathcal{Z} of a single holonomy u_i is subleading, provided that $m_{i < N} \in \{1, \dots, ab\}$ and $m_N \in \{1, \dots, 2ab\}$. In the building block Ψ defined in (2.22), the contribution of a single holonomy u_i consists of the two terms

$$\Phi_i^\pm \equiv \sum_{j(\neq i)}^N \log \tilde{\Gamma} \left(z_\pm \pm \omega \frac{j-i}{N}; ab\omega, ab\omega \right), \quad (\text{A.33})$$

where we have defined

$$z_\pm \equiv \Delta \pm \omega (m_j - m_i) + \omega (as + br). \quad (\text{A.34})$$

In particular, for the case $i = N$ we will use the shift property just proven to substitute m_N with \tilde{m}_N defined in (A.25).

We will now show that Φ_i^\pm is subleading. In the case $i = N$ this will allow us to choose \tilde{m}_N as in (2.21). In order to do this we want to bound the absolute value of the summand $\log \tilde{\Gamma}$ in Φ_i^\pm . What follows will be completely analogous to the argument used to show that (A.18) is subleading. After choosing a branch of the logarithm, the phases of $\tilde{\Gamma}$ can only contribute at order $\mathcal{O}(N)$ to Φ_i^\pm . As before, we exclude Stokes lines and note that for $\Delta \neq 0$ we can bound $|\log |\tilde{\Gamma}|$ with an N -independent constant so that $|\Phi_i^\pm| = \mathcal{O}(N)$. For $\Delta = 0$, z_\pm have the range

$$z_\pm \in \{-2ab + 1, \dots, 4ab - a - b - 1\}\omega, \quad (\text{A.35})$$

and the argument of $\tilde{\Gamma}$ may approach zeros or poles when $z_\pm = \bar{u}_{0,1}, \bar{u}_{0,2}, \bar{u}_{\infty,1}, \bar{u}_{\infty,2}, \bar{u}_{\infty,3}$, which are defined in (A.8). If this is the case, further inspection is required. Using again (A.9), we can write

$$\log \tilde{\Gamma} \left(z_\pm \pm \omega \frac{j-i}{N}; ab\omega, ab\omega \right) = \log \left[\frac{\prod_{m=1}^2 (z_\pm \pm \omega \frac{j-i}{N} - u_{0,m})^m s_{\tilde{\Gamma}}(z_\pm \pm \omega \frac{j-i}{N})}{\prod_{n=1}^3 (z_\pm \pm \omega \frac{j-i}{N} - u_{\infty,n})^n} \right] \quad (\text{A.36})$$

where $s_{\tilde{\Gamma}}$ is a function that is regular at $u_{\infty,1}, u_{\infty,2}, u_{\infty,3}$, and non-zero at $u_{0,1}, u_{0,2}$. This allows us to bound $|\log |s_{\tilde{\Gamma}}||$ with an N -independent constant, and its contribution to Φ_i^\pm is of order $\mathcal{O}(N)$. When $z_\pm = \bar{u}_{0,1}, \bar{u}_{0,2}, \bar{u}_{\infty,1}, \bar{u}_{\infty,2}, \bar{u}_{\infty,3}$ the logarithms of the other factors

²²This restriction is quite uninfluential, because the set of pairs $\{\tau + \mathbb{Z}, \sigma + \mathbb{Z}\}$ such that $\tau/\sigma = a/b \in \mathbb{Q}_{>0}$ with a, b both odd is still dense in \mathbb{H}^2 .

are either bounded by an N -independent constant, or are of the form

$$\sum_{j \neq i}^N \left| \log \left| x \pm \frac{j-i}{N} \right| \right| \leq (N-1) \log N, \quad (\text{A.37})$$

where $x = 0, \pm 1$. Notice that the use of the shift property previously proved plays a major role here. If we tried to apply this argument directly without first shifting m_N , we would have to consider an $\mathcal{O}(N)$ number of poles or zeros whose order is also $\mathcal{O}(N)$. This would lead to an $\mathcal{O}(N^3 \log N)$ bound, which does not help. What we did shows that a single Φ_i^\pm is of order $\mathcal{O}(N \log N)$ for any choice of the corresponding m_i . In particular this allows us to choose $\tilde{m}_N = N \bmod ab \in \{1, \dots, ab\}$ as we do in (2.21), without affecting the leading behavior of the building block Ψ .

A.3 Generic N

Here we generalize the computation done in Section 2.1 and consider a generic N which is not necessarily a multiple of ab . We will exploit many of the arguments in Section A.1. Let $N = ab\tilde{N} + q$, where $q \in \{0, \dots, ab-1\}$. We need to examine the leading order contribution of the building block

$$\Psi = \sum_{r=0}^{a-1} \sum_{s=0}^{b-1} \sum_{i \neq j}^N \log \tilde{\Gamma} \left(\Delta + \omega \frac{j-i}{N} + \omega(m_j - m_i + as + br); ab\omega, ab\omega \right). \quad (\text{A.38})$$

As shown in the final part of Section A.2, the contribution to the building block of a single holonomy u_i is subleading. Therefore the contribution of the last q holonomies $u_{ab\tilde{N}+1}, \dots, u_N$ is also subleading and can be discarded. Now, the sum over $i \neq j$ only goes up to $ab\tilde{N}$, and we can decompose the indices as in (2.26). Neglecting the $\gamma = \delta$ terms using the same argument as after (A.18), we get

$$\Psi \simeq \sum_{r=0}^{a-1} \sum_{s=0}^{b-1} \sum_{\gamma \neq \delta=0}^{\tilde{N}-1} \sum_{c,d=0}^{ab-1} \log \tilde{\Gamma} \left(\Delta + \omega \frac{\delta - \gamma}{\tilde{N} + \frac{q}{ab}} + \omega \frac{d-c}{N} + \omega(d-c + as + br); ab\omega, ab\omega \right). \quad (\text{A.39})$$

As in Section A.1, we want to drop $\omega(d-c)/N$ in the argument of the elliptic gamma function, and we can use the same reasoning given there, with the minor change that (A.13) takes the form

$$\tilde{R}_x = \frac{1}{N + \frac{q}{ab}} \sum_{\gamma \neq \delta}^N \frac{1}{\left| x + \frac{\delta - \gamma + \bar{c}}{N + q/ab} \right|}, \quad x = 0, \pm 1. \quad (\text{A.40})$$

The same bounds as for R_x can be used here, since one can show that

$$\tilde{R}_0 = R_0, \quad \tilde{R}_{\pm 1} \leq R_{\pm 1}. \quad (\text{A.41})$$

We can then use (2.23), as we did in Section 2.1, to change the moduli of the elliptic gamma function from $(ab\omega, ab\omega)$ to (ω, ω) :

$$\begin{aligned}
\Psi &\simeq \sum_{r=0}^{a-1} \sum_{s=0}^{b-1} \sum_{\substack{\tilde{N}-1 \\ \gamma \neq \delta = 0}} \sum_{c,d=0}^{ab-1} \log \tilde{\Gamma} \left(\Delta + \omega \frac{\delta - \gamma}{\tilde{N} + \frac{q}{ab}} + \omega(d - c + as + br); ab\omega, ab\omega \right) \\
&= \sum_{r=0}^{a-1} \sum_{s=0}^{b-1} \sum_{\substack{\tilde{N}-1 \\ \gamma \neq \delta = 0}} \log \tilde{\Gamma} \left(\Delta + \omega \frac{\delta - \gamma}{\tilde{N} + \frac{q}{ab}} + \omega(1 - ab + as + br); \omega, \omega \right) \\
&= \frac{1}{(ab)^2} \sum_{r=0}^{a-1} \sum_{s=0}^{b-1} \sum_{\substack{\tilde{N}-1 \\ \gamma \neq \delta = 0}} \sum_{c,d=0}^{ab-1} \log \tilde{\Gamma} \left(\Delta + \omega \frac{\delta - \gamma}{\tilde{N} + \frac{q}{ab}} + \omega(1 - ab + as + br); \omega, \omega \right).
\end{aligned} \tag{A.42}$$

In the last equality, to make future steps clearer, we added a sum over c, d even though nothing depends on c and d .

Now, in order to get the desired result we trace our steps backwards. First, we will reintroduce the term $\omega(d - c)/N$ into the argument of the elliptic gamma functions. Then we will add to the sum in (A.42) the $\gamma = \delta$ terms to form the sum over $i \neq j$ up to $ab\tilde{N}$. Finally we will add terms containing the last q holonomies $u_{ab\tilde{N}+1}, \dots, u_N$ in order to build the complete sum up to N . These are the exact same steps we just performed to express Ψ as in (A.42) up to subleading terms, with the only difference being that the moduli of $\tilde{\Gamma}$ are now (ω, ω) rather than $(ab\omega, ab\omega)$. Therefore the same arguments can be used, with only slight modifications involving the number and order of zeros and poles, but since these are parametrized here by r and s that are N -independent, this is of no consequence. At this point, Ψ at leading order is

$$\Psi \simeq \frac{1}{(ab)^2} \sum_{r=0}^{a-1} \sum_{s=0}^{b-1} \sum_{i \neq j}^N \log \tilde{\Gamma} \left(\Delta + \omega \frac{j - i}{N} + \omega(1 - ab + as + br); \omega, \omega \right), \tag{A.43}$$

and using the result of [6] (that is our equation (2.29)) we obtain

$$\Psi \simeq -\frac{\pi i N^2}{3(a\omega)(b\omega)} \frac{1}{ab} \sum_{r=0}^{a-1} \sum_{s=0}^{b-1} B_3 \left([\Delta]'_\omega + \omega(as + br - ab) \right). \tag{A.44}$$

Then, using the property of Bernoulli polynomials (2.33), we finally get (2.34).

B 5d $\mathcal{N} = 2$ Abelian gauged supergravity

We report here the general form of 5d $\mathcal{N} = 2$ Abelian gauged supergravity with n_V vector multiplets and n_H hypermultiplets [57–59].²³ The graviton multiplet contains a graviton, a

²³A more complete discussion was developed in [70].

gravitino and a vector; each vector multiplet contains a vector, a gaugino and a real scalar; each hypermultiplet contains four real scalars and a hyperino. All fermions are Dirac, but can conveniently be doubled with a symplectic Majorana condition. We follow the notation of [71]. We use indices

$$I, J, K = 1, \dots, n_V + 1, \quad i, j = 1, \dots, n_V, \quad u, v = 1, \dots, 4n_H \quad (\text{B.1})$$

for the gauge fields A_μ^I , for the scalars ϕ^i in vector multiplets, and for the scalars q^u in hypermultiplets, respectively. The data that define the theory are:

1. A very special real manifold \mathcal{SM} of real dimension n_V .
2. A quaternionic-Kähler manifold \mathcal{QM} of real dimension $4n_H$.
3. A set of $n_V + 1$ Killing vectors on \mathcal{QM} compatible with the quaternionic-Kähler structure (if $n_H = 0$, $n_V + 1$ FI parameters not all vanishing).

The Killing vectors could be linearly dependent or vanish.

The bosonic Lagrangian is given by

$$\begin{aligned} 8\pi G_N^{(5)} e^{-1} \mathcal{L}_{5d} = & \frac{R_s}{2} - \frac{1}{2} \mathcal{G}_{ij}(\phi) \partial_\mu \phi^i \partial^\mu \phi^j - \frac{1}{2} h_{uv}(q) \mathcal{D}_\mu q^u \mathcal{D}^\mu q^v - \frac{1}{4} G_{IJ}(\phi) F_{\mu\nu}^I F^{J\mu\nu} \\ & + \frac{e^{-1}}{48} C_{IJK} \epsilon^{\mu\nu\rho\sigma\lambda} F_{\mu\nu}^I F_{\rho\sigma}^J A_\lambda^K - g^2 V(\phi, q). \end{aligned} \quad (\text{B.2})$$

Here $G_N^{(5)}$ is the 5d Newton constant, $e d^5x$ is the spacetime volume form, R_s is the scalar curvature, $F_{\mu\nu}^I$ is the field strength of A_μ^I , g is a coupling constant, and V is the scalar potential. Let us explain the other terms.

Very special geometry. The scalars ϕ^i are real coordinates on the very special real manifold \mathcal{SM} [72]. The latter is specified by the totally symmetric tensor C_{IJK} (which, controlling also the Chern-Simons couplings, should be suitably quantized) as the submanifold

$$\mathcal{SM} = \left\{ \mathcal{V}(\Phi) \equiv \frac{1}{6} C_{IJK} \Phi^I \Phi^J \Phi^K = 1 \right\} \subset \mathbb{R}^{n_V+1}. \quad (\text{B.3})$$

Here Φ^I are coordinates on \mathbb{R}^{n_V+1} , and give rise to “sections” $\Phi^I(\phi^i)$ on \mathcal{SM} . The metrics G_{IJ} and \mathcal{G}_{ij} for vector fields and vector multiplet scalar fields are

$$G_{IJ}(\phi) = -\frac{1}{2} \frac{\partial}{\partial \Phi^I} \frac{\partial}{\partial \Phi^J} \log \mathcal{V} \Big|_{\mathcal{V}=1}, \quad \mathcal{G}_{ij}(\phi) = \partial_i \Phi^I \partial_j \Phi^J G_{IJ} \Big|_{\mathcal{V}=1} \quad (\text{B.4})$$

where $\partial_i \equiv \partial/\partial \phi^i$. We recognize that \mathcal{G} is the pull-back of G from \mathbb{R}^{n_V+1} to \mathcal{SM} . From (B.3) it immediately follows

$$C_{IJK} \Phi^I \Phi^J \partial_i \Phi^K \Big|_{\mathcal{V}=1} = 0. \quad (\text{B.5})$$

With a little bit of algebra one then obtains a more explicit expression for G :

$$G_{IJ} = -\frac{1}{2}C_{IJK}\Phi^K + \frac{1}{8}C_{IKL}C_{JMN}\Phi^K\Phi^L\Phi^M\Phi^N \Big|_{\nu=1}. \quad (\text{B.6})$$

It follows that the kinetic term for vector multiplet scalars can also be written as

$$-\frac{1}{2}\mathcal{G}_{ij}\partial_\mu\phi^i\partial^\mu\phi^j = \frac{1}{4}C_{IJK}\Phi^I\partial_\mu\Phi^J\partial^\mu\Phi^K \Big|_{\nu=1}. \quad (\text{B.7})$$

One can define on \mathcal{SM} the sections with lower indices:

$$\Phi_I \equiv \frac{2}{3}G_{IJ}\Phi^J \Big|_{\nu=1} = \frac{1}{6}C_{IJK}\Phi^J\Phi^K \Big|_{\nu=1} = \frac{1}{3}\frac{\partial\mathcal{V}}{\partial\Phi^I} \Big|_{\nu=1}. \quad (\text{B.8})$$

With simple algebra one can show the following identities:

$$\begin{aligned} \Phi_I\Phi^I &= 1, & G_{IJ} &= \frac{9}{2}\Phi_I\Phi_J - \frac{1}{2}C_{IJK}\Phi^K, \\ \partial_i\Phi_I &= -\frac{2}{3}G_{IJ}\partial_i\Phi^J, & \Phi_I\partial_i\Phi^I &= \Phi^I\partial_i\Phi_I = 0. \end{aligned} \quad (\text{B.9})$$

In particular, $\partial_i\Phi^I$ for $i = 1, \dots, n_V$ are the tangent vectors to \mathcal{SM} in \mathbb{R}^{n_V+1} while Φ_I is a 1-form orthogonal to \mathcal{SM} . Another identity (and similar ones obtained by lowering one or both of the indices I, J with the metric G) is

$$\mathcal{G}^{ij}\partial_i\Phi^I\partial_j\Phi^J = G^{IJ} - \frac{2}{3}\Phi^I\Phi^J, \quad (\text{B.10})$$

where G^{IJ} is the inverse of G_{IJ} . To prove it, one observes that the tensor on the LHS is the projector on \mathcal{SM} , and then verifies that the expression on the RHS has the same property.

When the manifold \mathcal{SM} is a locally symmetric space, one can find a constant symmetric tensor C^{IJK} with upper indices such that [57]

$$C^{IPQ}C_{P(JK}C_{LM)Q} = \frac{4}{3}\delta_{(J}^I C_{KLM)}. \quad (\text{B.11})$$

With some algebra, it follows that

$$\Phi^I = \frac{3}{2}G^{IJ}\Phi_J = \frac{9}{2}C^{IJK}\Phi_J\Phi^K, \quad G^{IJ} = 2\Phi^I\Phi^J - 6C^{IJK}\Phi^K, \quad (\text{B.12})$$

as well as

$$C^{IJK} = \frac{1}{8}G^{IL}G^{JM}G^{KN}C_{LMN}. \quad (\text{B.13})$$

Quaternionic-Kähler geometry. The scalars q^u are real coordinates on the quaternionic-Kähler manifold \mathcal{QM} with metric $h_{uv}(q)$ [73]. For $n_H \geq 2$,²⁴ this is a $4n_H$ -dimensional Riemannian manifold with holonomy $SU(2) \times Sp(n_H)/\mathbb{Z}_2$. To express this fact, it is convenient to introduce local “vielbeins” f_u^{iA} with $i = 1, 2$ (not to be confused with the index i of very special geometry) in the fundamental of $SU(2)$ and $A = 1, \dots, 2n_H$ in the fundamental of $Sp(n_H)$, such that

$$h_{uv} = f_u^{iA} f_v^{jB} \epsilon_{ij} \Omega_{AB}, \quad (\text{B.14})$$

where ϵ_{ij} and Ω_{AB} are the invariant tensors of $SU(2)$ and $Sp(n_H)$, respectively. Regarding (iA) as a composite index, the inverse of the matrix f_u^{iA} is $f_{iA}^u = h^{uv} f_v^{jB} \epsilon_{ji} \Omega_{BA}$. One can then construct a locally-defined triplet of almost complex structures

$$\vec{J}_u^v \equiv (J^x)_u^v = -i f_u^{iA} f_{jA}^v (\sigma^x)_i^j \quad (\text{B.15})$$

where $x = 1, 2, 3$ is in the adjoint of $SU(2)$ and $\vec{\sigma}$ are the Pauli matrices. The derived triplet of almost symplectic forms is $\vec{J}_{uv} = \vec{J}_u^t h_{tv}$. They are antisymmetric, using that $\vec{\sigma}_i^j \epsilon_{jk}$ is symmetric.²⁵ The almost complex structures automatically satisfy the quaternion relation

$$(J^x)_u^s (J^y)_s^t = -\delta^{xy} \delta_u^t + \epsilon^{xyz} (J^z)_u^t. \quad (\text{B.17})$$

The Levi-Civita connection takes values in $\mathfrak{su}(2) \times \mathfrak{sp}(n_H)$. Calling ω_{uj}^i and ρ_{uB}^A the two projections, respectively, they are determined by the requirement that f_u^{iA} be covariantly constant with respect to the full connection:

$$0 = \nabla_v f_u^{iA} + f_u^{jA} \omega_{vj}^i + f_u^{iB} \rho_{vB}^A. \quad (\text{B.18})$$

We can alternate between the vector and bispinor notations of $SU(2)$ with²⁶

$$\vec{\omega}_u = -i \omega_{ui}^j \vec{\sigma}_j^i, \quad \omega_{ui}^j = \frac{i}{2} \vec{\omega}_u \cdot \vec{\sigma}_i^j. \quad (\text{B.20})$$

²⁴The case $n_H = 1$ is special because $SU(2)^2/\mathbb{Z}_2 \cong SO(4)$ and so the holonomy condition does not impose any constraint on (orientable) Riemannian manifolds. However, supersymmetry requires (B.25) which we can take as the definition of a quaternionic-Kähler manifold of dimension 4. A 4-dimensional space satisfying (B.25) is Einstein with self-dual Weyl curvature.

²⁵Using the fact that a 2×2 matrix can be expanded in the basis $\{\mathbb{1}, \vec{\sigma}\}$, we also find

$$2f_u^{iA} f_{jA}^v = \delta_u^v \delta_j^i + i \vec{J}_u^v \cdot \vec{\sigma}_j^i. \quad (\text{B.16})$$

²⁶The $SU(2)$ connection satisfies $\epsilon^{jm} \omega_{um}^n \epsilon_{ni} = \omega_{ui}^j$, in particular $\omega_{uj}^j = 0$, and a similar condition is satisfied by ρ . This follows from the properties of the Pauli matrices. In going between the vector and bispinor notation one can use the identities

$$\vec{\sigma}_n^m \cdot \vec{\sigma}_i^j = \delta_n^j \delta_i^m - \epsilon^{mj} \epsilon_{ni}, \quad \vec{\sigma}_i^j \times \vec{\sigma}_\ell^m = i(\vec{\sigma}_i^m \delta_\ell^j - \delta_i^m \vec{\sigma}_\ell^j). \quad (\text{B.19})$$

The two connections are extracted from (B.18) through: $\omega_{ui}{}^j \delta_A^B + \delta_i^j \rho_{uA}{}^B = -f_{iA}{}^w \nabla_u f_w{}^{jB}$. From (B.18) it immediately follows

$$\tilde{\nabla}_w \vec{J}_u{}^v \equiv \nabla_w \vec{J}_u{}^v + \vec{\omega}_w \times \vec{J}_u{}^v = 0 . \quad (\text{B.21})$$

In other words, \vec{J} is covariantly constant with respect to its natural $SU(2)$ connection $\vec{\omega}$. From the integrability condition of (B.18) one also obtains (in bispinor and vector notation):

$$R_{uv}{}^s{}_t = \mathcal{R}_{uvi}{}^j f_{jA}{}^s f_t{}^{iA} + \mathcal{R}_{uvA}{}^B f_{jB}{}^s f_t{}^{jA} = -\frac{1}{2} \vec{\mathcal{R}}_{uv} \cdot \vec{J}_t{}^s + \mathcal{R}_{uvA}{}^B f_{jB}{}^s f_t{}^{jA} , \quad (\text{B.22})$$

where $R_{uv}{}^s{}_t$ is the Riemann tensor of h_{uv} and we defined

$$\begin{aligned} \mathcal{R}_{uvi}{}^j &\equiv 2\partial_{[u} \omega_{v]i}{}^j - 2\omega_{[u|i}{}^k \omega_{v]k}{}^j & \text{or} & & \vec{\mathcal{R}}_{uv} &\equiv 2\partial_{[u} \vec{\omega}_{v]} + \vec{\omega}_u \times \vec{\omega}_v \\ \mathcal{R}_{uvA}{}^B &\equiv 2\partial_{[u} \rho_{v]A}{}^B - 2\rho_{[u|A}{}^C \rho_{v]C}{}^B . \end{aligned} \quad (\text{B.23})$$

In particular

$$R_{uv}{}^s{}_t \vec{J}_s{}^t = 2n_H \vec{\mathcal{R}}_{uv} , \quad (\text{B.24})$$

i.e., the $SU(2)$ field strength $\vec{\mathcal{R}}_{uv}$ is the $\mathfrak{su}(2)$ projection of the Riemann curvature.

One can prove [74] (see also [73, 75]) that $SU(2) \times Sp(n_H)$ holonomy manifolds with $n_H \geq 2$ are automatically Einstein. In fact, they satisfy a stronger property: the Riemann curvature is the sum of the Riemann tensor of $\mathbb{H}\mathbb{P}^{n_H}$ and of a Weyl part,

$$\begin{aligned} R_{uvst} = \frac{R}{8n_H(n_H + 2)} &\left(h_{s[u} h_{v]t} + \vec{J}_{uv} \cdot \vec{J}_{st} - \vec{J}_{s[u} \cdot \vec{J}_{v]t} \right) + \\ &+ (f_u{}^{iA} f_v{}^{jB} \epsilon_{ij}) (f_s{}^{kC} f_t{}^{\ell D} \epsilon_{k\ell}) \mathcal{W}_{ABCD} . \end{aligned} \quad (\text{B.25})$$

The tensor \mathcal{W}_{ABCD} is totally symmetric and controls the Weyl curvature, which is contained in $Sp(n_H)$: it gives rise to a traceless (and thus Ricci flat) contribution to the Riemann curvature. From that expression we obtain

$$R_{vt} = \frac{R}{4n_H} h_{vt} , \quad \vec{\mathcal{R}}_{uv} = \frac{R}{4n_H(n_H + 2)} \vec{J}_{uv} . \quad (\text{B.26})$$

The first equation shows that the manifold is Einstein. The second equation shows that the $SU(2)$ part of the curvature is completely fixed in terms of the triplet of complex structures. The tensor \mathcal{W}_{ABCD} expresses the freedom in the $Sp(n_H)$ part.

While quaternionic-Kähler manifolds can have any size, local supersymmetry requires²⁷

$$\lambda \equiv \frac{R}{4n_H(n_H + 2)} = -1 , \quad (\text{B.27})$$

fixing the scalar curvature [73]. Hence the manifold of hypermultiplet scalars is a non-trivial quaternionic-Kähler manifold with negative scalar curvature.

²⁷Had we chosen a canonical normalization for the action of the hypermultiplet scalars, the scalar curvature would be fixed in terms of the Planck mass to $\lambda = -m_{\text{Pl}}^{-2}$ [73]. This reproduces the fact that the manifold of hypermultiplet scalars is hyper-Kähler in rigid supersymmetry.

Isometries and gauging. We consider gaugings of Abelian isometries of the quaternionic-Kähler manifold \mathcal{QM} by the vectors A_μ^I . The isometries are generated by (possibly vanishing or linearly dependent) Killing vectors $k_I^u(q)$ that also satisfy a quaternionic version of the triholomorphic condition:

$$h_{w(u}\nabla_v)k_I^w = 0, \quad \vec{J}_u{}^w(\nabla_w k_I^v) - (\nabla_u k_I^w)\vec{J}_w{}^v = \lambda \vec{J}_u{}^v \times \vec{P}_I. \quad (\text{B.28})$$

The second equation expresses the fact that the derivative of each Killing vector commutes with the triplet of complex structures, up to a rotation parametrized by the $SU(2)$ sections \vec{P}_I . Notice that the LHS can be written, after lowering v , as $2\tilde{\nabla}_{[u}(\vec{J}_{v]s}k_I^s)$, therefore in the hyper-Kähler case that $\lambda = 0$ and the $SU(2)$ bundle is trivial, this reduces to the familiar condition that the three symplectic forms \vec{J}_{uv} be preserved by the isometries. By taking the cross product of the second equation in (B.28) with $\vec{J}_v{}^u$ we obtain

$$2n_H\lambda\vec{P}_I = \vec{J}_u{}^v\nabla_v k_I^u. \quad (\text{B.29})$$

This shows that on quaternionic Kähler manifolds, the sections \vec{P}_I are completely fixed in terms of the Killing vectors. With a little bit of work²⁸ we obtain

$$\tilde{\nabla}_u\vec{P}_I = \vec{J}_{uv}k_I^w. \quad (\text{B.30})$$

This shows that \vec{P}_I are a triplet of moment maps for the action of k_I^u . Taking a derivative and using that $2\tilde{\nabla}_{[u}\tilde{\nabla}_{v]}\vec{P}_I = \vec{\mathcal{R}}_{uv} \times \vec{P}_I$ we get back the second equation in (B.28), showing that the correction term on the RHS is unavoidable. The divergence of (B.30) gives

$$\tilde{\nabla}^u\tilde{\nabla}_u\vec{P}_I = -2n_H\lambda\vec{P}_I, \quad (\text{B.31})$$

showing that the moment maps are eigenfunctions of the Laplacian.

Finally, let us consider for the moment the general case that the Killing vectors might form a non-Abelian group:

$$[k_I, k_J]^u = 2k_{[I}^s\nabla_s k_{J]}^u = f_{IJ}{}^K k_K^u, \quad (\text{B.32})$$

where on the LHS is the Lie bracket and $f_{IJ}{}^K$ are the structure constants. Multiplying (B.28) by $\nabla_v k_J^u$ and using (B.29), and then exploiting the derivative ∇_w of (B.32), we obtain

$$k_I^u\vec{J}_{uv}k_J^v = f_{IJ}{}^K\vec{P}_K + \lambda\vec{P}_I \times \vec{P}_J. \quad (\text{B.33})$$

This is called the equivariance relation. In the Abelian case we just set f to zero. In the special case $n_H = 0$ that there are no hypermultiplets, all Killing vectors vanish and the only

²⁸We take the derivative $\tilde{\nabla}$ of (B.29), recalling that \vec{J} is covariantly constant. From the algebraic Bianchi identity we have $R_{uvst}\vec{J}^{us} = \frac{1}{2}R_{vt}{}^s{}_u\vec{J}_s{}^u = n_H\vec{\mathcal{R}}_{vt} = n_H\lambda\vec{J}_{vt}$. Then we use that the vectors are Killing, as well as the properties of quaternionic-Kähler manifolds.

remnant of the quaternionic-Kähler structure is the condition $\vec{P}_I \times \vec{P}_J = 0$. The solution, up to $SU(2)$ rotations, is $P_I^x = \delta^{x3} \zeta_I$ where ζ_I are the so-called Fayet-Iliopoulos (FI) parameters, which in this case are extra parameters one needs to specify.

We now have all the ingredients to write the covariant derivative

$$\mathcal{D}_\mu q^u = \partial_\mu q^u + g A_\mu^I k_I^u, \quad (\text{B.34})$$

as well as the scalar potential

$$\begin{aligned} V &= P_I^x P_J^x \left(\frac{1}{2} \mathcal{G}^{ij} \partial_i \Phi^I \partial_j \Phi^J - \frac{2}{3} \Phi^I \Phi^J \right) + \frac{1}{2} h_{uv} k_I^u k_J^v \Phi^I \Phi^J \\ &= P_I^x P_J^x \left(\frac{1}{2} G^{IJ} - \Phi^I \Phi^J \right) + \frac{1}{2} h_{uv} k_I^u k_J^v \Phi^I \Phi^J \end{aligned} \quad (\text{B.35})$$

that couples the scalars on \mathcal{SM} and \mathcal{QM} . To go to the second line we used (B.10).

The covariant derivative of the supersymmetry parameter ϵ_i^{SUSY} (subject to symplectic-Majorana condition, with $i = 1, 2$) is

$$D_\mu \epsilon_i^{\text{SUSY}} = \left(\nabla_\mu \delta_i^j - \frac{i}{2} \vec{\mathcal{V}}_\mu \cdot \vec{\sigma}_i^j \right) \epsilon_j^{\text{SUSY}} \quad (\text{B.36})$$

with connection

$$\begin{aligned} \vec{\mathcal{V}}_\mu &= \mathcal{D}_\mu q^u \vec{\omega}_u - g A_\mu^I \vec{r}_I \quad \text{and} \quad \vec{r}_I = k_I^u \vec{\omega}_u - \lambda \vec{P}_I, \\ &= \partial_\mu q^u \vec{\omega}_u + g \lambda A_\mu^I \vec{P}_I \end{aligned} \quad (\text{B.37})$$

where λ is the constant (B.27). Under gauge transformations²⁹

$$\delta q^u = g \alpha^I k_I^u, \quad \delta A_\mu^I = -\partial_\mu \alpha^I \quad (\text{B.38})$$

with parameters α^I , using (B.26), (B.30) and (B.33) one can show that $\vec{\mathcal{V}}_\mu$ transforms as an $SU(2)$ connection:

$$\delta \vec{\mathcal{V}}_\mu = \partial_\mu \vec{\Lambda} + \vec{\mathcal{V}}_\mu \times \vec{\Lambda} \quad \text{with} \quad \vec{\Lambda} = g \alpha^I \vec{r}_I. \quad (\text{B.39})$$

Therefore, $D_\mu \epsilon_i^{\text{SUSY}}$ is covariant if ϵ_i^{SUSY} transforms as

$$\delta \epsilon_i^{\text{SUSY}} = \frac{i}{2} \vec{\Lambda} \cdot \vec{\sigma}_i^j \epsilon_j^{\text{SUSY}}. \quad (\text{B.40})$$

²⁹The covariant derivative transforms as $\delta \mathcal{D}_\mu q^u = g \alpha^I \mathcal{D}_\mu k_I^u$.

B.1 Conifold truncation in the general framework

Here we embed the consistent truncation of type IIB supergravity on $T^{1,1}$ to a 5d $\mathcal{N} = 2$ gauged supergravity with a so-called ‘‘Betti multiplet’’, described in Section 7 of [34] (called the ‘‘second model’’ in that paper), in the general framework. The model has $n_V = 2$ and $n_H = 2$. We identify the fields

$$\phi^i = \begin{pmatrix} u+v \\ w \end{pmatrix}_{\text{CF}}, \quad \Phi^I = \begin{pmatrix} e^{-4(u+v)/3} \\ -e^{2(u+v)/3} \cosh 2w \\ -e^{2(u+v)/3} \sinh 2w \end{pmatrix}_{\text{CF}}, \quad A^I = \begin{pmatrix} A \\ a_1^J \\ a_1^\Phi \end{pmatrix}_{\text{CF}}, \quad q^u = \begin{pmatrix} b_1^\Omega \\ b_2^\Omega \\ c_1^\Omega \\ c_2^\Omega \\ a \\ \phi \\ C_0 \\ u \end{pmatrix}_{\text{CF}} \quad (\text{B.41})$$

where ‘‘CF’’ indicates the notation of [34]. The scalar fields b^Ω, c^Ω are complex and we used $z_1 = \text{Re}(z), z_2 = \text{Im}(z)$ to indicate their real and imaginary parts, while u, v, w, a, ϕ, C_0 are real. The hypermultiplet scalars C_0 and ϕ together form the type IIB axiodilaton $C_0 + ie^{-\phi}$. Then we identify the Chern-Simons couplings

$$C_{122} = -C_{133} = 2 \quad (\text{B.42})$$

and symmetric permutations thereof, while all other components vanish, and the very special geometry of $SO(1,1) \times SO(1,1)$:

$$\mathcal{G}_{ij} = \begin{pmatrix} 4/3 & 0 \\ 0 & 4 \end{pmatrix}, \quad G_{IJ} = e^{-\frac{4}{3}(u+v)} \begin{pmatrix} \frac{1}{2}e^{4(u+v)} & 0 & 0 \\ 0 & \cosh(4w) & -\sinh(4w) \\ 0 & -\sinh(4w) & \cosh(4w) \end{pmatrix}. \quad (\text{B.43})$$

The tensor C^{IJK} has non-vanishing components $C^{122} = -C^{133} = 1/2$ and permutations.

The quaternionic-Kähler manifold is $\frac{SO(4,2)}{SO(4) \times SO(2)}$. Its metric is

$$h_{uv} dq^u dq^v = e^{-4u-\phi} db^\Omega d\bar{b}^\Omega + e^{-4u+\phi} (dc^\Omega - C_0 db^\Omega) (d\bar{c}^\Omega - C_0 d\bar{b}^\Omega) + \frac{1}{2} e^{-8u} \left(2da + \text{Re}(b^\Omega d\bar{c}^\Omega - c^\Omega d\bar{b}^\Omega) \right)^2 + \frac{1}{2} d\phi^2 + \frac{1}{2} e^{2\phi} dC_0^2 + 8du^2. \quad (\text{B.44})$$

In this normalization $R = -32$ and thus $\lambda = -1$. The $SU(2)$ connection is

$$\omega^1 - i\omega^2 = e^{-2u-\phi/2} db^\Omega + i e^{-2u+\phi/2} (dc^\Omega - C_0 db^\Omega) \\ \omega^3 = \frac{1}{2} e^{-4u} \left(2da + \text{Re}(b^\Omega d\bar{c}^\Omega - c^\Omega d\bar{b}^\Omega) \right) - \frac{1}{2} e^\phi dC_0. \quad (\text{B.45})$$

Finally, we identify the Killing vectors

$$k_1 = 3 \left(-b_2^\Omega \frac{\partial}{\partial b_1^\Omega} + b_1^\Omega \frac{\partial}{\partial b_2^\Omega} - c_2^\Omega \frac{\partial}{\partial c_1^\Omega} + c_1^\Omega \frac{\partial}{\partial c_2^\Omega} \right) + 2 \frac{\partial}{\partial a}, \quad k_2 = 2 \frac{\partial}{\partial a}, \quad k_3 = 0 \quad (\text{B.46})$$

and the corresponding moment maps

$$P_1^x = \begin{pmatrix} 3e^{\phi/2-2u}(c_1^\Omega - C_0 b_1^\Omega + e^{-\phi} b_2^\Omega) \\ 3e^{\phi/2-2u}(C_0 b_2^\Omega - c_2^\Omega + e^{-\phi} b_1^\Omega) \\ 3 - e^{-4u}(2 + 3b_2^\Omega c_1^\Omega - 3b_1^\Omega c_2^\Omega) \end{pmatrix}, \quad P_2^x = \begin{pmatrix} 0 \\ 0 \\ -2e^{-4u} \end{pmatrix}, \quad P_3^x = 0. \quad (\text{B.47})$$

The $SU(2)$ connection and the moment maps were given in [36] and can be translated into the notation of [34] (up to a conventional minus sign in the gauge fields) using the identifications

$$\phi^i = \begin{pmatrix} -3u_3 \\ u_2 \end{pmatrix}_{\text{HLS}}, \quad A^I = \begin{pmatrix} A_1 \\ \frac{k_{11} - k_{12}}{2} \\ \frac{k_{11} + k_{12}}{2} \end{pmatrix}_{\text{HLS}}, \quad q^u = \left(2 \operatorname{Re} b_0^1, 2 \operatorname{Im} b_0^1, 2 \operatorname{Re} b_0^2, 2 \operatorname{Im} b_0^2, \frac{k}{2}, \phi, a, u_1 \right)_{\text{HLS}}^\top \quad (\text{B.48})$$

where ‘‘HLS’’ indicates the notation of [36].

The theory has a supersymmetric AdS_5 vacuum at $u = v = w = b^\Omega = c^\Omega = 0$ and any value of a, C_0, ϕ (in particular, the axiodilaton can take any value). The potential is $V|_{\text{AdS}} = -6$ leading to AdS radius $\ell_5 = g^{-1}$. The spectrum therein was computed in [34] (see its Table 2). We are particularly interested in the spectrum of vector fields and the Killing vectors they couple to:

$$\begin{aligned} A^R &\equiv A^1 - 2A^2, & A^3 &: & m^2 = 0, & A^W &\equiv A^1 + A^2 : & m^2 = 24g^2. \\ k_R &= \frac{1}{3}(k_1 - k_2), & k_3 & & & k_W &= \frac{1}{3}(2k_1 + k_2) \end{aligned} \quad (\text{B.49})$$

The vector A^W acquires a mass by Higgs mechanism, eating the Stückelberg scalar a . The mass eigenstates are

$$\mathbb{B}^I{}_J A_\mu^J \quad \text{where} \quad \mathbb{B} = \begin{pmatrix} 1 & -2 & 0 \\ 0 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix} \quad (\text{B.50})$$

is the matrix that diagonalizes them (see also Appendix E).

C 4d $\mathcal{N} = 2$ Abelian gauged supergravity

We summarize the salient features of 4d $\mathcal{N} = 2$ Abelian gauged supergravity with n_V vector multiplets and n_H hypermultiplets, following [60, 61, 71]. The graviton multiplet contains a

graviton, two gravitini and a vector; each vector multiplet contains a vector, two gaugini and a complex scalar; each hypermultiplet contains four real scalars and two hyperini (all fermions can be taken Majorana). We use indices

$$\Lambda, \Sigma = 0, \dots, n_V, \quad i, j = 1, \dots, n_V, \quad u, v = 1, \dots, 4n_H \quad (\text{C.1})$$

for the gauge fields A_μ^Λ , for the complex scalars z^i in vector multiplets, and for the real scalars q^u in hypermultiplets, respectively. The data that define the theory are:

1. A special Kähler manifold \mathcal{KM} of complex dimension n_V .
2. A quaternionic-Kähler manifold \mathcal{QM} of real dimension $4n_H$.
3. A set of $n_V + 1$ Killing vectors on \mathcal{QM} compatible with the quaternionic-Kähler structure (if $n_H = 0$, $n_V + 1$ FI parameters not all vanishing).

The Killing vectors could be linearly dependent or vanish.

It is always possible to find a duality frame in which all gaugings are purely electric. In such frames the bosonic Lagrangian is

$$8\pi G_{\text{N}}^{(4)} e^{-1} \mathcal{L}_{\text{4d}} = \frac{R_s}{2} - g_{i\bar{j}}(z, \bar{z}) \partial_\mu z^i \partial^\mu \bar{z}^{\bar{j}} - \frac{1}{2} h_{uv}(q) \mathcal{D}_\mu q^u \mathcal{D}^\mu q^v + \frac{1}{8} \text{Im} \mathcal{N}_{\Lambda\Sigma}(z, \bar{z}) F_{\mu\nu}^\Lambda F^{\Sigma\mu\nu} + \frac{e^{-1}}{16} \text{Re} \mathcal{N}_{\Lambda\Sigma}(z, \bar{z}) F_{\mu\nu}^\Lambda F_{\rho\sigma}^\Sigma \epsilon^{\mu\nu\rho\sigma} - g^2 V(z, \bar{z}, q). \quad (\text{C.2})$$

The notation is mostly as in Appendix B. Let us explain the other terms.

Special Kähler geometry. The scalars z^i are complex coordinates on the special Kähler manifold \mathcal{KM} [61]. This is a Kähler-Hodge manifold — *i.e.*, a Kähler manifold with Kähler potential $\mathcal{K}(z, \bar{z})$ and metric $g_{i\bar{j}}(z, \bar{z}) = \partial_i \partial_{\bar{j}} \mathcal{K}$ as well as a line bundle (*i.e.*, a holomorphic vector bundle of rank 1) \mathcal{L} such that its first Chern class coincides (up to a constant) with the Kähler class $\omega = i\partial\bar{\partial}\mathcal{K}$ of the manifold³⁰ — further endowed with a flat $Sp(n_V + 1, \mathbb{R})$ symplectic bundle. The manifold comes equipped with a covariantly-holomorphic section of the tensor product of the symplectic bundle with the $U(1)$ -bundle \mathcal{U} associated to \mathcal{L} ,

$$\mathcal{V} = \begin{pmatrix} L^\Lambda \\ M_\Lambda \end{pmatrix} \quad \text{such that} \quad \begin{aligned} D_i \mathcal{V} &\equiv \partial_i \mathcal{V} + \frac{1}{2} (\partial_i \mathcal{K}) \mathcal{V} \\ D_{\bar{i}} \mathcal{V} &\equiv \partial_{\bar{i}} \mathcal{V} - \frac{1}{2} (\partial_{\bar{i}} \mathcal{K}) \mathcal{V} = 0, \end{aligned} \quad (\text{C.3})$$

obeying the constraints

$$\langle \mathcal{V}, \bar{\mathcal{V}} \rangle \equiv M_\Lambda \bar{L}^\Lambda - L^\Lambda \bar{M}_\Lambda = -i \quad (\text{C.4})$$

³⁰Because fermions are sections of the square root of \mathcal{L} , the Kähler class of \mathcal{KM} equal to the first Chern class of \mathcal{L} is required to be an even integer cohomology class.

and

$$\langle \mathcal{V}, D_i \mathcal{V} \rangle = 0, \quad (\text{C.5})$$

where we introduced the Sp -invariant antisymmetric form $i\langle \cdot, \cdot \rangle$. Equivalently, there is a holomorphic section of the tensor product of the symplectic bundle with \mathcal{L} ,³¹

$$v(z) = e^{-\mathcal{K}/2} \mathcal{V} \equiv \begin{pmatrix} X^\Lambda \\ F_\Lambda \end{pmatrix} \quad \text{such that} \quad \begin{aligned} D_i v &\equiv \partial_i v + (\partial_i \mathcal{K}) v \\ D_{\bar{i}} v &\equiv \partial_{\bar{i}} v = 0, \end{aligned} \quad (\text{C.6})$$

in terms of which the constraint (C.4) reads

$$\mathcal{K} = -\log(i \langle v, \bar{v} \rangle) = -\log \left[2 \operatorname{Im} (X^\Lambda \bar{F}_\Lambda) \right], \quad (\text{C.7})$$

while the constraint (C.5) becomes $\langle v, D_i v \rangle = \langle v, \partial_i v \rangle = 0$. From (C.3)–(C.5) it is easy to prove the following properties (or equivalent ones written in terms of v):

$$\begin{aligned} \langle D_i \mathcal{V}, \bar{\mathcal{V}} \rangle &= 0, & D_{\bar{j}} D_i \mathcal{V} &= g_{i\bar{j}} \mathcal{V}, & \langle D_i \mathcal{V}, D_{\bar{j}} \bar{\mathcal{V}} \rangle &= i g_{i\bar{j}} \\ \langle D_i \mathcal{V}, D_{\bar{j}} \mathcal{V} \rangle &= 0, & D_{[i} D_{\bar{j}]} \mathcal{V} &= 0 \end{aligned} \quad (\text{C.8})$$

from which the Kähler metric is extracted in a symplectic-invariant way.

The rescaling of X^Λ, F_Λ under Kähler transformations suggests to use X^Λ as homogeneous coordinates on \mathcal{KM} . It is always possible to find symplectic frames³² in which the Jacobian matrix $e^\lambda_i(z) = \partial_i (X^\lambda / X^0)$ (with $\lambda = 1, \dots, n_V$) is invertible. Notice that

$$\det(e^\lambda_i) = (X^0)^{n_V+1} \det(X^\Lambda, \partial_i X^\Lambda) = (X^0)^{n_V+1} \det(X^\Lambda, D_i X^\Lambda) \quad (\text{C.9})$$

where the two square matrices on the right have size $n_V + 1$, therefore the matrix $(X^\Lambda, \partial_i X^\Lambda)$ is invertible as well. Invertibility of the Jacobian ensures that we can use X^Λ as homogeneous coordinates, and regard $F_\Lambda(X)$ as homogeneous functions of degree 1, namely $X^\Sigma \partial_\Sigma F_\Lambda = F_\Lambda$. From (C.5) and (C.8), written as $\langle v, \partial_i v \rangle = \langle \partial_i v, \partial_{\bar{j}} v \rangle = 0$, one obtains the equations

$$(X^\Lambda, \partial_i X^\Lambda) \partial_{[\Lambda} F_{\Sigma]} (X^\Sigma, \partial_j X^\Sigma) = 0. \quad (\text{C.10})$$

Invertibility of the matrix implies $\partial_{[\Lambda} F_{\Sigma]} = 0$. Hence, in these frames, the sections F_Λ are the derivatives of a holomorphic homogeneous function $F(X)$ of degree 2, called the *prepotential*, namely $F_\Lambda = \partial_\Lambda F$. In such frames, the Kähler potential and thus the geometry are completely specified by the prepotential. The coordinates $t^i \equiv X^i / X^0$ with $i = 1, \dots, n_V$ are called *special coordinates*.

³¹In particular, $A = \partial \mathcal{K}$ is the Chern connection on \mathcal{L} . Moreover, $D_i \mathcal{V} = e^{\mathcal{K}/2} D_i v$ and $D_{\bar{i}} \mathcal{V} = e^{\mathcal{K}/2} D_{\bar{i}} v$.

³²See [76] for examples of frames in which, instead, a prepotential does not exist.

The couplings of vector fields to the scalars z^i are determined by the $(n_V + 1) \times (n_V + 1)$ period matrix \mathcal{N} , which is uniquely defined by the relations

$$M_\Lambda = \mathcal{N}_{\Lambda\Sigma} L^\Sigma, \quad D_{\bar{i}} \bar{M}_\Lambda = \mathcal{N}_{\Lambda\Sigma} D_{\bar{i}} \bar{L}^\Sigma. \quad (\text{C.11})$$

Explicitly, one needs to invert the matrix relation $(F_\Lambda, D_{\bar{i}} \bar{F}_\Lambda) = \mathcal{N}_{\Lambda\Sigma} (X^\Sigma, D_{\bar{i}} \bar{X}^\Sigma)$. The requirement that $g_{i\bar{j}}$ be positive definite guarantees that the rightmost matrix is invertible [61]. Indeed, introducing the square matrix $\mathcal{L}^\Lambda_I = (L^\Lambda, D_{\bar{i}} \bar{L}^\Lambda)$ of size $n_V + 1$, one can rewrite the scalar products in (C.4), (C.5) and (C.8) as

$$\mathcal{L}^\top (\mathcal{N} - \mathcal{N}^\top) \mathcal{L} = 0, \quad \mathcal{L}^\dagger (\mathcal{N} - \mathcal{N}^\dagger) \mathcal{L} = -i \text{diag}(1, g_{i\bar{j}}). \quad (\text{C.12})$$

The first equation shows that $\mathcal{N}_{\Lambda\Sigma}$ is a symmetric matrix, given the invertibility of \mathcal{L} . The second equation then, assuming that $g_{i\bar{j}}$ is positive definite, proves that \mathcal{L} is invertible and that $\mathbb{I}m \mathcal{N}_{\Lambda\Sigma}$ is negative definite. It also gives an expression for $\mathbb{I}m \mathcal{N}_{\Lambda\Sigma}$ that, after taking the inverse, reads

$$D_i L^\Lambda D_{\bar{j}} \bar{L}^\Sigma g^{i\bar{j}} + \bar{L}^\Lambda L^\Sigma = -\frac{1}{2} \left((\mathbb{I}m \mathcal{N})^{-1} \right)^{\Lambda\Sigma}. \quad (\text{C.13})$$

This relation, or the equivalent one in terms of the holomorphic section, will be used to rewrite the scalar potential below. When a prepotential exists, \mathcal{N} is obtained from

$$\mathcal{N}_{\Lambda\Sigma} = \bar{F}_{\Lambda\Sigma} + 2i \frac{(\mathbb{I}m F_{\Lambda\Gamma}) X^\Gamma (\mathbb{I}m F_{\Sigma\Delta}) X^\Delta}{X^\Omega (\mathbb{I}m F_{\Omega\Psi}) X^\Psi}, \quad (\text{C.14})$$

where $F_{\Lambda\Sigma} = \partial_\Lambda \partial_\Sigma F$. In this expression \mathcal{N} is manifestly symmetric.

Finally, one can define the tensor

$$\tilde{C}_{ijk} = \langle D_i D_j \mathcal{V}, D_k \mathcal{V} \rangle = \langle \mathcal{V}, D_k D_i D_j \mathcal{V} \rangle. \quad (\text{C.15})$$

Using (C.3)–(C.8) and the fact that the metric is Kähler, one easily proves that \tilde{C}_{ijk} is totally symmetric and covariantly holomorphic, $D_{\bar{\ell}} \tilde{C}_{ijk} = 0$ where \tilde{C} has twice the charge of \mathcal{V} . One can prove that $(\mathcal{V}, D_i \mathcal{V}, \bar{\mathcal{V}}, D_{\bar{i}} \bar{\mathcal{V}})$ pointwise form a basis for the symplectic bundle [61], hence

$$D_i D_j \mathcal{V} = i \tilde{C}_{ijk} g^{k\bar{k}} D_{\bar{k}} \bar{\mathcal{V}} \quad (\text{C.16})$$

follows by taking the product of the LHS with the basis. Among other things, \tilde{C} controls the curvature tensor: $R_{\bar{i}j\bar{k}\ell} = g_{\bar{j}\bar{i}} g_{\ell\bar{k}} + g_{j\bar{k}} g_{\ell\bar{i}} - \tilde{C}_{j\ell m} \tilde{C}_{\bar{i}\bar{k}\bar{n}} g^{m\bar{n}}$. In special coordinates the tensor \tilde{C} takes the particularly simple form

$$\tilde{C}_{ijk} = e^{\mathcal{K}} \partial_i \partial_j \partial_k \mathcal{F}(t) \quad \text{with} \quad \mathcal{F}(t) = (X^0)^{-2} F(X) \quad (\text{C.17})$$

and $t^i = X^i / X^0$.

Hypermultiplets and gauging. The part of the action involving the hypermultiplets has the same features as in the 5d case, summarized in Appendix B: the hypermultiplet scalars q^u (with $u = 1, \dots, 4n_H$) are coordinates on a quaternionic-Kähler manifold \mathcal{QM} with metric $h_{uv}(q)$. As before, we consider gauging of Abelian isometries of \mathcal{QM} , generated by $n_V + 1$ (possibly vanishing or linearly dependent) Killing vectors $k_\Lambda^u(q)$ that must be compatible with the quaternionic-Kähler structure, with associated triplets of moment maps $\vec{P}_\Lambda(q)$. In full generality one could consider both electric and magnetic gaugings, described by Killing vectors k_Λ^u and $k^{u\Lambda}$, respectively, and transforming as a vector under $Sp(n_V + 1, \mathbb{R})$ duality transformations. It is always possible to find a duality frame in which all gaugings are purely electric, and we will work in such a frame. Notice that there is no guarantee that in this frame a prepotential exists.

The scalar potential is

$$\begin{aligned} V &= 2 P_\Lambda^x P_\Sigma^x e^\mathcal{K} \left(g^{ij} D_i X^\Lambda D_{\bar{j}} \bar{X}^\Sigma - 3 X^\Lambda \bar{X}^\Sigma \right) + 4 e^\mathcal{K} h_{uv} k_\Lambda^u k_\Sigma^v X^\Lambda \bar{X}^\Sigma \\ &= -P_\Lambda^x P_\Sigma^x \left((\text{Im } \mathcal{N})^{-1\Lambda\Sigma} + 8 e^\mathcal{K} X^\Lambda \bar{X}^\Sigma \right) + 4 e^\mathcal{K} h_{uv} k_\Lambda^u k_\Sigma^v X^\Lambda \bar{X}^\Sigma . \end{aligned} \quad (\text{C.18})$$

To go to the second line we used (C.13).

The covariant derivative of the supersymmetry parameter ϵ_i^{SUSY} (subject to symplectic-Majorana condition, with $i = 1, 2$) is

$$D_\mu \epsilon_i^{\text{SUSY}} = \left(\nabla_\mu \delta_i^j - \frac{i}{2} \mathcal{A}_\mu \delta_i^j - \frac{i}{2} \vec{\mathcal{V}}_\mu \cdot \vec{\sigma}_i^j \right) \epsilon_j^{\text{SUSY}} \quad (\text{C.19})$$

with connections

$$\begin{aligned} \vec{\mathcal{V}}_\mu &= \partial_\mu q^u \vec{\omega}_u + g\lambda A_\mu^I \vec{P}_I \\ \mathcal{A}_\mu &= \frac{i}{2} \lambda \left[(\partial_\alpha \mathcal{K}) \partial_\mu z^\alpha - (\partial_{\bar{\alpha}} \mathcal{K}) \partial_\mu \bar{z}^{\bar{\alpha}} \right] . \end{aligned} \quad (\text{C.20})$$

Here $\vec{\mathcal{V}}_\mu$ is the $SU(2)$ connection that descends from the quaternionic-Kähler manifold \mathcal{QM} , as in the 5d case (B.37). Instead \mathcal{A}_μ descends from the connection on the $U(1)$ -bundle \mathcal{U} on the special Kähler manifold \mathcal{KM} .

D Reduction with background gauge fields

Following [65] we will now reduce, piece by piece, the bosonic Lagrangian (B.2) of 5d $\mathcal{N} = 2$ gauged supergravity down to 4d. We start in 5d with n_V vector multiplets and n_H hypermultiplets. We use indices

$$I, J = 1, \dots, n_V + 1 , \quad \Lambda, \Sigma = 0, \dots, n_V + 1 , \quad u, v = 1, \dots, 4n_H . \quad (\text{D.1})$$

We indicate the 5d vector fields as \widehat{A}_M^I (where $M, N = 0, \dots, 4$ are spacetime indices) and parametrize the vector multiplet scalars in terms of sections Φ^I subject to the cubic constraint $\mathcal{V}(\Phi) = 1$ in (B.3). The hypermultiplet scalars are q^u . We employ the rather standard Kaluza-Klein reduction ansatz (5.2) and (5.3):

$$\widehat{g}_{MN} = \begin{pmatrix} e^{2\widetilde{\phi}} g_{\mu\nu} + e^{-4\widetilde{\phi}} A_\mu^0 A_\nu^0 & -e^{-4\widetilde{\phi}} A_\mu^0 \\ -e^{-4\widetilde{\phi}} A_\nu^0 & e^{-4\widetilde{\phi}} \end{pmatrix}, \quad \widehat{g}^{MN} = \begin{pmatrix} e^{-2\widetilde{\phi}} g^{\mu\nu} & e^{-2\widetilde{\phi}} A^{0\mu} \\ e^{-2\widetilde{\phi}} A^{0\nu} & e^{4\widetilde{\phi}} + e^{-2\widetilde{\phi}} A_\rho^0 A^{0\rho} \end{pmatrix},$$

$$e_{(5)} = e^{2\widetilde{\phi}} e_{(4)}, \quad \Phi^I = -e^{2\widetilde{\phi}} z_2^I, \quad \widehat{A}_M^I = (A_\mu^I - z_1^I A_\mu^0, z_1^I + \xi^I). \quad (\text{D.2})$$

The last coordinate, that we call y and whose range Δy we leave generic for now, is compactified on a circle of length $e^{-2\widetilde{\phi}} \Delta y$, and no field depends on it. We indicated as \widehat{g}_{MN} and $e_{(5)}$ the 5d metric and the square root of its determinant, and as $g_{\mu\nu}$ and $e_{(4)}$ (with $\mu, \nu = 0, \dots, 3$ spacetime indices) their 4d counterparts. In 4d we end up with $n_V + 1$ vector multiplets, and we indicate as A_μ^Λ the vector fields. The physical scalars in 4d vector multiplets are the complex fields z^i . With a useful abuse of notation, we utilize the very same index I for 5d vector fields and 4d physical scalars, z^I , because in 4d we have one more vector field than in 5d. We also use the notation

$$z_1^I \equiv \text{Re } z^I, \quad z_2^I \equiv \text{Im } z^I. \quad (\text{D.3})$$

Notice that the real scalar $\widetilde{\phi}$ can be eliminated with the 5d constraint,

$$e^{-6\widetilde{\phi}} = -\mathcal{V}(z_2), \quad (\text{D.4})$$

then the scalars z^I can be treated as independent. The real parameters ξ^I represent background gauge fields along the circle, therefore, up to a gauge transformation, this ansatz is equivalent to a Scherk-Schwarz reduction.

The reduction of the Einstein term gives

$$8\pi G_N^{(4)} \mathcal{L}_1 = e_{(5)} \frac{\widehat{R}_s}{2} = e_{(4)} \left[\frac{R_s}{2} - 3 \partial_\mu \widetilde{\phi} \partial^\mu \widetilde{\phi} - \frac{e^{-6\widetilde{\phi}}}{8} F_{\mu\nu}^0 F^{0\mu\nu} \right] + \text{total derivatives}. \quad (\text{D.5})$$

Here \widehat{R}_s and R_s are the 5d and 4d Ricci scalars, respectively. The 4d and 5d Newton constants are related by

$$\frac{1}{G_N^{(4)}} = \frac{\Delta y}{G_N^{(5)}}. \quad (\text{D.6})$$

In the following, for clarity, we will omit the factor $8\pi G_N^{(4)}$. The reduction of the kinetic term of vector multiplet scalars gives

$$\mathcal{L}_2 = -e_{(5)} \frac{1}{2} G_{IJ} \widehat{g}^{MN} \partial_M \Phi^I \partial_N \Phi^J = e_{(4)} \left[-\frac{e^{4\widetilde{\phi}}}{2} G_{IJ} \partial_\mu z_2^I \partial^\mu z_2^J + 3 \partial_\mu \widetilde{\phi} \partial^\mu \widetilde{\phi} \right]. \quad (\text{D.7})$$

The last term exactly cancels the second term in \mathcal{L}_1 , therefore

$$\mathcal{L}_1 + \mathcal{L}_2 = e_{(4)} \left[\frac{R_s}{2} - \frac{e^{4\tilde{\phi}}}{2} G_{IJ} \partial_\mu z_2^I \partial^\mu z_2^J - \frac{e^{-6\tilde{\phi}}}{8} F_{\mu\nu}^0 F^{0\mu\nu} \right]. \quad (\text{D.8})$$

The reduction of the kinetic term of hypermultiplet scalars gives

$$\begin{aligned} \mathcal{L}_3 &= -e_{(5)} \frac{1}{2} h_{uv} \hat{g}^{MN} \hat{\mathcal{D}}_M q^u \hat{\mathcal{D}}_N q^v \\ &= e_{(4)} \left[-\frac{1}{2} h_{uv} \mathcal{D}_\mu q^u \mathcal{D}^\mu q^v - \frac{g^2 e^{6\tilde{\phi}}}{2} (k_0^u + z_1^I k_I^u) h_{uv} (k_0^v + z_1^J k_J^v) \right]. \end{aligned} \quad (\text{D.9})$$

Here $\hat{\mathcal{D}}_M q^u = \partial_M q^u + g \hat{A}_M^I k_I^u$ is the 5d covariant derivative in (B.34), while

$$\mathcal{D}_\mu q^u = \partial_\mu q^u + g A_\mu^I k_I^u + g A_\mu^0 \xi^I k_I^u = \partial_\mu q^u + g A_\mu^\Lambda k_\Lambda^u \quad (\text{D.10})$$

is the 4d covariant derivative, and we defined the new Killing vector

$$k_0^u \equiv \xi^I k_I^u. \quad (\text{D.11})$$

The reduction of the gauge kinetic term gives

$$\begin{aligned} \mathcal{L}_4 &= -e_{(5)} \frac{1}{4} G_{IJ} \hat{F}_{MN}^I \hat{F}^{JMN} \\ &= e_{(4)} \left[-\frac{e^{-2\tilde{\phi}}}{4} G_{IJ} (F_\mu^I - z_1^I F_{\mu\nu}^0) (F^{J\mu\nu} - z_1^J F^{0\mu\nu}) - \frac{e^{4\tilde{\phi}}}{2} G_{IJ} \partial_\mu z_1^I \partial^\mu z_1^J \right], \end{aligned} \quad (\text{D.12})$$

where \hat{F}_{MN} and $F_{\mu\nu}$ are the 5d and 4d field strengths, respectively. We used $\hat{F}_{\mu 4}^I = \partial_\mu z_1^I$ and $\hat{F}_{\mu\nu}^I = F_{\mu\nu}^I - z_1^I F_{\mu\nu}^0 + 2A_{[\mu}^0 \partial_{\nu]} z_1^I$.

In order to reduce the Chern-Simons term, we extend the geometry (5.2) to a 6d bulk whose boundary is the original 5d space. A convenient way to do that is to complete the circle parametrized by y into a unit disk with radius $\rho \in [0, 1]$. We extend the 5d connections \hat{A}^I in (5.3) to 6d connections \tilde{A}^I as follows:

$$\tilde{A}^I = A^I + \xi^I A^0 + \rho^2 (z_1^I + \xi^I) (dy - A^0). \quad (\text{D.13})$$

We then write the Chern-Simons action term as

$$\int_{5\text{d}} \mathcal{L}_5 = \int_{5\text{d}} \frac{1}{12} C_{IJK} \hat{F}^I \wedge \hat{F}^J \wedge \hat{A}^K = \int_{6\text{d}} \frac{1}{12} C_{IJK} \tilde{F}^I \wedge \tilde{F}^J \wedge \tilde{F}^K. \quad (\text{D.14})$$

Substituting $\tilde{F}^I = d\tilde{A}^I$ and performing the integrals over $d\rho^2 \wedge (dy - A^0)$, we extract the 4d reduced Lagrangian

$$\mathcal{L}_5 = \frac{1}{16} C_{IJK} \epsilon^{\mu\nu\rho\sigma} \left[(z_1^I + \xi^I) F_{\mu\nu}^J F_{\rho\sigma}^K - (z_1^I z_1^J - \xi^I \xi^J) F_{\mu\nu}^K F_{\rho\sigma}^0 + \frac{z_1^I z_1^J z_1^K + \xi^I \xi^J \xi^K}{3} F_{\mu\nu}^0 F_{\rho\sigma}^0 \right]. \quad (\text{D.15})$$

Notice that the contributions containing the ξ^I 's are standard theta terms.

Finally, the reduction of the scalar potential gives

$$\mathcal{L}_6 = -e_{(5)}g^2V = -e_{(4)}g^2 \left[P_I^x P_J^x \left(\frac{e^{2\tilde{\phi}}}{2} \mathcal{G}^{ij} \partial_i \Phi^I \partial_j \Phi^J - \frac{2e^{\delta\tilde{\phi}}}{3} z_2^I z_2^J \right) + \frac{e^{\delta\tilde{\phi}}}{2} h_{uv} k_I^u k_J^v z_2^I z_2^J \right]. \quad (\text{D.16})$$

We proceed now with recasting the various pieces of the action in the general form (C.2) of 4d $\mathcal{N} = 2$ gauged supergravity with $n_V + 1$ vector multiplets and n_H hypermultiplets. The Einstein term receives its contribution from \mathcal{L}_1 :

$$\mathcal{L}'_1 = e_{(4)} \frac{R_s}{2}. \quad (\text{D.17})$$

The kinetic term of vector multiplet scalars gets contributions from \mathcal{L}_2 and \mathcal{L}_4 :

$$\mathcal{L}'_2 = -e_{(4)} \frac{e^{4\tilde{\phi}}}{2} G_{IJ} \left(\partial_\mu z_2^I \partial^\mu z_2^J + \partial_\mu z_1^I \partial^\mu z_1^J \right) = -e_{(4)} g_{I\bar{J}} \partial z^I \partial^\mu \bar{z}^{\bar{J}}, \quad (\text{D.18})$$

where we defined the Hermitian metric

$$g_{I\bar{J}} = \frac{e^{4\tilde{\phi}}}{2} G_{I\bar{J}}. \quad (\text{D.19})$$

The kinetic term of hypermultiplet scalars gets its contribution from \mathcal{L}_3 ,

$$\mathcal{L}'_3 = -e_{(4)} \frac{1}{2} h_{uv} \mathcal{D}_\mu q^u \mathcal{D}^\mu q^v, \quad (\text{D.20})$$

with the covariant derivative \mathcal{D}_μ defined in (D.10)-(D.11). The gauge kinetic term receives contributions from \mathcal{L}_1 and \mathcal{L}_4 :

$$\mathcal{L}'_4 = -e_{(4)} \frac{e^{-6\tilde{\phi}}}{8} \left[F_{\mu\nu}^0 F^{0\mu\nu} + 4g_{IJ} (F_{\mu\nu}^I - z_1^I F_{\mu\nu}^0) (F^{J\mu\nu} - z_1^J F^{0\mu\nu}) \right] = e_{(4)} \frac{1}{8} \mathbb{I}m \mathcal{N}_{\Lambda\Sigma} F_{\mu\nu}^\Lambda F^{\Sigma\mu\nu} \quad (\text{D.21})$$

where we defined the field-dependent matrix of gauge couplings

$$\mathbb{I}m \mathcal{N}_{\Lambda\Sigma} = -e^{-6\tilde{\phi}} \begin{pmatrix} 1 + 4g_{KL} z_1^K z_1^L & -4g_{KJ} z_1^K \\ -4g_{IK} z_1^K & 4g_{IJ} \end{pmatrix} \quad (\text{D.22})$$

in which the indices Λ, Σ run over 0 and then the values of I, J . On the other hand, the field-dependent theta terms are contained in \mathcal{L}_5 :

$$\mathcal{L}'_5 = \mathcal{L}_5 = \frac{1}{16} \mathbb{R}e \mathcal{N}_{\Lambda\Sigma} \epsilon^{\mu\nu\rho\sigma} F_{\mu\nu}^\Lambda F_{\rho\sigma}^\Sigma \quad (\text{D.23})$$

where

$$\mathbb{R}e \mathcal{N}_{\Lambda\Sigma} = \begin{pmatrix} \frac{1}{3} C_{KLM} (z_1^K z_1^L z_1^M + \xi^K \xi^L \xi^M) & -\frac{1}{2} C_{JKL} (z_1^K z_1^L - \xi^K \xi^L) \\ -\frac{1}{2} C_{IKL} (z_1^K z_1^L - \xi^K \xi^L) & C_{IJK} (z_1^K + \xi^K) \end{pmatrix}. \quad (\text{D.24})$$

It turns out that $g_{I\bar{J}}$ and $\mathcal{N}_{\Lambda\Sigma}$ descend from the following prepotential:

$$\begin{aligned} F(X) &= \frac{1}{6} C_{IJK} \frac{\check{X}^I \check{X}^J \check{X}^K}{X^0} && \text{with } \check{X}^I \equiv X^I + \xi^I X^0 \\ &= \frac{1}{6} C_{IJK} \frac{X^I X^J X^K}{X^0} + \frac{1}{2} C_{IJK} \left(\xi^I X^J X^K + \xi^I \xi^J X^K X^0 + \frac{1}{3} \xi^I \xi^J \xi^K (X^0)^2 \right). \end{aligned} \quad (\text{D.25})$$

The terms in parenthesis involving the ξ^I 's only affect standard theta terms, which are topological and thus do not enter in the equations of motion. Indeed, using special coordinates $z^I = X^I/X^0$ and in the Kähler frame $|X^0|^2 = 1$, one derives the Kähler potential³³

$$\mathcal{K} = -\log \left(\frac{1}{6i} C_{IJK} (z^I - \bar{z}^I)(z^J - \bar{z}^J)(z^K - \bar{z}^K) \right) = -\log(8 e^{-6\tilde{\phi}}) \quad (\text{D.26})$$

from which the Kähler metric (D.19) with (B.6) follows. On the other hand

$$F_{\Lambda\Sigma} = \begin{pmatrix} \frac{1}{3} C_{KLM} (z^K z^L z^M + \xi^K \xi^L \xi^M) & -\frac{1}{2} C_{JKM} (z^K z^M - \xi^K \xi^L) \\ -\frac{1}{2} C_{IKM} (z^K z^M - \xi^K \xi^L) & C_{IJK} (z^K + \xi^K) \end{pmatrix} \quad (\text{D.27})$$

from which the matrix \mathcal{N} in (D.22) and (D.24) follows. It might be useful

$$\begin{aligned} (X^0)^{-2} X^\Lambda (\mathbb{I}m F_{\Lambda\Sigma}) X^\Sigma &= 4 C_{IJK} \left(\frac{1}{3} \mathbb{I}m(z^I z^J z^K) - \frac{1}{2} \mathbb{I}m(z^I z^J) \mathbb{R}e(z^K) \right) \\ &= -\frac{4}{3} C_{IJK} z_2^I z_2^J z_2^K = e^{-\mathcal{K}} = 8 e^{-6\tilde{\phi}}, \end{aligned} \quad (\text{D.28})$$

as well as $(\mathbb{I}m F_{I\Sigma}) X^\Sigma / X^0 = i C_{IKM} z_2^K z_2^M$.

Finally, the scalar potential gets contributions from \mathcal{L}_3 and \mathcal{L}_6 :

$$\begin{aligned} \mathcal{L}'_6 &= -e_{(4)} g^2 \left[P_I^x P_J^x \left(\frac{e^{2\tilde{\phi}}}{2} \mathcal{G}^{ij} \partial_i \Phi^I \partial_j \Phi^J - \frac{2e^{6\tilde{\phi}}}{3} z_2^I z_2^J \right) + \right. \\ &\quad \left. + \frac{e^{6\tilde{\phi}}}{2} h_{uv} \left(k_I^u k_J^v z_2^I z_2^J + (k_0^u + z_1^I k_I^u)(k_0^v + z_1^J k_J^v) \right) \right] \\ &= -e_{(4)} g^2 \left[-P_\Lambda^x P_\Sigma^x \left((\mathbb{I}m \mathcal{N})^{-1\Lambda\Sigma} + 8 e^{\mathcal{K}} X^\Lambda \bar{X}^\Sigma \right) + 4 e^{\mathcal{K}} h_{uv} k_\Lambda^u k_\Sigma^v X^\Lambda \bar{X}^\Sigma \right]. \end{aligned} \quad (\text{D.29})$$

To manipulate the first line we used (B.10) as well as

$$\left((\mathbb{I}m \mathcal{N})^{-1} \right)^{\Lambda\Sigma} + 8 e^{\mathcal{K}} X^{(\Lambda} \bar{X}^{\Sigma)} = -e^{6\tilde{\phi}} \begin{pmatrix} 0 & 0 \\ 0 & \frac{1}{4} g^{IJ} - z_2^I z_2^J \end{pmatrix}, \quad (\text{D.30})$$

which immediately follows from (D.22). Notice in particular that \vec{P}_0 drops out of the potential and cannot be extracted from it, but it is still determined as $\vec{P}_0 = \xi^I \vec{P}_I$ from (D.11). The action \mathcal{L}'_6 exactly reproduces the potential in (C.18).

³³The completely covariant expression for the Kähler potential is $e^{-\mathcal{K}} = 8 |X^0|^2 e^{-6\tilde{\phi}}$.

Summarizing, the compactification gives the following map from 5d to 4d data:

5d		4d	
n_V vector multiplets		$n_V + 1$ vector multiplets	
\mathcal{SM} with C_{IJK}	$\xrightarrow[\text{background fields}]{\text{reduction with } \xi^I}$	\mathcal{KM} with $F = \frac{1}{6}C_{IJK}\frac{\check{X}^I\check{X}^J\check{X}^K}{X^0}$	(D.31)
\mathcal{QM} with $h_{uv}(q)$		\mathcal{QM} with $h_{uv}(q)$	
gauging of k_I^u		gauging of $k_\Lambda^u = (\xi^J k_J^u, k_I^u)$	

where $\check{X}^I = X^I + \xi^I X^0$.

D.1 Reduction of the conifold truncation

The reduction of the 5d conifold truncation described in Appendix B.1 gives a 4d supergravity with the following data. The prepotential is

$$F = \frac{\check{X}^1((\check{X}^2)^2 - (\check{X}^3)^2)}{X^0}. \quad (\text{D.32})$$

It induces the vector multiplet scalar metric

$$g_{I\bar{J}} = \frac{1}{2} \begin{pmatrix} \frac{1}{2(z_2^1)^2} & 0 & 0 \\ \frac{(z_2^2)^2 + (z_2^3)^2}{((z_2^2)^2 - (z_2^3)^2)^2} & -\frac{2z_2^2 z_2^3}{((z_2^2)^2 - (z_2^3)^2)^2} \\ \text{Symmetrized} & \frac{(z_2^2)^2 + (z_2^3)^2}{((z_2^2)^2 - (z_2^3)^2)^2} \end{pmatrix} \quad (\text{D.33})$$

that depends on z_2^I , the theta terms (D.24) that depend on z_1^I and ξ^I , while the gauge coupling function $\text{Im}\mathcal{N}_{\Lambda\Sigma}$ takes a lengthier expression that depends on z_1^I and z_2^I and can be easily derived from (D.22). Since in 5d $k_3 = 0$, the 4d extra Killing vector is $k_0 = \xi^1 k_1 + \xi^2 k_2$.

E Black hole charges and their reduction

The electric black hole charges computed in [18] in our notation read

$$Q_{\check{\tau}} = -\frac{1}{8\pi G_N^{(5)} g} \int_{S_\infty^3} G_{\check{\tau}J} \star_5 \widehat{F}^J, \quad (\text{E.1})$$

where the integral is taken on the three-sphere at infinity, and are dimensionless. We recall that only a subspace $\widehat{A}_\mu^{\check{\tau}}$ of the vector fields are massless on the AdS₅ vacuum, and the index

\mathfrak{T} runs over them. The massless vectors are such that the hypermultiplet scalars sit at a fixed point of the gauged isometries, and are thus identified by the conditions

$$k_{\mathfrak{T}}^u(q) = 0. \quad (\text{E.2})$$

Indeed, let \mathbb{B}^I_J be a matrix of linear redefinitions such that $\mathbb{B}^I_J \widehat{A}^J_\mu$ are mass eigenstates. Such a matrix is characterized by $\mathbb{B}^I_J G^{JN} k_N^u h_{uv} k_L^v = \lambda_N^I \mathbb{B}^N_L$ where λ is the diagonal matrix of squared masses (in units of g^2). The corresponding linear transformation of charges is $Q_I \rightarrow Q_J (\mathbb{B}^{-1})^J_I$, while the Killing vectors corresponding to the mass eigenstates are $k_J^u (\mathbb{B}^{-1})^J_I$. Now consider a massless vector and let the index \mathfrak{T} be such that $\lambda_{\mathfrak{T}}^{\mathfrak{T}} = 0$ (not summed over \mathfrak{T}). Using non-degeneracy of the metrics G_{IJ} and h_{uv} , one easily proves that $k_J^u (\mathbb{B}^{-1})^J_{\mathfrak{T}} = 0$, which is (E.2).

Now, the equations of motion for the bosonic fields of 5d gauged supergravity that follow from (B.2) are

$$\begin{aligned} d(G_{IJ} \star_5 \widehat{F}^J) &= \frac{1}{4} C_{IJK} \widehat{F}^J \wedge \widehat{F}^K - g h_{uv} k_I^u \star_5 \widehat{\mathcal{D}}q^v \\ \widehat{R}_{MN} &= G_{IJ} \left(\widehat{F}^I_{MP} \widehat{F}^{JP}_N - \frac{1}{6} \widehat{g}_{MN} \widehat{F}^I_{PQ} \widehat{F}^{JPQ} \right) \\ &\quad + \mathcal{G}_{ij} \partial_M \phi^i \partial_N \phi^j + h_{uv} \widehat{\mathcal{D}}_M q^u \widehat{\mathcal{D}}_N q^v + \frac{2}{3} \widehat{g}_{MN} g^2 V. \end{aligned} \quad (\text{E.3})$$

Notice that (E.2) is just the condition not to have a source in the \mathfrak{T} -th component of Maxwell's equation from the hypermultiplets. We can express the charges $\mathcal{Q}_{\mathfrak{T}}$ in terms of integrals at the horizon [77] using the EOMs (E.3):

$$\mathcal{Q}_{\mathfrak{T}} = -\frac{1}{8\pi G_N^{(5)} g} \left[\int_{S^3_r} G_{\mathfrak{T}J} \star_5 \widehat{F}^J + \int_{S^3_r \times I[r, \infty]} \left(\frac{1}{4} C_{\mathfrak{T}JK} \widehat{F}^J \wedge \widehat{F}^K - g h_{uv} k_{\mathfrak{T}}^u \star_5 \widehat{\mathcal{D}}q^v \right) \right]. \quad (\text{E.4})$$

The first term is an integral evaluated at radius r , that we will take to be the horizon location. The second term is a correction, integrated on a cylinder $S^3 \times I$ where I is the interval from r to ∞ , that leads to a Page charge. Assuming that the condition $k_{\mathfrak{T}}^u(q) = 0$ remains true also on the black hole background,³⁴ the third term vanishes.

We can apply a similar manipulation to the angular momenta $J_{a=1,2}$. Given the spacetime Killing vectors $K_a \equiv K_a^M \partial_M$, the angular momenta are defined in [18] as

$$J_a = \frac{1}{16\pi G_N^{(5)}} \int_{S^3_\infty} \star_5 dK_a \quad (\text{E.5})$$

³⁴In the case of the conifold compactification discussed in Section 5.2, this assumption is true, see (5.28). We expect the assumption to be true in all cases.

where we have indicated with the same symbol $K_a \equiv K_{aM} dx^M$ the 1-forms dual to the Killing vectors, and the integral is evaluated once again at infinity. One can show that the Killing equation implies

$$d \star_5 dK = 2 \widehat{R}_{MN} K^M \star_5 dx^N . \quad (\text{E.6})$$

We can then use the EOMs (E.3) to replace the Ricci scalar \widehat{R}_{MN} . We assume that S^3 is invariant under the isometries generated by K_a , therefore, indicating as i_K the interior product, the integral of $\widehat{g}_{MN} K^M \star_5 dx^N = i_K(\star_5 1)$ vanishes. We also assume that $i_K d\phi^i = 0$. We obtain

$$J_a = \frac{1}{16\pi G_N^{(5)}} \left[\int_{S^3} \star_5 dK_a + 2 \int_{S^3 \times I} \left(G_{IJ} (i_{K_a} \widehat{F}^I) \wedge \star_5 \widehat{F}^J + h_{uv} (i_{K_a} \widehat{D}q^u) \star_5 \widehat{D}q^v \right) \right] . \quad (\text{E.7})$$

Now let us proceed and reduce the charges to 4d imposing the ansatz (D.2), in particular

$$\begin{aligned} \widehat{A}^I &= A^I + \xi^I A^0 + (z_1^I + \xi^I)(dy - A^0) \\ \widehat{F}^I &= F^I - z_1^I F^0 + dz_1^I \wedge (dy - A^0) , \end{aligned} \quad (\text{E.8})$$

and performing the integrals along the circle. Notice that because of (D.6) and since the horizon areas in 5d and 4d are related by $\text{Area}_{(5)} = \Delta y \text{Area}_{(4)}$, the black hole entropy is the same in 5d and 4d. We find

$$\begin{aligned} \int_{S^3} G_{IJ} \star_5 \widehat{F}^J &= \Delta y \int_{S^2} e^{-2\tilde{\phi}} G_{IJ} \star_4 (F^J - z_1^J F^0) \\ C_{IJK} \int_{S^3 \times I} \widehat{F}^J \wedge \widehat{F}^K &= -\Delta y C_{IJK} \int_{S^2} \left(2z_1^J F^K - z_1^J z_1^K F^0 \right) . \end{aligned} \quad (\text{E.9})$$

In the second equality we used that $z_1^I \rightarrow 0$ at infinity. The electric charges are thus

$$\mathcal{Q}_{\mathfrak{z}} = \frac{1}{g} \int_{S^2} \frac{\delta S_{4d}}{\delta F^{\mathfrak{z}}} - \frac{1}{8\pi G_N^{(4)} g} C_{\mathfrak{z}JK} \int_{S^2} \left(\frac{1}{2} \xi^J F^K + \frac{1}{4} \xi^J \xi^K F^0 \right) , \quad (\text{E.10})$$

where

$$\frac{\delta S_{4d}}{\delta F^\Lambda} = \frac{1}{16\pi G_N^{(4)}} \left(\text{Im} \mathcal{N}_{\Lambda\Sigma} \star_4 F^\Sigma + \text{Re} \mathcal{N}_{\Lambda\Sigma} F^\Sigma \right) \quad (\text{E.11})$$

are the derivatives of the action obtained from (C.2) with (D.22) and (D.24).

We define the 4d dimensionless magnetic charges as

$$p^\Lambda = \frac{g}{4\pi} \int_{S^2} F^\Lambda , \quad (\text{E.12})$$

where the integral can be done at any radius because of the Bianchi identities. On the other hand, the first Chern class of the circle fibration — that we take to be the Hopf fibration of S^3 — is $\frac{1}{\Delta y} \int dA^0 = 1$. Thus, we obtain a properly quantized $p^0 = 1$ if we set

$$\Delta y = \frac{4\pi}{g} . \quad (\text{E.13})$$

We will use this normalization from now on.

Let us now reduce the angular momentum. We consider the case $J_1 = J_2$, with $J_{1,2}$ normalized such that they generate orbits of length 2π , and define $J = (J_1 + J_2)/2$. The corresponding Killing vector and dual 1-form are

$$K^M \partial_M = \frac{\Delta y}{4\pi} \frac{\partial}{\partial y} = \frac{1}{g} \frac{\partial}{\partial y}, \quad K_M dx^M = \frac{1}{g} e^{-4\tilde{\phi}} (dy - A^0). \quad (\text{E.14})$$

The first term in (E.7) gives

$$\int_{S^3} \star_5 dK = -\frac{\Delta y}{g} \int_{S^2} e^{-6\tilde{\phi}} \star_4 F^0. \quad (\text{E.15})$$

To reduce the second term we use $i_K \widehat{F}^I = -\frac{1}{g} dz_1^I$, integrate by parts, and use the EOMs (E.3). To reduce the third term we use $i_K \widehat{\mathcal{D}}q^u = (z_1^I + \xi^I) k_I^u$ and $i_\omega(\star 1) = \star \omega$ for a 1-form ω . Eventually

$$\begin{aligned} J &= \frac{1}{8\pi G_N^{(4)} g} \left\{ \int_{S_r^2} \left[-\frac{1}{2} e^{-6\tilde{\phi}} \star_4 F^0 + e^{-2\tilde{\phi}} G_{IJ} z_1^I \star_4 (F^J - z_1^J F^0) \right. \right. \\ &\quad \left. \left. - C_{IJK} \left(\frac{1}{4} z_1^I z_1^J F^K - \frac{1}{6} z_1^I z_1^J z_1^K F^0 \right) \right] + \int_{S^2 \times I} \star_4 g k_0^u h_{uv} \mathcal{D}q^v \right\} \\ &= \frac{1}{g} \int_{S_r^2} \frac{\delta S_{4d}}{\delta F^0} - \frac{1}{8\pi G_N^{(4)} g} \left[C_{IJK} \xi^I \xi^J \int_{S_r^2} \left(\frac{1}{4} F^K + \frac{1}{6} \xi^K F^0 \right) + \int_{S^2 \times I} \star_4 g k_0^u h_{uv} \mathcal{D}q^v \right]. \end{aligned} \quad (\text{E.16})$$

The four-dimensional angular momentum of the black hole solution is proportional to $J_1 - J_2$, which vanishes in the case under consideration. This implies that we can impose spherical symmetry on S^2 . The section $\mathcal{D}q^v$ is charged under the Abelian vector fields A_μ^Λ , therefore the magnetic fluxes p^Λ give rise to an effective spin s on S^2 . However, the spin spherical harmonics [78, 79] have total angular momentum $j \geq |s|$, which should vanish in order to have a spherically-symmetric configuration. Since the Abelian symmetries are realized non-linearly on $\mathcal{D}q^v$ as soon as $k_\Lambda^u \neq 0$, we obtain the condition

$$p^\Lambda k_\Lambda^u(q) = 0 \quad (\text{E.17})$$

for spherically-symmetric black hole solutions. Without loss of generality, in Section 5 we have set $p^I = 0$ which implies $k_0^u = 0$. We then see that the last term in (E.16) vanishes.

The magnetic charges that appear in the attractor equations of [39], in our conventions, are (E.12) while the electric charges are

$$q_\Lambda = \frac{g}{4\pi} \int_{S_r^2} G_\Lambda \quad \text{with} \quad G_\Lambda = 16\pi G_N^{(4)} \frac{\delta S_{4d}}{\delta F^\Lambda}. \quad (\text{E.18})$$

Setting $p^I = 0$, we obtain the following dictionary between 5d and 4d charges:

$$\begin{aligned} q_0 &= 4G_N^{(4)} g^2 J + \frac{1}{3} C_{IJK} \xi^I \xi^J \xi^K p^0 \\ q_{\bar{x}} &= 4G_N^{(4)} g^2 Q_{\bar{x}} + \frac{1}{2} C_{\bar{x}JK} \frac{1}{2} \xi^J \xi^K p^0 . \end{aligned} \quad (\text{E.19})$$

E.1 Baryonic charge quantization in the conifold theory

In order to fix the exact relation between the supergravity charge Q_3 and the field theory baryonic charge Q_B , we deduce the Dirac quantization condition satisfied by A_μ^3 from the consistent reduction of [34].

The metric of $T^{1,1}$ is

$$ds^2 = \frac{1}{6} \sum_{i=1,2} \left(d\theta_i^2 + \sin^2 \theta_i d\varphi_i^2 \right) + \eta^2 \quad \text{with} \quad \eta = -\frac{1}{3} \left(d\psi + \sum_{i=1,2} \cos \theta_i d\varphi_i \right) . \quad (\text{E.20})$$

We define the 2-forms³⁵

$$\begin{aligned} J &= \frac{1}{6} \left(\sin \theta_1 d\theta_1 \wedge d\varphi_1 + \sin \theta_2 d\theta_2 \wedge d\varphi_2 \right) = \frac{1}{2} d\eta \\ \Phi &= \frac{1}{6} \left(\sin \theta_1 d\theta_1 \wedge d\varphi_1 - \sin \theta_2 d\theta_2 \wedge d\varphi_2 \right) . \end{aligned} \quad (\text{E.21})$$

The expansion of the 10d RR field strength F_5^{RR} in [34] around the $\text{AdS}_5 \times T^{1,1}$ vacuum (where $u = v = w = b^\Omega = c^\Omega = 0$), keeping only the dependence on the gauge fields and the Stückelberg scalar a , in our conventions reads

$$\begin{aligned} F_5^{\text{RR}} &= 4g \star_5 1 - 2g^{-1} (\star_5 Da) \wedge (\eta - g\widehat{A}^1) - g^{-2} (\star_5 d\widehat{A}^2) \wedge J + g^{-2} (\star_5 d\widehat{A}^3) \wedge \Phi \\ &\quad - g^{-3} d\widehat{A}^2 \wedge J \wedge (\eta - g\widehat{A}^1) - g^{-3} d\widehat{A}^3 \wedge \Phi \wedge (\eta - g\widehat{A}^1) \\ &\quad + g^{-4} J \wedge J \wedge (Da + 2(\eta - g\widehat{A}^1)) , \end{aligned} \quad (\text{E.22})$$

where \star_5 is the Poincaré dual in AdS_5 while $Da = da + 2g(\widehat{A}^1 + \widehat{A}^2)$. Dirac's quantization condition reads

$$\frac{1}{2\sqrt{\pi} \kappa_{10}} \int_{\mathcal{C}_5} F_5^{\text{RR}} \in \mathbb{Z} \quad (\text{E.23})$$

for any closed 5-cycle \mathcal{C}_5 . Here κ_{10} is the 10d gravitational coupling, related to the 5d Newton constant by

$$\frac{\text{Vol}(T^{1,1})}{g^5 \kappa_{10}^2} = \frac{1}{8\pi G_N^{(5)}} \quad (\text{E.24})$$

where $\text{Vol}(T^{1,1}) = 16\pi^3/27$. Applying (E.23) to $\mathcal{C}_5 = T^{1,1}$ and imposing that there are N units of 5-form flux, we recover (5.35). On the other hand, let us apply (E.23) to the 5-cycle

³⁵The 2-form J should not be confused with the angular momentum of the black hole.

$X_2 \times S^3$, where X_2 is the non-trivial 2-cycle of $T^{1,1}$ while S^3 is a spatial 3-sphere in AdS_5 . Using $\int_{X_2} J = 0$ and $\int_{X_2} \Phi = 4\pi/3$ as well as (5.35), we obtain

$$\frac{1}{2\sqrt{\pi} \kappa_{10}} \int_{X_2 \times S^3} F_5^{\text{RR}} = \frac{1}{6\pi G_N^{(5)} gN} \int_{S^3} \left(\star_5 \widehat{F}^3 + \widehat{F}^3 \wedge \widehat{A}^1 \right) = -\frac{4}{3N} Q_3 \in \mathbb{Z}, \quad (\text{E.25})$$

where $\widehat{F}^3 = d\widehat{A}^3$. According to (E.3) and using (B.42) and (B.43), the combination in parenthesis gives the Page charge Q_3 , which is conserved and quantized. Taking the 3-sphere to spatial infinity, it coincides with the charge defined in (E.1).

References

- [1] F. Benini, K. Hristov, and A. Zaffaroni, “Black hole microstates in AdS_4 from supersymmetric localization,” *JHEP* **05** (2016) 054, [arXiv:1511.04085 \[hep-th\]](#).
- [2] F. Benini and A. Zaffaroni, “Supersymmetric partition functions on Riemann surfaces,” *Proc. Symp. Pure Math.* **96** (2017) 13–46, [arXiv:1605.06120 \[hep-th\]](#).
- [3] F. Benini, K. Hristov, and A. Zaffaroni, “Exact microstate counting for dyonic black holes in AdS_4 ,” *Phys. Lett.* **B771** (2017) 462–466, [arXiv:1608.07294 \[hep-th\]](#).
- [4] A. Cabo-Bizet, D. Cassani, D. Martelli, and S. Murthy, “Microscopic origin of the Bekenstein-Hawking entropy of supersymmetric AdS_5 black holes,” *JHEP* **10** (2019) 062, [arXiv:1810.11442 \[hep-th\]](#).
- [5] S. Choi, J. Kim, S. Kim, and J. Nahmgoong, “Large AdS black holes from QFT,” [arXiv:1810.12067 \[hep-th\]](#).
- [6] F. Benini and P. Milan, “Black holes in 4d $\mathcal{N}=4$ Super-Yang-Mills,” *Phys. Rev. X* **10** (2020) 021037, [arXiv:1812.09613 \[hep-th\]](#).
- [7] S. Choi, C. Hwang, and S. Kim, “Quantum vortices, M2-branes and black holes,” [arXiv:1908.02470 \[hep-th\]](#).
- [8] N. Bobev and P. M. Cricigno, “Universal spinning black holes and theories of class \mathcal{R} ,” *JHEP* **12** (2019) 054, [arXiv:1909.05873 \[hep-th\]](#).
- [9] F. Benini, D. Gang, and L. A. Pando Zayas, “Rotating Black Hole Entropy from M5-branes,” *JHEP* **03** (2020) 057, [arXiv:1909.11612 \[hep-th\]](#).
- [10] A. Zaffaroni, “Lectures on AdS Black Holes, Holography and Localization,” [arXiv:1902.07176 \[hep-th\]](#).

- [11] C. Romelsberger, “Counting chiral primaries in $\mathcal{N}=1$, $d=4$ superconformal field theories,” *Nucl. Phys.* **B747** (2006) 329–353, [arXiv:hep-th/0510060 \[hep-th\]](#).
- [12] J. Kinney, J. M. Maldacena, S. Minwalla, and S. Raju, “An Index for 4 dimensional super conformal theories,” *Commun. Math. Phys.* **275** (2007) 209–254, [arXiv:hep-th/0510251 \[hep-th\]](#).
- [13] J. Bhattacharya, S. Bhattacharyya, S. Minwalla, and S. Raju, “Indices for Superconformal Field Theories in 3,5 and 6 Dimensions,” *JHEP* **02** (2008) 064, [arXiv:0801.1435 \[hep-th\]](#).
- [14] J. B. Gutowski and H. S. Reall, “Supersymmetric AdS_5 black holes,” *JHEP* **02** (2004) 006, [arXiv:hep-th/0401042 \[hep-th\]](#).
- [15] J. B. Gutowski and H. S. Reall, “General supersymmetric AdS_5 black holes,” *JHEP* **04** (2004) 048, [arXiv:hep-th/0401129 \[hep-th\]](#).
- [16] Z. W. Chong, M. Cvetič, H. Lu, and C. N. Pope, “Five-dimensional gauged supergravity black holes with independent rotation parameters,” *Phys. Rev.* **D72** (2005) 041901, [arXiv:hep-th/0505112 \[hep-th\]](#).
- [17] Z. W. Chong, M. Cvetič, H. Lu, and C. N. Pope, “General non-extremal rotating black holes in minimal five-dimensional gauged supergravity,” *Phys. Rev. Lett.* **95** (2005) 161301, [arXiv:hep-th/0506029 \[hep-th\]](#).
- [18] H. K. Kunduri, J. Lucietti, and H. S. Reall, “Supersymmetric multi-charge AdS_5 black holes,” *JHEP* **04** (2006) 036, [arXiv:hep-th/0601156 \[hep-th\]](#).
- [19] J. Markeviciute and J. E. Santos, “Evidence for the existence of a novel class of supersymmetric black holes with $\text{AdS}_5 \times S^5$ asymptotics,” *Class. Quant. Grav.* **36** (2019) 02LT01, [arXiv:1806.01849 \[hep-th\]](#).
- [20] J. Markeviciute, “Rotating Hairy Black Holes in $\text{AdS}_5 \times S^5$,” *JHEP* **03** (2019) 110, [arXiv:1809.04084 \[hep-th\]](#).
- [21] S. M. Hosseini, K. Hristov, and A. Zaffaroni, “An extremization principle for the entropy of rotating BPS black holes in AdS_5 ,” *JHEP* **07** (2017) 106, [arXiv:1705.05383 \[hep-th\]](#).
- [22] D. Cassani and L. Papini, “The BPS limit of rotating AdS black hole thermodynamics,” *JHEP* **09** (2019) 079, [arXiv:1906.10148 \[hep-th\]](#).

- [23] A. Cabo-Bizet and S. Murthy, “Supersymmetric phases of 4d $\mathcal{N}=4$ SYM at large N ,” [arXiv:1909.09597 \[hep-th\]](#).
- [24] F. Benini and P. Milan, “A Bethe Ansatz type formula for the superconformal index,” *Commun. Math. Phys.* **376** (2020) 1413–1440, [arXiv:1811.04107 \[hep-th\]](#).
- [25] C. Closset, H. Kim, and B. Willett, “ $\mathcal{N}=1$ supersymmetric indices and the four-dimensional A -model,” *JHEP* **08** (2017) 090, [arXiv:1707.05774 \[hep-th\]](#).
- [26] S. M. Hosseini, A. Nedelin, and A. Zaffaroni, “The Cardy limit of the topologically twisted index and black strings in AdS_5 ,” *JHEP* **04** (2017) 014, [arXiv:1611.09374 \[hep-th\]](#).
- [27] J. Hong and J. T. Liu, “The topologically twisted index of $\mathcal{N}=4$ super-Yang-Mills on $T^2 \times S^2$ and the elliptic genus,” *JHEP* **07** (2018) 018, [arXiv:1804.04592 \[hep-th\]](#).
- [28] A. Lanir, A. Nedelin, and O. Sela, “Black hole entropy function for toric theories via Bethe Ansatz,” *JHEP* **04** (2020) 091, [arXiv:1908.01737 \[hep-th\]](#).
- [29] A. Arabi Ardehali, J. Hong, and J. T. Liu, “Asymptotic growth of the 4d $\mathcal{N}=4$ index and partially deconfined phases,” [arXiv:1912.04169 \[hep-th\]](#).
- [30] A. González Lezcano and L. A. Pando Zayas, “Microstate counting via Bethe Ansatz in the 4d $\mathcal{N}=1$ superconformal index,” *JHEP* **03** (2020) 088, [arXiv:1907.12841v3 \[hep-th\]](#).
- [31] S. M. Hosseini, K. Hristov, and A. Zaffaroni, “A note on the entropy of rotating BPS $\text{AdS}_7 \times S^4$ black holes,” *JHEP* **05** (2018) 121, [arXiv:1803.07568 \[hep-th\]](#).
- [32] A. Amariti, I. Garozzo, and G. Lo Monaco, “Entropy function from toric geometry,” [arXiv:1904.10009 \[hep-th\]](#).
- [33] I. R. Klebanov and E. Witten, “Superconformal field theory on three-branes at a Calabi-Yau singularity,” *Nucl. Phys.* **B536** (1998) 199–218, [arXiv:hep-th/9807080 \[hep-th\]](#).
- [34] D. Cassani and A. F. Faedo, “A Supersymmetric consistent truncation for conifold solutions,” *Nucl. Phys.* **B843** (2011) 455–484, [arXiv:1008.0883 \[hep-th\]](#).
- [35] I. Bena, G. Giecold, M. Grana, N. Halmagyi, and F. Orsi, “Supersymmetric Consistent Truncations of IIB on $T^{1,1}$,” *JHEP* **04** (2011) 021, [arXiv:1008.0983 \[hep-th\]](#).
- [36] N. Halmagyi, J. T. Liu, and P. Szepietowski, “On $\mathcal{N}=2$ Truncations of IIB on $T^{1,1}$,” *JHEP* **07** (2012) 098, [arXiv:1111.6567 \[hep-th\]](#).

- [37] G. Dall’Agata and A. Gnecci, “Flow equations and attractors for black holes in $\mathcal{N}=2$ $U(1)$ gauged supergravity,” *JHEP* **03** (2011) 037, [arXiv:1012.3756 \[hep-th\]](#).
- [38] N. Halmagyi, M. Petrini, and A. Zaffaroni, “BPS black holes in AdS_4 from M-theory,” *JHEP* **08** (2013) 124, [arXiv:1305.0730 \[hep-th\]](#).
- [39] D. Klemm, N. Petri, and M. Rabbiosi, “Symplectically invariant flow equations for $\mathcal{N}=2$, $D=4$ gauged supergravity with hypermultiplets,” *JHEP* **04** (2016) 008, [arXiv:1602.01334 \[hep-th\]](#).
- [40] A. Cabo-Bizet, D. Cassani, D. Martelli, and S. Murthy, “The large- N limit of the 4d $\mathcal{N}=1$ superconformal index,” [arXiv:2005.10654 \[hep-th\]](#).
- [41] G. Felder and A. Varchenko, “Multiplication Formulas for the Elliptic Gamma Function,” [arXiv:math/0212155 \[math.QA\]](#).
- [42] G. Felder and A. Varchenko, “The elliptic gamma function and $SL(3, \mathbb{Z}) \ltimes \mathbb{Z}^3$,” *Adv. Math.* **156** (2000) 44–76, [arXiv:math/9907061 \[math.QA\]](#).
- [43] J. Kim, S. Kim, and J. Song, “A 4d $\mathcal{N}=1$ Cardy Formula,” [arXiv:1904.03455 \[hep-th\]](#).
- [44] A. Cabo-Bizet, D. Cassani, D. Martelli, and S. Murthy, “The asymptotic growth of states of the 4d $\mathcal{N}=1$ superconformal index,” *JHEP* **08** (2019) 120, [arXiv:1904.05865 \[hep-th\]](#).
- [45] D. Anselmi, D. Z. Freedman, M. T. Grisaru, and A. A. Johansen, “Nonperturbative formulas for central functions of supersymmetric gauge theories,” *Nucl. Phys.* **B526** (1998) 543–571, [arXiv:hep-th/9708042 \[hep-th\]](#).
- [46] S. Benvenuti, L. A. Pando Zayas, and Y. Tachikawa, “Triangle anomalies from Einstein manifolds,” *Adv. Theor. Math. Phys.* **10** (2006) 395–432, [arXiv:hep-th/0601054 \[hep-th\]](#).
- [47] D. R. Morrison and M. R. Plesser, “Nonspherical horizons. 1.,” *Adv. Theor. Math. Phys.* **3** (1999) 1–81, [arXiv:hep-th/9810201 \[hep-th\]](#).
- [48] A. Hanany and K. D. Kennaway, “Dimer models and toric diagrams,” [arXiv:hep-th/0503149 \[hep-th\]](#).
- [49] S. Franco, A. Hanany, K. D. Kennaway, D. Vegh, and B. Wecht, “Brane dimers and quiver gauge theories,” *JHEP* **01** (2006) 096, [arXiv:hep-th/0504110 \[hep-th\]](#).

- [50] S. Franco, A. Hanany, D. Martelli, J. Sparks, D. Vegh, and B. Wecht, “Gauge theories from toric geometry and brane tilings,” *JHEP* **01** (2006) 128, [arXiv:hep-th/0505211 \[hep-th\]](#).
- [51] B. Feng, Y.-H. He, K. D. Kennaway, and C. Vafa, “Dimer models from mirror symmetry and quivering amoebae,” *Adv. Theor. Math. Phys.* **12** (2008) 489–545, [arXiv:hep-th/0511287 \[hep-th\]](#).
- [52] A. Butti and A. Zaffaroni, “R-charges from toric diagrams and the equivalence of a -maximization and Z -minimization,” *JHEP* **11** (2005) 019, [arXiv:hep-th/0506232 \[hep-th\]](#).
- [53] S. Kim and K.-M. Lee, “1/16-BPS Black Holes and Giant Gravitons in the $\text{AdS}_5 \times S^5$ Space,” *JHEP* **12** (2006) 077, [arXiv:hep-th/0607085 \[hep-th\]](#).
- [54] K. Hristov, “Dimensional reduction of BPS attractors in AdS gauged supergravities,” *JHEP* **12** (2014) 066, [arXiv:1409.8504 \[hep-th\]](#).
- [55] K. Hristov, S. Katmadas, and C. Toldo, “Rotating attractors and BPS black holes in AdS_4 ,” *JHEP* **01** (2019) 199, [arXiv:1811.00292 \[hep-th\]](#).
- [56] K. Hristov, S. Katmadas, and C. Toldo, “Matter-coupled supersymmetric Kerr-Newman- AdS_4 black holes,” *Phys. Rev. D* **100** (2019) 066016, [arXiv:1907.05192 \[hep-th\]](#).
- [57] M. Gunaydin, G. Sierra, and P. K. Townsend, “The Geometry of $\mathcal{N}=2$ Maxwell-Einstein Supergravity and Jordan Algebras,” *Nucl. Phys.* **B242** (1984) 244–268.
- [58] M. Gunaydin, G. Sierra, and P. K. Townsend, “Gauging the $d=5$ Maxwell-Einstein Supergravity Theories: More on Jordan Algebras,” *Nucl. Phys.* **B253** (1985) 573.
- [59] A. Ceresole and G. Dall’Agata, “General matter coupled $\mathcal{N}=2$, $D=5$ gauged supergravity,” *Nucl. Phys.* **B585** (2000) 143–170, [arXiv:hep-th/0004111 \[hep-th\]](#).
- [60] L. Andrianopoli, M. Bertolini, A. Ceresole, R. D’Auria, S. Ferrara, P. Fre, and T. Magri, “ $\mathcal{N}=2$ supergravity and $\mathcal{N}=2$ superYang-Mills theory on general scalar manifolds: Symplectic covariance, gaugings and the momentum map,” *J. Geom. Phys.* **23** (1997) 111–189, [arXiv:hep-th/9605032 \[hep-th\]](#).
- [61] B. Craps, F. Roose, W. Troost, and A. Van Proeyen, “What is special Kahler geometry?,” *Nucl. Phys.* **B503** (1997) 565–613, [arXiv:hep-th/9703082 \[hep-th\]](#).

- [62] L. Andrianopoli, S. Ferrara, and M. A. Lledo, “Scherk-Schwarz reduction of $D=5$ special and quaternionic geometry,” *Class. Quant. Grav.* **21** (2004) 4677–4696, [arXiv:hep-th/0405164](#) [hep-th].
- [63] K. Behrndt, G. Lopes Cardoso, and S. Mahapatra, “Exploring the relation between 4D and 5D BPS solutions,” *Nucl. Phys.* **B732** (2006) 200–223, [arXiv:hep-th/0506251](#) [hep-th].
- [64] G. Lopes Cardoso, J. M. Oberreuter, and J. Perz, “Entropy function for rotating extremal black holes in very special geometry,” *JHEP* **05** (2007) 025, [arXiv:hep-th/0701176](#) [hep-th].
- [65] H. Looyestijn, E. Plaushinn, and S. Vandoren, “New potentials from Scherk-Schwarz reductions,” *JHEP* **12** (2010) 016, [arXiv:1008.4286](#) [hep-th].
- [66] D. Klemm, N. Petri, and M. Rabbiosi, “Black string first order flow in $\mathcal{N}=2$, $d=5$ Abelian gauged supergravity,” *JHEP* **01** (2017) 106, [arXiv:1610.07367](#) [hep-th].
- [67] K. Hristov and A. Rota, “6d-5d-4d reduction of BPS attractors in flat gauged supergravities,” *Nucl. Phys. B* **897** (2015) 213–228, [arXiv:1410.5386](#) [hep-th].
- [68] F. Benini, H. Khachatryan, and P. Milan, “Black hole entropy in massive Type IIA,” *Class. Quant. Grav.* **35** (2018) 035004, [arXiv:1707.06886](#) [hep-th].
- [69] S. M. Hosseini, K. Hristov, and A. Passias, “Holographic microstate counting for AdS_4 black holes in massive IIA supergravity,” *JHEP* **10** (2017) 190, [arXiv:1707.06884](#) [hep-th].
- [70] E. Bergshoeff, S. Cucu, T. De Wit, J. Gheerardyn, R. Halbersma, S. Vandoren, and A. Van Proeyen, “Superconformal $\mathcal{N}=2$, $D=5$ matter with and without actions,” *JHEP* **10** (2002) 045, [arXiv:hep-th/0205230](#) [hep-th].
- [71] D. Z. Freedman and A. Van Proeyen, *Supergravity*. Cambridge Univ. Press, Cambridge, UK, 2012.
- [72] B. de Wit and A. Van Proeyen, “Special geometry, cubic polynomials and homogeneous quaternionic spaces,” *Commun. Math. Phys.* **149** (1992) 307–334, [arXiv:hep-th/9112027](#) [hep-th].
- [73] J. Bagger and E. Witten, “Matter Couplings in $\mathcal{N}=2$ Supergravity,” *Nucl. Phys.* **B222** (1983) 1–10.

- [74] M. Berger, “Sur les groupes d’holonomie homogènes de variétés à connexion affine et des variétés Riemanniennes,” *Bull. Soc. Math. France* **83** (1955) 279–330.
- [75] S. Salamon, “Quaternionic Kähler Manifolds,” *Invent. Math.* **67** (1982) 143–171.
- [76] A. Ceresole, R. D’Auria, S. Ferrara, and A. Van Proeyen, “Duality transformations in supersymmetric Yang-Mills theories coupled to supergravity,” *Nucl. Phys.* **B444** (1995) 92–124, [arXiv:hep-th/9502072](#) [[hep-th](#)].
- [77] K. Hanaki, K. Ohashi, and Y. Tachikawa, “Comments on charges and near-horizon data of black rings,” *JHEP* **12** (2007) 057, [arXiv:0704.1819](#) [[hep-th](#)].
- [78] E. Newman and R. Penrose, “Note on the Bondi-Metzner-Sachs group,” *J. Math. Phys.* **7** (1966) 863–870.
- [79] T. T. Wu and C. N. Yang, “Dirac Monopole Without Strings: Monopole Harmonics,” *Nucl. Phys. B* **107** (1976) 365.