



SCUOLA INTERNAZIONALE SUPERIORE DI STUDI AVANZATI

SISSA Digital Library

Self-consistent harmonic approximation in presence of non-local couplings

*Original*

Self-consistent harmonic approximation in presence of non-local couplings / Giachetti, G., Defenu, N., Ruffo, S., Trombettoni, A.. - In: EUROPHYSICS LETTERS. - ISSN 0295-5075. - 133:5(2021), pp. 1-11. [10.1209/0295-5075/133/57004]

*Availability:*

This version is available at: 20.500.11767/127025 since: 2022-03-11T14:01:46Z

*Publisher:*

*Published*

DOI:10.1209/0295-5075/133/57004

*Terms of use:*

Testo definito dall'ateneo relativo alle clausole di concessione d'uso

*Publisher copyright*  
EDP Sciences

This version is available for education and non-commercial purposes.

note finali coverpage

(Article begins on next page)

# Self-consistent harmonic approximation with non-local couplings

Guido Giachetti<sup>1</sup>, Nicolò Defenu<sup>2</sup>, Stefano Ruffo<sup>1,3</sup>, and Andrea Trombettoni<sup>1,4</sup>

<sup>1</sup>*SISSA and INFN, Sezione di Trieste, Via Bonomea 265, I-34136 Trieste, Italy*

<sup>2</sup>*Institute for Theoretical Physics, ETH Zürich Wolfgang-Pauli-Str. 27, 8093 Zurich, Switzerland*

<sup>3</sup>*Istituto dei Sistemi Complessi, Consiglio Nazionale delle Ricerche, Via Madonna del Piano 10, I-50019 Sesto Fiorentino, Italy*

<sup>4</sup>*Department of Physics, University of Trieste, Strada Costiera*

January 1, 2021

## Abstract

We derive the self-consistent harmonic approximation for the  $2D$  XY model with non-local interactions. The resulting equation for the variational couplings holds for any form of the spin-spin coupling as well as for any dimension. Our analysis is then specialized to power-law couplings decaying with the distance  $r$  as  $\propto 1/r^{2+\sigma}$  in order to investigate the robustness, at finite  $\sigma$ , of the Berezinskii-Kosterlitz-Thouless (BKT) transition, which occurs in the short-range limit  $\sigma \rightarrow \infty$ . We propose an ansatz for the functional form of the variational couplings and show that for any  $\sigma > 2$  the BKT mechanism occurs. The present investigation provides an upper bound for the lower critical threshold  $\sigma^* = 2$ , above which the traditional BKT transition persists in spite of the LR couplings.

## 1 Introduction

Among the different strategies commonly employed to study interacting systems, one that is often used – in its simplest form – is based on the determination of the non-interacting model that better approximates, in a variational sense, the initial, interacting problem. This method, referred to as the Self-Consistent Harmonic Approximation (SCHA), can be improved by considering non-quadratic, but solvable, approximations of the problem at hand or by an integration of the quantum fluctuations to determine an optimized classical potential [1, 2]. Then, the SCHA and its variants may take different forms both for classical or quantum models, and for equilibrium or dynamical properties. They are in general based on the variational principle of the minimization of the energy (or free energy) difference between the interacting model and the approximating one, calculated using the state of the latter [3]. When the approximating model is non-interacting, this approach cannot be used to determine non-trivial correlation functions or interference effects, but also in this case SCHA provides estimates for the equilibrium free energy and the properties of the phases of the model under study. SCHA and its variants, including the so-called pure quantum SCHA, have been as well used to provide an estimate of the quantum corrections to the free energy of nonlinear systems [4, 5] and it has proven useful in several contexts, ranging from the study of phonon spectra in metals and insulators (see [6] and refs. therein) to quantum antiferromagnets [7] and arrays of Josephson junctions [8].

A context in which SCHA has been widely employed along the years is provided by the two-dimensional XY model, where the celebrated Berezinskii-Kosterlitz-Thouless (BKT) transition takes place [9–11]. In the BKT transition, the low temperature phase features vortex-antivortex pairs and power-law correlation functions (with temperature-dependent exponents), while in the high temperature phase topological charges can freely propagate leading to exponential decay in the correlation functions [12–15]. The superfluid density exhibits a universal jump at the critical temperature  $T_{BKT}$  [16] and, correspondingly, both the critical exponent  $\eta$  [10, 11] and the power law exponent for the finite size scaling of the largest eigenvalue of the one-body density matrix [17], jump at  $T_{BKT}$  from a finite value, respectively  $1/4$  and  $7/8$ , to zero.

The Hamiltonian of the short-range, nearest-neighbor XY model reads

$$H = -\frac{J_0}{2} \sum_{\mathbf{i}, \mathbf{j}} \cos(\theta_{\mathbf{i}} - \theta_{\mathbf{j}}), \quad (1)$$

where  $J_0$  is the coupling constant and the sum is on all the pairs of nearest-neighbor sites of a  $2D$  square lattice, on which the variables  $\theta_i$  are defined. The temperature  $T_{BKT}$  of the XY model (1) has been the subject of considerable work, and the value determined by Monte Carlo simulations is given by  $k_B T_{BKT}/J_0 = 0.893 \pm 0.001$  [18–23], with  $k_B$  the Boltzmann constant. We will put, as usual,  $\beta = 1/k_B T$ , with  $T$  the temperature.

The SCHA for the short-range Hamiltonian (1) applies in a particularly transparent way. One introduces the quadratic Hamiltonian  $H_0 = \frac{\tilde{J}_0}{4} \sum_{\mathbf{i}, \mathbf{j}} (\theta_{\mathbf{i}} - \theta_{\mathbf{j}})^2$  and then variationally determines  $\tilde{J}_0$  as a function of  $J_0$  and  $T$ , or – equivalently – the dimensionless coupling  $\beta \tilde{J}_0$  as a function of  $\beta J_0$  [24–27]. It is found that  $\tilde{J}_0$  is different from zero for  $T$  smaller than a temperature, denoted by  $T_c$ , at which the effective  $\tilde{J}_0$  drops to zero [24]. An improved determination of the BKT critical temperature  $T_{BKT}$  can be obtained inserting  $\tilde{J}_0(T)$  in the Nelson-Kosterlitz condition [16] at the BKT critical point [25, 26]. An advantage of this approach is that it can be extended to the quantum phase model [8, 28, 29] describing arrays of Josephson junctions [30, 31] and ultracold bosons in optical lattices in the large filling limit [32].

When the couplings are non-local, i.e. the spins in sites  $\mathbf{i}$  and  $\mathbf{j}$  are coupled with a strength  $J(\mathbf{i}, \mathbf{j})$ , the application of the SCHA with the introduction of a variational coupling matrix  $\tilde{J}(\mathbf{i}, \mathbf{j})$  faces with the practical problem of solving the full set of conditions for the  $\tilde{J}$ 's. Here, we focus on the case of Hamiltonians with non-local couplings, proposing and discussing the consequences of a functional ansatz for the variational couplings  $\tilde{J}(\mathbf{i}, \mathbf{j})$ . In particular, we will consider the case of the  $d = 2$  XY with power-law couplings  $J(\mathbf{i}, \mathbf{j})$  decaying asymptotically as  $J \sim 1/|\mathbf{i} - \mathbf{j}|^{2+\sigma}$ . The reason for this choice is three-fold.

*i)* The nearest-neighbor  $2D$  XY model has been extensively studied with SCHA [24–27] providing a benchmark for the short-range limit of the theory. *ii)* The study of statistical mechanics models with long-range interactions attracted considerable attention along the last few decades (see the reviews [33–35]). The general result is that exists a value of the exponent of the power-law decay, denoted by  $\sigma^*$ , such that for  $\sigma > \sigma^*$  the universality class is the same of the short-range limit  $\sigma \rightarrow \infty$ . A first derivation by Sak, focusing on  $O(n)$  models [36], provided the result  $\sigma^* = 2 - \eta_{SR}$ , where  $\eta_{SR}$  is the anomalous dimension of the short-range ( $\sigma \rightarrow \infty$ ) limit. The validity of this result has been thoroughly investigated using a variety of techniques [37–47], yielding very strong evidences in its favor. Nevertheless, Sak's result only concerns second-order phase transitions, while the BKT transition is an infinite order one, so that the traditional  $\sigma^* = 2 - \eta_{SR}$  threshold does not apply to the XY model with non-local couplings. *iii)* Results on the BKT transition for the  $2D$  XY model with non-local couplings are very rare, to the best of our knowledge, with exceptions coming from related models such as  $1D$  quantum XXZ spin models with long-range interactions [48–50] and random-dilute graphs [51]. Then, qualitative information coming from the SCHA may provide insights for further investigations.

## 2 SCHA with non-local couplings

We consider the XY model on a 2-dimensional square lattice of  $N$  sites with non-local couplings

$$\beta H \equiv -\frac{1}{2} \sum_{\mathbf{i}, \mathbf{j}} J(r) \cos(\theta_{\mathbf{i}} - \theta_{\mathbf{j}}). \quad (2)$$

where  $r = |\mathbf{r}|$  and  $\mathbf{r} = \mathbf{i} - \mathbf{j}$  and, as previously mentioned, we choose

$$J(r) = \frac{J}{r^{2+\sigma}}, \quad (3)$$

with  $\sigma > 0$ . In Eq. (2) we are setting  $J \equiv \beta J_0$  and we take energy in units of  $J_0$  (restoring  $J_0$  when useful for clarity).

However, for the moment we do not need to fix a specific form for the couplings  $J(r)$ , and our results till Eq. (20) hold for any non-local couplings  $J(r)$ .

Proceeding according to the SCHA, we replace the cosine in the original Hamiltonian (2) with a quadratic term

$$\beta H_0 = \frac{1}{4} \sum_{\mathbf{i}, \mathbf{j}} \tilde{J}(\mathbf{r})(\theta_{\mathbf{i}} - \theta_{\mathbf{j}})^2 \quad (4)$$

where  $\tilde{J}(\mathbf{r})$  is a generic function that has to be determined in a self-consistent way, in order to approximate the original Hamiltonian at best. One may wonder whether the asymptotic behavior of  $\tilde{J}(\mathbf{i})$  can be different from that of  $J(r)$ , thus it is convenient to derive the self-consistent equations in the most general setting.

To further proceed, we introduce the partition function of the quadratic model

$$Z_0 = \int \prod_{\mathbf{j}} d\theta_{\mathbf{j}} e^{-\beta H_0}, \quad (5)$$

and the corresponding free energy  $F_0$  defined by  $Z_0 = e^{-\beta F_0}$ . The average on the corresponding Boltzmann measure is defined as

$$\langle \cdot \rangle_0 = \frac{1}{Z_0} \int \prod_{\mathbf{j}} d\theta_{\mathbf{j}} e^{-\beta H_0}. \quad (6)$$

In agreement with the variational principle [3] the best possible result for the couplings  $\tilde{J}$  is obtained by minimizing the quantity

$$\mathcal{F} = \beta F_0 + \beta \langle H \rangle_0 - \beta \langle H_0 \rangle_0 \quad (7)$$

with respect to  $\tilde{J}(\mathbf{r})$ . The equipartition theorem implies that  $\langle H_0 \rangle_0 = \frac{N}{2\beta}$ , so that this term can be ignored. On the other hand:

$$\begin{aligned} \beta \langle H \rangle_0 &= -\frac{1}{2} \sum_{\mathbf{i}, \mathbf{j}} J(|\mathbf{i} - \mathbf{j}|) \langle \cos(\theta_{\mathbf{j}} - \theta_{\mathbf{i}}) \rangle_0 \\ &= -\frac{N}{2} \sum_{\mathbf{r}} J(r) e^{-\frac{1}{2} \langle (\theta_0 - \theta_{\mathbf{r}})^2 \rangle_0} \end{aligned} \quad (8)$$

where we made use of the identity  $\langle \cos A \rangle_0 = e^{-\frac{1}{2} \langle A^2 \rangle_0}$ , valid for every Gaussian measure, and of the translational invariance of the system. Finally, in order to find  $F_0$ , and to compute the correlation functions, we have to diagonalize  $H_0$  by means of the Fourier transform. Defining

$$\theta_{\mathbf{q}} = \frac{1}{\sqrt{N}} \sum_{\mathbf{j}} e^{-i\mathbf{q}\cdot\mathbf{j}} \theta_{\mathbf{j}}; \quad \theta_{\mathbf{j}} = \frac{1}{\sqrt{N}} \sum_{\mathbf{q} \in BZ} e^{i\mathbf{q}\cdot\mathbf{j}} \theta_{\mathbf{q}} \quad (9)$$

(where the sum on the wavevectors  $\mathbf{q}$  is on the first Brillouin zone) we then have

$$\beta H_0 = \frac{1}{2} \sum_{\mathbf{q} \in BZ} K(\mathbf{q}) |\theta_{\mathbf{q}}|^2 \quad (10)$$

where

$$K(\mathbf{q}) = \sum_{\mathbf{r}} \tilde{J}(\mathbf{r}) (1 - \cos(\mathbf{q} \cdot \mathbf{r})) \quad (11)$$

It follows  $Z_0 = (2\pi)^{N/2} \prod_{\mathbf{q} \in BZ} K(\mathbf{q})^{-1/2}$  and

$$\beta F_0 = \frac{1}{2} \sum_{\mathbf{q} \in BZ} \ln K(\mathbf{q}) \quad (12)$$

From Eq. (10) follows that  $\langle \theta_{\mathbf{q}} \theta_{\mathbf{q}'} \rangle_0 = \delta_{\mathbf{q}+\mathbf{q}'} K(\mathbf{q})^{-1}$ . Then

$$\begin{aligned} \langle (\theta_0 - \theta_{\mathbf{r}})^2 \rangle_0 &= \frac{1}{N} \sum_{\mathbf{q}, \mathbf{q}' \in BZ} \langle \theta_{\mathbf{q}} \theta_{\mathbf{q}'} \rangle_0 (1 - e^{i\mathbf{q}\cdot\mathbf{r}})(1 - e^{i\mathbf{q}'\cdot\mathbf{r}}) \\ &= \frac{2}{N} \sum_{\mathbf{q} \in BZ} \frac{1 - \cos(\mathbf{q} \cdot \mathbf{r})}{K(\mathbf{q})} \end{aligned} \quad (13)$$

### 3 Minimization of the free energy

We finally find

$$\mathcal{F} = \frac{1}{2} \sum_{\mathbf{q} \in BZ} \ln K(\mathbf{q}) - \frac{N}{2} \sum_{\mathbf{r}} J(r) e^{-G(\mathbf{r})} \quad (14)$$

with

$$G(\mathbf{r}) = \frac{1}{N} \sum_{\mathbf{q} \in BZ} \frac{1 - \cos(\mathbf{q} \cdot \mathbf{r})}{K(\mathbf{q})}. \quad (15)$$

In the lattice case and at generic distance  $\mathbf{r}$ , the quantity  $G(\mathbf{r})$  also depends on the direction of the vector  $\mathbf{r}$ , but for large distances one may see that it only depends on the modulus  $r$ , as expected.

A close inspection of Eq. (14) reveals that the couplings  $\tilde{J}(\mathbf{r})$  appear in  $\mathcal{F}$  only through the quantities  $K(\mathbf{q})$ . Then, to proceed with the minimization of  $\mathcal{F}$ , it is sufficient to derive with respect to the  $K(\mathbf{q})$ 's, obtaining

$$\frac{\delta \mathcal{F}}{\delta K(\mathbf{q})} = \frac{1}{2K(\mathbf{q})} + \frac{N}{2} \sum_{\mathbf{r}} J(r) \frac{\delta G(\mathbf{r})}{\delta K(\mathbf{q})} e^{-G(\mathbf{r})} \quad (16)$$

and in turn

$$\frac{\delta G(\mathbf{r})}{\delta K(\mathbf{q})} = -\frac{1}{N} \frac{1 - \cos(\mathbf{q} \cdot \mathbf{r})}{K(\mathbf{q})^2}, \quad (17)$$

so that

$$\frac{\delta \mathcal{F}}{\delta K(\mathbf{q})} = \frac{K(\mathbf{q}) - \sum_{\mathbf{r}} J(r) [1 - \cos(\mathbf{q} \cdot \mathbf{r})] e^{-G(\mathbf{r})}}{2K(\mathbf{q})^2}. \quad (18)$$

Exploiting the definition (11) for  $K(\mathbf{q})$  it follows

$$\frac{\delta \mathcal{F}}{\delta K(\mathbf{q})} = \frac{1}{2K(\mathbf{q})^2} \sum_{\mathbf{r}} \mathcal{A}(\mathbf{r}) (1 - \cos(\mathbf{q} \cdot \mathbf{r})), \quad (19)$$

where  $\mathcal{A}(\mathbf{r}) \equiv \tilde{J}(\mathbf{r}) - J(r) e^{-G(\mathbf{r})}$ , Eq. (19) being valid for each value of  $\mathbf{q} \in BZ$  and implying:

$$J(r) = \tilde{J}(\mathbf{r}) e^{G(\mathbf{r})}. \quad (20)$$

Eq. (20) is the desired relation between the couplings  $\tilde{J}(\mathbf{r})$  of the optimizing model and the couplings  $J(r)$  of the initial model. In solving it, one can actually look for a solution such that the couplings  $\tilde{J}$  depend only on  $r$ . Eq. (20) has been derived for the  $2D$  XY model, but the same calculations can be extended for different dimensions. The same structure of Eq. (20) is found for  $O(n)$  models as well. Finally, we notice that Eq. (20) is valid for any non-local form of the couplings  $J(r)$ , such as the exponential one, and, therefore, it is not limited to the power-law decaying form in Eq. (3).

### 4 Ansatz for the variational couplings

We now come back to the specific problem of the  $2D$  XY model with power-law decaying couplings, as defined in Eq. (3):  $J(r) = \frac{J}{r^{2+\sigma}}$ . We remind that we are setting  $J = \beta J_0$  and energy units so that  $J_0 = 1$  (unless differently stated).

Within this choice, when the parameter  $\sigma \rightarrow \infty$ , only nearest-neighbor couplings are present, and the Hamiltonian (1) with coupling  $J_0$  between pairs of nearest-neighbors is retrieved. Moreover, we assume  $\sigma > 0$  so that the additivity of the thermodynamic quantities is preserved [33, 34]. As recalled in the Introduction, in the short-range limit  $\sigma \rightarrow \infty$ , the system undergoes the BKT transition and this phenomenology is qualitatively captured by the SCHA approximation, in which the original Hamiltonian is replaced with a variational quadratic ansatz. The variational coupling jumps discontinuously to zero at a temperature  $T_c$  (see Fig. 1) and provides an estimate for the superfluid stiffness, which can be used in the BKT renormalization group flow equations [24–27].

Our main goal is to discuss the corresponding behavior for finite  $\sigma$ . Without being able to clarify the nature of the universality class and the value of critical exponents, SCHA is nevertheless giving first




Figure 1: Ratio  $\tilde{J}/J$  as a function of  $T = 1/J$  for different values of  $\sigma$ . For each  $\sigma$  we find a jump of  $\tilde{J}$  to zero at a temperature denoted in the text and in Fig. 2 as  $T_c$ .




Figure 2:  $T_c(\sigma)$  at which  $\tilde{J}$  jumps, in units of the  $\sigma \rightarrow \infty$  value  $T_c(\infty) = \frac{4}{e}$ . We see that the temperature is finite for  $\tilde{\sigma} \rightarrow 2$ .




Figure 3: Behavior of  $\tilde{\sigma} - \sigma$  as a function of  $\sigma$  for different values of the temperature  $T$ . It is seen that  $\tilde{\sigma} > \sigma$  everywhere.




Figure 4: Estimate of  $T_{BKT}(\sigma)$  in units of the  $\sigma \rightarrow \infty$  value  $T_{BKT}(\infty)$ .

information whether the BKT phenomenology is stable for large  $\sigma$ , as one would expect, and if one can perform an estimate, or – better – put an upper bound for the value of  $\sigma^*$  (defined in this  $d = 2$   $O(2)$  case such that for  $\sigma > \sigma^*$  one has a BKT transition).

For large  $\sigma$  (i.e. in the short-range limit), the quantity  $\langle \cos(\theta_{\mathbf{j}+\mathbf{r}} - \theta_{\mathbf{j}}) \rangle_0 = e^{-G(\mathbf{r})}$  has a temperature-dependent power law behavior. In order to study such temperature dependence in the power-law case, one needs to extract the large distance properties of the variational couplings  $\tilde{J}(\mathbf{r})$ . Therefore, it is natural to study how a different asymptotic power-law behavior in  $\tilde{J}(\mathbf{r})$  with respect to  $J(r)$  can actually arise from (20), as anticipated. This lead to the ansatz

$$\tilde{J}(\mathbf{r}) \equiv \frac{\tilde{J}}{r^{2+\tilde{\sigma}}} \quad (21)$$

where  $\tilde{\sigma}$  may, in general, be different from  $\sigma$ . Inserting the ansatz (21) in (20), we have to determine  $\tilde{\sigma}$  and  $\tilde{J}$  as a function of  $\sigma$  and  $J$ . We know that in the short-range limit ( $\tilde{\sigma}, \sigma \rightarrow \infty$ )  $\tilde{J}$  as a function of  $T$  has a jump from a finite value to zero [24].

An advantage of the ansatz (21) is to give first information about the robustness of the BKT transition from the knowledge of  $\tilde{\sigma}$ . Indeed, in the short-range nearest-neighbor case, a textbook calculation gives an estimate of  $T_{BKT}$  by calculating the energetic and entropic contributions to the free energy  $\Delta F$  of a free vortex, see e.g. Chapter 4 of [52]. By putting  $N = L^2$  and the lattice spacing  $\equiv 1$ , one has that  $\Delta F = \pi J_0 \ln L - k_B T \ln L^2$ , giving  $k_B T_{BKT} = \pi J_0/2$ . A similar situation of competition between the energetic and entropic contributions occurs for the 1D Ising model with power-law interactions ( $\propto |r|^{-1-\sigma}$ ), where the excitations are magnetization kinks rather than vortices [53]. In our 2D case, one can see that if  $\tilde{\sigma} > 2$  then BKT behavior is expected. This can be understood by observing that for  $q \rightarrow 0$  the scaling of the propagator due to the effective interactions ( $\propto 1/q^{\tilde{\sigma}}$ ) is irrelevant with respect to the one of the free propagator ( $\propto 1/q^2$ ), from which one can conclude the irrelevance of the non-local interaction when  $\tilde{\sigma} > 2$ . We notice that this is in agreement with the known rigorous result of a low-temperature phase with spontaneous symmetry breaking and magnetization for  $XY$  couplings decaying faster than  $1/r^4$  in 2D [54]. In conclusion, if one finds

$$\tilde{\sigma}(\sigma) > 2, \quad (22)$$

then it is possible to conclude that SCHA is indicating persistence of BKT at that  $\sigma$ .

## 5 Results

Let us start from the case  $\sigma > 2$ , deferring our comments on the case  $\sigma < 2$  to the end of the present Section. By plugging the ansatz (21) in Eq. (20), we numerically solve for  $\tilde{J}$  and  $\tilde{\sigma}$  in a square lattice. We considered growing values of  $N$  to solve Eq. (20), and we observed that convergence of the result is obtained for  $N \sim 10^4$ .

In Fig. 1 we show  $\tilde{J}/J$  as a function of the temperature for different values of  $\sigma \geq 2$ . The temperature  $T_c$  corresponding to the jump is  $\sigma$ -dependent (see below for a discussion of the difference between  $T_c$  and  $T_{BKT}$ ). In the  $\sigma \rightarrow \infty$  limit it converges to the short-range value  $k_B T_c(\infty) = \frac{4}{e} J_0$  [24]. The behavior of  $T_c(\sigma)$  in units of  $T_c(\infty)$  is shown in Fig. 2. We found that  $T_c$  is a decreasing function of  $\sigma$  which remains finite as  $\sigma$  approaches 2.

Our main result is that one has  $\tilde{\sigma} > 2$ . This is apparent from Fig 3, where we show the behavior of  $\tilde{\sigma} - \sigma$  as a function of  $\sigma$  for different values of the temperature  $T$ . This quantity is of course only defined as long as  $T < T_c(\sigma)$ . In particular, since  $T_c(\sigma) \rightarrow T_c(\infty)$  as  $\sigma \rightarrow \infty$ , for  $T < T_c(\infty)$  the curve is defined for every  $\sigma$ . As expected,  $\tilde{\sigma} - \sigma$  is temperature-dependent and  $\tilde{\sigma} \rightarrow \sigma$  as  $T \rightarrow 0$ . For  $T < T_c(\infty)$ , as  $\sigma \rightarrow \infty$ ,  $\tilde{\sigma} - \sigma$  goes to a constant, in agreement with the natural expectation of recovering the short-range result. It is important to notice that, for each value of the parameters, we have that  $\tilde{\sigma} > \sigma$  and therefore the validity of the condition (22).

Going to the  $\sigma < 2$  case, since  $T_c$  is finite for  $\sigma \rightarrow 2^+$  and also  $\tilde{\sigma} > \sigma$ , one may be tempted to investigate whether the BKT phase can survive in the  $\sigma < 2$  regime, and ascertain if the value of  $\sigma$  for which the long-range tail of the couplings modifies the critical behavior is actually smaller than 2. However, for  $\sigma < 2$  the solution of Eq. (20) with  $\tilde{J}(\mathbf{r})$  in the form of Eq. (21) is no longer unique. Then, since our ansatz considers only a subspace of the possible functional forms of  $\tilde{J}(r)$  and the difference in the energy of the solutions can be of the same order of the error introduced by the ansatz (21), we cannot safely draw any conclusion for  $\sigma < 2$ . To further explore the  $\sigma < 2$  region one has to resort both to the numerical solution of Eq. (20) without the use of the ansatz (21) and to the application of more sophisticated techniques. It is anyway fair to conclude that 2 is an upper bound for  $\sigma^*$ , i.e. for  $\sigma > 2$  one has a BKT transition at finite temperature.

## 6 Estimate of $T_{BKT}$

$T_c(\sigma)$  can be considered only a very crude estimate of the temperature at which the universal jump occurs. For the short-range limit, a better estimate of  $T_{BKT}$  can be obtained if we define  $T_{BKT}$  as the temperature at which the variational  $\tilde{J}$  reaches the value  $\tilde{J}_{BKT} = \frac{2}{\pi}$  [26]. This value, restoring the temperature, is obtained by considering the Nelson-Kosterlitz condition [16] for the short-range XY model, reading  $J_{superfl}/k_B T = 2/\pi$ , and using the SCHA value  $\tilde{J}_0$  as estimate for the superfluid stiffness  $J_{superfl}$ . In this way, one can obtain  $k_B T_{BKT} = 1.06 J_0$ , improving with respect to the value  $T_c(\sigma \rightarrow \infty)$  (given by  $k_B T_c = (4/e) J_0 = 1.47 J_0$ ). A further improvement could be obtained by using estimates of the dielectric constant in the Nelson-Kosterlitz transition, giving  $k_B T_{BKT} = 0.96 J_0$  [26]. These estimates can be compared with the functional renormalization group estimate,  $k_B T_{BKT} = 0.94 J_0$  [55], with a (functional) renormalization group calculation using a renormalized initial condition,  $k_B T_{BKT} = 0.89 J_0$  [56], with the analytic calculation based on the mapping on the 1D quantum XXZ model,  $k_B T_{BKT} = 0.883 J_0$  [57], and with the previously mentioned Monte Carlo value  $k_B T_{BKT} = 0.893 J_0$ . We notice that the Nelson-Kosterlitz condition can be used to extract  $T_{BKT}$  with high precision from Monte Carlo data [23].

The condition  $\tilde{J}_{BKT} = \frac{2}{\pi}$  can be thought as corresponding to the value at which the Gaussian low-temperature theory becomes unstable due to the presence of the topological excitations [10, 11]. This improves the estimate of  $T_{BKT}$ , since the plain SCHA does not account for the presence of these excitations, i.e. the real mechanism underlying the transition.

Then, we expect that a better estimate of  $T_{BKT}(\sigma)$  can be obtained if we apply the same line of reasoning for finite  $\sigma$ , i.e. using a two-step approach in which SCHA is used to estimate the superfluid stiffness and then the latter is used in the Nelson-Kosterlitz condition. However, with general non-local couplings, and power-law couplings in particular, a microscopic relation between the coupling  $J(r)$  and the superfluid stiffness is not known. One can think to use  $\tilde{J}$  instead of  $J$ , but this would heavily underestimate the effect of the tails of the variational couplings  $\tilde{J}$ .

We propose to proceed defining the analog of the nearest-neighbors coupling in the context of our variational quadratic Hamiltonian (21). We then make use once again of the variational method to determine the best value of  $J_{nn}$  in

$$\beta H_{nn} = \frac{J_{nn}}{2} \sum_{\mathbf{i}, \mathbf{j}} (\theta_{\mathbf{i}} - \theta_{\mathbf{j}})^2 \quad (23)$$

which best approximates  $H_0$ . We find the result:

$$J_{nn}(\tilde{J}, \tilde{\sigma}) = \frac{1}{2N} \sum_{\mathbf{q} \in BZ} \frac{K(\mathbf{q})}{2 - \cos q_x - \cos q_y} \quad (24)$$

where  $K(\mathbf{q})$  is computed from (11) with  $\tilde{J}(r) = \frac{\tilde{J}}{r^{2+\sigma}}$  and  $\tilde{J}, \tilde{\sigma}$  corresponding to the solutions of the variational equation (20).

We can then find our estimate of  $T_{BKT}$  by choosing  $\tilde{J} = \frac{1}{T}$  such that  $J_{nn} = \frac{2}{\pi}$ . The behavior of  $T_{BKT}(\sigma)$  in units of the short-range value  $T_{BKT}(\infty)$  are shown in Fig. 4. The dependence on  $\sigma$  is stronger than those of  $T_c(\sigma)$ , even if the qualitative behavior is the same. In particular,  $T_{BKT}(\sigma)$  remains finite as well as  $\sigma \rightarrow 2^+$ . Whether this suggests or not the scenario in which the BKT transition survives for  $\sigma < 2$  remains open.

## 7 Conclusions

In this paper we studied the self-consistent harmonic approximation (SCHA) for the 2D XY model with non-local couplings. The approach relies on the well known procedure of approximating the interacting model under study with a quadratic model, whose coefficients are optimized minimizing the free energy difference. We derived an equation for the variational couplings which is valid for general non-local interactions and in a form that can be used for other models, such as long-range  $O(n)$  models. Then, we focused on power-law couplings  $J(r)$  decaying as  $J(r) = J/r^{2+\sigma}$ .

The short-range limit  $\sigma \rightarrow \infty$  exhibits the Berezinskii-Kosterlitz-Thouless (BKT) transition. To extract information about the robustness of the BKT transition at finite  $\sigma$  we propose an ansatz for the functional form of the variational couplings  $\tilde{J}(r)$  reading  $\tilde{J}(r) = \tilde{J}/r^{2+\tilde{\sigma}}$ , where  $\tilde{J}$  and  $\tilde{\sigma}$  have to be determined as a function of  $J$  and  $\sigma$ .

The study of this dependence revealed that for  $\sigma > 2$  BKT occurs. The critical temperature at which  $\tilde{J}$  drops to zero has been determined. We then used the output of the SCHA calculation in the Nelson-Kosterlitz condition at the critical point to obtain an improved estimate of the BKT critical temperature.

Our results suggest that, once the value  $\sigma^*$  is defined in such a way that for  $\sigma > \sigma^*$  one has a BKT transition, then one has an upper bound for  $\sigma^*$  given by  $\sigma^* = 2$ . For  $\sigma < 2$  the SCHA has two solutions for the variational parameters. Therefore further work is needed to clarify the structure of the phase diagram and the critical points for  $\sigma < 2$ .

*Acknowledgments:* We thank N. Dupuis, T. Enss, G. Gori and I. Nandori for fruitful discussions. We also thank R. Vaia for useful correspondence. This work is supported by the CNR / HAS (Italy-Hungary) project ‘‘Strongly interacting systems in confined geometries’’ and by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany’s Excellence Strategy ‘‘EXC-2181/1- 390900948’’ (the Heidelberg STRUCTURES Excellence Cluster). This work is part of MUR-PRIN2017 project ‘‘Coarse-grained description for non- equilibrium systems and transport phenomena (CO-NEST)’’ No. 201798CZL whose partial financial support is acknowledged.

- [1] Thomas R. Koehler. ‘‘Theory of the Self-Consistent Harmonic Approximation with Application to Solid Neon’’. In: *Phys. Rev. Lett.* 17 (2 1966), pp. 89–91.
- [2] A. Cuccoli et al. ‘‘The effective potential and effective Hamiltonian in quantum statistical mechanics’’. In: *Journal of Physics: Condensed Matter* 7.41 (1995), pp. 7891–7938.
- [3] R. P. Feynman. *Statistical mechanics*. Reading, W. A. Benjamin, 1972.
- [4] R. Giachetti and V. Tognetti. ‘‘Variational Approach to Quantum Statistical Mechanics of Nonlinear Systems with Application to Sine-Gordon Chains’’. In: *Phys. Rev. Lett.* 55 (9 1985), pp. 912–915.
- [5] R. P. Feynman and H. Kleinert. ‘‘Effective classical partition functions’’. In: *Phys. Rev. A* 34 (6 1986), pp. 5080–5084.
- [6] I. Errea, M. Calandra, and F. Mauri. ‘‘Anharmonic free energies and phonon dispersions from the stochastic self-consistent harmonic approximation: Application to platinum and palladium hydrides’’. In: *Phys. Rev. B* 89 (6 2014), p. 064302.

- [7] A. Cuccoli et al. “Two-dimensional quantum Heisenberg antiferromagnet: Effective-Hamiltonian approach to the thermodynamics”. In: *Phys. Rev. B* 56 (22 1997), pp. 14456–14468.
- [8] E. Šimánek. *Inhomogeneous superconductors: granular and quantum effects*. New York, Oxford University Press, 1994.
- [9] V. L. Berezinskii. “Destruction of long-range order in one-dimensional and two-dimensional systems having a continuous symmetry group i. classical systems”. In: *Sov. Phys. JETP* 34 (1972), p. 610.
- [10] J.M. Kosterlitz and D.J. Thouless. “Ordering, metastability and phase transitions in two-dimensional systems”. In: *J. Phys. C* 6 (1973), pp. 1181–1203.
- [11] J.M. Kosterlitz. “The critical properties of the two-dimensional xy model”. In: *J. Phys. C* 7.6 (1974), p. 1046.
- [12] P. Minnhagen and G. G. Warren. “Superfluid density of a two-dimensional fluid”. In: *Phys. Rev. B* 24 (5 1981), pp. 2526–2532.
- [13] R. Kenna and A.C. Irving. “The Kosterlitz-Thouless universality class”. In: *Nuclear Physics B* 485.3 (1997), pp. 583–612. ISSN: 0550-3213.
- [14] Z. Gulácsi and M. Gulácsi. “Theory of phase transitions in two-dimensional systems”. In: *Adv. Phys.* 47.1 (1998), pp. 1–89. eprint: <http://dx.doi.org/10.1080/000187398243564>.
- [15] J. M. Kosterlitz. “Kosterlitz–Thouless physics: a review of key issues”. In: *Rep. Progr. Phys.* 79.2 (2016), p. 026001.
- [16] David R. Nelson and J. M. Kosterlitz. “Universal Jump in the Superfluid Density of Two-Dimensional Superfluids”. In: *Phys. Rev. Lett.* 39 (19 1977), pp. 1201–1205.
- [17] A. Colcelli et al. “Finite temperature off-diagonal long-range order for interacting bosons”. In: *Phys. Rev. B* 102 (18 2020), p. 184510.
- [18] R. Gupta et al. “Phase Transition in the 2DXY Model”. In: *Phys. Rev. Lett.* 61 (17 1988), pp. 1996–1999.
- [19] R. Gupta and C. F. Baillie. “Critical behavior of the two-dimensional XY model”. In: *Phys. Rev. B* 45 (6 1992), pp. 2883–2898.
- [20] M. Hasenbusch. “The two-dimensionalXYmodel at the transition temperature: a high-precision Monte Carlo study”. In: *Journal of Physics A: Mathematical and General* 38.26 (2005), pp. 5869–5883.
- [21] N. Schultka and E. Manousakis. “Finite-size scaling in two-dimensional superfluids”. In: *Phys. Rev. B* 49 (17 1994), pp. 12071–12077.
- [22] Y. Komura and Y. Okabe. “Large-Scale Monte Carlo Simulation of Two-Dimensional Classical XY Model Using Multiple GPUs”. In: *Journal of the Physical Society of Japan* 81.11 (2012), p. 113001. eprint: <https://doi.org/10.1143/JPSJ.81.113001>.
- [23] Y. Hsieh, Y. Kao, and Anders W Sandvik. “Finite-size scaling method for the Berezinskii–Kosterlitz–Thouless transition”. In: *Journal of Statistical Mechanics: Theory and Experiment* 2013.09 (2013), P09001.
- [24] V. L Pokrovskii and G. V. Uimin. “Magnetic properties of plane and layer systems”. In: *Journal of Experimental and Theoretical Physics* 38.4 (1974), p. 847.
- [25] Yu E Lozovik and S G Akopov. “On phase diagram of granular superconductor”. In: *Journal of Physics C: Solid State Physics* 14.2 (1981), pp. L31–L35.
- [26] A. S. T. Pires and M. E. Gouvêa. “Phase transition and spin dynamics in the two-dimensional easy-plane ferromagnet”. In: *Phys. Rev. B* 48 (17 1993), pp. 12698–12703.
- [27] A. S. T. Pires. “Low-temperature thermodynamic study of the XY model using a self-consistent harmonic approximation”. In: *Phys. Rev. B* 53 (1 1996), pp. 235–239.
- [28] A. Cuccoli et al. “Quantum effects on the BKT phase transition of two-dimensional Josephson arrays”. In: *Phys. Rev. B* 61 (17 2000), pp. 11289–11292.

- [29] A. Smerzi, P. Sodano, and A. Trombettoni. “Finite-temperature renormalization group analysis of interaction effects in 2D lattices of Bose–Einstein condensates”. In: *Journal of Physics B: Atomic, Molecular and Optical Physics* 37.7 (2004), S265–S274.
- [30] P. Martinoli and C. Leemann. “Two Dimensional Josephson Junction Arrays”. In: *J. of Low Temp. Phys.* 118.5 (2000), pp. 699–731. ISSN: 1573-7357.
- [31] R. Fazio and H. van der Zant. “Quantum phase transitions and vortex dynamics in superconducting networks”. In: *Phys. Rep.* 355.4 (2001), pp. 235 –334. ISSN: 0370-1573.
- [32] A. Trombettoni, A. Smerzi, and P. Sodano. “Observable signature of the Berezinskii–Kosterlitz–Thouless transition in a planar lattice of Bose–Einstein condensates”. In: *New Journal of Physics* 7 (2005), pp. 57–57.
- [33] A. Campa, T. Dauxois, and S. Ruffo. “Statistical mechanics and dynamics of solvable models with long-range interactions”. In: *Physics Reports* 480.3 (2009), pp. 57 –159. ISSN: 0370-1573.
- [34] A. Campa et al. *Physics of Long-Range Interacting Systems*. Oxford Univ. Press, 2014.
- [35] N. Defenu et al. “Criticality of spin systems with weak long-range interactions”. In: *Journal of Physics A: Mathematical and Theoretical* 53.14 (2020), p. 143001.
- [36] J. Sak. “Recursion Relations and Fixed Points for Ferromagnets with Long-Range Interactions”. In: *Phys. Rev. B* 8 (1 1973), pp. 281–285.
- [37] E. Luijten and Henk W. J. Blöte. “Classical critical behavior of spin models with long-range interactions”. In: *Phys. Rev. B* 56 (14 1997), pp. 8945–8958.
- [38] E. Luijten and H. W. J. Blöte. “Boundary between Long-Range and Short-Range Critical Behavior in Systems with Algebraic Interactions”. In: *Phys. Rev. Lett.* 89.2 (June 2002), p. 1.
- [39] T. Blanchard, M. Picco, and M.A. Rajabpour. “Influence of long-range interactions on the critical behavior of the Ising model”. In: *EPL (Europhysics Letters)* 101.5 (Mar. 2013), p. 56003.
- [40] E. Brézin, G. Parisi, and F. Ricci-Tersenghi. “The Crossover Region Between Long-Range and Short-Range Interactions for the Critical Exponents”. In: *J. Stat. Phys.* 157.4-5 (Aug. 2014), pp. 855–868.
- [41] M.C. Angelini, G. Parisi, and F. Ricci-Tersenghi. “Relations between short-range and long-range Ising models”. In: *Phys. Rev. E* 89 (6 2014), p. 062120.
- [42] N. Defenu, A. Trombettoni, and A. Codello. “Fixed-point structure and effective fractional dimensionality for O(N) models with long-range interactions”. In: *Phys. Rev. E* 92.5 (2015), pp. 1–11. ISSN: 15502376. arXiv: 1211.3991.
- [43] N. Defenu, A. Trombettoni, and S. Ruffo. “Anisotropic long-range spin systems”. In: *Phys. Rev. B* 94.22 (2016), p. 224411. ISSN: 2469-9950.
- [44] C. Behan et al. “Long-Range Critical Exponents near the Short-Range Crossover”. In: *Phys. Rev. Lett.* 118 (24 2017), p. 241601.
- [45] C. Behan et al. “A scaling theory for the long-range to short-range crossover and an infrared duality”. In: *J. Phys. A: Math. Th.* 50.35, 354002 (Sept. 2017), p. 354002.
- [46] G. Gori et al. “One-dimensional long-range percolation: A numerical study”. In: *Phys. Rev. E* 96.1 (July 2017), p. 012108.
- [47] T. Horita, H. Suwa, and S. Todo. “Upper and lower critical decay exponents of Ising ferromagnets with long-range interaction”. In: *Phys. Rev. E* 95.1 (Jan. 2017), p. 012143.
- [48] A. Bermudez et al. “Long-range Heisenberg models in quasiperiodically driven crystals of trapped ions”. In: *Phys. Rev. B* 95.2 (Jan. 2017), p. 024431.
- [49] M. F. Maghrebi, Z. X. Gong, and A. V. Gorshkov. “Continuous Symmetry Breaking in 1D Long-Range Interacting Quantum Systems”. In: *Phys. Rev. Lett.* 119 (2 2017), p. 023001.
- [50] T. Botzung et al. *Effects of energy extensivity on the quantum phases of long-range interacting systems*. 2019. arXiv: 1909.12105 [cond-mat.str-el].

- [51] F. Cescatti et al. “Analysis of the low-temperature phase in the two-dimensional long-range diluted XY model”. In: *Phys. Rev. B* 100 (5 2019), p. 054203.
- [52] M. Le Bellac. *Quantum and statistical field theory*. Clarendon Press, 1991.
- [53] D. J. Thouless. “Long-Range Order in One-Dimensional Ising Systems”. In: *Phys. Rev.* 187 (2 1969), pp. 732–733.
- [54] H. Kunz and C.-E. Pfister. “First order phase transition in the plane rotator ferromagnetic model in two dimensions”. In: *Comm. Math. Phys.* 46.3 (1976), pp. 245–251.
- [55] N. Defenu et al. “Nonperturbative renormalization group treatment of amplitude fluctuations for  $-\varphi$ -4topological phase transitions”. In: *Phys. Rev. B* 96.17 (Nov. 2017), p. 174505.
- [56] I. Maccari et al. “Interplay of spin waves and vortices in the two-dimensional XY model at small vortex-core energy”. In: *Physical Review B* 102.10 (2020). ISSN: 2469-9969.
- [57] D. C. Mattis. “Transfer matrix in plane-rotator model”. In: *Physics Letters A* 104.6 (1984), pp. 357–360. ISSN: 0375-9601.