



SCUOLA INTERNAZIONALE SUPERIORE DI STUDI AVANZATI

SISSA Digital Library

Quantum Hitchin Systems via β -Deformed Matrix Models

Original

Quantum Hitchin Systems via β -Deformed Matrix Models / Bonelli, G., Maruyoshi, K., Tanzini, A.. - In: COMMUNICATIONS IN MATHEMATICAL PHYSICS. - ISSN 0010-3616. - 358:3(2018), pp. 1041-1064. [10.1007/s00220-017-3053-0]

Availability:

This version is available at: 20.500.11767/86954 since: 2019-02-11T15:31:30Z

Publisher:

Published

DOI:10.1007/s00220-017-3053-0

Terms of use:

Testo definito dall'ateneo relativo alle clausole di concessione d'uso

Publisher copyright
Springer

This version is available for education and non-commercial purposes.

note finali coverpage

(Article begins on next page)

Quantum Hitchin Systems via β -deformed Matrix Models

Giulio Bonelli, Kazunobu Maruyoshi and Alessandro Tanzini

International School of Advanced Studies (SISSA)
via Bonomea 265, 34136 Trieste, Italy and INFN, Sezione di Trieste

Abstract

We study the quantization of Hitchin systems in terms of β -deformations of generalized matrix models related to conformal blocks of Liouville theory on punctured Riemann surfaces. We show that in a suitable limit, corresponding to the Nekrasov-Shatashvili one, the loop equations of the matrix model reproduce the Hamiltonians of the *quantum* Hitchin system on the sphere and the torus with marked points. The eigenvalues of these Hamiltonians are shown to be the ϵ_1 -deformation of the chiral observables of the corresponding $\mathcal{N} = 2$ four dimensional gauge theory. Moreover, we find the exact wave-functions in terms of the matrix model representation of the conformal blocks with degenerate field insertions.

1 Introduction

$\mathcal{N} = 2$ supersymmetric gauge theories in four dimensions display very interesting mathematical structures in their supersymmetrically saturated sectors. These structures allow an exact characterization of several important physical aspects, such as their low energy behavior and stable spectra. These data are encoded in the celebrated Seiberg-Witten solution [1]. It was soon realized that the Seiberg-Witten data can be recovered from integrable systems in terms of their spectral curves [2]. In this context the Hitchin integrable system has emerged as the fundamental geometric structure underlying the M-theory description of $\mathcal{N} = 2$ theories [3, 4, 5].

On the other hand the Seiberg-Witten solution can also be recovered, at least in the case of linear and elliptic quiver $\mathcal{N} = 2$ theories, via equivariant localization on the instanton moduli space [6, 7]. Indeed, this approach contains further information encoded in the expansion in the equivariant parameters of the Ω -background. This opens the issue of relating the full Nekrasov partition function to a suitable quantization of the Hitchin system.

A crucial result in this context is provided by the AGT correspondence [8] relating the Nekrasov partition function to conformal blocks of Liouville/Toda field theories in two dimensions. In [9, 10] it was proposed that this correspondence should be regarded as a two parameter quantization of the Hitchin system itself, or, in field theory language, as its second quantization. Here we will address these issues in the particular limiting case in which one of the two equivariant parameters is vanishing. This was identified by Nekrasov and Shatashvili [11] to provide the first quantization of the integrable system*. In the context of AGT correspondence the instanton partition function in the Nekrasov-Shatashvili limit can be related to the insertion of degenerate fields in the Liouville theory [28, 29], which corresponds to the insertion of surface operators in the gauge theory side [30, 31, 32, 33, 34, 35, 36].

In our approach, we will make use of the matrix model perspective on AGT correspondence developed in [37]. This was further elaborated for Liouville theory on the sphere in [38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53], on the torus in [54, 55] and in [56] at all genera. In this context the equivariant parameters of the Nekrasov partition function are encoded in the β -deformation of the standard Van-der-Monde measure [37]. Notice that for β -deformed matrix models the algebraic equation defining the spectral curve gets deformed into a *differential* equation which can be interpreted as a Schrödinger equation [57, 58, 59]. Our proposal identifies this differential equation, in the Nekrasov-Shatashvili limit, as providing the quantum Hamiltonians of the associated Hitchin integrable system. Moreover the associated

*See also [12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27] for the relation between the gauge theory and the quantized integrable system.

wave-function is described in terms of the β -deformed generalized matrix model corresponding to degenerate field insertions in the Liouville theory [30, 31, 29].

Let us notice also that the quantization of Hitchin systems plays a vital rôle in the context of Langlands duality [60].

The organization of this paper is as follows. In section 2, we review the M-theory perspective on the Hitchin system and some basic facts on its quantization at low genera, namely the sphere and the torus with marked points. We derive the loop equation for the generalized β -deformed matrix model both in the sphere and torus case in sections 3 and 4 respectively and we show that in the Nekrasov-Shatashvili limit these reproduces the Hamiltonians of the quantized Hitchin system. Moreover, we provide a description of the associated wave-functions in terms of the β -deformed generalized matrix model describing degenerate field insertions in the Liouville theory [30, 31, 29]. In section 5 we present some final comments and further directions.

2 Quantum Hitchin systems *in nuce*

In the M-theory framework, the Hitchin system arises by considering the geometry of a system of N M5-branes wrapped on a manifold Y_6 with topology $\mathcal{C} \times \mathbf{R}^4 \times \{pt\}$ in $T^*\mathcal{C} \times \mathbf{R}^4 \times \mathbf{R}^3$ where \mathcal{C} is a Riemann surface. This [37] should be equipped with a non trivial fibration of \mathbf{R}^4 over \mathcal{C} which specifies the Ω -background of Nekrasov [6]. The geometry of the M5-branes bound state is described by an N -fold branched covering of \mathcal{C} given by the algebraic equation

$$x^N = \sum_{j=2}^N \phi_j(z) x^{N-j} \tag{2.1}$$

where x is a section of $T^*\mathcal{C}$ and ϕ_j are $(j, 0)$ -holomorphic differentials on \mathcal{C} whose singularity structure at the punctures identifies the matter content of the gauge theory. The quantum mechanics of this latter structure passes by interpreting (2.1) as the spectral curve of the associated classical system encoding the Seiberg-Witten solution (2.1), and then quantizing via a suitable deformation.

In what follows, we focus on the case with two M5-branes and the corresponding Hitchin systems. In this case, the M-theory curve is specified by the quadratic differential: $x^2 = W_2(z)$. We further focus on the situation where the singularities of the quadratic differential are all of *regular* type meaning the poles of at most degree 2. There is one kind of regular punctures in this two M5-branes case, so the Riemann surface \mathcal{C} is just specified by genus and the number of punctures. Thus, we denote this by $\mathcal{C}_{g,n}$. Under the particular marking of $\mathcal{C}_{g,n}$, the worldvolume low energy theory is the weakly coupled $SU(2)^{n+3g-3}$ superconformal quiver gauge theory [61, 4]. The Hitchin system is associated with this gauge theory.

In order to have explicit parametrization of the Hamiltonians, let's review and specify Hitchin systems and their quantization at low genera, that is sphere and torus with an arbitrary number of punctures, following the approach of [62].

Let us start from some general features of the system. The Higgs field satisfies the condition

$$\bar{\partial}_A \Phi = \sum_k \lambda_k \delta_{w_k}$$

which can be simplified by $\bar{A} = h^{-1} \bar{\partial} h$ and $\Phi = h^{-1} \tilde{\Phi} h$ to

$$\bar{\partial} \tilde{\Phi} = \sum_k \nu_k \delta_{w_k}, \quad (2.2)$$

where $\nu_k = h^{-1}(w_k) \lambda_k h(w_k)$. Eq.(2.2) admits a unique solution iff $\sum_k \nu_k = 0$, given by

$$\tilde{\Phi} = \sum_k \nu_k \omega_{w_k, z^*} + \Phi^0 \quad (2.3)$$

where ω_{PQ} is the unique normalized abelian differential of third kind, i.e. holomorphic on $\Sigma \setminus \{P, Q\}$ with simple poles at P and Q with residues respectively $+1$ and -1 and vanishing A-periods, while $\Phi^0 = \phi_I^0 \omega_I$ is a Lie algebra valued holomorphic differential and ω_I is a basis of normalized holomorphic differentials on Σ .

As explained in [62], the Poisson brackets are induced by the Lie algebra of the complexified gauge group. At every puncture the residues are expanded as $\nu_k = \sum_a \nu_k^a t^a$, where t^a is a basis of the Lie algebra and the Poisson brackets are $\{\nu_k^a, \nu_l^b\}_{PB} = i \delta_{kl} f^{abc} \nu_k^c$.

The Hamiltonians are the Chern polynomials of the Higgs field, namely the coefficients of the expansion of the spectral curve (2.1) as

$$\det(\Phi - x \cdot \mathbf{1}) = 0.$$

The quantization of the integrable system, as proposed in [62], is induced by the quantization of the Poisson brackets above.

The case of our interest is a projection to the Cartan degrees of freedom of the general integrable system specified to $SL(2, \mathbf{C})$.

In the sphere case, there are no holomorphic differentials and the Higgs field reads (see also [10])

$$\tilde{\Phi} = \sum_k \frac{\nu_k}{z - w_k} \frac{dz}{2\pi i}$$

where ν_k is the only Cartan element. The corresponding relevant Hamiltonians are generated by

$$\text{Tr} \Phi^2 = \sum_k \left(\frac{\mathbf{J}_k^2}{(z - w_k)^2} + \frac{\mathbf{H}_k^{(0)}}{z - w_k} \right) \left(\frac{dz}{2\pi i} \right)^2 \quad (2.4)$$

where

$$\mathbf{J}_k^2 = \text{Tr } \nu_k^2 \quad \mathbf{H}_k^{(0)} = 2 \sum_{l \neq k} \frac{1}{w_k - w_l} \text{Tr } \nu_l \nu_k \quad (2.5)$$

According to the general discussion above, the quantization of these operators is provided by replacing ν_k at each puncture with the corresponding spin operators.

Analogously, in the torus case, the Higgs field is

$$\tilde{\Phi} = \left(\sum_k \nu_k \frac{\vartheta_1'(z - w_k)}{\vartheta_1(z - w_k)} + 2\pi i p \right) \frac{dz}{2\pi i} \quad (2.6)$$

from which it follows that

$$\text{Tr } \Phi^2 = \sum_k \left(\mathcal{P}(z - w_k) \mathbf{J}_k^2 + \frac{\vartheta_1'(z - w_k)}{\vartheta_1(z - w_k)} \mathbf{H}_k^{(1)} + \mathbf{H}_0^{(1)} \right) \left(\frac{dz}{2\pi i} \right)^2, \quad (2.7)$$

where[†]

$$\begin{aligned} \mathbf{H}_k^{(1)} &= 2 \sum_{l \neq k} \text{Tr } \nu_k \nu_l \frac{\vartheta_1'(w_k - w_l)}{\vartheta_1(w_k - w_l)} + 4\pi i \text{Tr } \nu_k p, \\ \mathbf{H}_0^{(1)} &= -4\pi^2 \text{Tr } p^2 - \eta_1 \sum_k \mathbf{J}_k^2 + \frac{1}{2} \sum_{k,l;k \neq l} \text{Tr } \nu_k \nu_l \frac{\vartheta_1''(w_k - w_l)}{\vartheta_1(w_k - w_l)}. \end{aligned} \quad (2.8)$$

See Appendix A for the definition of the theta functions. In (2.7), \mathcal{P} is the Weierstrass \mathcal{P} -function. To obtain (2.7) we used the identity (A.8) relating the Weierstrass \mathcal{P} -function and its primitive $\zeta'(z) = -\mathcal{P}(z)$.

In what follows, we will see the appearing of the above Hitchin Hamiltonians via the generalized beta-deformed matrix model.

3 Matrix model: genus zero

The AGT relation associated with a sphere is the one between the Nekrasov partition function of $\mathcal{N} = 2$ superconformal $SU(2)^{n-3}$ linear quiver gauge theory and the n -point conformal block on the sphere. Both of them are specified by the marking of $\mathcal{C}_{0,n}$. We can obtain the beta-deformed matrix model starting from the Dotsenko-Fateev integral representation of the conformal block [63] as follows. (See [37, 47]). In terms of the free field $\phi(z)$ (whose OPE is

[†]Notice that with respect to (4.10) in [62], in $\mathbf{H}_0^{(1)}$ we find also a term proportional to η_1 , which will reveal to be crucial in comparing with the quantum Seiberg-Witten curve. This term was invisible to the authors of [62], being the absolute normalization of the differential which they admittedly do not check.

$\phi(z)\phi(\omega) \sim -\frac{1}{2} \log(z - \omega)$), the n -point conformal block is described by inserting the screening operators

$$Z^{\mathcal{C}_{0,n}} = \left\langle \left(\int d\lambda_I : e^{2b\phi(\lambda_I)} : \right)^N \prod_{k=0}^{n-1} V_{m_k}(w_k) \right\rangle_{\text{free on } \mathcal{C}_0}, \quad (3.1)$$

where the vertex operator $V_{m_k}(w_k)$ is given by $: e^{2m_k\phi(w_k)} :$. The momentum conservation condition relates the external momenta and the number of integrals as $\sum_{k=0}^{n-1} m_k = bN - Q$. By evaluating the OPEs, it is easy to obtain

$$Z^{\mathcal{C}_{0,n}} = C(m_k, w_k) \tilde{Z}^{\mathcal{C}_{0,n}} \equiv e^{F^{\mathcal{C}_{0,n}}/g_s^2}, \quad (3.2)$$

where $\tilde{Z}^{\mathcal{C}_{0,n}}$ is the beta-deformation of one matrix model

$$\tilde{Z}^{\mathcal{C}_{0,n}} = \int \prod_{I=1}^N d\lambda_I \prod_{I < J} (\lambda_I - \lambda_J)^{-2b^2} e^{\frac{b}{g_s} \sum_I W(\lambda_I)} \equiv e^{\tilde{F}^{\mathcal{C}_{0,n}}/g_s^2}, \quad (3.3)$$

and

$$W(z) = \sum_{k=0}^{n-2} 2m_k \log(z - w_k), \quad C(m_k, w_k) = \prod_{k < \ell \leq n-2} (w_k - w_\ell)^{-\frac{2m_k m_\ell}{g_s^2}}. \quad (3.4)$$

We have introduced the parameter g_s by rescaling $m_k \rightarrow \frac{m_k}{g_s}$. We relate m_k with the mass parameters of the gauge theory. Also, we have chosen three insertion points as $w_0 = 0$, $w_1 = 1$ and $w_{n-1} = \infty$. The remaining parameters are identified with the gauge theory coupling constants $q_i = e^{2\pi i \tau_i}$ ($i = 1, \dots, n-3$) as follows:

$$w_2 = q_1, \quad w_3 = q_1 q_2, \quad \dots, \quad w_{n-2} = q_1 q_2 \dots q_{n-3}. \quad (3.5)$$

While the dependence on m_{n-1} disappeared in the potential, this is recovered by the momentum conservation condition

$$\sum_{k=0}^{n-2} m_k + m_{n-1} = b g_s N - g_s Q. \quad (3.6)$$

We will refer to F_m as free energy.

The identification of the parameter b with the Nekrasov's deformation parameters is given by

$$\epsilon_1 = b g_s, \quad \epsilon_2 = \frac{g_s}{b}. \quad (3.7)$$

Note that, in the case of $b = i$ ($c = 1$), this reduces to the usual hermitian matrix model and this case corresponds to the self-dual background $\epsilon_1 = -\epsilon_2$.

Here we define the resolvent of the matrix model as

$$R(z_1, \dots, z_k) = (b g_s)^k \sum_{I_1} \frac{1}{z_1 - \lambda_{I_1}} \dots \sum_{I_k} \frac{1}{z_k - \lambda_{I_k}}. \quad (3.8)$$

For $k = 1$, this reduces to the usual resolvent.

3.1 Wave-function and conformal block

In the following, we mainly concentrate on the limit where $\epsilon_2 \rightarrow 0$ with ϵ_1 and the other parameters keeping fixed. In other words, the limit is $b \rightarrow \infty$ and $g_s \rightarrow 0$ with bg_s and N keeping finite. This is the limit by Nekrasov and Shatashvili [11]. In [10, 29], it was shown that the conformal blocks on a sphere with the additional insertion of the degenerate fields $V_{\frac{1}{2b}}(z) = e^{-\frac{\phi(z)}{b}}$ capture the quantization of the integrable systems.

In this limit, the beta-deformed partition function can be written as $\int \prod_{I=1}^N d\lambda_I \exp(-\frac{1}{\epsilon_2} \widetilde{W})$ where

$$\widetilde{W} = \sum_I W(\lambda_I) + 2\epsilon_1 \sum_{I < J} \log(\lambda_I - \lambda_J). \quad (3.9)$$

Thus the leading order part of the free energy can be obtained from the value of the critical points which solve the equations of motion:

$$W'(\lambda_I) + 2\epsilon_1 \sum_{J \neq I} \frac{1}{\lambda_I - \lambda_J} = 0. \quad (3.10)$$

We note that these two terms are of the same order in the limit because N and ϵ_1 are kept finite.

In this section, we will show that under the identification the beta-deformed matrix model $Z^{C_{0,n}}$ with the n -point conformal block, the integral representation of the degenerate conformal block can be written in terms of the resolvent of the original matrix model (3.8), in the $\epsilon_2 \rightarrow 0$ limit. More explicitly, we will show

$$\frac{Z_{deg}^{C_{0,n+n}}}{Z^{C_{0,n}}} (z_1, z_2, \dots, z_n) = \prod_{i=1}^n \Psi_i(z_i), \quad \Psi_i(z_i) = \exp \left(\frac{1}{\epsilon_1} \int^{z_i} x(z') dz' \right), \quad (3.11)$$

in the $\epsilon_2 \rightarrow 0$ limit, where $Z_{deg}^{C_{0,n+n}}$ is the matrix model partition function corresponding to the $2n$ -point conformal block with n degenerate fields inserted at $z = z_i$. This property of “separation of variables” agrees with the corresponding result of the Virasoro conformal block as in [32, 10]. The differential $x(z)dz$ is identified with the “quantized” Seiberg-Witten differential which is given, in terms of matrix model language, by

$$x(z) = \langle R(z) \rangle - \frac{W'(z)}{2}, \quad (3.12)$$

where $\langle \dots \rangle = \frac{1}{Z^{C_{0,n}}} \int \prod d\lambda \prod (\lambda_I - \lambda_J)^{-2b^2} e^{b \sum W/g_s} \dots$. This relation (3.11) is a simple generalization of the one obtained in [20] for the single degenerate field insertion.

First of all, we consider a more generic expression: the $(n + \ell)$ -point conformal block where ℓ degenerate fields are inserted

$$\begin{aligned}
Z_{deg}^{\mathcal{C}_{0,n+\ell}} &= \left\langle \prod_{i=1}^{\ell} V_{\frac{1}{2b}}(z_i) \left(\int d\lambda e^{2b\phi(\lambda)} \right)^N \prod_{k=0}^{n-1} V_{\frac{m_k}{g_s}}(w_k) \right\rangle \\
&= \prod_{i<j} (z_i - z_j)^{-\frac{1}{2b^2}} \prod_{0 \leq k < \ell \leq n-2} (w_k - w_\ell)^{-\frac{2m_k m_\ell}{g_s^2}} \prod_{i=1}^{\ell} \prod_{k=0}^{n-2} (z_i - w_k)^{-\frac{m_k}{bg_s}} \\
&\quad \times \int \prod_{I=1}^N d\lambda_I \prod_{I<J} (\lambda_I - \lambda_J)^{-2b^2} \prod_I \prod_{k=0}^{n-2} (\lambda_I - w_k)^{\frac{2bm_k}{g_s}} \prod_{i=1}^{\ell} (z_i - \lambda_I), \quad (3.13)
\end{aligned}$$

where the potential $W(z)$ is the same as (3.4). We have taken w_{n-1} to infinity and omitted the factor including this, as we have done above. The momentum conservation is however modified by the degenerate field insertion as

$$\sum_{k=0}^{n-1} m_k + \frac{\ell g_s}{2b} = bg_s N - g_s Q. \quad (3.14)$$

By dividing by $Z^{\mathcal{C}_{0,n}}$ and taking a log, we obtain

$$\log \frac{Z_{deg}^{\mathcal{C}_{0,n+\ell}}}{Z^{\mathcal{C}_{0,n}}} = -\frac{1}{2b^2} \sum_{i<j} \log(z_i - z_j) - \sum_i \frac{W(z_i)}{2bg_s} + \log \left\langle \prod_{i,I} (z_i - \lambda_I) \right\rangle. \quad (3.15)$$

Notice that the expectation value is defined as above, but with the modified momentum conservation (3.14). By defining $e^L = \prod_{i,I} (z_i - \lambda_I)$, we notice that

$$L = \sum_{i,I} \log(z_i - \lambda_I) = \sum_{i,I} \int^{z_i} \frac{dz'_i}{z'_i - \lambda_I}, \quad (3.16)$$

where we have ignored irrelevant terms due to the end points of the integrations. Then, we use that the expectation value of e^L can be written as $\log \langle e^L \rangle = \sum_{k=1}^{\infty} \frac{1}{k!} \langle L^k \rangle_{conn}$ [20], where $\langle \dots \rangle_{conn}$ means the connected part of the correlator, $\langle L^2 \rangle_{conn} = \langle L^2 \rangle - \langle L \rangle^2$, etc. Thus, (3.15) can be expressed, by using (3.16), as

$$\log \frac{Z_{deg}^{\mathcal{C}_{0,n+\ell}}}{Z^{\mathcal{C}_{0,n}}} = -\sum_i \frac{W(z_i)}{2\epsilon_1} + \sum_{k=1}^{\infty} \frac{1}{k!} \left\langle \left(\sum_{i,I} \int^{z_i} \frac{dz'_i}{z'_i - \lambda_I} \right)^k \right\rangle_{conn}. \quad (3.17)$$

We will now consider the limit where $\epsilon_2 \rightarrow 0$. In this limit, the terms with $k > 1$ are subleading contributions compared with the $k = 1$ terms since the connected part of the expectation value can be ignored in this limit. Thus, we obtain

$$\log \frac{Z_{deg}^{\mathcal{C}_{0,n+\ell}}}{Z^{\mathcal{C}_{0,n}}} \rightarrow \frac{1}{\epsilon_1} \sum_i \int^{z_i} x(z') dz', \quad (3.18)$$

where we have used (3.12). Thus, by setting $\ell = n$, we have obtained (3.11). This indicates that the properties of the conformal block with degenerate field insertions are build in the resolvent of the matrix model in the $\epsilon_2 \rightarrow 0$ limit.

3.2 Loop equations

The argument in the previous section shows that the relation with the integrable system can be seen by analyzing the resolvent, in particular, the loop equations. Thus, we derive it here with finite β . First of all, we keep the potential arbitrary and obtain

$$\begin{aligned} 0 &= \frac{1}{\tilde{Z}^{c_{0,n}}} \int \prod_{I=1}^N d\lambda_I \sum_K \frac{\partial}{\partial \lambda_K} \left[\frac{1}{z - \lambda_K} \prod_{I < J} (\lambda_I - \lambda_J)^{-2b^2} e^{\frac{b}{g_s} \sum_I W(\lambda_I)} \right] \\ &= -\frac{1}{g_s^2} \langle R(z, z) \rangle - \frac{b + \frac{1}{b}}{g_s} \langle R(z)' \rangle + \frac{1}{g_s^2} W'(z) \langle R(z) \rangle - \frac{f(z)}{g_s^2}, \end{aligned} \quad (3.19)$$

where R' is the z -derivative of the resolvent and we have defined

$$f(z) = bg_s \left\langle \sum_I \frac{W'(z) - W'(\lambda_I)}{z - \lambda_I} \right\rangle. \quad (3.20)$$

The expectation value is defined as the matrix model average. By multiplying (3.19) by g_s^2 , we obtain

$$0 = -\langle R(z, z) \rangle - (\epsilon_1 + \epsilon_2) \langle R(z)' \rangle + \langle R(z) \rangle W'(z) - f(z). \quad (3.21)$$

In the case of the hermitian matrix model $b = i$, the second term vanishes and the equation reduces to the well-known one.

Let us then analyze $f(z)$ by specifying the potential to the Penner type one (3.4). In this case,

$$f(z) = \sum_{k=0}^{n-2} \frac{c_k}{z - w_k}, \quad (3.22)$$

where for $k \geq 2$

$$c_k = -bg_s \left\langle \sum_I \frac{2m_k}{\lambda_I - w_k} \right\rangle = g_s^2 \frac{\partial \log \tilde{Z}^{c_{0,n}}}{\partial w_k} = \frac{\partial \tilde{F}_m}{\partial w_k}. \quad (3.23)$$

While we cannot write c_0 and c_1 as above because we have chosen $w_0 = 0$ and $w_1 = 1$, we can see that they are written in terms of c_k with $k \geq 2$. First of all, due to the equations of motion: $\langle \sum_I W'(\lambda_I) \rangle = 0$, the sum of c_k is constrained to vanish $\sum_{k=0}^{n-2} c_k = 0$. In order to find another

constraint, we consider the asymptotic at large z of the loop equation. The asymptotic of the resolvent is $\langle R(z) \rangle \sim \frac{bg_s N}{z}$, so that the leading terms at large z in the loop equations satisfy

$$-(bg_s N)^2 + (\epsilon_1 + \epsilon_2)bg_s N + bg_s N \sum_{k=0}^{n-2} 2m_k - \sum_{k=0}^{n-2} w_k c_k = 0. \quad (3.24)$$

The leading term of order $1/z$ in $f(z)$ vanishes via the first constraint. Thus, we obtain

$$\sum_{k=0}^{n-2} w_k c_k = \left(\sum_{k=0}^{n-2} m_k + m_{n-1} + \frac{ng_s}{2b} + g_s Q \right) \left(\sum_{k=0}^{n-2} m_k - m_{n-1} + \frac{ng_s}{2b} \right) \equiv M^2, \quad (3.25)$$

where we have used the momentum conservation[‡] (3.14) with $\ell = n$. Therefore, c_0 and c_1 can be written in terms of c_k (3.23). These constraints are related to the Virasoro constraints [64].

$\epsilon_2 \rightarrow 0$ limit

As above, in the $\epsilon_2 \rightarrow 0$ limit, the connected part of (3.8) can be ignored in this limit: $\langle R(z, z) \rangle \rightarrow \langle R(z) \rangle^2$. Taking this into account, the loop equation (3.21) becomes

$$0 = -\langle \tilde{R}(z) \rangle^2 - \epsilon_1 \langle \tilde{R}(z)' \rangle + \langle \tilde{R}(z) \rangle W'(z) - \tilde{f}(z), \quad (3.26)$$

where \tilde{R} and \tilde{f} are $R|_{\epsilon_2 \rightarrow 0}$ and $f|_{\epsilon_2 \rightarrow 0}$ respectively. In the following, we will omit the tildes of R and f . Then, in terms of $x = \langle R(z) \rangle - \frac{W'(z)}{2}$, the equation becomes

$$0 = -x^2 - \epsilon_1 x' + U(z), \quad (3.27)$$

where

$$U(z) = \frac{W'(z)^2}{4} - \frac{\epsilon_1}{2} W''(z) - f(z). \quad (3.28)$$

This equation is similar to the one obtained in [57]. It is easy to see that this can be written as the Schrödinger-type equation:

$$0 = -\epsilon_1^2 \frac{\partial^2}{\partial z^2} \Psi(z) + U(z) \Psi(z), \quad (3.29)$$

where $\Psi(z)$ is defined in (3.11).

The above argument is applicable for an arbitrary potential $W(z)$. Here we return to the Penner-type one (3.4) and see the relation with the Gaudin Hamiltonian[§]. (3.28) becomes in this case

$$U(z) = \sum_{k=0}^{n-2} \frac{m_k(m_k + \epsilon_1)}{(z - w_k)^2} + \sum_k \frac{H_k}{z - w_k} - \sum_{k=0}^{n-2} \frac{c_k}{z - w_k} \quad (3.30)$$

[‡]We are using the modified momentum conservation to apply this to the argument in the previous section. However, the difference will disappear in the $\epsilon_2 \rightarrow 0$ limit.

[§]A relation with Gaudin system at finite ϵ_2 has been noticed also in [57]

where

$$H_k = \sum_{\ell(\neq k)} \frac{2m_k m_\ell}{w_k - w_\ell}. \quad (3.31)$$

$U(z)$ is the vacuum expectation value of (2.4). In particular, notice that the residue of the quadratic pole in (3.30) corresponds to the eigenvalue of \mathbf{J}_k^2 quantized in ϵ_1 units and that $H_k - c_k$ are the vacuum energies of the quantum Hamiltonians $\mathbf{H}_k^{(0)}$ (2.5).

Let us rewrite c_k in terms of the gauge theory variables. Since the moduli of the sphere are related with the gauge coupling constants of the linear quiver gauge theory as in (3.5), the derivatives with respect to w_k can be written as $2\pi i w_k \frac{\partial}{\partial w_k} = \frac{\partial}{\partial \tau_{k-1}} - \frac{\partial}{\partial \tau_k}$, where the second term vanished when $k = n - 2$. Therefore, for $k = 2, \dots, n - 2$, by using (3.23), we obtain

$$c_k = \frac{1}{2\pi i w_k} \left(\frac{\partial \tilde{F}^{C_{0,n}}}{\partial \tau_{k-1}} - \frac{\partial \tilde{F}^{C_{0,n}}}{\partial \tau_k} \right) = \frac{1}{2\pi i w_k} (u_{k-1} - u_k). \quad (3.32)$$

where u_k are closely related with the gauge theory variables $\langle \text{tr} \phi_k^2 \rangle$, ϕ_k being the vector multiplet scalar of the k -th gauge group. Indeed, supposing that the free energy $F^{C_{0,n}}$ is identified with the prepotential (with ϵ_1 and ϵ_2) of the gauge theory (this has indeed been checked for $n = 4$ in [55, 44, 45, 46] in some orders in the moduli), we can use the ϵ -deformed version [65, 66, 9] of the Matone relation [67, 68, 69] to relate u_k with $\langle \text{tr} \phi_k^2 \rangle$. Note that there is still difference between $\tilde{F}^{C_{0,n}}$ and $F^{C_{0,n}}$, we will explicitly consider this in an example below. Instead, for c_0 and c_1 , we can use the two constraints derived above and obtain

$$\begin{aligned} c_0 &= \sum_{k=2}^{n-2} (w_k - 1)c_k - M^2 = \frac{1}{2\pi i} \sum_{k=2}^{n-2} \frac{w_k - 1}{w_k} (u_{k-1} - u_k) - M^2, \\ c_1 &= - \sum_{k=2}^{n-2} w_k c_k + M^2 = - \frac{1}{2\pi i} \sum_{k=2}^{n-2} (u_{k-1} - u_k) + M^2. \end{aligned} \quad (3.33)$$

We have obtained the differential equations for $\Psi = \prod_i \Psi_i(z_i)$ each of which is (3.29) satisfied by $\Psi_i(z_i)$. These are similar to the differential equations satisfied by the $(n+n)$ -point Virasoro conformal block where n vertex operators are chosen to be degenerate $V(z) = e^{-\frac{\phi(z)}{b}}$, as in [10]. As an example, we will explicitly see in the subsequent section the differential equation for $n = 4$ is the same as that obtained from the Virasoro conformal block.

3.3 Sphere with four punctures

We now consider the case corresponding to a sphere with four puncture where the matrix model potential is given by

$$W(z) = \sum_{k=0}^2 2m_k \log(z - w_k), \quad w_0 = 0, \quad w_1 = 1, \quad w_2 = q. \quad (3.34)$$

This corresponds to $SU(2)$ gauge theory with four flavors and the Gaudin model on a sphere with four punctures where the number of commuting Hamiltonian is just one. Thus we simply consider the wave-function $\Psi(z) = e^{\frac{1}{\epsilon_1} \int^z x dz'}$ and the differential equation satisfied by it.

In this case, $U(z)$ can be evaluated in terms of c_2 as

$$U(z) = \sum_{k=0}^2 \frac{m_k^2 + \epsilon_1 m_k}{(z - w_k)^2} + \frac{m_3^2 - \sum_{i=0}^2 m_i^2 + \epsilon_1 (m_3 - \sum_{k=0}^2 m_k)}{z(z-1)} + \frac{-q(q-1)c_2 + 2(qm_1m_2 + (q-1)m_2m_0)}{z(z-1)(z-q)}, \quad (3.35)$$

Let us relate this with the one obtained from the Virasoro conformal block. We consider the last line of $U(z)$. Let us recall the definition of the free energy (3.2) and (3.3). Since the difference of them is expressed by $C = q^{-\frac{2m_2m_0}{g_s^2}} (1-q)^{-\frac{m_1m_2}{g_s^2}}$ in this case, the free energies are related by

$$F^{\mathcal{C}_{0,4}} = \tilde{F}^{\mathcal{C}_{0,4}} - 2m_1m_2 \log(1-q) - 2m_2m_0 \log q. \quad (3.36)$$

Therefore, its derivative is

$$q(1-q) \frac{\partial F^{\mathcal{C}_{0,4}}}{\partial q} = q(1-q) \frac{\partial \tilde{F}^{\mathcal{C}_{0,4}}}{\partial q} + 2qm_1m_2 - 2(1-q)m_2m_0. \quad (3.37)$$

We notice that the right hand side is the numerator of the last line of $U(z)$.

Here let us redefine the mass parameters as

$$\tilde{m}_0 = m_0 + \frac{\epsilon_1}{2}, \quad \tilde{m}_3 = m_3 - \frac{\epsilon_1}{2}. \quad (3.38)$$

In this notation, U can be written as

$$U(z) = \frac{\tilde{m}_0^2 - \frac{\epsilon_1^2}{4}}{z^2} + \frac{m_1(m_1 + \epsilon_1)}{(z-1)^2} + \frac{m_2(m_2 + \epsilon_1)}{(z-q)^2} - \frac{-\tilde{m}_3^2 + \tilde{m}_0^2 + m_1(m_1 + \epsilon_1) + m_2(m_2 + \epsilon_1)}{z(z-1)} + \frac{q(1-q)}{z(z-1)(z-q)} \frac{\partial F}{\partial q}. \quad (3.39)$$

Note here that the first four terms are exactly the potential which considered in [29] $V(z)$. Also, the last term might correspond to the ‘‘eigenvalue’’ in [29]. The Schrödinger-like equation becomes

$$-\epsilon_1^2 \frac{\partial^2}{\partial z^2} \Psi(z) + V(z) \Psi(z) = -\frac{q(1-q)}{z(z-1)(z-q)} \frac{\partial F}{\partial q} \Psi(z). \quad (3.40)$$

Note that this differential equation has also been derived in [20] from the free field expression, the first line of (3.13).

4 Matrix model: genus one

In this section, we consider the matrix model corresponding to the conformal block on a torus with punctures [37, 54, 55, 56]. We will derive the loop equations of the matrix model and relate it with the differential equations of the corresponding Hitchin system.

We consider the n -point conformal block on a torus whose integral description is

$$\begin{aligned} Z^{\mathcal{C}_{1,n}} &= e^{\tilde{F}^{\mathcal{C}_{1,n}}/g_s^2} = \int \prod_I d\lambda_I \prod_{I<J} \left[\frac{\vartheta_1(\lambda_I - \lambda_J)}{\eta(\tau)^3} \right]^{-2b^2} \prod_I \prod_{k=1}^n \left[\frac{\vartheta_1(\lambda_I - w_k)}{\eta(\tau)^3} \right]^{\frac{2bm_k}{g_s}} e^{\frac{4\pi bp}{g_s} \sum_I \lambda_I} \\ &\quad \times \prod_{k<\ell} \left[\frac{\vartheta_1(w_k - w_\ell)}{\eta(\tau)^3} \right]^{-\frac{2m_k m_\ell}{g_s^2}} e^{-\frac{4\pi p \sum_k m_k w_k}{g_s^2}} \\ &= C(w_k, m_k, p) \tilde{Z}^{\mathcal{C}_{1,n}}, \end{aligned} \quad (4.1)$$

where $C(w_k, m_k, p)$ is the λ independent coefficient

$$C(w_k, m_k, p) = \eta^{-3(-b^2 N(N-1) + \frac{2bN}{g_s} \sum_k m_k + \frac{2}{g_s^2} \sum_{k<\ell} m_k m_\ell)} \prod_{k<\ell} \vartheta_1(w_k - w_\ell)^{-\frac{2m_k m_\ell}{g_s^2}} e^{-\frac{4\pi p \sum_k m_k w_k}{g_s^2}} \quad (4.2)$$

and thus,

$$\tilde{Z}^{\mathcal{C}_{1,n}} = e^{\tilde{F}^{\mathcal{C}_{1,n}}/g_s^2} = \int \prod_I d\lambda_I \prod_{I<J} \vartheta_1(\lambda_I - \lambda_J)^{-2b^2} e^{\frac{b}{g_s} \sum_I W(\lambda_I)} \quad (4.3)$$

with the potential

$$W(z) = \sum_{k=1}^n 2m_k \log \vartheta_1(z - w_k) + 4\pi p z. \quad (4.4)$$

While we do not know the representation of this in terms of a matrix, we refer to this as (generalized) matrix model. As the matrix model in the case of the sphere, this expression has been obtained [54, 56] from the free field expression, similar to (3.1) but on the torus, following from the Liouville correlator by the method in [71]. The parameters must satisfy the momentum conservation

$$\sum_{k=1}^n m_k = b g_s N. \quad (4.5)$$

See Appendix A for the definition of the elliptic theta functions.

This matrix model is related with the $\mathcal{N} = 2$ elliptic $SU(2)$ quiver gauge theory [61] which is obtained from two M5-branes on $\mathcal{C}_{1,n}$ with specifying its marking. The gauge theory coupling constants $q_i = e^{2\pi i \tau_i}$ ($i = 1, \dots, n$) are identified with the moduli of the torus as

$$e^{2\pi i w_k} = \prod_{i=k}^{n-1} q_i, \quad q \equiv e^{2\pi i \tau} = \prod_{i=1}^n q_i. \quad (4.6)$$

The parameters m_k are directly identified with the mass parameters of the bifundamentals. The remaining parameters of the matrix model, the filling fractions and momentum p in the potential determine the Coulomb moduli of the gauge theory.

4.1 Wave-function and conformal block

As in the previous section, we can show the relation between the resolvent of this matrix model and the conformal block with the degenerate fields $\Phi_{1,2} = e^{-\frac{1}{b}\phi(z)}$. The integral representation of the latter is

$$Z_{deg}^{\mathcal{C}_{1,n+\ell}}(z_i) = C(w_k, m_k, p) \int \prod_I d\lambda_I \prod_{I < J} \vartheta_1(\lambda_I - \lambda_J)^{-2b^2} \prod_I \vartheta_1(\lambda_I - w_1)^{\frac{2bm_1}{g_s}} e^{\frac{4\pi bp}{g_s} \sum_I \lambda_I} \quad (4.7)$$

$$\times \prod_{i=1}^{\ell} \left(\prod_k \left[\frac{\vartheta_1(z_i - w_k)}{\eta(\tau)^3} \right]^{-\frac{m_k}{bg_s}} \prod_I \left[\frac{\vartheta_1(z_i - \lambda_I)}{\eta(\tau)^3} \right] e^{-\frac{2\pi p}{bg_s} z_i} \right) \prod_{i < j} \left[\frac{\vartheta_1(z_i - z_j)}{\eta(\tau)^3} \right]^{-\frac{1}{2b^2}},$$

where z_i are the insertion points of the degenerate fields and the momentum conservation is slightly deformed from the original one to

$$\sum_{k=1}^n m_k + \frac{\ell g_s}{2b} = bg_s N. \quad (4.8)$$

As before, we consider $\log \frac{Z_{deg}^{\mathcal{C}_{1,n+\ell}}}{Z^{\mathcal{C}_{1,n}}}$. The main object we want to know is $\log \left\langle \prod_{I,i} \vartheta_1(z_i - \lambda_I) \right\rangle$, where the expectation value is defined in the same way as in the sphere case. Note that we are using the momentum conservation (4.8). Then, if we define $e^L = \prod_{I,i} \vartheta_1(z_i - \lambda_I)$, we can rewrite it as $L = \sum_{I,i} \int^{z_i} \frac{\vartheta_1'(z'_i - \lambda_I)}{\vartheta_1(z'_i - \lambda_I)} dz'_i$. By using the same argument as the sphere case: $\log \langle e^L \rangle = \sum_k \frac{1}{k!} \langle L^k \rangle_{conn}$, we therefore obtain

$$\log \frac{Z_{deg}^{\mathcal{C}_{1,n+\ell}}}{Z^{\mathcal{C}_{1,n}}} = - \sum_i \frac{W(z_i)}{2bg_s} - \frac{3\ell(\ell+1)}{4b^2} \log \eta - \frac{1}{2b^2} \sum_{i < j} \log \vartheta(z_i - z_j)$$

$$+ \sum_{k=1}^{\infty} \frac{1}{k!} \left\langle \left(\sum_{I,i} \int^{z_i} \frac{\vartheta_1'(z'_i - \lambda_I)}{\vartheta_1(z'_i - \lambda_I)} dz'_i \right)^k \right\rangle_{conn}. \quad (4.9)$$

where we have used the momentum conservation in the second term.

We introduce the deformation parameters and take the limit where $\epsilon_2 \rightarrow 0$ while ϵ_1 keeping fixed. In this limit, the path integral is dominated by the solutions of the equations of motion

$$W'(\lambda_I) - 2bg_s \prod_{J \neq I} \frac{\vartheta_1'(\lambda_I - \lambda_J)}{\vartheta_1(\lambda_I - \lambda_J)} = 0, \quad (4.10)$$

as we have seen in the previous section, and the connected part with $k \geq 2$ in (4.9) is negligible. Thus we obtain

$$\log \frac{Z_{deg}^{\mathcal{C}_{1,n+\ell}}}{Z^{\mathcal{C}_{1,n}}} \rightarrow \frac{1}{\epsilon_1} \sum_i \int^{z_i} \left(\langle R(z'_i) \rangle - \frac{W'(z'_i)}{2} \right) dz'_i \equiv \log \prod_i \Psi(z_i). \quad (4.11)$$

where R is an analog of the resolvent of the sphere case

$$R(z_1, \dots, z_k) = (bg_s)^k \sum_{I_1} \frac{\vartheta'_1(z_1 - \lambda_{I_1})}{\vartheta_1(z_1 - \lambda_{I_1})} \dots \sum_{I_k} \frac{\vartheta'_1(z_k - \lambda_{I_k})}{\vartheta_1(z_k - \lambda_{I_k})}. \quad (4.12)$$

Thus, we have related $Z_{deg}^{\mathcal{C}_{1,n+\ell}}$ with the resolvent or Ψ at least in the $\epsilon_2 \rightarrow 0$ limit.

One simple consequence which follows from the above formulas is about the monodromies of Ψ or $Z_{deg}^{\mathcal{C}_{1,n+1}}$. (For simplicity, we consider the $\ell = 1$ case. The generalization to $\ell \geq 2$ might be straightforward.) Along the A cycle, $Z_{deg}^{\mathcal{C}_{1,n+1}}$ behaves as

$$Z_{deg}^{\mathcal{C}_{1,n+1}}(z + \pi) = (-1)^{N - \frac{\sum_k m_k}{bg_s}} e^{-\frac{2\pi^2}{bg_s} p} Z_{deg}^{\mathcal{C}_{1,n+1}}(z) = (-1)^{\frac{1}{2b^2}} e^{-\frac{2\pi^2}{bg_s} p} Z_{deg}^{\mathcal{C}_{1,n+1}}(z), \quad (4.13)$$

where we have used the momentum conservation. Similarly, we can evaluate the B -cycle monodromy

$$\frac{Z_{deg}^{\mathcal{C}_{1,n+1}}(z + \pi\tau)}{Z^{\mathcal{C}_{1,n}}} = e^{-\frac{W(z)}{2bg_s}} \eta^{-\frac{3}{2b^2}} \left\langle \prod_I \vartheta_1(z - \lambda_I) e^{2i \sum_I \lambda_I} \right\rangle \exp \left(-\frac{iz}{b^2} - \frac{\pi i\tau}{2b^2} - \frac{2\pi^2 \tau p}{bg_s} \right) \quad (4.14)$$

On the other hand, let us consider the shift of $p \rightarrow p - \frac{g_s}{2\pi ib}$ in $Z_{deg}^{\mathcal{C}_{1,n+1}}$, which gives the additional factor $\exp \left(-\frac{iz}{b^2} + 2i \sum_I \lambda_I \right)$ in the integrals. Therefore, we obtain

$$Z_{deg}^{\mathcal{C}_{1,n+1}}(z + \pi\tau; p) = Z_{deg}^{\mathcal{C}_{1,n+1}}(z; p - \frac{g_s}{2\pi ib}) e^{-\frac{2\pi^2 \tau p}{bg_s} - \frac{\pi i\tau}{2b^2}}. \quad (4.15)$$

We note that these are indeed the same monodromies as those of the conformal block with the degenerate field [30, 29].

We can further proceed to derive the special geometry (ϵ_1 -deformed Seiberg-Witten) relation for the resolvent. In the $\epsilon_2 \rightarrow 0$ limit, (4.11) shows that $Z_{deg}^{\mathcal{C}_{1,n+1}}$ can be expanded as

$$Z_{deg}^{\mathcal{C}_{1,n+1}}(z) = \exp \left(\frac{\mathcal{F}(\epsilon_1)}{\epsilon_1 \epsilon_2} + \frac{1}{\epsilon_1} \int^z x(z') dz' + \mathcal{O}(\epsilon_2) \right), \quad (4.16)$$

where the first term is $\mathcal{F}(\epsilon_1) = \lim_{\epsilon_2 \rightarrow 0} F^{\mathcal{C}_{1,n}}$. The above A and B cycle monodromies indicate that for the second term

$$\oint_A x(z) dz = -2\pi^2 p, \quad \oint_B x(z) dz = -2\pi^2 \tau p - \frac{1}{2\pi i} \frac{\partial \mathcal{F}(\epsilon_1)}{\partial p}. \quad (4.17)$$

This corresponds to the Seiberg-Witten relation in the presence of the ϵ_1 dependence [12, 13]. For other independent cycles which correspond to the legs of the pants decomposition of the torus, it is natural to expect that the similar relation is satisfied.

4.2 Loop equations

Let us derive the loop equations of the matrix model for the torus. As discussed above, in order to relate with the conformal block, we will use the momentum conservation (4.8). From the Schwinger-Dyson equation for an arbitrary transformation $\delta\lambda_K = \frac{\vartheta_1'(z-\lambda_K)}{\vartheta_1(z-\lambda_K)}$, we derive

$$0 = g_s^2 \left\langle \sum_I \left(\frac{\vartheta_1'(z-\lambda_I)}{\vartheta_1(z-\lambda_I)} \right)^2 \right\rangle - g_s^2 \left\langle \sum_I \frac{\vartheta_1''(z-\lambda_I)}{\vartheta_1(z-\lambda_I)} \right\rangle + bg_s W'(z) \left\langle \sum_I \frac{\vartheta_1'(z-\lambda_I)}{\vartheta_1(z-\lambda_I)} \right\rangle + t(z) - 2b^2 g_s^2 \left\langle \sum_{I<J} \frac{\vartheta_1'(\lambda_I-\lambda_J)}{\vartheta_1(\lambda_I-\lambda_J)} \left(\frac{\vartheta_1'(z-\lambda_I)}{\vartheta_1(z-\lambda_I)} - \frac{\vartheta_1'(z-\lambda_J)}{\vartheta_1(z-\lambda_J)} \right) \right\rangle, \quad (4.18)$$

where we have multiplied the both sides by g_s^2 and defined

$$t(z) = bg_s \left\langle \sum_I \frac{\vartheta_1'(z-\lambda_I)}{\vartheta_1(z-\lambda_I)} (W'(\lambda_I) - W'(z)) \right\rangle. \quad (4.19)$$

We then use the formula (A.9) to calculate the last term and, after some algebra, we obtain

$$0 = -\langle R(z, z) \rangle - \left(b + \frac{1}{b} \right) g_s \langle R'(z) \rangle + W'(z) \langle R(z) \rangle + b^2 g_s^2 N \left\langle \sum_I \frac{\vartheta_1''(z-\lambda_I)}{\vartheta_1(z-\lambda_I)} \right\rangle + t(z) + b^2 g_s^2 \left\langle \sum_{I<J} \frac{\vartheta_1''(\lambda_I-\lambda_J)}{\vartheta_1(\lambda_I-\lambda_J)} \right\rangle + 3b^2 g_s^2 \eta_1 N(N-1). \quad (4.20)$$

This equation is valid for an arbitrary potential.

From now on, let us consider the potential corresponding to the toric conformal block (4.4). In this case, $t(z)$ can be evaluated by using (A.9) again as

$$t(z) = 2bg_s \sum_{k=1}^n m_k \frac{\vartheta_1'(z-w_k)}{\vartheta_1(z-w_k)} \left\langle \sum_I \frac{\vartheta_1'(\lambda_I-w_k)}{\vartheta_1(\lambda_I-w_k)} \right\rangle - bg_s \sum_{k=1}^n m_k \left\langle \frac{\vartheta_1''(\lambda_I-w_k)}{\vartheta_1(\lambda_I-w_k)} \right\rangle - bg_s N \sum_{k=1}^n m_k \frac{\vartheta_1''(z-w_k)}{\vartheta_1(z-w_k)} - bg_s \sum_{k=1}^n m_k \left\langle \sum_I \frac{\vartheta_1''(z-\lambda_I)}{\vartheta_1(z-\lambda_I)} \right\rangle - 6bg_s N \eta_1 \sum_{k=1}^n m_k. \quad (4.21)$$

By substituting this into (4.20) and using the momentum conservation, we obtain

$$0 = -\langle R(z, z) \rangle - \left(b + \frac{1}{b} \right) g_s \langle R'(z) \rangle + W'(z) \langle R(z) \rangle - 3bg_s(N+1)\eta_1 \sum_k m_k + 2bg_s \sum_{k=1}^n m_k \frac{\vartheta_1'(z-w_k)}{\vartheta_1(z-w_k)} \left\langle \sum_I \frac{\vartheta_1'(\lambda_I-w_k)}{\vartheta_1(\lambda_I-w_k)} \right\rangle - bg_s N \sum_k m_k \frac{\vartheta_1''(z-w_k)}{\vartheta_1(z-w_k)} - bg_s \sum_k m_k \left\langle \sum_I \frac{\vartheta_1''(\lambda_I-w_k)}{\vartheta_1(\lambda_I-w_k)} \right\rangle + b^2 g_s^2 \left\langle \sum_{I<J} \frac{\vartheta_1''(\lambda_I-\lambda_J)}{\vartheta_1(\lambda_I-\lambda_J)} \right\rangle + \frac{g_s^2 \ell}{2} \left\langle \sum_I \frac{\vartheta_1''(z-\lambda_I)}{\vartheta_1(z-\lambda_I)} \right\rangle + \frac{3\ell}{2} g_s^2 \eta_1 (N-1). \quad (4.22)$$

where the last two terms come from the deformation of the momentum conservation. Let us note that the τ -derivative of the partition function $\tilde{Z}^{\mathcal{C}_{1,n}}$

$$\frac{4g_s^2}{\tilde{Z}^{\mathcal{C}_{1,n}}} \frac{\partial \tilde{Z}^{\mathcal{C}_{1,n}}}{\partial \ln q} = -bg_s \sum_k m_k \left\langle \sum_I \frac{\vartheta_1''(\lambda_I - w_k)}{\vartheta_1(\lambda_I - w_k)} \right\rangle + b^2 g_s^2 \left\langle \sum_{I < J} \frac{\vartheta_1''(\lambda_I - \lambda_J)}{\vartheta_1(\lambda_I - \lambda_J)} \right\rangle, \quad (4.23)$$

where we have used $-8 \frac{\partial}{\partial \ln q} \vartheta_1(a) = \vartheta_1''(a)$. This is the third line in (4.22). Similarly, the derivatives with respect to w_k produce the first term in the second line of (4.22):

$$-\frac{g_s^2}{\tilde{Z}^{\mathcal{C}_{1,n}}} \frac{\partial \tilde{Z}^{\mathcal{C}_{1,n}}}{\partial w_k} = 2bg_s m_k \left\langle \sum_I \frac{\vartheta_1'(\lambda_I - w_k)}{\vartheta_1(\lambda_I - w_k)} \right\rangle. \quad (4.24)$$

We will fix one of the moduli of the torus as $w_n = 0$ below. In this case, the derivative with respect to w_n is understood as the right hand side of (4.24). Putting all these together, we finally obtain

$$\begin{aligned} 0 = & -\langle R(z, z) \rangle - \left(b + \frac{1}{b}\right) g_s \langle R'(z) \rangle + W'(z) \langle R(z) \rangle - 3bg_s(N+1)\eta_1 \sum_k m_k \\ & -g_s^2 \sum_{k=1}^n \frac{\vartheta_1'(z - w_k)}{\vartheta_1(z - w_k)} \frac{\partial \ln \tilde{Z}^{\mathcal{C}_{1,n}}}{\partial w_k} - bg_s N \sum_k m_k \frac{\vartheta_1''(z - w_k)}{\vartheta_1(z - w_k)} + 4g_s^2 \frac{\partial \ln \tilde{Z}^{\mathcal{C}_{1,n}}}{\partial \ln q} \\ & + \frac{g_s^2 \ell}{2} \left\langle \sum_I \frac{\theta_1''(z - \lambda_I)}{\theta_1(z - \lambda_I)} \right\rangle + \frac{3\ell}{2} g_s^2 \eta_1 (N-1). \end{aligned} \quad (4.25)$$

$\epsilon_2 \rightarrow 0$ limit

We consider the $\epsilon_2 \rightarrow 0$ limit. We can use under this limit the equations of motion (4.10). Furthermore, as discussed in the previous section, $\langle R(z, z) \rangle \sim \langle R(z) \rangle^2$. Also, the last two terms in (4.25) disappears in this limit. Naively, the terms with the derivative of $\ln \tilde{Z}^{\mathcal{C}_{1,n}}$ also disappear due to the factor $g_s^2 (= \epsilon_1 \epsilon_2)$. However, they does not because $\ln \tilde{Z}^{\mathcal{C}_{1,n}}$ behaves as $\tilde{F}^{\mathcal{C}_{1,n}}/g_s^2$. Thus, the loop equation (4.25) becomes

$$\begin{aligned} 0 = & -\langle R(z) \rangle^2 \Big|_{\epsilon_2 \rightarrow 0} - \epsilon_1 \langle R'(z) \rangle \Big|_{\epsilon_2 \rightarrow 0} + W'(z) \langle R(z) \rangle \Big|_{\epsilon_2 \rightarrow 0} - 3\eta_1 \left(\sum_\ell m_\ell \right) \left(\sum_k m_k + \epsilon_1 \right) \\ & - \left(\sum_\ell m_\ell \right) \sum_k m_k \frac{\vartheta_1''(z - w_k)}{\vartheta_1(z - w_k)} - \sum_{k=1}^n \frac{\vartheta_1'(z - w_k)}{\vartheta_1(z - w_k)} \frac{\partial \tilde{F}^{\mathcal{C}_{1,n}}}{\partial w_k} \Big|_{\epsilon_2 \rightarrow 0} + 4 \frac{\partial \tilde{F}^{\mathcal{C}_{1,n}}}{\partial \ln q} \Big|_{\epsilon_2 \rightarrow 0}. \end{aligned} \quad (4.26)$$

We will omit $\Big|_{\epsilon_2 \rightarrow 0}$ in what follows. After some algebra (by introducing $x = \langle R(z) \rangle - W'/2$ and by using the formula of the theta function), we obtain

$$\begin{aligned} 0 = & -x^2 - \epsilon_1 x' + \sum_k m_k (m_k + \epsilon_1) \mathcal{P}(z - w_k) \\ & + \sum_k \frac{\vartheta_1'(z - w_k)}{\vartheta_1(z - w_k)} \left(H_k - \frac{\partial \tilde{F}^{\mathcal{C}_{1,n}}}{\partial w_k} \right) + H_0 + 4 \frac{\partial \tilde{F}^{\mathcal{C}_{1,n}}}{\partial \ln q}, \end{aligned} \quad (4.27)$$

where $\mathcal{P}(z)$ is the Weierstrass elliptic function (A.3) and

$$\begin{aligned} H_k &= 4\pi p m_k + 2 \sum_{\ell(\neq k)} m_k m_\ell \frac{\vartheta'_1(w_k - w_\ell)}{\vartheta_1(w_k - w_\ell)}, \\ H_0 &= 4\pi^2 p^2 - \eta_1 \sum_k m_k(m_k + \epsilon_1) + \frac{1}{2} \sum_{k \neq \ell} m_k m_\ell \frac{\vartheta''_1(w_k - w_\ell)}{\vartheta_1(w_k - w_\ell)}. \end{aligned} \quad (4.28)$$

Therefore, we obtain that $H_k - \frac{\partial \tilde{F}^{\mathcal{C}_{1,n}}}{\partial w_k}$ and $H_0 + 4 \frac{\partial \tilde{F}^{\mathcal{C}_{1,n}}}{\partial \ln q}$ are the vacuum energies of the quantum Hitchin Hamiltonians $\mathbf{H}_k^{(1)}$ and $\mathbf{H}_0^{(1)}$ in (2.7). This shows that this matrix model captures the quantization of the Hitchin system associated with the torus.

We can write this equation in the form of the differential equation satisfied by the wavefunction $\Psi(z) = e^{\frac{1}{\epsilon_1} \int^z x(z') dz'}$:

$$\begin{aligned} &\left(-\epsilon_1^2 \frac{\partial^2}{\partial z^2} + \sum_k m_k(m_k + \epsilon_1) \mathcal{P}(z - w_k) + \sum_k \frac{\vartheta'_1(z - w_k)}{\vartheta_1(z - w_k)} H_k + H_0 \right) \Psi(z) \\ &= \left(\sum_k \frac{\vartheta'_1(z - w_k)}{\vartheta_1(z - w_k)} \frac{\partial \tilde{F}^{\mathcal{C}_{1,n}}}{\partial w_k} - 4 \frac{\partial \tilde{F}^{\mathcal{C}_{1,n}}}{\partial \ln q} \right) \Psi(z) \end{aligned} \quad (4.29)$$

By considering $\prod_{i=1}^n \Psi(z_i)$, this satisfies the above differential equations for each z_i . This is related with the KZB equation [72, 73, 74] as discussed in [28] (see also [75]).

It is easy to translate the terms in the right hand side in (4.29) to the gauge theory variables. By the identification of the moduli, the derivatives with respect to w_k are written as $\frac{\partial}{\partial w_k} = \frac{\partial}{\partial \tau_k} - \frac{\partial}{\partial \tau_{k-1}}$ (for $k = 1, \dots, n-1$), where $\tau_0 = \tau_n$, and also $\frac{\partial}{\partial \ln q} = \frac{1}{2\pi i} \frac{\partial}{\partial \tau_n}$. Therefore, we obtain

$$\frac{\partial \tilde{F}^{\mathcal{C}_{1,n}}}{\partial w_k} = u_k - u_{k-1}, \quad (4.30)$$

for $k = 1, \dots, n-1$, where we have defined $u_k = \frac{\partial \tilde{F}^{\mathcal{C}_{1,n}}}{\partial \tau_k}$ and $u_0 \equiv u_n$. For $k = n$, we calculate

$$\frac{\partial \tilde{F}^{\mathcal{C}_{1,n}}}{\partial w_n} = - \sum_{k=1}^{n-1} \frac{\partial \tilde{F}^{\mathcal{C}_{1,n}}}{\partial w_k} + 4\pi p \sum_{k=1}^n m_k = -u_{n-1} + u_n + 4\pi p \sum_{k=1}^n m_k \quad (4.31)$$

where we have used the equations of motion and the momentum conservation. (We also ignored the term depending on ϵ_2 .) Also, for the derivative with respect to q we obtain

$$-4 \frac{\partial \tilde{F}^{\mathcal{C}_{1,n}}}{\partial \ln q} = -\frac{2}{\pi i} u_n. \quad (4.32)$$

As in the sphere case, these u_k are related with $\langle \text{tr} \phi_k^2 \rangle$. Note however that there could be a difference of them since u_k here are the derivatives of \tilde{F} , which will be seen below.

For completeness, let us rewrite the above equation in terms of the original partition function $Z^{\mathcal{C}_{1,n}}$. The difference between $Z^{\mathcal{C}_{1,n}}$ and $\tilde{Z}^{\mathcal{C}_{1,n}}$ is given by C (4.2), which gives rise to

$$\begin{aligned}
4\frac{\partial\tilde{F}^{\mathcal{C}_{1,n}}}{\partial\ln q} &= 4\frac{\partial F^{\mathcal{C}_{1,n}}}{\partial\ln q} + 3\sum_k m_k(m_k + \epsilon_1)\eta_1 - \sum_{k<\ell} m_k m_\ell \frac{\vartheta_1''(w_k - w_\ell)}{\vartheta_1(w_k - w_\ell)}, \\
-\frac{\partial\tilde{F}^{\mathcal{C}_{1,n}}}{\partial w_k} &= -\frac{\partial F^{\mathcal{C}_{1,n}}}{\partial w_k} - 2\sum_{\ell(\neq k)} m_k m_\ell \frac{\vartheta_1'(w_k - w_\ell)}{\vartheta_1(w_k - w_\ell)} - 4\pi p m_k, \quad (k = 1, \dots, n-1) \\
-\frac{\partial\tilde{F}^{\mathcal{C}_{1,n}}}{\partial w_n} &= \sum_{k=1}^{n-1} \frac{\partial F^{\mathcal{C}_{1,n}}}{\partial w_k} + \sum_{\ell(\neq n)} m_n m_\ell \frac{\vartheta_1'(w_\ell)}{\vartheta_1(w_\ell)} - 4\pi p m_n,
\end{aligned} \tag{4.33}$$

where we have used (4.31). Thus, we obtain from (4.27)

$$\begin{aligned}
0 &= -x^2 - \epsilon_1 x' + \sum_k m_k(m_k + \epsilon_1)\mathcal{P}(z - w_k) + 2\sum_k m_k(m_k + \epsilon_1)\eta_1 \\
&\quad - \sum_{k=1}^{n-1} \frac{\vartheta_1'(z - w_k)}{\vartheta_1(z - w_k)} \frac{\partial F^{\mathcal{C}_{1,n}}}{\partial w_k} + \frac{\vartheta_1'(z)}{\vartheta_1(z)} \left(\sum_{k=1}^{n-1} \frac{\partial F^{\mathcal{C}_{1,n}}}{\partial w_k} + \sum_{\ell(\neq n)} m_n m_\ell \frac{\vartheta_1'(w_\ell)}{\vartheta_1(w_\ell)} \right) + 4\pi^2 p^2 + 4\frac{\partial F^{\mathcal{C}_{1,n}}}{\partial\ln q}.
\end{aligned} \tag{4.34}$$

4.3 One-punctured torus

In the $n = 1$ case, we can see the relation with the elliptic Calogero-Moser model. We will take $w_1 = 0$. The potential is

$$W(z) = 2m_1 \log \vartheta_1(z) + 4\pi p z. \tag{4.35}$$

In this case, it is easy to calculate the loop equation (4.34):

$$0 = -x(z)^2 - \epsilon_1 x'(z) + m_1(m_1 + \epsilon_1)\mathcal{P}(z) - 4u(\epsilon_1), \tag{4.36}$$

where

$$\begin{aligned}
u(\epsilon_1) &= -\pi^2 p^2 - g_s^2 \frac{\partial \ln Z^{\mathcal{C}_{1,1}}}{\partial \ln q} - \frac{m_1(m_1 + \epsilon_1)\eta_1}{2} \\
&= -\pi^2 p^2 + \frac{\partial}{\partial \ln q} (\mathcal{F}^{\mathcal{C}_{1,1}} - 2m_1(m_1 + \epsilon_1) \ln \eta),
\end{aligned} \tag{4.37}$$

where we have used that $\eta_1 = 4\frac{\partial \ln \eta}{\partial \ln q}$ (See Appendix A) and defined the free energy as $\mathcal{F}^{\mathcal{C}_{1,1}} = \lim_{\epsilon_2 \rightarrow 0} F^{\mathcal{C}_{1,1}}$. Note that the free energy is the one evaluated in the $\epsilon_2 \rightarrow 0$ limit. Eq.(4.37) is the ϵ_1 -deformed version of the relation between $\langle \text{tr} \phi^2 \rangle$ and the Coulomb modulus u [66] which coincides with the one found in [24].

By introducing the “wave-function” $\Psi = e^{\frac{1}{\epsilon_1} \int^z dz' x(z')}$, we finally obtain

$$\left[-\epsilon_1^2 \frac{\partial^2}{\partial z^2} + m_1(m_1 + \epsilon_1)\mathcal{P}(z) \right] \Psi = 4u\Psi. \tag{4.38}$$

The left hand side is the Calogero-Moser Hamiltonian and u in the right hand side can be considered as the eigenvalue of the Hamiltonian. Note that similar equations have been derived from the Virasoro conformal block [29] and affine \hat{sl}_2 conformal block [28]. Indeed, by assuming the equivalence of the partition function of the matrix model $Z^{\mathcal{C}_{1,1}}$ with the one-point conformal block on a torus, (4.38) becomes the exactly same equation as the one obtained from the conformal block with the degenerate field. (See Section 3.1.2 in [29]. The identification of the parameter is $a = i\pi p$.)

We also emphasize that the differential $x dz$ in the wave-function satisfies the special geometry relation (4.17). This is equivalent to the proposal in [12] stating that the ϵ_1 -deformed prepotential can be obtained from the ϵ_1 -deformed special geometry relation for the $\mathcal{N} = 2^*$ theory, by using the same argument as in [29].

4.3.1 Large N limit and prepotential

Before going to next, let us consider the loop equation in the large N limit which can be obtain by taking $\epsilon_1 \rightarrow 0$ further in (4.36):

$$x^2 = m^2 \mathcal{P}(z) - 4u(\epsilon_1 = 0), \quad (4.39)$$

which is the Seiberg-Witten curve of the $\mathcal{N} = 2^*$ gauge theory [3]. Indeed, the parameter u can be written as

$$u(\epsilon_1 = 0) = -\pi^2 p^2 + \frac{\partial}{\partial \ln q} \left(\mathcal{F}_0^{\mathcal{C}_{1,1}} - 2m_1^2 \ln \eta \right). \quad (4.40)$$

where $\mathcal{F}_0^{\mathcal{C}_{1,1}}$ is the leading contribution of the full free energy in the limit where $\epsilon_{1,2} \rightarrow 0$. The first term corresponds the classical contribution to the prepotential. The last term denotes the shift of the Coulomb moduli parameter from the value of the physical expectation value $\langle \text{Tr } \phi^2 \rangle$ [1, 76, 66].

Indeed, we can be more precise. Under the identification $i\pi p$ with the vev of the vector multiplet scalar a , it is easy to show from (4.13) that

$$a = i\pi p = \frac{1}{2\pi i} \int_0^\pi x dz. \quad (4.41)$$

This and the fact that the form of x here is the same as the Seiberg-Witten differential of the $\mathcal{N} = 2^*$ gauge theory where $\mathcal{F}_0^{\mathcal{C}_{1,1}}$ is changed to the prepotential (see [77, 29]) lead to the conclusion that the free energy in the large N limit of this matrix model is exactly the same as the prepotential of the gauge theory (under the identification above).

5 Conclusions

In this paper we proposed that the β -deformation of matrix models provides, in a suitable limit, the quantization of the associated integrable system. In particular we have shown that the loop equations for the β -deformed generalized matrix models [54, 56] reproduce in the Nekrasov-Shatashvili limit the Hamiltonians of the quantum Hitchin system associated to the sphere and torus with marked points. Moreover, we have shown how to obtain the wave function from degenerate field insertions.

It would be interesting to understand if this procedure could provide a general quantization prescription of integrable systems which can be linked to specific matrix models. To this end it would be very useful to provide further evidence and examples. For instance, it would be interesting to investigate the β -deformed Chern-Simons matrix model [49] in this direction. Furthermore, the extension of our approach to q -deformed conformal blocks, along the lines of [78], would be worth to be analyzed with the aim of connecting our results with topological strings.

On a more specific side, a natural extension of our analysis concerns Hitchin systems on curves of higher genera, the point being a generalization of the identity (A.8). A further explorable direction would be the extension to higher rank gauge groups with a multi-matrix model approach.

The problem of understanding the proper quantization of the Seiberg-Witten geometry has been explored recently also from a different view point consisting in a saddle point analysis of the instanton partition in the Nekrasov-Shatashvili limit [19, 24, 27]. In Section 4 we have seen that for the $\mathcal{N} = 2^*$ theory the eigenvalues of the Hamiltonian match. It would be interesting to further explore the relation between the two quantizations.

Acknowledgements

We thank A. Brini, B. Dubrovin, H. Itoyama, S. Pasquetti, F. Yagi, for interesting discussions and comments.

G.B. and K.M. are partially supported by the INFN project TV12. A.T. is partially supported by PRIN “Geometria delle varietà algebriche e loro spazi di moduli” and the INFN project PI14 “Nonperturbative dynamics of gauge theories”.

Appendix

A Elliptic functions

The elliptic theta function is defined by

$$\vartheta_1(z|\tau) = 2q^{1/8} \sin z \prod_{n=1}^{\infty} (1 - q^n)(1 - 2q^n \cos 2z + q^{2n}) \quad (\text{A.1})$$

which has pseudo periodicity

$$\vartheta_1(z + \pi|\tau) = -\vartheta_1(z|\tau), \quad \vartheta_1(z + \pi\tau|\tau) = e^{-i(2z + \pi\tau)} \vartheta_1(z|\tau). \quad (\text{A.2})$$

This function satisfies $\vartheta_1''(z|\tau) = -8 \frac{\partial}{\partial \ln q} \vartheta_1(z|\tau)$ where $\vartheta_1'(z|\tau) = \frac{\partial}{\partial z} \vartheta_1(z|\tau)$.

The Weierstrass elliptic function \mathcal{P} is double periodic with periods π and $\pi\tau$ and is expressed as

$$\mathcal{P}(z) = -\zeta'(z), \quad \zeta(z) = \frac{\vartheta_1'(z|\tau)}{\vartheta_1(z|\tau)} + 2\eta_1 z, \quad \eta_1 = -\frac{1}{6} \frac{\vartheta_1'''(z|\tau)|_{z=0}}{\vartheta_1'(z|\tau)|_{z=0}}, \quad (\text{A.3})$$

where $\vartheta_1(z|\tau)$ is elliptic theta function. The Weierstrass function satisfies

$$\mathcal{P}(z)' = 4\mathcal{P}(z)^3 - g_2\mathcal{P}(z) - g_3, \quad (\text{A.4})$$

where

$$g_2 = \frac{4}{3} \left(1 + 240 \sum_{n=1}^{\infty} \frac{n^3 q^n}{1 - q^n} \right),$$

$$g_3 = \frac{8}{27} \left(1 - 504 \sum_{n=1}^{\infty} \frac{n^5 q^n}{1 - q^n} \right), \quad (\text{A.5})$$

We also define g_1 whose expansion is

$$g_1 = -\frac{1}{3} \left(1 - 24 \sum_{n=1}^{\infty} \frac{nq^n}{1 - q^n} \right), \quad (\text{A.6})$$

which is related with η_1 as $g_1 = -2\eta_1$:

$$\eta_1 = \frac{1}{6} \left(1 - 24 \sum_{n=1}^{\infty} \frac{nq^n}{1 - q^n} \right) = 4 \frac{\partial}{\partial \ln q} \ln \eta, \quad (\text{A.7})$$

where $\eta = q^{1/24} \prod_{n=1}^{\infty} (1 - q^n)$ is the Dedekind eta function.

The Weierstrass and zeta functions satisfies the identity

$$\mathcal{P}(a + b) + \mathcal{P}(a) + \mathcal{P}(b) = (\zeta(a + b) - \zeta(a) - \zeta(b))^2. \quad (\text{A.8})$$

This leads to the following formula of the theta function

$$\frac{\vartheta_1'(b - a)}{\vartheta_1(b - a)} \left(\frac{\vartheta_1'(a)}{\vartheta_1(a)} - \frac{\vartheta_1'(b)}{\vartheta_1(b)} \right) = \frac{\vartheta_1'(a) \vartheta_1'(b)}{\vartheta_1(a) \vartheta_1(b)} - \frac{1}{2} \left(\frac{\vartheta_1''(a)}{\vartheta_1(a)} + \frac{\vartheta_1''(b)}{\vartheta_1(b)} + \frac{\vartheta_1''(b - a)}{\vartheta_1(b - a)} \right) - 3\eta_1. \quad (\text{A.9})$$

References

- [1] N. Seiberg and E. Witten, “Electric - magnetic duality, monopole condensation, and confinement in $N=2$ supersymmetric Yang-Mills theory,” Nucl. Phys. B **426** (1994) 19 [Erratum-ibid. B **430** (1994) 485] [arXiv:hep-th/9407087].
N. Seiberg and E. Witten, “Monopoles, duality and chiral symmetry breaking in $N=2$ supersymmetric QCD,” Nucl. Phys. B **431**, 484 (1994) [arXiv:hep-th/9408099].
- [2] A. Gorsky, I. Krichever, A. Marshakov, A. Mironov and A. Morozov, “Integrability and Seiberg-Witten exact solution,” Phys. Lett. B **355**, 466 (1995) [arXiv:hep-th/9505035].
E. J. Martinec and N. P. Warner, “Integrable systems and supersymmetric gauge theory,” Nucl. Phys. B **459**, 97 (1996) [arXiv:hep-th/9509161].
T. Nakatsu and K. Takasaki, “Whitham-Toda hierarchy and $N = 2$ supersymmetric Yang-Mills theory,” Mod. Phys. Lett. A **11**, 157 (1996) [arXiv:hep-th/9509162].
H. Itoyama and A. Morozov, “Integrability and Seiberg-Witten Theory: Curves and Periods,” Nucl. Phys. B **477**, 855 (1996) [arXiv:hep-th/9511126]; “Prepotential and the Seiberg-Witten Theory,” Nucl. Phys. B **491**, 529 (1997) [arXiv:hep-th/9512161].
A. Gorsky, A. Marshakov, A. Mironov and A. Morozov, “ $N=2$ Supersymmetric QCD and Integrable Spin Chains: Rational Case $N_f < 2N_c$,” Phys. Lett. B **380**, 75 (1996) [arXiv:hep-th/9603140].
- [3] R. Donagi and E. Witten, “Supersymmetric Yang-Mills Theory And Integrable Systems,” Nucl. Phys. B **460**, 299 (1996) [arXiv:hep-th/9510101].
- [4] D. Gaiotto, “ $N=2$ dualities,” arXiv:0904.2715 [hep-th].
- [5] D. Gaiotto, G. W. Moore and A. Neitzke, “Four-dimensional wall-crossing via three-dimensional field theory,” Commun. Math. Phys. **299** (2010) 163 [arXiv:0807.4723 [hep-th]].
- [6] N. A. Nekrasov, “Seiberg-Witten prepotential from instanton counting,” arXiv:hep-th/0306211.
N. Nekrasov and A. Okounkov, “Seiberg-Witten theory and random partitions,” arXiv:hep-th/0306238.
- [7] R. Flume and R. Poghossian, “An Algorithm for the microscopic evaluation of the coefficients of the Seiberg-Witten prepotential,” Int. J. Mod. Phys. A **18**, 2541 (2003) [arXiv:hep-th/0208176].

- U. Bruzzo, F. Fucito, J. F. Morales and A. Tanzini, “Multiinstanton calculus and equivariant cohomology,” *JHEP* **0305**, 054 (2003) [arXiv:hep-th/0211108].
- F. Fucito, J. F. Morales and R. Poghossian, “Instantons on quivers and orientifolds,” *JHEP* **0410**, 037 (2004) [arXiv:hep-th/0408090].
- [8] L. F. Alday, D. Gaiotto and Y. Tachikawa, “Liouville Correlation Functions from Four-dimensional Gauge Theories,” *Lett. Math. Phys.* **91**, 167 (2010) [arXiv:0906.3219 [hep-th]].
- [9] G. Bonelli and A. Tanzini, “Hitchin systems, N=2 gauge theories and W-gravity,” *Phys. Lett. B* **691**, 111 (2010) [arXiv:0909.4031 [hep-th]].
- [10] J. Teschner, “Quantization of the Hitchin moduli spaces, Liouville theory, and the geometric Langlands correspondence I,” arXiv:1005.2846 [hep-th].
- [11] N. A. Nekrasov and S. L. Shatashvili, “Quantization of Integrable Systems and Four Dimensional Gauge Theories,” arXiv:0908.4052 [hep-th].
- [12] A. Mironov and A. Morozov, “Nekrasov Functions and Exact Bohr-Sommerfeld Integrals,” *JHEP* **1004**, 040 (2010) [arXiv:0910.5670 [hep-th]].
- [13] A. Mironov and A. Morozov, “Nekrasov Functions from Exact BS Periods: the Case of SU(N),” *J. Phys. A* **43**, 195401 (2010) [arXiv:0911.2396 [hep-th]].
- [14] A. Popolitov, “On relation between Nekrasov functions and BS periods in pure SU(N) case,” arXiv:1001.1407 [hep-th].
- [15] N. Nekrasov and E. Witten, “The Omega Deformation, Branes, Integrability, and Liouville Theory,” arXiv:1002.0888 [hep-th].
- [16] D. Orlando and S. Reffert, “Relating Gauge Theories via Gauge/Bethe Correspondence,” arXiv:1005.4445 [hep-th].
- [17] W. He and Y. G. Miao, “On the magnetic expansion of Nekrasov theory: the SU(2) pure gauge theory,” arXiv:1006.1214 [hep-th].
- [18] K. K. Kozłowski and J. Teschner, “TBA for the Toda chain,” arXiv:1006.2906 [math-ph].
- [19] R. Poghossian, “Deforming SW curve,” arXiv:1006.4822 [hep-th].
- [20] A. Marshakov, A. Mironov and A. Morozov, “On AGT Relations with Surface Operator Insertion and Stationary Limit of Beta-Ensembles,” *Teor. Mat. Fiz.* **164**, 1:3 (2010) [arXiv:1011.4491 [hep-th]].

- [21] Y. Yamada, “A quantum isomonodromy equation and its application to $N=2$ $SU(N)$ gauge theories,” *J. Phys. A* **44**, 055403 (2011) [arXiv:1011.0292 [hep-th]].
- [22] M. Piatek, “Classical conformal blocks from TBA for the elliptic Calogero-Moser system,” [arXiv:1102.5403 [hep-th]].
- [23] N. Nekrasov, A. Rosly, S. Shatashvili, “Darboux coordinates, Yang-Yang functional, and gauge theory,” [arXiv:1103.3919 [hep-th]].
- [24] F. Fucito, J. F. Morales, R. Poghossian and D. R. Pacifici, “Gauge theories on Ω -backgrounds from non commutative Seiberg-Witten curves,” arXiv:1103.4495 [hep-th].
- [25] Y. Zenkevich, “Nekrasov prepotential with fundamental matter from the quantum spin chain,” arXiv:1103.4843 [math-ph].
- [26] N. Dorey, T. J. Hollowood and S. Lee, “Quantization of Integrable Systems and a 2d/4d Duality,” arXiv:1103.5726 [hep-th].
- [27] H. -Y. Chen, N. Dorey, T. J. Hollowood, S. Lee, “A New 2d/4d Duality via Integrability,” [arXiv:1104.3021 [hep-th]].
- [28] L. F. Alday and Y. Tachikawa, “Affine $SL(2)$ conformal blocks from 4d gauge theories,” *Lett. Math. Phys.* **94**, 87-114 (2010). [arXiv:1005.4469 [hep-th]].
- [29] K. Maruyoshi and M. Taki, “Deformed Prepotential, Quantum Integrable System and Liouville Field Theory,” arXiv:1006.4505 [hep-th].
- [30] L. F. Alday, D. Gaiotto, S. Gukov, Y. Tachikawa and H. Verlinde, “Loop and surface operators in $N=2$ gauge theory and Liouville modular geometry,” *JHEP* **1001**, 113 (2010) [arXiv:0909.0945 [hep-th]].
- [31] T. Dimofte, S. Gukov and L. Hollands, “Vortex Counting and Lagrangian 3-manifolds,” arXiv:1006.0977 [hep-th].
- [32] C. Kozcaz, S. Pasquetti and N. Wyllard, “A & B model approaches to surface operators and Toda theories,” *JHEP* **1008** (2010) 042 [arXiv:1004.2025 [hep-th]].
- [33] M. Taki, “Surface Operator, Bubbling Calabi-Yau and AGT Relation,” arXiv:1007.2524 [hep-th].

- [34] H. Awata, H. Fuji, H. Kanno, M. Manabe and Y. Yamada, “Localization with a Surface Operator, Irregular Conformal Blocks and Open Topological String,” arXiv:1008.0574 [hep-th].
- [35] C. Kozcaz, S. Pasquetti, F. Passerini and N. Wyllard, “Affine $sl(N)$ conformal blocks from $N=2$ $SU(N)$ gauge theories,” arXiv:1008.1412 [hep-th];
- [36] G. Bonelli, A. Tanzini and J. Zhao, “Vertices, Vortices and Interacting Surface Operators,” arXiv:1102.0184 [hep-th].
- [37] R. Dijkgraaf and C. Vafa, “Toda Theories, Matrix Models, Topological Strings, and $N=2$ Gauge Systems,” arXiv:0909.2453 [hep-th].
- [38] H. Itoyama, K. Maruyoshi and T. Oota, “The Quiver Matrix Model and 2d-4d Conformal Connection,” *Prog. Theor. Phys.* **123**, 957 (2010) [arXiv:0911.4244 [hep-th]].
- [39] T. Eguchi and K. Maruyoshi, “Penner Type Matrix Model and Seiberg-Witten Theory,” *JHEP* **1002**, 022 (2010) [arXiv:0911.4797 [hep-th]].
- [40] R. Schiappa and N. Wyllard, “An A_r threesome: Matrix models, 2d CFTs and 4d $N=2$ gauge theories,” arXiv:0911.5337 [hep-th];
- [41] A. Mironov, A. Morozov and S. Shakirov, “Matrix Model Conjecture for Exact BS Periods and Nekrasov Functions,” *JHEP* **1002**, 030 (2010) [arXiv:0911.5721 [hep-th]];
- [42] M. Fujita, Y. Hatsuda, T. -S. Tai, “Genus-one correction to asymptotically free Seiberg-Witten prepotential from Dijkgraaf-Vafa matrix model,” *JHEP* **1003**, 046 (2010). [arXiv:0912.2988 [hep-th]].
- [43] P. Sulkowski, “Matrix models for beta-ensembles from Nekrasov partition functions,” *JHEP* **1004**, 063 (2010). [arXiv:0912.5476 [hep-th]].
- [44] A. Mironov, A. Morozov and S. Shakirov, “Conformal blocks as Dotsenko-Fateev Integral Discriminants,” *Int. J. Mod. Phys. A* **25**, 3173 (2010) [arXiv:1001.0563 [hep-th]];
- [45] H. Itoyama and T. Oota, “Method of Generating q -Expansion Coefficients for Conformal Block and $N=2$ Nekrasov Function by beta-Deformed Matrix Model,” *Nucl. Phys. B* **838**, 298 (2010) [arXiv:1003.2929 [hep-th]];
- [46] A. Mironov, A. Morozov and A. Morozov, “Matrix model version of AGT conjecture and generalized Selberg integrals,” arXiv:1003.5752 [hep-th];

- [47] T. Eguchi and K. Maruyoshi, “Seiberg-Witten theory, matrix model and AGT relation,” JHEP **1007**, 081 (2010) [arXiv:1006.0828 [hep-th]].
- [48] H. Itoyama, T. Oota, N. Yonezawa, “Massive Scaling Limit of beta-Deformed Matrix Model of Selberg Type,” [arXiv:1008.1861 [hep-th]].
- [49] A. Brini, M. Marino, S. Stevan, “The Uses of the refined matrix model recursion,” [arXiv:1010.1210 [hep-th]].
- [50] M. C. N. Cheng, R. Dijkgraaf and C. Vafa, “Non-Perturbative Topological Strings And Conformal Blocks,” arXiv:1010.4573 [hep-th].
- [51] O. P. Santillan, “Geometric transitions, double scaling limits and gauge theories,” [arXiv:1103.1422 [hep-th]].
- [52] A. Mironov, A. Morozov, A. Popolitov, S. Shakirov, “Resolvents and Seiberg-Witten representation for Gaussian beta-ensemble,” [arXiv:1103.5470 [hep-th]].
- [53] H. Itoyama, N. Yonezawa, “ ϵ -Corrected Seiberg-Witten Prepotential Obtained From Half Genus Expansion in beta-Deformed Matrix Model,” [arXiv:1104.2738 [hep-th]].
- [54] K. Maruyoshi and F. Yagi, “Seiberg-Witten curve via generalized matrix model,” JHEP **1101**, 042 (2011) [arXiv:1009.5553 [hep-th]].
- [55] A. Mironov, A. Morozov, S. Shakirov, “On ‘Dotsenko-Fateev’ representation of the toric conformal blocks,” J. Phys. A **A44**, 085401 (2011). [arXiv:1010.1734 [hep-th]].
- [56] G. Bonelli, K. Maruyoshi, A. Tanzini and F. Yagi, “Generalized matrix models and AGT correspondence at all genera,” arXiv:1011.5417 [hep-th].
- [57] B. Eynard and O. Marchal, “Topological expansion of the Bethe ansatz, and non-commutative algebraic geometry,” JHEP **0903**, 094 (2009) [arXiv:0809.3367 [math-ph]].
- [58] L. Chekhov, B. Eynard, O. Marchal, “Topological expansion of the Bethe ansatz, and quantum algebraic geometry,” [arXiv:0911.1664 [math-ph]].
- [59] L. O. Chekhov, B. Eynard, O. Marchal, “Topological expansion of β -ensemble model and quantum algebraic geometry in the sectorwise approach,” Theor. Math. Phys. **166**, 141-185 (2011). [arXiv:1009.6007 [math-ph]].
- [60] E. Frenkel, “Lectures on the Langlands program and conformal field theory,” arXiv:hep-th/0512172.

- [61] E. Witten, “Solutions of four-dimensional field theories via M theory,” Nucl. Phys. **B500**, 3-42 (1997). [hep-th/9703166].
- [62] K. Gawedzki, P. Tran-Ngoc-Bich, “Hitchin systems at low genera,” J. Math. Phys. **41**, 4695-4712 (2000). [hep-th/9803101].
- [63] V. S. Dotsenko and V. A. Fateev, “Conformal algebra and multipoint correlation functions in 2D statistical models,” Nucl. Phys. B **240**, 312 (1984);
V. S. Dotsenko and V. A. Fateev, “Four Point Correlation Functions And The Operator Algebra In The Two-Dimensional Conformal Invariant Theories With The Central Charge $C < 1$,” Nucl. Phys. B **251**, 691 (1985).
- [64] A. A. Belavin, A. M. Polyakov and A. B. Zamolodchikov, “Infinite conformal symmetry in two-dimensional quantum field theory,” Nucl. Phys. B **241**, 333 (1984).
- [65] R. Flume, F. Fucito, J. F. Morales and R. Poghossian, “Matone’s relation in the presence of gravitational couplings,” JHEP **0404**, 008 (2004) [arXiv:hep-th/0403057].
- [66] F. Fucito, J. F. Morales, R. Poghossian and A. Tanzini, “ $N = 1$ superpotentials from multi-instanton calculus,” JHEP **0601**, 031 (2006) [arXiv:hep-th/0510173].
- [67] M. Matone, “Instantons and recursion relations in $N=2$ SUSY gauge theory,” Phys. Lett. B **357**, 342 (1995) [arXiv:hep-th/9506102].
- [68] J. Sonnenschein, S. Theisen and S. Yankielowicz, “On the Relation Between the Holomorphic Prepotential and the Quantum Moduli in SUSY Gauge Theories,” Phys. Lett. B **367**, 145 (1996) [arXiv:hep-th/9510129].
- [69] T. Eguchi and S. K. Yang, “Prepotentials of $N = 2$ Supersymmetric Gauge Theories and Soliton Equations,” Mod. Phys. Lett. A **11**, 131 (1996) [arXiv:hep-th/9510183].
- [70] V. G. Knizhnik and A. B. Zamolodchikov, “Current algebra and Wess-Zumino model in two dimensions,” Nucl. Phys. B **247**, 83 (1984).
- [71] M. Goulian and M. Li, “Correlation functions in Liouville theory,” Phys. Rev. Lett. **66** (1991) 2051.
- [72] D. Bernard, “On The Wess-Zumino-Witten Models On The Torus,” Nucl. Phys. B **303**, 77 (1988).

- [73] P. I. Etingof and A. A. Kirillov, “Representation of affine Lie algebras, parabolic differential equations and Lamé functions,” arXiv:hep-th/9310083.
- [74] G. Felder, C. Weiczerkowski, “Conformal blocks on elliptic curves and the Knizhnik-Zamolodchikov-Bernard equations,” Commun. Math. Phys. **176**, 133-162 (1996). [hep-th/9411004].
- [75] T. Eguchi and H. Ooguri, “Conformal and Current Algebras on General Riemann Surface,” Nucl. Phys. B **282**, 308 (1987).
- [76] N. Dorey, V. V. Khoze and M. P. Mattis, “On mass-deformed $N = 4$ supersymmetric Yang-Mills theory,” Phys. Lett. B **396**, 141 (1997) [arXiv:hep-th/9612231].
- [77] V. A. Fateev and A. V. Litvinov, “On AGT conjecture,” JHEP **1002**, 014 (2010) [arXiv:0912.0504 [hep-th]].
- [78] H. Awata and Y. Yamada, “Five-dimensional AGT Relation and the Deformed beta-ensemble,” Prog. Theor. Phys. **124** (2010) 227 [arXiv:1004.5122 [hep-th]].