



# A Note on Non-local Sobolev Spaces and Non-local Perimeters

Konstantinos Bessas<sup>1</sup> · Giuseppe Cosma Brusca<sup>2</sup>

Received: 11 April 2025 / Revised: 24 July 2025 / Accepted: 26 August 2025 /

Published online: 20 November 2025

© The Author(s) 2025

## Abstract

We investigate the space of non-local Sobolev functions associated with an integral kernel. We prove an extension result, Sobolev and Poincaré inequalities and an isoperimetric inequality for the non-local perimeter restricted to a set. Finally, we remark on non-local isoperimetric problems, even when the underlying kernel is not necessarily radially symmetric.

**Keywords** Non-local Sobolev spaces · Functions of non-local bounded variation · Non-local interactions · Non-local perimeters · Isoperimetric problems

**Mathematics Subject Classification** 49Q20 · 26B30 · 46E35

## 1 Introduction

The aim of this note is presenting a result of continuous extension for non-local Sobolev spaces in which interactions are weighted by a non-negative kernel  $K$ . Letting  $K$  be a kernel, i.e., a measurable function from  $\mathbb{R}^d$  to  $[0, +\infty]$  such that  $K \not\equiv 0$  up to negligible sets, and letting  $\Omega$  denote an open subset of  $\mathbb{R}^d$ , we consider the seminorm

---

✉ Konstantinos Bessas  
konstantinos.bessas@unipv.it

Giuseppe Cosma Brusca  
gbrusca@sissa.it

<sup>1</sup> Dipartimento di Matematica, Università di Pavia, Via Adolfo Ferrata 5, 27100 Pavia, Italy

<sup>2</sup> SISSA, Via Bonomea 265, 34136 Trieste, Italy

$$[u]_{W^{K,p}(\Omega)} := \left( \int_{\Omega \times \Omega} |u(x) - u(y)|^p K(x-y) \, dx dy \right)^{\frac{1}{p}} \in [0, +\infty], \quad (1.1)$$

for  $p \in [1, +\infty)$  and  $u : \mathbb{R}^d \rightarrow [-\infty, +\infty]$  measurable and finite a.e., and introduce the Banach space

$$W^{K,p}(\Omega) := \left\{ u \in L^p(\Omega) : [u]_{W^{K,p}(\Omega)} < +\infty \right\}$$

equipped with the norm  $\|u\|_{W^{K,p}(\Omega)} := \|u\|_{L^p(\Omega)} + [u]_{W^{K,p}(\Omega)}$ .  $W^{K,p}(\Omega)$  is a generalization of the usual *fractional Gagliardo-Slobodeckij-Sobolev space*  $W^{s,p}(\Omega)$  (see also [1, Section 3], where similar spaces are studied).

We prove that, under fairly mild assumptions on the kernel,  $W^{K,p}(\Omega)$  is continuously embedded in  $W^{K,p}(\mathbb{R}^d)$  through an extension operator, provided that  $\Omega$  is a bounded, open set with Lipschitz boundary. The proof of this result is obtained by adapting the arguments employed in [2] for fractional Sobolev spaces, i.e., in the case where  $K$  is the *fractional kernel*  $K(z) = |z|^{-d-sp}$  for  $s \in (0, 1)$ . In order to treat a rather general class of kernels, we detect a set of assumptions that are needed to us, but that may possibly be weakened, in order to carry out our analysis. Some of these hypotheses have been discussed by the first-named author and Stefani in [3] for studying the properties of the above function space when  $\Omega = \mathbb{R}^d$ . In that work, it is remarked that certain assumptions are necessary in order to obtain a non-trivial function space: as an example, a simple computation shows that if  $K \in L^1(\mathbb{R}^d)$ , then  $W^{K,p}(\mathbb{R}^d) = L^p(\mathbb{R}^d)$ , and, for this reason, we may always assume that the kernel is *non-integrable*,

$$K \notin L^1(\mathbb{R}^d). \quad (\text{Nint})$$

In this work, we prove that another natural assumption is the integrability of the kernel *far from the origin*, i.e.,

$$K \in L^1(\mathbb{R}^d \setminus B_r) \text{ for all } r > 0, \quad (\text{Far})$$

where  $B_r$  denotes the Euclidean ball centred at the origin with radius  $r$ . Indeed, we show that, if the above is not in force, then  $W^{K,p}(\mathbb{R}^d) = \{0\}$ .

The main difficulties in mimicking the arguments presented in [2] are due to the general structure of the kernel  $K$ , as radial symmetry and homogeneity are usually lacking. Here, we present the main assumptions that we need, additionally to the above mentioned (Nint) and (Far), in order to obtain our result.

We let  $|\cdot|_*$  denote a generic norm on  $\mathbb{R}^d$ , and, for the sake of clarity, we let  $|\cdot|_2$  denote the standard Euclidean norm on  $\mathbb{R}^d$ . Given a measurable function  $K : \mathbb{R}^d \rightarrow [0, +\infty]$ , we will consider the following conditions:

there exists  $c_0 \in (0, 1]$  such that

$$|x|_* \leq |y|_* \implies K(x) \geq c_0 K(y); \quad (\text{Dec}(c_0, |\cdot|_*))$$

there exist positive constants  $D$  and  $C_D$  such that

$$|y|_* = 2|x|_*, |x|_* \leq D \implies K(x) \leq C_D K(y); \quad (\text{Dou}(D, |\cdot|_*))$$

$$\int_{\mathbb{R}^d} K(z) \min\{1, |z|_2^p\} dz < +\infty. \quad (\text{Nts}_p)$$

We comment on the above hypotheses. The assumption  $(\text{Dec}(c_0, |\cdot|_*))$  tells that  $K$  is (almost) *decreasing* with respect to the distance induced by the norm  $|\cdot|_*$  and up to a multiplicative constant  $c_0$ . More precisely, we observe that if  $c_0 = 1$ , then the kernel is of the form  $K(z) = \kappa(|z|_*)$  for some  $\kappa : [0, +\infty) \rightarrow [0, +\infty]$  non-increasing. Our analysis takes into account also kernels which are not necessarily of this form, indeed if  $c_0 \in (0, 1)$ , such assumption allows to deal with kernels that are not described by a one-dimensional, non-increasing profile. For instance, we may consider

$$\kappa : [0, +\infty) \rightarrow [0, +\infty]$$

a non-increasing function, and

$$\phi : \mathbb{R}^d \rightarrow [\alpha, \beta]$$

a measurable function with  $0 < \alpha < \beta$ . Then, the kernel

$$K(z) = \kappa(|z|_*)\phi(z),$$

satisfies  $(\text{Dec}(c_0, |\cdot|_*))$  with  $c_0 = \alpha/\beta$ .

The doubling assumption  $(\text{Dou}(D, |\cdot|_*))$  gives quantitative information on the growth of the kernel close to the origin. We will take advantage of this assumption, combined with  $(\text{Dec}(c_0, |\cdot|_*))$ , in order to overcome the lack of homogeneity of  $K$ . Indeed, this is one of the main issues that requires some adaptations compared to the fractional case.

We also observe that if a kernel satisfies both  $(\text{Dou}(D, |\cdot|_*))$  and  $(\text{Dec}(c_0, |\cdot|_*))$ , then necessarily  $c_0 \leq C_D$ .

Finally, the assumption  $(\text{Nts}_p)$  implies that the kernel  $K$  is *not too singular*: namely, it is integrable away from the origin and it is allowed to have a singularity at the origin which cannot be stronger than the singularity of the kernel  $1/|\cdot|_2^{p+d}$ . This hypothesis is satisfied for instance by the aforementioned fractional kernel  $K(z) = |z|^{-d-sp}$  with  $s \in (0, 1)$  and by all the integrable kernels. In the literature  $(\text{Nts}_p)$  is also known as the *p-Lévy integrability condition*.  $(\text{Nts}_p)$  is stronger than  $(\text{Far})$  and it implies that the usual local Sobolev space  $W^{1,p}(\mathbb{R}^d)$  is contained in  $W^{K,p}(\mathbb{R}^d)$ , with continuous injection. We point out that, as a consequence of the equivalence of the norms on  $\mathbb{R}^d$ ,  $(\text{Nts}_p)$  could be equivalently stated in terms of the generic norm  $|\cdot|_*$ .

A prototypical kernel that satisfies all the above assumptions is the one piecewise defined by fractional kernels:

$$K(z) = \begin{cases} \alpha_1 |z|_*^{-d-s_1 p} & \text{if } |z|_* \leq R_1, \\ \alpha_k |z|_*^{-d-s_k p} & \text{if } R_{k-1} < |z|_* \leq R_k, \quad k \in \{2, \dots, M-1\}, \\ \alpha_M |z|_*^{-d-s_M p} & \text{if } |z|_* > R_{M-1}, \end{cases}$$

where  $M \in \mathbb{N}^+$  is fixed and it holds that  $\alpha_1 \geq \dots \geq \alpha_M \geq 0$ ,  $1 \leq R_1 < \dots < R_M$ , and  $0 < s_1 \leq \dots \leq s_M < 1$ .

Also mixed fractional-logarithmic kernels can be taken into account in our analysis. For instance, the kernel

$$K(z) = \begin{cases} (1 - \log(|z|_*))^{-\alpha} |z|_*^{-d-s} & \text{if } |z|_* \leq 1, \\ 0 & \text{if } |z|_* > 1 \end{cases}$$

always satisfies (Far),  $(\text{Dou}(D, |\cdot|_*))$  and  $(\text{Nts}_1)$ ; while  $(\text{Nint})$  holds for

$$(s, \alpha) \in (0, 1) \times \mathbb{R} \cup \{0\} \times (-\infty, 1],$$

and  $(\text{Dec}(c_0, |\cdot|_*))$  holds for every  $c_0 \in (0, 1]$  whenever  $\alpha \in (-\infty, 0]$ .

Finally, we observe that assumption  $(\text{Dec}(c_0, |\cdot|_*))$  allows to consider oscillating kernels when  $c_0 \in (0, 1)$ . Indeed, an example of admissible kernel is obtained starting from the one-dimensional profile

$$\kappa(t) = \begin{cases} \beta t^{-d-s} & \text{if } t \in (0, 1], \\ \alpha \sin(2\pi t) + \beta & \text{if } t \in (1, M], \\ \beta(t - M + 1)^{-d-s} & \text{if } t \in (M, +\infty), \end{cases}$$

where  $M \in \mathbb{N}^+$ ,  $0 < \alpha < \beta$ , and it is given by

$$K(z) = \kappa(|z|_*),$$

which satisfies  $(\text{Dec}(c_0, |\cdot|_*))$  with  $c_0 = \frac{\beta-\alpha}{\alpha+\beta}$ .

We point out that an extension result is also conjectured in [4, Theorem 3.78] under the further assumption that the kernel is radial, without requiring a doubling property. In our proof, radial symmetry of the kernel is not required, but, in order to employ the arguments of [2], we also rely on  $(\text{Dou}(D, |\cdot|_*))$ .

As an application of the extension result, we prove continuous and compact embeddings of  $W^{K,p}(\Omega)$  in  $L^q(\Omega)$  for some  $q$  possibly strictly larger than  $p$ . At that end, we resort to some results that are already present in the literature, see again [3] or [5] by Cesaroni and Novaga. To some extent, the reason that leads us to study the extension property of non-local Sobolev spaces is that it allows us to prove Poincaré-Wirtinger and Sobolev inequalities, which, in turn, can be used to obtain a non-local isoperimetric inequality relative to Lipschitz domains that may be employed for the study of (non-local) isoperimetric problems on the whole  $\mathbb{R}^d$ .

We finally point out that [6, Sections 6 and 7] contains results connected to non-local compact embeddings and Poincaré type inequalities.

### 1.1 Non-local Geometric Variational Problems

Several non-local geometric variational problems can be studied applying the theory of non-local Sobolev functions.

A first example is contained in [3], where the space of functions of non-local bounded variation (which coincides with  $W^{K,1}(\mathbb{R}^d)$ ) was investigated and its theory was applied to generalize results for a denoising model with  $L^1$  fidelity and a non-local total variation associated to a singular kernel. These results are related to the fidelity of the model and were first proved in [7] for the fractional kernel. In recent years there has been interest for non-local image denoising problems, see for instance [8–10].

Another class of geometric variational problems involving the fractional kernel can be found in [11, 12] where the main object is a model to describe the shape of a liquid under the action of a bulk energy. In these models the structure of the fractional kernel seems to play a significant role, due to the strong (although partial) regularity theory in this context, which is not clear for more general kernels. We also recall recent non-local models dealing with the study of liquids subjected to non-local interactions [13–17].

Another application is related to the study of the non-local isoperimetric problem, which will be discussed in more detail in Sect. 6. We anticipate that under strong symmetry and monotonicity assumptions on the kernel, the theory is well-known. Nevertheless, as soon as one drops these hypotheses, the situation becomes significantly more complex. Some partial results are contained in [5] for integrable kernels and in [18, 19] for the fractional anisotropic perimeters.

To conclude, we mention the recent work [20] on symmetrization results for general non-local linear elliptic and parabolic problems.

### 1.2 Plan of the Paper

The plan of the paper is the following. In Sect. 2 we first fix some notations and recall some preliminary results.

In Sect. 3 we prove some properties of the space of non-local Sobolev functions. First, the density of smooth functions in non-local Sobolev spaces is proved in Proposition 3.1, then the triviality of such spaces if (Far) is not in force is proved in Proposition 3.3.

Section 4 is devoted to the proof of the extension result (Theorem 4.5) and also contains, as an immediate Corollary, a density result of functions that are smooth up to the boundary of the regular domain  $\Omega$ .

In Sect. 5 we recall some known facts about continuous and compact embeddings for  $W^{K,p}(\mathbb{R}^d)$  and apply our extension theorem to obtain analogous results for  $W^{K,p}(\Omega)$ , see Corollary 5.2 and Corollary 5.4, together with a Poincaré-Wirtinger inequality (Proposition 5.6).

In Sect. 6, the monotonicity with respect to the radius of the non-local perimeter of Euclidean balls is addressed in Proposition 6.7. An explicit formula to compute, in dimension one, the non-local perimeter of balls (i.e., intervals) is the object of Proposition 6.9. An explicit example showing that in general Euclidean balls are not solutions of the non-local isoperimetric problem with volume constraint (even if the kernel is radially symmetric) is contained in Proposition 6.12. As a consequence of the previously obtained Poincaré–Wirtinger inequality, in Corollary 6.13 we furnish a non-local isoperimetric inequality relative to a regular domain  $\Omega$ .

## 2 Notation and Preliminary Results

We fix some basic notations and recall some well-known facts, whose proof can be found, for instance, in [3].

Let  $d \geq 1$ . We put  $E^c := \mathbb{R}^d \setminus E$  for every  $E \subset \mathbb{R}^d$ . For  $E \subset \mathbb{R}^d$  we denote by  $\mathbb{1}_E$  the characteristic function of  $E$ , which is defined by  $\mathbb{1}_E(x) := 1$  if  $x \in E$  and  $\mathbb{1}_E(x) := 0$  if  $x \in E^c$ . For any  $E, F \subset \mathbb{R}^d$ , we put  $\text{dist}(E, F) := \inf \{|x - y|_2 : x \in E, y \in F\}$ .

We let the symbol  $\mathcal{L}^d$  denote the Lebesgue measure of  $\mathbb{R}^d$  and, unless otherwise specified, the term *measurable* is used as a shorthand of  $\mathcal{L}^d$ -measurable and *a.e.* as a shorthand of  $\mathcal{L}^d$ -almost everywhere. If  $E \subset \mathbb{R}^d$  is a measurable set we denote  $|E| := \mathcal{L}^d(E)$ .  $\mathcal{H}^{d-1}$  denotes the Hausdorff measure of dimension  $d - 1$ .

We write  $E \sqcup F$  to denote  $E \cup F$  whenever  $E, F \subset \mathbb{R}^d$  are measurable sets such that  $|E \cap F| = 0$ .

We denote by  $B_r(x) := \{y \in \mathbb{R}^d : |y - x|_2 < r\}$  the open Euclidean ball of radius  $r > 0$  centred at  $x \in \mathbb{R}^d$  and  $B_r := B_r(0)$ . Moreover, we let  $B^m(x) := B_{r(m)}(x)$ , where  $r(m) = (m/|B_1|)^{1/d}$ , be the open Euclidean ball of volume  $m > 0$  centred at  $x \in \mathbb{R}^d$  and  $B^m := B^m(0)$ .

We let the symbol  $\partial E$  denote the topological boundary of a set  $E \subset \mathbb{R}^d$ .

We define the forward difference operator  $\Delta_h : L^1_{\text{loc}}(\mathbb{R}^d) \rightarrow L^1_{\text{loc}}(\mathbb{R}^d)$  as,

$$\Delta_h u(x) := u(x + h) - u(x),$$

for every  $x, h \in \mathbb{R}^d$ .

We then observe that, if  $K$  is a kernel,

$$[u]_{W^{K,p}(\mathbb{R}^d)} = \left( \int_{\mathbb{R}^d} \|\Delta_h u\|_{L^p(\mathbb{R}^d)}^p K(h) \, dh \right)^{\frac{1}{p}}, \quad (2.1)$$

for all  $p \in [1, +\infty)$ .

We say that  $K$  is *symmetric* if

$$K(x) = K(-x) \text{ for all } x \in \mathbb{R}^d, \quad (\text{Sym})$$

$K$  is *strictly positive* if

$$K(x) > 0 \text{ for all } x \in \mathbb{R}^d, \quad (\text{Pos})$$

$K$  has *positive infimum around the origin* if

$$\text{there exist } r, \mu > 0 \text{ such that } K(x) \geq \mu \text{ for all } x \in B_r. \quad (\text{Inf})$$

**Remark 2.1** Let  $K$  be a kernel,  $p \in [1, +\infty)$  and  $\Omega \subset \mathbb{R}^d$  open. We define  $K_{\text{sym}}(x) := (K(x) + K(-x))/2$  for every  $x \in \mathbb{R}^d$  and we observe that  $K_{\text{sym}}$  is a kernel satisfying (Sym), and

$$[u]_{W^{K,p}(\Omega)} = [u]_{W^{K_{\text{sym}},p}(\Omega)},$$

for every  $u : \Omega \rightarrow [-\infty, +\infty]$  measurable and finite a.e.. Furthermore, passing from  $K$  to  $K_{\text{sym}}$ , the assumptions (Far), (Nint), (Nts<sub>p</sub>), (Pos) and (Inf) are preserved.

In view of Remark 2.1, we observe that the hypothesis (Sym) is not restrictive if one is interested in computing (1.1), provided that  $K_{\text{sym}}$  preserves all the initial assumptions satisfied by  $K$ , up to replace  $K$  with the kernel  $K_{\text{sym}}$  defined in Remark 2.1.

### 3 Properties of Non-local Sobolev Functions

In this section we first prove a density result for non-local Sobolev functions and then show that (Far) is essential to make the spaces  $W^{K,p}(\mathbb{R}^d)$  non-trivial. For a general density result we also mention [6, Theorem 3.11].

**Proposition 3.1**  $C^\infty(\mathbb{R}^d) \cap W^{K,p}(\mathbb{R}^d)$  is dense in  $W^{K,p}(\mathbb{R}^d)$  for every  $p \in [1, +\infty)$ .

**Proof** We follow the same strategy as in [21, Theorem 6.62].

Let  $K$  be a kernel and let  $p \in [1, +\infty)$ . Let  $\{\rho_\varepsilon\}_{\varepsilon>0}$  be a family of standard mollifiers and for every  $\varepsilon > 0$  define  $u_\varepsilon := \rho_\varepsilon * u$ . We claim that for every  $\varepsilon > 0$ ,

$$[u_\varepsilon]_{W^{K,p}(\mathbb{R}^d)} \leq [u]_{W^{K,p}(\mathbb{R}^d)}, \quad (3.1)$$

and that

$$\lim_{\varepsilon \rightarrow 0^+} [u_\varepsilon - u]_{W^{K,p}(\mathbb{R}^d)} = 0. \quad (3.2)$$

We define  $D_\varepsilon(h) := \|\Delta_h u_\varepsilon - \Delta_h u\|_{L^p(\mathbb{R}^d)}^p$ ,  $U(h) := \|\Delta_h u\|_{L^p(\mathbb{R}^d)}^p$  and  $U_\varepsilon(h) := \|\Delta_h u_\varepsilon\|_{L^p(\mathbb{R}^d)}^p$  for every  $h \in \mathbb{R}^d$  and  $\varepsilon > 0$ . Thanks to (2.1), for every  $\varepsilon > 0$  we get,

$$[u_\varepsilon - u]_{W^{K,p}(\mathbb{R}^d)}^p = \int_{\mathbb{R}^d} D_\varepsilon(h) K(h) \, dh. \quad (3.3)$$

We note that  $\Delta_h u_\varepsilon = \rho_\varepsilon * \Delta_h u$  for every  $h \in \mathbb{R}^d$  and  $\varepsilon > 0$ , which implies that  $U_\varepsilon(h) \leq U(h)$  for every  $\varepsilon > 0$ . Thus, invoking (2.1) again, (3.1) immediately follows. Moreover,  $\lim_{\varepsilon \rightarrow 0^+} D_\varepsilon(h) = 0$  and  $D_\varepsilon(h) \leq 2^{p-1}(U_\varepsilon(h) + U(h)) \leq 2^p U(h)$  for every  $h \in \mathbb{R}^d$ . Since  $2^p K U \in L^1(\mathbb{R}^d)$ , because of (2.1), it is a domination for the family  $\{K D_\varepsilon\}_{\varepsilon > 0}$  and so we can apply Lebesgue dominated convergence theorem to pass to the limit in (3.3) to get (3.2).  $\square$

If a kernel does not satisfy (Far), it necessarily develops singularities away from the origin (at finite distances or at infinity), as explained in the following:

**Lemma 3.2** *Let  $K$  be a kernel.  $K$  does not satisfy assumption (Far) if and only if either*

$$\int_{B_r^c} K(h) \, dh = +\infty, \quad (3.4)$$

*for every  $r > 0$ , or there exists  $h_0 \in \mathbb{R}^d \setminus \{0\}$  such that*

$$\int_{\Omega} K(h) \, dh = +\infty, \quad (3.5)$$

*for every open neighborhood  $\Omega \subset \mathbb{R}^d$  of  $h_0$ .*

*In particular, if  $K$  satisfies (Far), then  $K \in L^1_{\text{loc}}(\mathbb{R}^d \setminus \{0\})$ .*

**Proof** Let  $K$  be a kernel which does not satisfy assumption (Far).

If  $K \notin L^1(\mathbb{R}^d \setminus B_r)$  for every  $r > 0$ , then (3.4) holds. Therefore, we can assume that there exist  $0 < r < r'$  s.t.  $K \notin L^1(B_{r'} \setminus B_r)$ . By contradiction, assume that for every  $h_0 \in \overline{B_{r'} \setminus B_r}$  there exists  $\Omega$  neighborhood of  $h_0$  such that (3.5) does not hold. By compactness of  $\overline{B_{r'} \setminus B_r}$  we can find a finite open cover of  $B_{r'} \setminus B_r$



such that the integral of  $K$  on each element of the cover is finite. This contradicts the fact that  $K$  is not integrable on  $B_{r'} \setminus B_r$ . Therefore, the thesis holds for some  $h_0 \in \overline{B_{r'}} \setminus \overline{B_r} \subset \mathbb{R}^d \setminus \{0\}$ .

Conversely, if there exists  $h_0 \in \mathbb{R}^d \setminus \{0\}$  such that (3.5) holds for every  $\Omega \subset \mathbb{R}^d$  neighborhood of  $h_0$ , we put  $r := |h_0| > 0$  and observe that  $K \notin L^1(\mathbb{R}^d \setminus B_r)$ . To conclude we simply recall that if we assume (3.4), then (Far) cannot hold by definition.  $\square$

**Proposition 3.3** *Let  $K$  be a kernel which does not satisfy assumption (Far). Then,  $W^{K,p}(\mathbb{R}^d) = \{0\}$  for every  $p \in [1, +\infty)$ .*

**Proof** Let  $K$  be a kernel which does not satisfy assumption (Far) and let  $p \in [1, +\infty)$ .

*Case 1.* Let  $u \in C^0(\mathbb{R}^d) \cap W^{K,p}(\mathbb{R}^d)$ . Having in mind (2.1), we define  $U(h) := \|\Delta_h u\|_{L^p(\mathbb{R}^d)}^p$  for every  $h \in \mathbb{R}^d$  and  $U(\infty) := \int_{\mathbb{R}^d} \liminf_{h \rightarrow \infty} |\Delta_h u(x)|^p dx$ , so that

$$[u]_{W^{K,p}(\mathbb{R}^d)}^p = \int_{\mathbb{R}^d} U(h) K(h) dh.$$

Observe that  $0 \leq U(h) \leq 2^p \|u\|_{L^p(\mathbb{R}^d)}^p$  for all  $h \in \mathbb{R}^d \cup \{\infty\}$  and thanks to Fatou's Lemma it is lower semicontinuous.

By Lemma 3.2 either there exists  $h_0 \in \mathbb{R}^d \setminus \{0\}$  such that (3.5) holds for every  $\Omega$  neighborhood of  $h_0$ , or (3.4) holds for every  $r > 0$ . In the first case, we claim that  $U(h_0) = 0$ . Indeed, if  $U(h_0) > 0$ , by lower semicontinuity, we can find a neighborhood  $\Omega$  of  $h_0$  such that  $U > U(h_0)/2$  on  $\Omega$  and consequently obtain

$$+\infty > 2[u]_{W^{K,p}(\mathbb{R}^d)}^p \geq \int_{\Omega} U(h) K(h) dh \geq \frac{U(h_0)}{2} \int_{\Omega} K(h) dh,$$

which contradicts (3.5). This implies that  $U(h_0) = 0$ . In the second case, we claim that  $U(\infty) = 0$ . Indeed, if  $U(\infty) > 0$ , by lower semicontinuity, we can find  $r > 0$  such that  $U > U(\infty)/2$  on  $B_{r^c}$  and consequently obtain

$$+\infty > 2[u]_{W^{K,p}(\mathbb{R}^d)}^p \geq \int_{B_{r^c}} U(h) K(h) dh \geq \frac{U(\infty)}{2} \int_{B_{r^c}} K(h) dh,$$

which contradicts (3.4)

In the first case, the condition  $U(h_0) = 0$  implies that  $u(x + h_0) = u(x)$  for every  $x \in \mathbb{R}^d$ , in other words  $u$  is periodic with respect to translations of vectors in  $h_0\mathbb{Z}$ . Since  $u \in L^p(\mathbb{R}^d) \cap C^0(\mathbb{R}^d)$ ,  $u \equiv 0$ .

In the second case, the condition  $U(\infty) = 0$  implies that  $\liminf_{h \rightarrow \infty} |\Delta_h u(x)| = 0$  for a.e.  $x \in \mathbb{R}^d$ . If we assume that  $u \not\equiv 0$  then, thanks to the continuity of  $u$  we can find  $x_0 \in \mathbb{R}^d$  such that  $u(x_0) \neq 0$  and  $\liminf_{h \rightarrow \infty} |u(x_0 + h) - u(x_0)| = 0$ . In particular,  $\liminf_{h \rightarrow \infty} |u(x_0 + h)| \geq |u(x_0)| > 0$ , which contradicts the fact that  $u(\cdot + x_0) \in L^p(\mathbb{R}^d)$ .

*Case 2.* Let  $u \in W^{K,p}(\mathbb{R}^d)$ . Thanks to Proposition 3.1 we can find a sequence  $\{u_k\}_{k \in \mathbb{N}} \subset C^\infty(\mathbb{R}^d) \cap W^{K,p}(\mathbb{R}^d)$  such that  $u_k \rightarrow u$  in  $W^{K,p}(\mathbb{R}^d)$ . Since by the previous case we know that  $u_k \equiv 0$  for all  $k \in \mathbb{N}$ , we deduce that  $u \equiv 0$  a.e. in  $\mathbb{R}^d$ .  $\square$

## 4 An Extension Result

In this section we prove the extension result; that is, the continuous embedding of  $W^{K,p}(\Omega)$  in  $W^{K,p}(\mathbb{R}^d)$ , provided that the boundary of  $\Omega$  is regular enough and that the kernel  $K$  satisfies suitable assumptions. Our proof is obtained by adapting the arguments contained in Section 5 of [2] and follows by three preliminary lemmas. The first one furnishes the natural extension of a function in  $W^{K,p}(\Omega)$  having compact support in  $\Omega$ . In the following, we will let  $C$  denote a positive constant that may change from line to line.

**Lemma 4.1** *Let  $K$  be a kernel satisfying (Far),  $\Omega \subset \mathbb{R}^d$  an open set, and  $u \in W^{K,p}(\Omega)$  with  $p \in [1, +\infty)$ . Assume that there exists a compact set  $V \subset \Omega$  such that  $u = 0$  a.e. in  $\Omega \setminus V$ , then the function  $\tilde{u}$  defined as*

$$\tilde{u}(x) := \begin{cases} u(x) & \text{if } x \in \Omega, \\ 0 & \text{if } x \in \Omega^c \end{cases} \quad (4.1)$$

*belongs to  $W^{K,p}(\mathbb{R}^d)$  and there exists a positive constant  $C = C(K, V, \Omega)$  such that*

$$\|\tilde{u}\|_{W^{K,p}(\mathbb{R}^d)} \leq C \|u\|_{W^{K,p}(\Omega)}.$$

**Proof** It holds that  $\|\tilde{u}\|_{L^p(\mathbb{R}^d)} = \|u\|_{L^p(\Omega)}$ . Moreover, there exists  $r > 0$  such that the sets  $x - \Omega^c$  and  $\Omega^c - y$  are contained in  $B_r^c(0)$  for all  $x, y \in V$ , which entails

$$\max \left\{ \int_{x - \Omega^c} K(z) \, dz, \int_{\Omega^c - y} K(z) \, dz \right\} \leq \int_{B_r^c(0)} K(z) \, dz. \quad (4.2)$$

As for the seminorm, we have

$$\begin{aligned}
\int_{\mathbb{R}^d} \int_{\mathbb{R}^d} |\tilde{u}(x) - \tilde{u}(y)|^p K(x-y) \, dx dy &= \int_{\Omega} \int_{\Omega} |u(x) - u(y)|^p K(x-y) \, dx dy \\
&\quad + \int_{\Omega} |u(x)|^p \left\{ \int_{\Omega^c} K(x-y) \, dy \right\} \, dx \\
&\quad + \int_{\Omega} |u(y)|^p \left\{ \int_{\Omega^c} K(x-y) \, dx \right\} \, dy \\
&= [u]_{W^{K,p}(\Omega)}^p + \int_V |u(x)|^p \left\{ \int_{x-\Omega^c} K(z) \, dz \right\} \, dx \\
&\quad + \int_V |u(y)|^p \left\{ \int_{\Omega^c-y} K(z) \, dz \right\} \, dy \\
&\leq [u]_{W^{K,p}(\Omega)}^p + 2\|u\|_{L^p(\Omega)}^p \int_{B_r^c(0)} K(z) \, dz,
\end{aligned}$$

where we performed two times the change of variables  $z := x - y$ , and the last inequality follows by (4.2). Finally, the thesis follows by (Far).  $\square$

The following lemma allows us to extend functions to a larger set  $\Omega$  provided that this can be obtained by reflecting the initial domain with respect to a hyperplane. We will say that an open set  $\Omega \subset \mathbb{R}^d$  is symmetric with respect to the coordinate  $x_d$  if for every  $x = (x_1, \dots, x_d) \in \Omega$ , then  $(x_1, \dots, -x_d) \in \Omega$ .

**Lemma 4.2** *Let  $\Omega \subset \mathbb{R}^d$  be an open set symmetric with respect to the coordinate  $x_d$  and let  $K$  be a kernel satisfying (Far),  $(\text{Dec}(c_0, |\cdot|_*))$ , and  $(\text{Dou}(D, |\cdot|_*))$ . Consider the sets  $\Omega_+ := \{x \in \Omega : x_d > 0\}$  and  $\Omega_- := \Omega \setminus \Omega_+$ , and let  $u \in W^{K,p}(\Omega_+)$  with  $p \in [1, +\infty)$ . Then the function  $\bar{u}$  defined as*

$$\bar{u}(x) := \begin{cases} u(x', x_d) & \text{if } x \in \Omega_+ \\ u(x', -x_d) & \text{if } x \in \Omega_- \end{cases}, \quad (4.3)$$

*belongs to  $W^{K,p}(\Omega)$  and there exists a positive constant  $C = C(c_0, C_D, |\cdot|_*, K)$  such that*

$$\|\bar{u}\|_{W^{K,p}(\Omega)} \leq C \|u\|_{W^{K,p}(\Omega_+)}.$$

**Proof** It is immediate to observe that  $\|u\|_{L^p(\Omega)}^p = 2\|u\|_{L^p(\Omega_+)}^p$ . As for the seminorms, it holds

$$\begin{aligned}
& \int_{\Omega} \int_{\Omega} |\bar{u}(x) - \bar{u}(y)|^p K(x - y) \, dx dy \\
&= [u]_{W^{\kappa,p}(\Omega_+)}^p + \int_{\Omega_+} \int_{\Omega_-} |u(x) - u(y', -y_d)|^p K(x - y) \, dx dy \\
&\quad + \int_{\Omega_-} \int_{\Omega_+} |u(x', -x_d) - u(y)|^p K(x - y) \, dx dy \\
&\quad + \int_{\Omega_-} \int_{\Omega_-} |u(x', -x_d) - u(y', -y_d)|^p K(x - y) \, dx dy.
\end{aligned}$$

We estimate separately the last three addends.

By the change of variables  $z := (y', -y_d)$ , we have

$$\begin{aligned}
& \int_{\Omega_+} \int_{\Omega_-} |u(x) - u(y', -y_d)|^p K(x - y) \, dx dy \\
&= \int_{\Omega_+} \int_{\Omega_+} |u(x) - u(z)|^p K((x_1 - z_1, \dots, x_{d-1} - z_{d-1}, x_d + z_d)) \, dx dz.
\end{aligned}$$

Note that, since  $y = (z', -z_d)$ ,

$$|x - y|_2 = |(x_1 - z_1, \dots, x_{d-1} - z_{d-1}, x_d + z_d)|_2 \geq |x - z|_2,$$

hence, by the equivalence of the norms, there exists a positive constant  $c_*$  such that

$$c_* |(x_1 - z_1, \dots, x_{d-1} - z_{d-1}, x_d + z_d)|_* \geq |x - z|_*,$$

and then, by  $(\text{Dec}(c_0, |\cdot|_*))$ , we obtain

$$\begin{aligned}
& \int_{\Omega_+} \int_{\Omega_+} |u(x) - u(z)|^p K((x_1 - z_1, \dots, x_{d-1} - z_{d-1}, x_d + z_d)) \, dx dz \\
&\leq \frac{1}{c_0} \int_{\Omega_+} \int_{\Omega_+} |u(x) - u(z)|^p K\left(\frac{x - z}{c_*}\right) \, dx dz.
\end{aligned}$$

Now we argue differently for short-range and long-range interactions. Let  $m \in \mathbb{N}^+$  be such that

$$\frac{2^m}{c_*} \geq 1.$$

Applying  $m$  times  $(\text{Dou}(D, |\cdot|_*))$ , we deduce that for every  $x, z \in \mathbb{R}^d$  s.t.  $|x - z|_* \leq \frac{c_* D}{2^{m-1}}$ ,

$$K\left(\frac{x - z}{c_*}\right) \leq C_D^m K\left(\frac{2^m(x - z)}{c_*}\right), \quad (4.4)$$

where the iterated application of  $(\text{Dou}(D, |\cdot|_*))$  is made possible by the fact that

$$\frac{2^k}{c_*} |x - z|_* \leq D$$

for all  $k \in \{0, \dots, m-1\}$ .

Applying (4.4), and then applying  $(\text{Dec}(c_0, |\cdot|_*))$ , we obtain

$$\begin{aligned} & \iint_{\Omega_+^2 \cap \{|x-z|_* \leq \frac{c_* D}{2^{m-1}}\}} |u(x) - u(z)|^p K\left(\frac{x - z}{c_*}\right) dx dz \\ & \leq C_D^m \int_{\Omega_+} \int_{\Omega_+} |u(x) - u(z)|^p K\left(\frac{2^m(x - z)}{c_*}\right) dx dz \\ & \leq \frac{C_D^m}{c_0} \int_{\Omega_+} \int_{\Omega_+} |u(x) - u(z)|^p K(x - z) dx dz. \end{aligned} \quad (4.5)$$

As for long-range interactions, by the change of variables  $w := \frac{x-z}{c_*}$ , we have

$$\begin{aligned} & \iint_{\Omega_+^2 \cap \{|x-z|_* > \frac{c_* D}{2^{m-1}}\}} |u(x) - u(z)|^p K\left(\frac{x - z}{c_*}\right) dx dz \\ & \leq \int_{\Omega_+} 2^{p-1} |u(x)|^p \int_{\{z \in \mathbb{R}^d: |x-z|_* > \frac{c_* D}{2^{m-1}}\}} K\left(\frac{x - z}{c_*}\right) dz dx \\ & \quad + \int_{\Omega_+} 2^{p-1} |u(z)|^p \int_{\{x \in \mathbb{R}^d: |x-z|_* > \frac{c_* D}{2^{m-1}}\}} K\left(\frac{x - z}{c_*}\right) dx dz \\ & \leq 2^p \|u\|_{L^p(\Omega_+)}^p c_*^d \int_{\{w \in \mathbb{R}^d: |w|_* > \frac{D}{2^{m-1}}\}} K(w) dw. \end{aligned} \quad (4.6)$$

Gathering (4.5) and (4.6), and using (Far) together with the equivalence of the norms, we obtain

$$\int_{\Omega_+} \int_{\Omega_-} |u(x) - u(y', -y_d)|^p K(x - y) dx dy \leq C \|u\|_{W^{K,p}(\Omega_+)}, \quad (4.7)$$

$C$  being a positive constant depending on  $K, c_*, C_D$ , and  $c_0$ .

With the same argument, we get

$$\int_{\Omega_-} \int_{\Omega_+} |u(x', -x_d) - u(y)|^p K(x - y) \, dx dy \leq C \|u\|_{W^{K,p}(\Omega_+)}^p. \quad (4.8)$$

Finally, we treat the last summand. By the changes of variables  $z := (x', -x_d)$  and  $w := (y', -y_d)$ , we have

$$\begin{aligned} & \int_{\Omega_-} \int_{\Omega_-} |u(x', -x_d) - u(y', -y_d)|^p K(x - y) \, dx dy \\ &= \int_{\Omega_+} \int_{\Omega_+} |u(z) - u(w)|^p K((z_1 - w_1, \dots, z_{d-1} - w_{d-1}, -z_d + w_d)) \, dz dw. \end{aligned}$$

As before, we obtain that

$$c_* |(z_1 - w_1, \dots, z_{d-1} - w_{d-1}, -z_d + w_d)|_* \geq |z - w|_*,$$

and then, by  $(\text{Dec}(c_0, |\cdot|_*))$ , we get

$$\begin{aligned} & \int_{\Omega_+} \int_{\Omega_+} |u(z) - u(w)|^p K((z_1 - w_1, \dots, z_{d-1} - w_{d-1}, -z_d + w_d)) \, dz dw \\ & \leq \frac{1}{c_0} \int_{\Omega_+} \int_{\Omega_+} |u(z) - u(w)|^p K\left(\frac{z - w}{c_*}\right) \, dz dw. \end{aligned}$$

Reasoning as for the first two addends, we obtain

$$\int_{\Omega_-} \int_{\Omega_-} |u(x', -x_d) - u(y', -y_d)|^p K(x - y) \, dx dy \leq C \|u\|_{W^{K,p}(\Omega_+)}^p, \quad (4.9)$$

$C$  being a positive constant depending on  $K, c_*, C_D$ , and  $c_0$ . Putting together (4.7), (4.8), and (4.9) we obtain the thesis.  $\square$

**Remark 4.3** If the norm  $|\cdot|_*$  is symmetric with respect to the hyperplane  $\{x_d = 0\}$  in the sense that

$$|(x_1, \dots, x_d)|_* = |(x_1, \dots, -x_d)|_* \quad (4.10)$$

for all  $x \in \mathbb{R}^d$ , we do not need the doubling assumption in Lemma 4.2. Indeed, let  $x \in \Omega_+, y \in \Omega_-$ , and  $z := (y', -y_d) \in \Omega_+$ . There exists  $\lambda \in [0, 1]$  such that

$$\begin{aligned} z - x &= (y_1 - x_1, \dots, y_{d-1} - x_{d-1}, -y_d - x_d) \\ &= (y_1 - x_1, \dots, y_{d-1} - x_{d-1}, \lambda(y_d - x_d) + (1 - \lambda)(x_d - y_d)); \end{aligned}$$

then, by the convexity of the norm, we obtain

$$\begin{aligned} |z - x|_* &\leq \lambda|(y_1 - x_1, \dots, y_{d-1} - x_{d-1}, y_d - x_d)|_* \\ &\quad + (1 - \lambda)|(y_1 - x_1, \dots, y_{d-1} - x_{d-1}, x_d - y_d)|_* \\ &= |y - x|_*, \end{aligned}$$

where the last equality follows by (4.10). The above inequality allows to carry out the proof of the previous lemma without making use of  $(\text{Dou}(D, |\cdot|_*))$ . Moreover, this observation simplifies the argument as it allows to not reasoning differently for short-range and long-range interactions. Note that, if the norm  $|\cdot|_*$  is symmetric with respect to a different hyperplane  $\Pi$ , then the above lemma is still valid without requiring  $(\text{Dou}(D, |\cdot|_*))$  provided that the set  $\Omega$  can be obtained by reflecting the set  $\Omega_+$  with respect to  $\Pi$ .

The last lemma is needed to resort to an argument involving partition of unity that we will employ in the proof of the extension result.

**Lemma 4.4** *Let  $K$  be a kernel satisfying  $(\text{Nts}_p)$ ,  $\Omega \subset \mathbb{R}^d$  an open set,  $u \in W^{K,p}(\Omega)$  with  $p \in [1, +\infty)$ , and  $\psi$  a Lipschitz continuous function compactly supported in  $\Omega$  such that  $0 \leq \psi \leq 1$ . Then the function  $\psi u$  belongs to  $W^{K,p}(\Omega)$  and, letting  $\text{Lip } \psi$  denote the Lipschitz constant of  $\psi$ , there exists a positive constant  $C = C(\text{Lip } \psi, K)$  such that*

$$\|\psi u\|_{W^{K,p}(\Omega)} \leq C \|u\|_{W^{K,p}(\Omega)}.$$

**Proof** Since  $0 \leq \psi \leq 1$ , it holds that  $\|\psi u\|_{L^p(\Omega)} \leq \|u\|_{L^p(\Omega)}$ . By adding and subtracting  $\psi(x)u(y)$ , we evaluate the seminorm as follows,

$$\begin{aligned} &\int_{\Omega} \int_{\Omega} |\psi u(x) - \psi u(y)|^p K(x - y) \, dx dy \\ &\leq 2^{p-1} \left\{ \int_{\Omega} \int_{\Omega} |\psi(x)|^p |u(x) - u(y)|^p K(x - y) \, dx dy \right. \\ &\quad \left. + \int_{\Omega} \int_{\Omega} |u(y)|^p |\psi(x) - \psi(y)|^p K(x - y) \, dx dy \right\} \\ &\leq 2^{p-1} [u]_{W^{K,p}(\Omega)}^p \\ &\quad + 2^{p-1} \int_{\Omega} \int_{\Omega} |u(y)|^p |\psi(x) - \psi(y)|^p K(x - y) \, dx dy. \end{aligned}$$

By treating differently short-range and long-range interactions, we obtain

$$\begin{aligned} & \iint_{\Omega^2 \cap \{|x-y|_2 \leq 1\}} |u(y)|^p |\psi(x) - \psi(y)|^p K(x-y) \, dx dy \\ & \leq (\text{Lip } \psi)^p \|u\|_{L^p(\Omega)}^p \int_{B_1(0)} K(z) |z|_2^p \, dz, \end{aligned}$$

and, since  $|\psi(x) - \psi(y)|^p \leq 1$  for all  $x, y \in \Omega$  thanks to the assumption  $0 \leq \psi \leq 1$ ,

$$\begin{aligned} & \iint_{\Omega^2 \cap \{|x-y|_2 > 1\}} |u(y)|^p |\psi(x) - \psi(y)|^p K(x-y) \, dx dy \\ & \leq \int_{\Omega} |u(y)|^p \int_{B_1^c(y)} K(x-y) \, dx dy = \|u\|_{L^p(\Omega)}^p \int_{B_1^c(0)} K(z) \, dz. \end{aligned}$$

Putting together these estimates, and making use of  $(\text{Nts}_p)$ , which implies (Far), the proof is concluded.  $\square$

Finally, we prove our main result.

**Theorem 4.5** *Let  $K$  be a kernel satisfying  $(\text{Dec}(c_0, |\cdot|_*))$ ,  $(\text{Dou}(D, |\cdot|_*))$ , and  $(\text{Nts}_p)$ ,  $\Omega \subset \mathbb{R}^d$  a bounded, open set with Lipschitz boundary and  $p \in [1, +\infty)$ . Then  $W^{K,p}(\Omega)$  is continuously embedded in  $W^{K,p}(\mathbb{R}^d)$ ; that is, for every  $u \in W^{K,p}(\Omega)$  there exists  $\tilde{u} \in W^{K,p}(\mathbb{R}^d)$  such that the restriction of  $\tilde{u}$  in  $\Omega$  equals  $u$  and there exists a positive constant  $C = C(c_0, C_D, K, \Omega, |\cdot|_*)$  such that*

$$\|\tilde{u}\|_{W^{K,p}(\mathbb{R}^d)} \leq C \|u\|_{W^{K,p}(\Omega)}.$$

**Proof** Consider  $\{B_j\}_{j=1}^M$  a finite cover of  $\partial\Omega$  made up of open Euclidean balls,  $\partial\Omega \subset \bigcup_{j=1}^M B_j$ . Let  $\{\psi_j\}_{j=0}^{M+1}$  be a smooth partition of unity such that  $\text{supp } \psi_0 \subset \Omega$ ,  $\text{supp } \psi_j \subset B_j$  for every  $j \in \{1, \dots, M\}$ ,  $\text{supp } \psi_{M+1} \subset \mathbb{R}^d \setminus \Omega$  and  $\sum_{j=0}^{M+1} \psi_j = \sum_{j=0}^M \psi_j = 1$  in  $\Omega$ .

By Lemma 4.4,  $\psi_0 u \in W^{K,p}(\Omega)$  and, by Lemma 4.1, the extension  $\widetilde{\psi_0 u}$  belongs to  $W^{K,p}(\mathbb{R}^d)$  with

$$\|\widetilde{\psi_0 u}\|_{W^{K,p}(\mathbb{R}^d)} \leq C \|u\|_{W^{K,p}(\Omega)}, \quad (4.11)$$

where  $C$  depends on  $\psi_0, \Omega, K$ .

Now, we consider the restriction of  $u$  to  $\Omega \cap B_j$ . We consider a bi-Lipschitz change of variables  $T_j : Q \rightarrow B_j$  mapping  $Q_+$  into  $\Omega \cap B_j$ , where we set

$$Q := \left(-\frac{1}{2}, \frac{1}{2}\right)^d \quad \text{and} \quad Q_+ := Q \cap \{x_d > 0\},$$



and we define  $v_j(y) := u(T_j(y))$ ,  $y \in Q_+$ . We aim at proving that  $v_j \in W^{K,p}(Q_+)$ . By the change of variables  $x := T_j^{-1}(y)$ , we have that  $\|v_j\|_{L^p(Q_+)} \leq C\|u\|_{L^p(\Omega \cap B_j)}$ , with depending on  $T_j$ . As for the seminorms, we observe that there exists  $L_j > 0$  such that

$$\frac{1}{L_j} |T_j^{-1}(x) - T_j^{-1}(y)|_2 \geq |x - y|_2,$$

that, by the equivalence of the norms, implies

$$\frac{c_*}{L_j} |T_j^{-1}(x) - T_j^{-1}(y)|_* \geq |x - y|_*. \quad (4.12)$$

The changes of variables  $x := T_j(\hat{x})$  and  $y := T_j(\hat{y})$ , (4.12), and  $(\text{Dec}(c_0, |\cdot|_*))$ , lead to

$$\begin{aligned} & \int_{Q_+} \int_{Q_+} |v_j(\hat{x}) - v_j(\hat{y})|^p K(\hat{x} - \hat{y}) \, d\hat{x} d\hat{y} \\ &= \int_{Q_+} \int_{Q_+} |u(T_j(\hat{x})) - u(T_j(\hat{y}))|^p K(\hat{x} - \hat{y}) \, d\hat{x} d\hat{y} \\ &= \int_{\Omega \cap B_j} \int_{\Omega \cap B_j} |u(x) - u(y)|^p K(T_j^{-1}(x) - T_j^{-1}(y)) |\det T_j^{-1}|^2 \, dx dy \\ &\leq \frac{C}{c_0} \int_{\Omega \cap B_j} \int_{\Omega \cap B_j} |u(x) - u(y)|^p K\left(\frac{L_j}{c_*}(x - y)\right) \, dx dy, \end{aligned}$$

for some positive constant depending on  $T_j$ . Now we argue similarly to the proof of Lemma 4.2. We fix  $m_j \in \mathbb{N}$  such that

$$\frac{2^{m_j} L_j}{c_*} \geq 1,$$

and we apply  $m_j$  times assumption  $(\text{Dou}(D, |\cdot|_*))$  to estimate the contributions of the short-range interactions by

$$\begin{aligned}
& \iint_{(\Omega \cap B_j)^2 \cap \{|x-y|_* \leq \frac{c_* D}{L_j 2^{m_j-1}}\}} |u(x) - u(y)|^p K\left(\frac{L_j}{c_*}(x-y)\right) dx dy \\
& \leq C_D^{m_j} \iint_{(\Omega \cap B_j)^2 \cap \{|x-y|_* \leq \frac{c_* D}{L_j 2^{m_j-1}}\}} |u(x) - u(y)|^p K\left(2^{m_j} \frac{L_j}{c_*}(x-y)\right) dx dy \\
& \leq \frac{C_D^{m_j}}{c_0} \iint_{(\Omega \cap B_j)^2 \cap \{|x-y|_* \leq \frac{c_* D}{L_j 2^{m_j-1}}\}} |u(x) - u(y)|^p K(x-y) dx dy \\
& \leq \frac{C_D^{m_j}}{c_0} [u]_{W^{K,p}(\Omega)}^p.
\end{aligned} \tag{4.13}$$

Analogously, long-range interactions can be estimated as follows:

$$\begin{aligned}
& \iint_{(\Omega \cap B_j)^2 \cap \{|x-y|_* > \frac{c_* D}{L_j 2^{m_j-1}}\}} |u(x) - u(y)|^p K\left(\frac{L_j}{c_*}(x-y)\right) dx dy \\
& \leq 2^{p-1} \|u\|_{L^p(\Omega)}^p \int_{\{z \in \mathbb{R}^d : |z|_* > \frac{c_* D}{L_j 2^{m_j-1}}\}} K\left(\frac{L_j}{c_*} z\right) dz \\
& = 2^{p-1} \left(\frac{c_*}{L_j}\right)^d \|u\|_{L^p(\Omega)}^p \int_{\{z \in \mathbb{R}^d : |z|_* > \frac{D}{2^{m_j-1}}\}} K(z) dz.
\end{aligned} \tag{4.14}$$

Gathering (4.13) and (4.14), and using (Far), we infer that  $v_j \in W^{K,p}(Q_+)$  with

$$\|v_j\|_{W^{K,p}(Q_+)} \leq C \|u\|_{W^{K,p}(\Omega)}, \tag{4.15}$$

where  $C$  depends on  $T_j$ ,  $c_0$ ,  $c_*$ ,  $C_D$  and  $K$ .

We are in position to apply Lemma 4.2 in order to obtain a function  $\bar{v}_j \in W^{K,p}(Q)$  with

$$\|\bar{v}_j\|_{W^{K,p}(Q)} \leq C \|v_j\|_{W^{K,p}(Q_+)}. \tag{4.16}$$

We define  $w_j(x) := \bar{v}_j(T_j^{-1}(x))$  for  $x \in B_j$  and, arguing as for the function  $v_j$ , we obtain that  $w_j \in W^{K,p}(B_j)$ , with

$$\|w_j\|_{W^{K,p}(B_j)} \leq C \|\bar{v}_j\|_{W^{K,p}(Q)}, \tag{4.17}$$

for some  $C$  depending on  $T_j$ ,  $c_0$ ,  $c_*$ ,  $C_D$ , and  $K$ .

Therefore, combining Lemmas 4.4 and 4.1, we have that  $\widetilde{\psi_j w_j} \in W^{K,p}(\mathbb{R}^d)$  with

$$\|\widetilde{\psi_j w_j}\|_{W^{K,p}(\mathbb{R}^d)} \leq C \|u\|_{W^{K,p}(\Omega)}, \tag{4.18}$$

with the constant  $C$  that, also by (4.15), (4.16) and (4.17), depends on  $T_j$ ,  $c_0$ ,  $c_*$ ,  $C_D$ ,  $\text{Lip } \psi_j$ ,  $K$ , and  $\Omega$ . In conclusion, the function

$$\tilde{u} := \widetilde{\psi_0 u} + \sum_{j=1}^M \widetilde{\psi_j w_j}$$

is the desired extension as it coincides with  $u$  on  $\Omega$  and, gathering (4.11) and (4.18), it satisfies

$$\|\tilde{u}\|_{W^{K,p}(\mathbb{R}^d)} \leq C \|u\|_{W^{K,p}(\Omega)},$$

where the positive constant depends on  $C_D$ ,  $c_0$ ,  $c_*$ , the kernel  $K$ , and the set  $\Omega$  through the Lipschitz continuous changes of variables  $\{T_j\}_{j=1}^M$  and the smooth functions  $\{\psi_j\}_{j=0}^M$ .  $\square$

As a first application of Theorem 4.5, we combine it with Proposition 3.1 and immediately obtain the following result.

**Corollary 4.6** *Let  $K$  be a kernel satisfying  $(\text{Dec}(c_0, |\cdot|_*))$ ,  $(\text{Dou}(D, |\cdot|_*))$ , and  $(\text{Nts}_p)$ ,  $\Omega \subset \mathbb{R}^d$  a bounded, open set with Lipschitz boundary and  $p \in [1, +\infty)$ . Then,  $C^\infty(\overline{\Omega}) \cap W^{K,p}(\Omega)$  is dense in  $W^{K,p}(\Omega)$  for every  $p \in [1, +\infty)$ .*

## 5 Sobolev and Poincaré Inequalities

In this section, we present a compactness result in  $W^{K,p}(\Omega)$  together with some estimates in the spirit of Poincaré–Wirtinger and Sobolev inequalities on bounded, open, regular subsets of  $\mathbb{R}^d$ . At that end, we recall some results that are already present in the literature starting with a result of local compactness which can be found in [3, Theorem 2.11] and [6, Theorem 6.2]. In the following, we will let  $C$  denote a positive constant that may change from line to line.

**Theorem 5.1** *Let  $K$  be a kernel satisfying  $(\text{Nint})$  and  $(\text{Far})$  and let  $\{u_k\}_k \subset W^{K,p}(\mathbb{R}^d)$  with  $p \in [1, +\infty)$  be a sequence such that*

$$\sup_k \|u_k\|_{W^{K,p}(\mathbb{R}^d)} < +\infty.$$

*Then there exist a subsequence  $\{k_j\}_j$  and  $u \in W^{K,p}(\mathbb{R}^d)$  such that*

$$u_{k_j} \rightarrow u \text{ in } L^p_{\text{loc}}(\mathbb{R}^d) \text{ as } j \rightarrow +\infty.$$

Having at disposal our extension theorem, we prove the following.

**Corollary 5.2** *Let be a kernel satisfying (Nint),  $(\text{Dec}(c_0, |\cdot|_*))$ ,  $(\text{Dou}(D, |\cdot|_*))$ , and  $(\text{Nts}_p)$ ,  $\Omega \subset \mathbb{R}^d$  a bounded, open set with Lipschitz boundary, and  $p \in [1, +\infty)$ . Then  $W^{K,p}(\Omega)$  is compactly embedded in  $L^p(\Omega)$ ; that is, if  $\{u_k\}_k \subset W^{K,p}(\Omega)$  is such that*

$$\sup_k \|u_k\|_{W^{K,p}(\Omega)} < +\infty,$$

*there exist a subsequence  $\{k_j\}_j$  and  $u \in W^{K,p}(\Omega)$  such that*

$$u_{k_j} \rightarrow u \text{ in } L^p(\Omega) \text{ as } j \rightarrow +\infty.$$

**Proof** Consider a sequence  $\{u_k\}_k$  which is bounded in  $W^{K,p}(\Omega)$ . First, we apply Theorem 4.5 to obtain a sequence  $\{\tilde{u}_k\}_k \subset W^{K,p}(\mathbb{R}^d)$  such that

$$\sup_k \|\tilde{u}_k\|_{W^{K,p}(\mathbb{R}^d)} < +\infty,$$

then, by Theorem 5.1 we find a subsequence  $\{k_j\}_j$  and  $u \in W^{K,p}(\mathbb{R}^d)$  such that

$$\tilde{u}_{k_j} \rightarrow u \text{ in } L^p_{\text{loc}}(\mathbb{R}^d) \text{ as } j \rightarrow +\infty.$$

This, in turn, implies  $u_{k_j} \rightarrow u$  in  $L^p(\Omega)$ , which concludes the proof.  $\square$

If  $\Omega \subset \mathbb{R}^d$  is a measurable set of finite measure we put  $\Omega^* = B^{|\Omega|}$ . If  $u : \mathbb{R}^d \rightarrow [-\infty, +\infty]$  is a measurable function such that  $|\{|u| > t\}| < +\infty$  for all  $t > 0$ , we define  $u^*$ , the *symmetric-decreasing rearrangement* of  $u$ , by

$$u^*(x) := \int_0^{+\infty} \mathbb{1}_{\{|u|>t\}^*}(x) dt,$$

for every  $x \in \mathbb{R}^d$ .

We illustrate a Sobolev-type inequality that is presented in [5] and we immediately deduce an analogous result on bounded, regular domains through the extension result.

**Proposition 5.3** *Assume that there exist  $q \in [1, +\infty)$  and a positive constant such that,*

$$\mathcal{P}_{K^*}(B^m) \geq \frac{m^{\frac{1}{q}}}{C} \quad \text{for every } m > 0. \quad (5.1)$$

*Then for every  $u \in W^{K,1}(\mathbb{R}^d)$  it holds that*

$$\|u\|_{L^q(\mathbb{R}^d)} \leq C[u]_{W^{K,1}(\mathbb{R}^d)}.$$

**Corollary 5.4** *Let  $K$  be a kernel satisfying  $(\text{Dec}(c_0, |\cdot|_*))$ ,  $(\text{Dou}(D, |\cdot|_*))$ , and  $(\text{Nts}_1)$ ,  $\Omega \subset \mathbb{R}^d$  a bounded, open set with Lipschitz boundary, and assume that (5.1) holds for some  $q \in [1, +\infty)$ . Then  $W^{K,1}(\Omega)$  is continuously embedded in  $L^q(\Omega)$ ; that is, there exists a positive constant such that*

$$\|u\|_{L^q(\Omega)} \leq C\|u\|_{W^{K,1}(\Omega)}$$

for every  $u \in W^{K,1}(\Omega)$ .

**Proof** Given  $u \in W^{K,1}(\Omega)$ , by Theorem 4.5 there exists an extension  $\tilde{u} \in W^{K,1}(\mathbb{R}^d)$  such that

$$\|\tilde{u}\|_{W^{K,1}(\mathbb{R}^d)} \leq C\|u\|_{W^{K,1}(\Omega)},$$

then, by Proposition 5.3 we obtain

$$\|\tilde{u}\|_{L^q(\mathbb{R}^d)} \leq C\|u\|_{W^{K,1}(\Omega)}.$$

Since  $\tilde{u}$  coincides with  $u$  on  $\Omega$ , the thesis follows.  $\square$

In view of the study of non-local isoperimetric problems related to the kernel  $K$ , a useful tool is a relative isoperimetric inequality. Such estimate is a particular case of a Poincaré-Wirtinger inequality, that in turn, can be obtained through the embedding results that we already proved. We premise a useful remark on constant functions.

**Remark 5.5** We observe that, in general,  $[u]_{W^{K,p}(\Omega)} = 0$  does not imply that  $u$  is constant a.e. on  $\Omega$ . For instance, if  $\Omega$  is made up of two disjoint balls whose distance is larger than the diameter of the support of  $K$ , then  $[u]_{W^{K,p}(\Omega)} = 0$  for every locally constant function  $u$ . However, if  $\Omega$  is connected and  $K$  satisfies (Nint), (Far), and  $(\text{Dec}(c_0, |\cdot|_*))$ , such property holds true. To see this, it suffices to show that  $[u]_{W^{K,p}(\Omega)} = 0$  implies  $u$  is locally constant. Let  $x_0 \in \Omega$  and let  $r > 0$  such that  $B_{2r}(x_0) \subset \subset \Omega$ , then

$$0 = [u]_{W^{K,p}(\Omega)} \geq \int_{B_r(0)} K(h) \left\{ \int_{B_r(x_0)} |u(x+h) - u(x)|^p dx \right\} dh.$$

As a consequence of (Nint), (Far), and  $(\text{Dec}(c_0, |\cdot|_*))$ , we deduce that (Inf) holds. This fact, combined with the above inequality, implies (if  $r > 0$  is sufficiently small)

$$0 = \int_{B_r(0)} \left\{ \int_{B_r(x_0)} |u(x+h) - u(x)|^p dx \right\} dh,$$

which in turn yields  $u(x+h) = u(x)$  for a.e.  $h \in B_r(0)$  and  $x \in B_r(x_0)$ ; i.e.,  $u(x) = u(y)$  for a.e.  $x, y \in B_{2r}(x_0)$ . Therefore,  $u$  is a.e. equal to a constant in a neighborhood of  $x_0$ .

In order to simplify the notations in the following, we set

$$u_\Omega := \frac{1}{|\Omega|} \int_{\Omega} u \, dx,$$

for every  $u \in W^{K,1}(\Omega)$ .

**Proposition 5.6** *Let  $\Omega \subset \mathbb{R}^d$  a bounded, connected, open set with Lipschitz boundary, and  $p \in [1, +\infty)$ . Let  $K$  be a kernel satisfying (Nint), (Dec( $c_0, |\cdot|_*$ )), (Dou( $D, |\cdot|_*$ )), and (Nts $_p$ ). Then there exists a positive constant  $C$  such that*

$$\left( \int_{\Omega} |u - u_\Omega|^p \, dx \right)^{\frac{1}{p}} \leq C[u]_{W^{K,p}(\Omega)} \quad (5.2)$$

for every  $u \in W^{K,p}(\Omega)$ . Moreover, if (5.1) holds for some  $q \in [1, +\infty)$ , then there exists a positive constant  $C$  such that

$$\left( \int_{\Omega} |u - u_\Omega|^q \, dx \right)^{\frac{1}{q}} \leq C[u]_{W^{K,1}(\Omega)} \quad (5.3)$$

for every  $u \in W^{K,1}(\Omega)$ .

**Proof** The proof of (5.2) is obtained by the usual contradiction-compactness argument. If (5.2) fails, there exists a sequence  $\{u_k\}_k \subset W^{K,p}(\Omega)$  such that for all  $k \in \mathbb{N}$  it holds  $(u_k)_\Omega = 0$ ,  $\|u_k\|_{L^p(\Omega)} = 1$ , and moreover

$$[u_k]_{W^{K,p}(\Omega)} \rightarrow 0 \quad \text{as } k \rightarrow +\infty.$$

Applying Corollary 5.2 we deduce there exists  $u \in W^{K,p}(\Omega)$  such that, up to a not relabelled subsequence,  $u_k \rightarrow u$  in  $L^p(\Omega)$ , and  $u_k \rightarrow u$  a.e. in  $\Omega$ . We claim that  $u$  is constant; indeed, by Fatou's Lemma,

$$0 = \liminf_{k \rightarrow +\infty} \int_{\Omega} \int_{\Omega} |u_k(x) - u_k(y)|^p K(x-y) \, dx dy \geq \int_{\Omega} \int_{\Omega} |u(x) - u(y)|^p K(x-y) \, dx dy$$

so that Remark 5.5 and the connectedness of  $\Omega$  imply the claim. Since  $u_\Omega = 0$ , we infer  $u = 0$ , which contradicts  $\|u\|_{L^p(\Omega)} = 1$ .

The proof of (5.3) is similar. Arguing again by contradiction, there exists a sequence  $\{u_k\}_k \subset W^{K,1}(\Omega)$  such that for all  $k \in \mathbb{N}$  it holds  $(u_k)_\Omega = 0$  and  $\|u_k\|_{L^q(\Omega)} = 1$ , and moreover

$$[u_k]_{W^{K,1}(\Omega)} \rightarrow 0 \quad \text{as } k \rightarrow +\infty.$$

Note that, by (5.2), this implies  $u_k \rightarrow 0$  in  $L^1(\Omega)$ , which in turn yields  $u_k \rightarrow 0$  in  $W^{K,1}(\Omega)$  as  $k \rightarrow +\infty$ . Applying Corollary 5.4, we infer that  $u_k \rightarrow 0$  in  $L^q(\Omega)$  as  $k \rightarrow +\infty$ , which contradicts the fact that  $\|u_k\|_{L^q(\Omega)} = 1$  for all  $k \in \mathbb{N}$ , concluding the proof.  $\square$

**Remark 5.7** If  $K$  satisfies (Pos) and  $(\text{Dec}(c_0, |\cdot|_*))$ , the proof of (5.2) follows by a straightforward computation without requiring any additional assumption either on  $K$  or on  $\partial\Omega$ . Indeed, we put  $d = \text{diam } \Omega$  and obtain

$$\begin{aligned} \int_{\Omega} |u - u_{\Omega}|^p dx &\leq \frac{1}{|\Omega|} \int_{\Omega} \int_{\Omega} |u(x) - u(y)|^p dx dy \\ &\leq \frac{1}{|\Omega| \inf\{K(z) : z \in B_d(0)\}} \int_{\Omega} \int_{\Omega} |u(x) - u(y)|^p K(x - y) dx dy, \end{aligned}$$

which is the claimed inequality since, by the assumptions, it holds  $\inf\{K(z) : z \in B_d(0)\} > 0$ .

We also remark that an improved version of this inequality is obtained in [1] in the case  $\Omega = B_{\rho}(x_0)$  and  $K(z) = [|z|_2^d f^p(|z|_2)]^{-1}$ ,  $f$  being continuous, increasing, and such that  $f(0) = 0$ . In that work, exploiting the structure of the kernel and a further doubling assumption, inequality (5.2) is obtained in a quantitative version so that the multiplicative constant appearing on the right-hand side explicitly depends on  $\rho$ .

## 6 Remarks on the Non-local Isoperimetric Problem

Let  $K$  be a kernel and let  $\Omega, E \subset \mathbb{R}^d$  be measurable sets. We define the  $K$ -perimeter of  $E$  relative to  $\Omega$  as

$$\mathcal{P}_K(E; \Omega) := \left( \frac{1}{2} \int_{\Omega \times \Omega} + \int_{\Omega \times \Omega^c} \right) |\mathbb{1}_E(x) - \mathbb{1}_E(y)| K(x - y) dx dy,$$

the  $K$ -perimeter of  $E$  by  $\mathcal{P}_K(E) := \mathcal{P}_K(E; \mathbb{R}^d) = \frac{1}{2}[\mathbb{1}_E]_{W^{K,1}(\mathbb{R}^d)}$  and finally we put

$$\mathcal{V}_K(E) := \int_{E \times E} K(x - y) dx dy.$$

For every  $m > 0$  we consider the following variational problems,

$$p_K(m) := \inf_{E \subset \mathbb{R}^d, |E|=m} \mathcal{P}_K(E), \quad (P_{K,m})$$

$$\sup_{E \subset \mathbb{R}^d, |E|=m} \mathcal{V}_K(E), \quad (P'_{K,m})$$

where the function  $p_K$  is usually referred to as the *isoperimetric profile* or the *isovolumetric function*. We recall that, by Remark 2.1, it would not be restrictive to replace the kernel  $K$  by its symmetrized kernel  $K_{\text{sym}}(x) = (K(x) + K(-x))/2$ . Therefore, we will often assume that (Sym) is satisfied.

**Remark 6.1** We observe that if  $K \in L^1(\mathbb{R}^d)$  and (Sym) holds, then

$$\mathcal{P}_K(E) := \|K\|_{L^1(\mathbb{R}^d)}|E| - \mathcal{V}_K(E).$$

Therefore, under this assumptions,  $(P_{K,m})$  is equivalent to  $(P'_{K,m})$  for every  $m > 0$ .

## 6.1 Non-local Perimeter of Balls

In this section, given a kernel  $K$ , we are interested in studying the function

$$(0, +\infty) \ni r \mapsto \mathcal{P}_K(B_r),$$

which in general is bounded from above by the function  $(0, +\infty) \ni r \mapsto p_K(|B_r|)$ . We start introducing the non-local curvature.

**Definition 6.2** (*Non-local curvature*) Let  $K$  be a kernel satisfying (Sym) and (Far) and let  $E \subset \mathbb{R}^d$  be a measurable set. We define

$$H_{K,\partial E}(x) := \limsup_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}^d \setminus B_\varepsilon(x)} (\mathbb{1}_{E^c}(y) - \mathbb{1}_E(y)) K(x-y) dy \in [-\infty, +\infty] \quad (6.1)$$

for every  $x \in \mathbb{R}^d$ .

**Remark 6.3** Let  $E \subset \mathbb{R}^d$  be a measurable set and  $x \in \mathbb{R}^d$ . We observe that (6.1) is well-defined, being,

$$\int_{\mathbb{R}^d \setminus B_\varepsilon(x)} |\mathbb{1}_{E^c}(y) - \mathbb{1}_E(y)| K(x-y) dy \leq 2 \int_{\mathbb{R}^d \setminus B_\varepsilon} K(z) dz < +\infty,$$



for every  $\varepsilon > 0$ . We also observe that,

$$H_{K, \partial E}(x) = \limsup_{\varepsilon \rightarrow 0^+} H_{K_\varepsilon, \partial E}(x) \in [-\infty, +\infty],$$

where  $\{K_\varepsilon\}_{\varepsilon > 0}$  is a family of kernels defined by  $K_\varepsilon := \mathbb{1}_{B_\varepsilon^c} K$  for every  $\varepsilon > 0$ .

**Lemma 6.4** *Let  $K$  be a kernel satisfying (Sym) and (Far) and  $C \subset \mathbb{R}^d$  be a convex set s.t.  $\partial C \neq \emptyset$ . Then,  $H_{K, \partial C}(x) \in [0, +\infty]$  for every  $x \in \partial C$ .*

**Proof** Let  $x \in \partial C$ . Let  $H^+$  be a half-space which does not contain  $C$  and whose boundary is a supporting hyperplane of containing  $x$ . Let  $H^- := \mathbb{R}^d \setminus H^+$ . For every  $\varepsilon > 0$  we define the kernel  $K_\varepsilon := \mathbb{1}_{B_\varepsilon^c} K$ . Then, being  $C^c = H^+ \sqcup (H^- \setminus C)$  and  $H^- = C \sqcup (H^- \setminus C)$ , we can write

$$\begin{aligned} H_{K_\varepsilon, \partial C}(x) &= \int_{C^c} K_\varepsilon(x-y) dy - \int_C K_\varepsilon(x-y) dy \\ &= \int_{H^+} K_\varepsilon(x-y) dy + \int_{H^- \setminus C} K_\varepsilon(x-y) dy \\ &\quad - \int_{H^-} K_\varepsilon(x-y) dy + \int_{H^- \setminus C} K_\varepsilon(x-y) dy. \end{aligned}$$

Since  $K_\varepsilon$  satisfies (Sym), by a change of variables we get,

$$\int_{H^+} K_\varepsilon(x-y) dy = \int_{H^-} K_\varepsilon(x-y) dy.$$

This implies that,

$$H_{K_\varepsilon, \partial C}(x) = 2 \int_{H^- \setminus C} K_\varepsilon(x-y) dy \geq 0,$$

which allows us to deduce that,

$$H_{K, \partial C}(x) = \limsup_{\varepsilon \rightarrow 0^+} H_{K_\varepsilon, \partial C}(x) \geq 0.$$

This concludes the proof.  $\square$

**Lemma 6.5** *Let  $K$  be a kernel satisfying (Sym) and s.t.  $K \in C^1(\mathbb{R}^d \setminus \{0\})$  and  $\sup_{z \in \mathbb{R}^d \setminus \{0\}} K(z) |z|^{\frac{d+s}{2}} < +\infty$  for some  $s \in (0, 1)$ . Then, the function  $(0, +\infty) \ni r \mapsto \mathcal{P}_K(B_r)$  is monotone non-decreasing.*

**Proof** We observe that the assumptions imply that  $K$  satisfies (Far). Let  $f(r) := \mathcal{P}_K(B_r)$  for every  $r > 0$ . Let  $\nu_{B_r}(x) := x/r$  for every  $x \in \partial B_r$  be the outer unit normal to  $\partial B_r$ . Fix  $r > 0$  and let  $X \in C_c^\infty(\mathbb{R}^d; \mathbb{R}^d)$  s.t.  $X(x) = \nu_{B_r}(x)$  for every  $x \in \partial B_r$ . Let  $\Phi_t(x) := x + tX(x)$  for every  $t \in \mathbb{R}$  and  $x \in \mathbb{R}^d$ .

Thanks to [22, Theorem 6.1],  $f$  is differentiable and

$$\frac{df}{dr}(r) = \frac{d}{dt} \bigg|_{t=0} \mathcal{P}_K(\Phi_t(B_r)) = \int_{\partial B_r} H_{K, \partial B_r} X \cdot \nu_{B_r} d\mathcal{H}^{d-1} = \int_{\partial B_r} H_{K, \partial B_r} d\mathcal{H}^{d-1}.$$

We conclude that  $\frac{df}{dr}(r) \geq 0$  by Lemma 6.4.  $\square$

**Lemma 6.6** *Let  $\{K_k\}_{k \in \mathbb{N}} \subset L^1(\mathbb{R}^d)$  be a sequence of kernels and  $K \in L^1(\mathbb{R}^d)$  be a kernel. If  $K_k \rightharpoonup K$  in  $L^1(\mathbb{R}^d)$ , then  $\lim_k [u]_{W^{K_k, 1}(\mathbb{R}^d)} = [u]_{W^{K, 1}(\mathbb{R}^d)}$  for every  $u \in L^1(\mathbb{R}^d)$ .*

**Proof** We define  $U(h) := \|\Delta_h u\|_{L^1(\mathbb{R}^d)}$  for every  $h \in \mathbb{R}^d$ , so that

$$[u]_{W^{K_k, 1}(\mathbb{R}^d)} = \int_{\mathbb{R}^d} U(h) K_k(h) dh, \quad (6.2)$$

for every  $k \in \mathbb{N}$ .

Since  $U \in L^\infty(\mathbb{R}^d)$ , with  $\|U\|_{L^\infty(\mathbb{R}^d)} \leq 2\|u\|_{L^1(\mathbb{R}^d)}$ , we can pass to the limit in (6.2) and conclude the proof.  $\square$

**Proposition 6.7** *Let  $K$  be a kernel. Then, the function  $(0, +\infty) \ni r \mapsto \mathcal{P}_K(B_r)$  is monotone non-decreasing.*

**Proof** Without loss of generality, recalling Remark 2.1, we may assume that (Sym) holds. If  $K$  does not satisfy (Far), then by Proposition 3.3  $\mathcal{P}_K(B_r) = +\infty$  for every  $r > 0$ , so that there is nothing to prove. Therefore, we assume that (Far) holds and observe that Lemma 3.2 guarantees then that  $K \in L^1_{\text{loc}}(\mathbb{R}^d \setminus \{0\})$ . Let  $0 < r < R$ .

*Step 1.* Assume that  $K \in L^1(\mathbb{R}^d)$ . Then we can find a sequence of kernels  $\{K_k\}_{k \in \mathbb{N}} \subset C_c^\infty(\mathbb{R}^d)$  s.t.  $K_k \rightarrow K$  in  $L^1(\mathbb{R}^d)$ . For every  $k \in \mathbb{N}$  we can also assume that  $K_k$  satisfies (Sym) up to replacing it with the kernel defined by  $(K_k(x) + K_k(-x))/2$  for every  $x \in \mathbb{R}^d$  (see Remark 2.1). In particular,

$K_k$  satisfies the hypotheses of Lemma 6.5 for every  $k \in \mathbb{N}$ , which implies that  $\mathcal{P}_{K_k}(B_r) \leq \mathcal{P}_{K_k}(B_R)$  for every  $k \in \mathbb{N}$ . Thanks to Lemma 6.6 we conclude that  $\mathcal{P}_K(B_r) \leq \mathcal{P}_K(B_R)$ .

*Step 2.* Let  $K \in L^1_{\text{loc}}(\mathbb{R}^d \setminus \{0\})$  satisfying (Sym) and define  $K_k := \min\{k, K\}$  for every  $k \in \mathbb{N}$ . Then,  $K_k \in L^1(\mathbb{R}^d)$  and it satisfies (Sym) for every  $k \in \mathbb{N}$ . By the previous step and since  $K_k \leq K$  for every  $k \in \mathbb{N}$ , we obtain

$$\mathcal{P}_{K_k}(B_r) \leq \mathcal{P}_{K_k}(B_R) \leq \mathcal{P}_K(B_R). \quad (6.3)$$

By the monotone convergence theorem (or by Fatou's lemma) from (6.3) we conclude that  $\mathcal{P}_K(B_r) \leq \mathcal{P}_K(B_R)$ .  $\square$

**Remark 6.8** If  $K$  is a kernel satisfying (Nts<sub>1</sub>), recalling [3, eq. (2.8)], we can obtain the following bound

$$\mathcal{P}_K(B_r) \leq \max \left\{ |B_1| r^d, \frac{1}{2} \mathcal{H}^{d-1}(\partial B_1) r^{d-1} \right\} \int_{\mathbb{R}^d} \min\{1, |x|_2\} K(x) \, dx,$$

for every  $r > 0$ .

We now provide an explicit formula to compute the non-local perimeter of one-dimensional balls.

**Proposition 6.9** *Let  $d = 1$  and let  $K$  be a kernel satisfying (Sym) and (Far). Let  $x_0 \in (0, +\infty)$ ,  $c_0 \in \mathbb{R}$  and  $G \in W^{1,1}_{\text{loc}}((0, +\infty))$  defined by  $G(x) := \int K(y) \, dy + c_0$  for every  $x > 0$ . Let  $H \in C^1((0, +\infty))$  s.t.  $\frac{dH}{dx} = G$ . Then, for every  $r > 0$ ,*

$$\frac{1}{2} \mathcal{P}_K((-r, r)) = 2G(+\infty)r + H(0^+) - H(2r), \quad (6.4)$$

where  $H(0^+) := \lim_{x \rightarrow 0^+} H(x) \in (-\infty, +\infty]$  and  $G(+\infty) := \lim_{x \rightarrow +\infty} G(x) \in (-\infty, +\infty)$ .

**Proof** Recalling Remark 3.2, we observe that  $K \in L^1_{\text{loc}}(\mathbb{R} \setminus \{0\})$ , so that  $G \in W^{1,1}_{\text{loc}}((0, +\infty))$  is well-defined.

We observe that  $G$  is monotone non-decreasing and that  $H$  is convex. Furthermore, either  $H$  is monotone (either non-decreasing, or non-increasing) in  $(0, +\infty)$ , or  $H$  is monotone non-increasing in  $(0, x']$  and monotone non-decreasing in  $[x', +\infty)$  for some  $x' > 0$ . In any case,  $\lim_{x \rightarrow 0^+} H(x) \in (-\infty, +\infty]$  and  $\lim_{x \rightarrow +\infty} G(x) \in (-\infty, +\infty)$  ( $G(x) < +\infty$  thanks to (Far)). We can then fix  $r > 0$  and compute,

$$\begin{aligned}
\frac{1}{2} \mathcal{P}_K((-r, r)) &= \frac{1}{2} \left( \int_{-\infty}^{-r} dy + \int_r^{+\infty} dy \right) \int_{-r}^r K(x-y) dx \\
&= \int_{-\infty}^{-r} \int_{-r}^r K(x-y) dx dy \\
&= \int_{-\infty}^{-r} (G(r-y) - G(-r-y)) dy \\
&\stackrel{(*)}{=} \lim_{M \rightarrow +\infty} \int_{-M}^{-r} \frac{d}{dy} [-H(r-y) + H(-r-y)] dy \\
&= -H(2r) + H(0^+) + \lim_{M \rightarrow +\infty} (H(r+M) - H(-r+M))
\end{aligned}$$

where the second equality is a consequence of (Sym), while the equality marked by  $\circledast$  is justified by the monotone convergence theorem, being

$$G(r-y) - G(-r-y) = \int_{-r}^r K(x-y) dx \geq 0 \text{ for any } y \in (-\infty, -r).$$

By Lagrange's mean value theorem for every  $M > 0$  we can find  $c_{r,M} \in (-r+M, r+M)$  s.t.  $H(r+M) - H(-r+M) = G(c_{r,M}) 2r$ . Therefore,

$$\lim_{M \rightarrow +\infty} (H(r+M) - H(-r+M)) = G(+\infty) 2r,$$

which completes the proof of (6.4).  $\square$

**Remark 6.10** Under the assumptions of Proposition 6.9, if for some  $r > 0$  the kernel satisfies  $\mathcal{P}_K((-r, r)) < +\infty$ , then the function  $(0, +\infty) \ni r \mapsto \mathcal{P}_K((-r, r))$  is everywhere finite and, taking the first derivative in (6.4), we observe that it is monotone non-decreasing, concave and at least of class  $C^1((0, +\infty))$ .

**Example 6.11** As an application of Proposition 6.9, we can obtain explicit formulas for the non-local perimeter of balls of some one-dimensional kernels. This is useful in order to ascertain the validity of (5.1) for  $d = 1$ , which is needed to obtain Propositions 5.3 and 5.6 and Corollary 5.4 (see also Corollary 6.13).

If we consider the fractional kernel  $K_s(x) = |x|^{-1-s}$  with  $s \in (0, 1)$  in dimension 1, we can choose  $G(x) = -x^{-s}/s$  and  $H(x) = -x^{1-s}/((1-s)s)$  for  $x > 0$  to find  $\mathcal{P}_{K_s}((-r, +r)) = -2H(2r)$  for every  $r > 0$ . This tells that (5.1) holds if the exponent  $1/q$  equals  $1-s$ ; i.e., for  $q = \frac{1}{1-s}$ .

If we consider the kernel  $K(x) = |x|^{-1}(-\log|x|)^{\gamma-1} \mathbb{1}_{(-1/3, 1/3)}(x)$ , where  $\gamma > 0$ , cfr. [3, Remark 2.20] and [23], we can choose

$$G(x) = \begin{cases} -\frac{(-\log(x))^\gamma}{\gamma} & \text{for } 0 < x < \frac{1}{3} \\ -\frac{(\log(3))^\gamma}{\gamma} & \text{for } x \geq \frac{1}{3} \end{cases},$$

$$H(x) = \begin{cases} -\frac{\int_0^x (-\log(t))^\gamma dt}{\gamma} & \text{for } 0 < x < \frac{1}{3} \\ -\frac{(\log(3))^\gamma}{\gamma} \left(x - \frac{1}{3}\right) - \frac{\int_0^{1/3} (-\log(t))^\gamma dt}{\gamma} & \text{for } x \geq \frac{1}{3} \end{cases},$$

so

that

$\phi(r) := \frac{1}{2} \mathcal{P}_K((-r/2, +r/2)) = G(+\infty)r - H(r) = -\frac{(\log(3))^\gamma}{\gamma}r - H(r)$ , for every  $r > 0$ .

We first observe that  $\phi$  is constant on  $[1/3, +\infty)$ , so that (5.1) cannot hold. This is not surprising, since  $K$  has compact support.

Nevertheless, we can consider a variant of  $K$  whose behavior in  $[1/3, +\infty)$  is, for instance, of fractional type, so that  $\phi$  restricted to  $[1/3, +\infty)$  is bounded from below by  $r^\alpha$  for some  $\alpha \in (0, 1]$ . We then deduce that  $\lim_{r \rightarrow 0^+} \phi(r)/r = +\infty$  and  $\lim_{r \rightarrow 0^+} \phi(r)/r^\alpha = 0$  for every  $\alpha \in (0, 1)$ . This, tells that if (5.1) holds, then necessarily the power of the volume in (5.1) must be 1.

## 6.2 Existence of Non-local Isoperimetric Sets

We now discuss some results in the literature related to the existence of minimizers of the problem  $(P_{K,m})$ .

Let  $m > 0$ . [5, Theorem 5.6 and Proposition 5.4] guarantee that if  $K \in L^1(\mathbb{R}^d)$  is a generic kernel, then a suitable relaxation of  $(P_{K,m})$  admits at least a solution and every solution of this relaxed problem has compact support. Moreover, in [5, Theorem 5.7 and Remark 5.8] it is shown that if  $K \in L^1(\mathbb{R}^d)$  satisfies a further technical condition (which is satisfied for instance if the kernel is positive definite) then  $(P_{K,m})$  admits at least a solution, which is compact. A key ingredient for these results is Remark 6.1 together with suitable first and second variations formulas.

On the other hand, if  $K$  is a sufficiently regular kernel (satisfying the assumptions of Proposition 6.9) and if  $E$  is a solution of  $(P_{K,m})$ , then it is a volume-constrained stationary set for the perimeter  $\mathcal{P}_K$  and as a consequence of [22, Theorem 6.1] the non-local curvature  $H_{K,\partial E}$  (recall Definition 6.2) is constant on  $\partial E$ .

Under symmetry assumptions, [3, Theorem 2.19] (cfr. also [5, Proposition 3.1]) tells us that  $(P_{K,m})$  is solved by Euclidean balls of volume  $m$  (which are the unique solutions) if  $K$  is a radial and decreasing kernel with respect to the Euclidean norm, i.e. if  $K$  satisfies  $(\text{Dec}(1, |\cdot|_2))$ . This proof is based on the *Riesz's rearrangement inequality*, which is a crucial tool no more available once the kernel presents a different symmetry. In fact, in [24] it is proved that Riesz's inequality fails if the rearrangement is performed by means of a general convex body rather than the Euclidean ball.

We finally stress that this is not simply a technical issue:  $(\text{Dec}(1, |\cdot|_2))$  is essential to obtain the optimality of Euclidean balls as Proposition 6.12 shows.

**Proposition 6.12** *Let  $K$  be a kernel satisfying (Sym) and (Far). Assume that there exists  $\delta > 0$  such that  $\{K_\delta > 0\}$  has non-empty interior, where  $K_\delta := \mathbb{1}_{B_\delta^c} K$ . Then, Euclidean balls of volume  $m_r := |B_r|$  do not solve  $(P_{K_\delta, m_r})$  for every  $r \in (0, \delta/2)$ .*

**Proof** Let  $\delta > 0$ ,  $r \in (0, \delta/2)$  and  $m_r := |B_r|$ . Thanks to Remark 6.1, we observe that  $(P_{K_\delta, m_r})$  is equivalent to  $(P'_{K_\delta, m_r})$ , where  $K_\delta := \mathbb{1}_{B_\delta^c} K \in L^1(\mathbb{R}^d)$  thanks to (Far).

Let  $x_0$  be a point in the interior of  $\{K_\delta > 0\}$  and let  $\varepsilon_0 > 0$  s.t.  $B_{\varepsilon_0}(x_0) \subset \{K_\delta > 0\} \subset B_\delta^c$ . Let  $\Omega_1 = B_{r/\sqrt[d]{2}}(x_0/2)$ ,  $\Omega_2 = B_{r/\sqrt[d]{2}}(-x_0/2)$  and observe that  $\text{dist}(\Omega_1, \Omega_2) = |x_0|_2 - 2r/\sqrt[d]{2} > \delta - 2r/\sqrt[d]{2} > 0$ .

We fix  $0 < \varepsilon < \min\{\varepsilon_0/2, r/\sqrt[d]{2}\}$  and put  $\Omega := \Omega_1 \sqcup \Omega_2$ , noting that  $|\Omega| = m_r$ . Then,

$$\mathcal{V}_{K_\delta}(\Omega) = \mathcal{V}_{K_\delta}(\Omega_1) + \mathcal{V}_{K_\delta}(\Omega_2) + 2 \int_{\Omega_1} \int_{\Omega_2} K_\delta(x-y) \, dy \, dx.$$

In particular, using the fact that for every  $x \in B_\varepsilon(x_0/2) \subset \Omega_1$  and every  $y \in B_\varepsilon(-x_0/2) \subset \Omega_2$  we have  $x-y \in B_{\varepsilon_0}(x_0) \subset \{K_\delta > 0\}$ ,

$$\begin{aligned} \mathcal{V}_{K_\delta}(\Omega) &\geq 2 \int_{\Omega_1} \int_{\Omega_2} K_\delta(x-y) \, dy \, dx \\ &\geq 2 \int_{B_\varepsilon(x_0/2)} \int_{B_\varepsilon(-x_0/2)} K_\delta(x-y) \, dy \, dx > 0 = \mathcal{V}_{K_\delta}(B_r), \end{aligned}$$

where the last equality follows from the restriction  $r \in (0, \delta/2)$  combined with the inclusion  $B_\delta \subset \{K_\delta = 0\}$ . This shows that  $B_r$  is not a solution of  $(P'_{K_\delta, m_r})$ , which concludes the proof.  $\square$

We conclude this note addressing to a possible application of the extension result in Sect. 4 and of the inequalities previously obtained in Sect. 5. More precisely, inequality (5.3) can be employed to infer the following non-local isoperimetric inequality relative to a regular set  $\Omega$ .

**Corollary 6.13** *Let  $\Omega \subset \mathbb{R}^d$  a bounded, open set with Lipschitz boundary, and assume that (5.1) holds for some  $q \in [1, +\infty)$ . Let  $K$  be a kernel satisfying (Nint),  $(\text{Dec}(c_0, |\cdot|_*))$ ,  $(\text{Dou}(D, |\cdot|_*))$ , and  $(\text{Nts}_1)$ . Then there exists a positive constant  $C$  such that*

$$\min\{|E \cap \Omega|, |\Omega \setminus E|\}^{\frac{1}{q}} \leq C \mathcal{P}_K(E; \Omega) \quad (6.5)$$

for every  $E \subseteq \mathbb{R}^d$  measurable.

**Proof** Without loss of generality, assume  $\mathcal{P}_K(E; \Omega) < +\infty$ . We prove the statement under the further assumption that  $\min\{|E \cap \Omega|, |\Omega \setminus E|\} = |E \cap \Omega|$ , the other case being analogous. Applying Proposition 5.6 with  $u = \mathbb{1}_E$  and noting that we are supposing  $|\Omega \setminus E| \geq |\Omega|/2$ , we obtain

$$\begin{aligned} \mathcal{P}_K(E; \Omega) &\geq \frac{1}{2} [\chi_E]_{W^{K,1}(\Omega)} \\ &\geq \frac{1}{2C} \left( \int_{\Omega} \left| \chi_E - \frac{|E \cap \Omega|}{|\Omega|} \right|^q dx \right)^{\frac{1}{q}} \\ &= \frac{1}{2C} \left[ |E \cap \Omega| \left( \frac{|\Omega \setminus E|}{|\Omega|} \right)^q + |\Omega \setminus E| \left( \frac{|E \cap \Omega|}{|\Omega|} \right)^q \right]^{\frac{1}{q}} \\ &\geq \frac{1}{2C} |E \cap \Omega|^{\frac{1}{q}} \frac{|\Omega \setminus E|}{|\Omega|} \geq \frac{1}{4C} |E \cap \Omega|^{\frac{1}{q}}, \end{aligned}$$

where the constant  $C > 0$  is the one appearing in Proposition 5.6. This concludes the proof.  $\square$

**Acknowledgements** KB has been supported by Fondazione Cariplo, grant n° 2023-0873. KB is member of the INdAM-GNAMPA and his work is partially supported by the INdAM -GNAMPA 2024 Project "Proprietà geometriche e regolarità in problemi variazionali locali e non locali", codice CUP E53C23001670001. GCB is member of the INdAM-GNAMPA.

**Author Contributions** KB and GCB equally contributed to the manuscript.

**Funding** Open access funding provided by Università degli Studi di Pavia within the CRUI-CARE Agreement. KB has been supported by Fondazione Cariplo, grant n° 2023-0873. GCB did not receive support from any organization for the submitted work.

## Declarations

**Conflict of interest** The authors declare that they have not financial or non-financial interests related to the work submitted for publication.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

1. Duzaar, F., Gastel, A., Mingione, G.: Elliptic systems, singular sets and Dini continuity. *Comm. Partial Differ. Equ.* **29**(7–8), 1215–1240 (2004)

2. Di Nezza, E., Palatucci, G., Valdinoci, E.: Hitchhiker's guide to the fractional Sobolev spaces. *Bull. Sci. Math.* **136**(5), 521–573 (2012)
3. Bessas, K., Stefani, G.: Non-local BV functions and a denoising model with  $L^1$  fidelity. *Adv. Calc. Var.* **18**(1), 189–217 (2025)
4. Gounoue, GFF.:  $L^2$ -theory for nonlocal operators on domains, Ph.D. thesis, Universität Bielefeld, (2020)
5. Cesaroni, A., Novaga, M.: The isoperimetric problem for nonlocal perimeters. *Discrete Contin. Dyn. Syst. Ser. S* **11**(3), 425–440 (2018)
6. Foghem, G.: Stability of complement value problems for  $p$ -Lévy operators. *NoDEA Nonlinear Diff. Equ. Appl.* **32**(1), 1 (2025)
7. Bessas, K.: Fractional total variation denoising model with  $L^1$  fidelity. *Nonlinear Anal.* **222**, 112926 (2022)
8. Antil, H., Díaz, H., Jing, T., Schikorra, A.: Nonlocal bounded variations with applications. *SIAM J. Math. Anal.* **56**(2), 1903–1935 (2024)
9. José, M.: Mazón, Marcos Solera, and Julián Toledo,  $(BV, L^p)$ -decomposition,  $p = 1, 2$ , of functions in metric random walk spaces. *Adv. Calc. Var.* **15**(3), 515–550 (2022)
10. Novaga, M., Onoue, F.: Local Hölder regularity of minimizers for nonlocal variational problems. *Commun. Contemp. Math.* **25**(10), 2250058 (2023)
11. Bessas, K., Novaga, M., Onoue, F.: On the shape of small liquid drops minimizing nonlocal energies. *ESAIM Control Optim. Calc. Var.* **29**, 86 (2023)
12. Cesaroni, A., Novaga, M.: Volume constrained minimizers of the fractional perimeter with a potential energy. *Discrete Contin. Dyn. Syst. Ser. S* **10**(4), 715–727 (2017)
13. Maggi, F., Valdinoci, E.: Capillarity problems with nonlocal surface tension energies. *Comm. Partial Differ. Equ.* **42**(9), 1403–1446 (2017)
14. Dipierro, S., Maggi, F., Valdinoci, E.: Asymptotic expansions of the contact angle in nonlocal capillarity problems. *J. Nonlinear Sci.* **27**(5), 1531–1550 (2017)
15. Dipierro, S., Maggi, F., Valdinoci, E.: Minimizing cones for fractional capillarity problems. *Rev. Mat. Iberoam.* **38**(2), 635–658 (2022)
16. De Luca, A., Dipierro, S., Valdinoci, E.: Nonlocal capillarity for anisotropic kernels. *Math. Ann.* **388**(4), 3785–3846 (2024)
17. Indrei, E.: The nonlocal Almgren problem. *Math.* **13**(17), 2716 (2025)
18. Ludwig, M.: Anisotropic fractional perimeters. *J. Differ. Geom.* **96**(1), 77–93 (2014)
19. Kreuml, A.: The anisotropic fractional isoperimetric problem with respect to unconditional unit balls. *Commun. Pure Appl. Anal.* **20**(2), 783–799 (2021)
20. Ferone, V., Piscitelli, G., Volzone, B.: Symmetrization results for general nonlocal linear elliptic and parabolic problems. *J. Math. Pures Appl.* **190**, 103597 (2024)
21. Leoni, G.: A First Course in Fractional Sobolev Spaces, Graduate Studies in Mathematics, vol. 229, p. 2023. American Mathematical Society, Providence (2023)
22. Figalli, A., Fusco, N., Maggi, F., Millot, V., Morini, M.: Isoperimetry and stability properties of balls with respect to nonlocal energies. *Comm. Math. Phys.* **336**(1), 441–507 (2015)
23. Brué, E., Nguyen, Q.-H.: On the Sobolev space of functions with derivative of logarithmic order. *Adv. Nonlinear Anal.* **9**(1), 836–849 (2020)
24. Schaftringen, J.: Anisotropic symmetrization. *Ann. Inst. H. Poincaré C Anal. Non Linéaire* **23**(4), 539–565 (2006)

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.