



SCUOLA INTERNAZIONALE SUPERIORE DI STUDI AVANZATI

SISSA Digital Library

Algebraic groups in non-commutative probability theory revisited

Original

Algebraic groups in non-commutative probability theory revisited / Chevyrev, I., Ebrahimi-Fard, K., Patras, F.. - In: INFINITE DIMENSIONAL ANALYSIS QUANTUM PROBABILITY AND RELATED TOPICS. - ISSN 0219-0257. - 27:3(2024). [10.1142/s021902572450005x]

Availability:

This version is available at: 20.500.11767/148772 since: 2025-10-21T09:02:49Z

Publisher:

Published

DOI:10.1142/s021902572450005x

Terms of use:

Testo definito dall'ateneo relativo alle clausole di concessione d'uso

Publisher copyright

World Scientific Publishing Co.

This version is available for education and non-commercial purposes.

note finali coverpage

(Article begins on next page)

ALGEBRAIC GROUPS IN NON-COMMUTATIVE PROBABILITY THEORY REVISITED

ILYA CHEVYREV, KURUSCH EBRAHIMI-FARD, AND FRÉDÉRIC PATRAS

ABSTRACT. The role of coalgebras as well as algebraic groups in non-commutative probability has long been advocated by the school of von Waldenfels and Schürmann. Another algebraic approach was introduced more recently, based on shuffle and pre-Lie calculus, and results in another construction of groups of characters encoding the behaviour of states. Comparing the two, the first approach, recast recently in a general categorical language by Manzel and Schürmann, can be seen as largely driven by the theory of universal products, whereas the second construction builds on Hopf algebras and a suitable algebraization of the combinatorics of noncrossing set partitions. Although both address the same phenomena, moving between the two viewpoints is not obvious. We present here an attempt to unify the two approaches by making explicit the Hopf algebraic connections between them. Our presentation, although relying largely on classical ideas as well as results closely related to Manzel and Schürmann's aforementioned work, is nevertheless original on several points and fills a gap in the non-commutative probability literature. In particular, we systematically use the language and techniques of algebraic groups together with shuffle group techniques to prove that two notions of algebraic groups naturally associated with free, respectively Boolean and monotone, probability theories identify. We also obtain explicit formulas for various Hopf algebraic structures and detail arguments that had been left implicit in the literature.

Key words: universal products; free product; states; cumulants; additive convolution; shuffle algebra; pre-Lie algebra; Hopf algebra.

MSC classes: 16T05, 16T10, 16T30, 17A30, 46L53, 46L54

CONTENTS

1. Introduction	1
2. General notation and conventions	2
3. Group structures from convolutions: the classical case	5
4. Universal products: the non-commutative case	7
5. Products of states	12
6. The monotone group of states	14
7. The Boolean group of states	18
8. The free group of states	22
References	25

1. INTRODUCTION

The school of von Waldenfels and Schürmann has advocated the use of coalgebras as well as algebraic groups and their Hopf algebras of coordinate functions in the context of non-commutative probability theory [3, 4, 5, 22, 31]. More recently, two of the present authors

introduced a different approach, which is mainly driven by shuffle and pre-Lie calculus, leading to another construction of groups of characters [10, 11, 12, 13, 14]. The first construction can be understood as largely determined by the theory of universal products and additive free, monotone and boolean convolutions, whereas the second builds on a suitable algebraization of the combinatorics of lattices of set partitions and is encoded into a Hopf algebra structure on $T(T^+(\mathcal{A}))$, the double tensor algebra over an algebra \mathcal{A} of non-commutative random variables. Although both approaches address the same phenomena, moving from one to the other has no obvious mathematical formulation.

The goal of the work at hand is to present a unified Hopf algebraic point of view that permits to compare the different approaches and to make the connections transparent by including explicit formulas on the Hopf algebraic structures. The key object appears to be $S(T^+(\mathcal{A}))$, the Hopf algebra of polynomials over the tensor algebra over \mathcal{A} . Our exposition of results, while relying on classical ideas, is original on several points. Moreover, we obtain explicit formulas for the Hopf algebraic structures and detail arguments that have been sometimes left implicit in the literature.

Although we have tried to give a self-contained presentation, we need to assume some familiarity with the foundations of non-commutative probability theory as well as the theory of Hopf algebras. The reader who would like to see more details on non-commutative probability and related topics is referred to [6, 18, 20, 29, 30]. Detailed studies of the notion of universal product in non-commutative probability can be found in [4, 17, 19, 28]. Regarding background and details on Hopf and pre-Lie algebras, in particular of combinatorial nature, we refer the reader to the book [8].

The paper is organised as follows. The next section introduces a collection of notations and conventions used throughout the paper. In Section 3, we consider the group structure coming from convolution in the case of classical probability. Section 4 presents the notion of universal product in the context of non-commutative probability. Section 5 recalls the shuffle algebra approach to moment-cumulant relations in non-commutative probability. Sections 6, 7 and 8, discuss the monotone, free and Boolean group structures for states.

Acknowledgements: The authors would like to thank the anonymous referee for their detailed comments that helped to improve the presentation of the paper.

This work was partially supported by the project ‘‘Pure Mathematics in Norway’’, funded by Trond Mohn Foundation and Tromsø Research Foundation. IC acknowledges support by the EPSRC via the New Investigator Award EP/X015688/1. KEF was supported by the Research Council of Norway through project 302831 ‘‘Computational Dynamics and Stochastics on Manifolds’’ (CODYSMA). FP acknowledges support from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation program (Dual project, grant agreement No. 670624), from the ANR project Algebraic Combinatorics, Renormalization, Free probability and Operads – CARPLO (Project-ANR-20-CE40-0007) and from the ANR–FWF project PAGCAP (ANR-21-CE48-0020, Austrian Science Fund (FWF), grant I 5788).

2. GENERAL NOTATION AND CONVENTIONS

Let \mathbb{N}^* denote the positive integers. For any $n \in \mathbb{N}^*$, we define the sets $[n] := \{1, \dots, n\}$ and $\langle n \rangle := \{0, \dots, n - 1\}$. Throughout the article, (\mathcal{A}, φ) stands for a non-commutative

probability space. This means that \mathcal{A} is an associative algebra with unit $1_{\mathcal{A}}$ over a field \mathbb{K} of characteristic zero (usually taken to be the complex numbers in concrete applications). See the standard reference [20] for more details. It is important to distinguish the unit of \mathcal{A} from the unit of the ground field: we denote the former $1_{\mathcal{A}}$ (or 1) and the latter $\mathbf{1}$. The map φ is an unital \mathbb{K} -valued linear form on \mathcal{A} , i.e., $\varphi(1_{\mathcal{A}}) = \mathbf{1}$. The elements of \mathcal{A} play the role of random variables, while the map φ should be considered as the expectation.

Classical cumulants are defined in probability where the algebra \mathcal{A} of random variables is commutative, as a family of multilinear functionals $\{c_n : \mathcal{A}^n \rightarrow \mathbb{C}\}_{n \geq 1}$ iteratively determined by classical moment-cumulant relations

$$(1) \quad \varphi(a_1 \cdots a_n) = \sum_{\pi \in P_n} \prod_{\pi_i \in \pi} c_{|\pi_i|}(a_1, \dots, a_n | \pi_i),$$

see [21, Sec. 3.2]. The sum on the right hand side runs over all set partitions of $\{1, \dots, n\}$. Here and below we write for a block $\pi_i = \{i_1, \dots, i_{|\pi_i|}\}$ in a set partitions $\pi = \{\pi_1, \dots, \pi_m\} \in P_n$ and a multilinear functional f

$$f_{|\pi_i|}(a_1, \dots, a_n | \pi_i) := f_{|\pi_i|}(a_{i_1}, \dots, a_{i_{|\pi_i|}}).$$

Free, Boolean, and monotone cumulants in non-commutative probability provide effective combinatorial tools for studying non-commutative distributions of random variables. Free cumulants [26, 20] form a family of multilinear functionals $\{k_n : \mathcal{A}^n \rightarrow \mathbb{C}\}_{n \geq 1}$ which are recursively defined by free moment-cumulant relations

$$(2) \quad \varphi(a_1 \cdots a_n) = \sum_{\pi \in NC_n} \prod_{\pi_i \in \pi} k_{|\pi_i|}(a_1, \dots, a_n | \pi_i).$$

Boolean cumulants [27] form another family of multilinear functionals $\{b_n : \mathcal{A}^n \rightarrow \mathbb{C}\}_{n \geq 1}$ recursively defined by Boolean moment-cumulant relations

$$(3) \quad \varphi(a_1 \cdots a_n) = \sum_{\pi \in \text{Int}_n} \prod_{\pi_i \in \pi} b_{|\pi_i|}(a_1, \dots, a_n | \pi_i).$$

Monotone cumulants [16] form a family of multilinear functionals $\{h_n : \mathcal{A}^n \rightarrow \mathbb{C}\}_{n \geq 1}$ recursively defined by the monotone moment-cumulant relations

$$(4) \quad \varphi(a_1 \cdots a_n) = \sum_{\pi \in NC_n} \frac{1}{t(\pi)!} \prod_{\pi_i \in \pi} h_{|\pi_i|}(a_1, \dots, a_n | \pi_i).$$

The sums on the right hand sides of (2) and (4) run over the lattice NC_n of non-crossing partitions of the set $\{1, \dots, n\}$. The sum on the right hand side of (3) on the other hand is defined over the lattice $\text{Int}_n \subset NC_n$ of interval partitions (i.e., where each block is an interval).

Tensor algebras: The non-unital tensor algebra $\bigoplus_{n > 0} \mathcal{A}^{\otimes n}$ over \mathcal{A} is denoted $T^+(\mathcal{A})$. On the other hand, the unital tensor algebra is obtained by adding a copy of the ground field, that is, $T(\mathcal{A}) := \mathbb{K} \oplus T^+(\mathcal{A})$. The same conventions apply for the tensor (resp. non-unital tensor) algebra $T(V)$ (resp. $T^+(V)$) over a vector space V .

In general, when $B = \bigoplus_{n \in \mathbb{N}} B_n$ is a graded connected algebra (meaning that the degree zero component of the graded algebra is isomorphic to the ground field), we will write B^+ for its augmentation ideal, that is, $\bigoplus_{n \in \mathbb{N}^*} B_n$.

Tensors $a_1 \otimes \cdots \otimes a_n \in T(\mathcal{A})$ are written as words $a_1 \cdots a_n$. They have to be distinguished from the product of the a_i seen as elements in the algebra \mathcal{A} , which is written $a_1 \cdot_{\mathcal{A}} \cdots \cdot_{\mathcal{A}} a_n$. The unit $\mathbf{1}$ of the ground field is identified with the empty word \emptyset . The word $a \cdots a \in \mathcal{A}^{\otimes n}$ is denoted a^n – and should not be confused with the n -th power of a in \mathcal{A} , which is written $a^{\mathcal{A}n}$.

The natural algebra structure on $T^+(\mathcal{A})$ refers to the concatenation product of words, which turns it into a non-commutative graded algebra with the natural degree of a word $w = a_1 \cdots a_n$ being its length, i.e., its number of letters, $\deg(w) := n$. In the unital case the empty word is of length zero and, as we saw already, plays the role of the unit for the concatenation product.

We use here the convention that identifies the algebra of polynomials $S(V)$ (also known as the symmetric tensor algebra) over a vector space V with the space of covariants $\bigoplus_{n \geq 0} T^n(V)/S_n$, where S_n is the n -th symmetric group acting on tensors of length n by permutation:

$$\forall \sigma \in S_n, v_1, \dots, v_n \in V, \quad \sigma(v_1 \cdots v_n) := v_{\sigma^{-1}(1)} \cdots v_{\sigma^{-1}(n)}.$$

The augmentation ideal $S^+(V)$ is $\bigoplus_{n > 0} T^n(V)/S_n$. The same approach is used for the algebra of polynomials $S(X)$ over a set X .

We define then $S(T^+(\mathcal{A}))$ and $S^+(T^+(\mathcal{A}))$ to be the polynomial algebra, respectively the non-unital polynomial algebra over $T^+(\mathcal{A})$. We again identify $\mathbf{1} \in \mathbb{K}$ with the unit of $S(T^+(\mathcal{A}))$ to distinguish it from the unit of \mathcal{A} and thus treat \mathbb{K} as a subspace of $S(T^+(\mathcal{A}))$. Notice that, as the ground field is of characteristic zero, there is a canonical bijection between invariants and covariants, and $S(T^+(\mathcal{A}))$ identifies with the space of symmetric tensors over $T^+(\mathcal{A})$. See [8, Sect. 2.4] for a general discussion of polynomial (Hopf) algebras.

Elements in $S(T^+(\mathcal{A}))$ are represented using a parenthesized bar-notation, i.e., $(w_1 | \cdots | w_n) \in S(T^+(\mathcal{A}))$, where the words $w_i \in T^+(\mathcal{A})$, $i = 1, \dots, n$. The algebra $S(T^+(\mathcal{A}))$ is commutative with the bar-product $a|b := (w_1 | \cdots | w_n | v_1 | \cdots | v_m)$ for $a = (w_1 | \cdots | w_n)$ and $b = (v_1 | \cdots | v_m)$. We denote by $\deg(a) := \deg(w_1) + \cdots + \deg(w_n)$ the degree of an element $a \in S(T^+(\mathcal{A}))$.

Analogous conventions hold for both the non-unital and unital double tensor algebras, defined to be $T^+(T^+(\mathcal{A})) := \bigoplus_{n > 0} T^+(\mathcal{A})^{\otimes n}$, respectively $T(T^+(\mathcal{A})) := \bigoplus_{n \geq 0} T^+(\mathcal{A})^{\otimes n}$. Elements are denoted using the bracket-bar notation, $[w_1 | \cdots | w_n] \in T^+(T^+(\mathcal{A}))$, for words $w_i \in T^+(\mathcal{A})$, $i = 1, \dots, n$.

Set theoretic conventions: Consider now a canonically ordered subset $S := \{s_1, \dots, s_m\}$ of the set $[n]$. For a word $a_1 \cdots a_n \in T^+(\mathcal{A})$, we set $a_S := a_{s_1} \cdots a_{s_m} \in T^+(\mathcal{A})$ and $a_{\emptyset} := \mathbf{1}$. A connected component $S' := \{s_i, \dots, s_{i+j}\} \subseteq S \subseteq [n]$ is a maximal sequence of successive elements in S , i.e., $s_{i+k} = s_i + k$ for all $1 \leq k \leq j$, and $s_{i-1} < s_i - 1$ and $s_{i+j+1} > s_{i+j} + 1$ whenever $i > 1$ and $i + j < m$ respectively. For any $S \subseteq [n]$ we denote by S_1, \dots, S_p and J_1^S, \dots, J_k^S the connected components of S and of $[n] - S$, respectively. Both are naturally ordered in terms of their minimal (or maximal) elements. For example, if $n = 7$ and $S = \{2, 3, 6\}$, we get $p = 2$ connected components of S , $S_1 = \{2, 3\}$, $S_2 = \{6\}$ and $k = 3$ connected components of $[n] - S$, $J_1^S = \{1\}$, $J_2^S = \{4, 5\}$, $J_3^S = \{7\}$.

For two partitions π_1, π_2 of $[n]$, we say that π_1 is finer than π_2 and write $\pi_1 \leq \pi_2$ if every block of π_1 is a subset of one block in π_2 . The relation \leq turns the set of partitions into a lattice. With this notation, $\pi_{\text{Int}}(S) := \{S_1, \dots, S_p, J_1^S, \dots, J_k^S\}$ is the maximal interval partition which is finer than the two-block partition $\{S, [n] - S\}$. We note that this is a dual closure map in the sense of [7, 1.4 Closures].

3. GROUP STRUCTURES FROM CONVOLUTIONS: THE CLASSICAL CASE

Let us start with classical probability as a template for the non-commutative constructions which are to come in the next section. The following presentation revisits ideas from classical probability in an algebraic language recasting them in the setting presented in the recent work [15], which formulates moment-cumulant relations as well as Wick polynomials in the framework of algebraic groups.

We assume that all distributions of random variables have determinate moment problem, that is, they are entirely characterized by the sequence of all their moments (think for example to random variables with compact support, Gaussian variables, or alike; see [1] as a standard reference on these questions). This hypothesis arises from a parallel assumption made within the combinatorial approach to free probability, where the distribution of a random variable is identified with the sequence of its moments at all orders. See for example the definition of distributions in free probability in the standard reference [20, Chapter 1].

Consider now a commutative unital algebra \mathcal{A} of classical random variables with expectation value operator \mathbb{E} . It is convenient to encode distributions of random variables by the linear form still written $\mathbb{E} : S(\mathcal{A}) \rightarrow \mathbb{K}$,

$$a_1 \cdots a_n \longmapsto \mathbb{E}(a_1 \cdots a_n) = \mathbb{E}[a_1 \underset{\mathcal{A}}{\cdot} \cdots \underset{\mathcal{A}}{\cdot} a_n],$$

where $\mathbb{E}(\mathbf{1}) := 1$ and $a_1 \cdots a_n$ stands for a monomial of degree n in the polynomial algebra $S(\mathcal{A})$ (not to be confused with $a_1 \underset{\mathcal{A}}{\cdot} \cdots \underset{\mathcal{A}}{\cdot} a_n$, the product of the a_i in \mathcal{A}). The distribution of an element $a \in \mathcal{A}$ is then obtained as the sequence of its moments $\mathbb{E}(a^n)$. The (additive) convolution of two distributions of scalar-valued random variables X, Y belonging to \mathcal{A} is then defined as the distribution of the sum $X' + Y'$, where X' and Y' are independent and have the same laws as X and Y . The simplest way to make this process effective is to construct the tensor product algebra $\mathcal{A} \otimes \mathcal{A}$ equipped with the expectation value operator $\mathbb{E} \otimes \mathbb{E}$. Given two integrable functions $f(X)$ and $g(Y)$, of X respectively Y , one gets

$$\mathbb{E} \otimes \mathbb{E}(f(X) \otimes g(Y)) = \mathbb{E}(f(X))\mathbb{E}(g(Y)),$$

which shows that $X' := X \otimes \mathbf{1}$ and $Y' := \mathbf{1} \otimes Y$ are two independent copies in $\mathcal{A} \otimes \mathcal{A}$ of the random variables X and Y .

This process dualizes as follows.

Definition 1. *Let ϕ and ψ be two unital linear forms on $S(\mathcal{A})$. The (additive) convolution product of ϕ and ψ is the unital linear form on $S(\mathcal{A})$ defined by:*

$$\phi * \psi(\mathbf{1}) := 1,$$

$$\phi * \psi(a_1 \cdots a_n) := \phi \otimes \psi((a_1 \otimes \mathbf{1} + \mathbf{1} \otimes a_1) \cdots (a_n \otimes \mathbf{1} + \mathbf{1} \otimes a_n)).$$

Corollary 2. *If X and X' are two independent identically distributed (i.i.d.) random variables, then the distribution of $X + X'$ with respect to \mathbb{E} identifies with the distribution of X with respect to $\mathbb{E} * \mathbb{E}$.*

Furthermore, as $\phi * \psi(a_1 \cdots a_n) = \phi(a_1 \cdots a_n) + \psi(a_1 \cdots a_n) + \text{l.o.t.}$, where l.o.t. refers to a sum of products of evaluations of ϕ and ψ on tensors of length strictly less than n , a standard inductive argument (on the length of tensors) that we omit shows that the $*$ -product defines a group structure on the set of unital linear forms on $S(\mathcal{A})$.

Definition 3. We call the set \mathcal{G} of unital linear forms on $S(\mathcal{A})$, equipped with the convolution product, the group of generalized measures on \mathcal{A} .

Let us assume now that \mathcal{A} has a basis $\mathcal{B} = (b_i)_{i \in \mathbb{N}}$. Think for example of a Gaussian family $\{\Omega_1, \dots, \Omega_n\}$ and the algebra spanned by monomials in the $\Omega_1, \dots, \Omega_n$ (which is a dense subalgebra of the square integrable functions).

Given a sequence $I = (i_1, \dots, i_k)$ with $0 \leq i_1 \leq \dots \leq i_k$, we set $b_I := b_{i_1} \cdots b_{i_k} \in S(\mathcal{A})$. As usual, b_I should *not* be confused with the product $b_{i_1} \cdot_{\mathcal{A}} \cdots \cdot_{\mathcal{A}} b_{i_k}$ in \mathcal{A} . The set \mathcal{X} of all monomials b_I defines then a (complete) system of coordinates on \mathcal{G} , where, for an element $\phi \in \mathcal{G}$, we specify

$$\langle \phi | b_I \rangle := \phi(b_I).$$

Here complete refers to the fact that this set \mathcal{X} of coordinate functions induces a bijective correspondence between elements of \mathcal{G} and families of scalars indexed by sequences of integers:

$$\phi \longmapsto \{\phi(b_I)\}_I.$$

The definition of the group law on \mathcal{G} implies

$$\langle \phi * \psi | b_I \rangle = \sum_{A \amalg B = [I]} \langle \phi | b_{A,I} \rangle \langle \psi | b_{B,I} \rangle,$$

where, if $A = \{j_1, \dots, j_k\}$, $b_{A,I} := b_{i_{j_1}} \cdots b_{i_{j_k}}$. This shows that the b_I are representative functions on the group \mathcal{G} . See [8, 2.7 Representative functions] for details. By standard arguments, the polynomial algebra $S(\mathcal{X})$ is a Hopf algebra whose coproduct δ is given on generators by

$$(5) \quad \delta(b_I) = \sum_{A \amalg B = [I]} b_{A,I} \otimes b_{B,I},$$

and the following theorem holds (see [8, 3.5 Algebraic groups]).

Theorem 4. *The (commutative) group \mathcal{G} of generalized measures is, up to a canonical isomorphism, the group of characters (algebra maps to the ground field) of the (commutative and cocommutative) Hopf algebra $S(\mathcal{X})$, with coproduct given by (5).*

Remark 5. *A terminological precision: it follows from general arguments (namely the fundamental theorem for coalgebras), but is easy to check directly here, that the Hopf algebra $S(\mathcal{X})$ is a direct limit of commutative Hopf subalgebras which are finitely generated as algebras. The group \mathcal{G} is therefore an inverse limit of affine algebraic groups (groups of characters on a finitely generated commutative Hopf algebra), that is, a pro-affine algebraic group.*

Let us show briefly why these constructions are meaningful for probability. Assume again that we consider a given expectation value functional \mathbb{E} on \mathcal{A} and view it as an element of \mathcal{G} . Let us write \star for the convolution product of linear forms on $S(\mathcal{X})$: given $\lambda, \beta \in S(\mathcal{X})^*$,

$$\lambda \star \beta(b_{I_1} \cdots b_{I_k}) := (\lambda \otimes \beta) \circ \delta(b_{I_1} \cdots b_{I_k}).$$

The product map $*$ on \mathcal{G} is the restriction of \star to the set of characters – so, the two products agree on \mathcal{G} , but we prefer to keep a notational distinction between them for the sake of clarity.

Proposition 6. *For elements $a_1, \dots, a_n \in \mathcal{A}$, we have*

$$\log^*(\mathbb{E})(a_1 \cdots a_n) = c_n(a_1, \dots, a_n),$$

where c_n stands for the multivariate classical cumulant map as in (1).

Proof. Iterating the coproduct formula (5) gives

$$\begin{aligned} \log^*(\mathbb{E})(a_1 \cdots a_n) &= \sum_{k=1}^n \frac{(-1)^{k-1}}{k} \sum_{P \in OP_n^k} \mathbb{E}[a_{P_1}] \cdots \mathbb{E}[a_{P_k}] \\ &= \sum_{k=1}^n (-1)^{k-1} (k-1)! \sum_{P \in P_n^k} \mathbb{E}[a_{P_1}] \cdots \mathbb{E}[a_{P_k}], \end{aligned}$$

where OP_n^k and P_n^k stand for the set of ordered partitions of length k respectively the set of partitions of length k of $[n]$. The last term provides the Leonov–Shiryaev definition of multivariate cumulants, see [21, Sec. 3.2]. The combinatorial study of the moments/cumulants relations actually goes back at least to Schützenberger’s thesis [23]; see for instance [15, Eq. (12)] for details on an Hopf algebraic account). \square

However, these ideas can be recast differently. Consider a basis element $b_I = b_{i_1} \cdots b_{i_k}$ in \mathcal{X} . We use Proposition 6 to introduce another system of coordinates on \mathcal{G} . Let \mathcal{Y} be the set of the c_I , where, setting $n := |I|$,

$$c_I := \sum_{k=1}^n (-1)^{k-1} (k-1)! \sum_{P \in P_n^k} b_{P_1, I} \cdots b_{P_k, I}.$$

As $c_I = b_I + \text{l.o.t.}$, the c_I indeed define another complete system of coordinates on \mathcal{G} . Furthermore, the same argument as for \mathbb{E} gives for $\alpha \in \mathcal{G}$ (viewed as a character on $S(\mathcal{X})$),

$$\log^*(\alpha)(b_I) = \langle \alpha | c_I \rangle.$$

As the products $*$ and \star are commutative and both agree on \mathcal{G} , we obtain for given $\alpha, \beta \in \mathcal{G}$:

$$\begin{aligned} \langle \alpha * \beta | c_I \rangle &= \log^*(\alpha * \beta)(b_I) \\ &= \log^*(\alpha)(b_I) + \log^*(\beta)(b_I) \\ &= \langle \alpha | c_I \rangle + \langle \beta | c_I \rangle. \end{aligned}$$

This formula can be interpreted as follows. As the \star product is dual to the coproduct on $S(\mathcal{X})$, the formula expresses the fact that the c_I are primitive elements in $S(\mathcal{X})$. It follows that, in the system of coordinates \mathcal{Y} , the group \mathcal{G} obeys the additive group law: the c_I -coordinate of $\alpha * \beta$ is the sum of the c_I -coordinates of α and β .

Several of the phenomena we have accounted for in the classical commutative case will resurface in the non-commutative context, although some differences will emerge as well, particularly when examining the so-called monotone and anti-monotone cases, which we will introduce in the upcoming section.

4. UNIVERSAL PRODUCTS: THE NON-COMMUTATIVE CASE

Consider now a non-commutative probability space (\mathcal{A}, φ) . The distributions of random variables (seen as elements of \mathcal{A}) are then encoded by the linear form (see again [20] for details), $\phi : T(\mathcal{A}) \rightarrow \mathbb{K}$,

$$a_1 \cdots a_n \longmapsto \phi(a_1 \cdots a_n) = \varphi(a_1 \mathbin{\vphantom{a}} \mathcal{A} \cdots \mathcal{A} a_n),$$

where $a_1 \cdots a_n$ stands now for a non-commutative monomial of degree n in the tensor algebra over \mathcal{A} (once again, not to be confused with the product of the a_i in \mathcal{A}). The distribution of $a \in \mathcal{A}$ is then obtained as the sequence of its moments $\phi(a^n)$.

Analogously to the commutative case, the (additive) convolution of two distributions of random variables X, Y belonging to \mathcal{A} could then be defined as the distribution of the sum $X' + Y'$, where X' and Y' are independent, having the same laws as X and Y . However, the analogy stops here. Indeed, whereas in the commutative case there is a unique notion of independence, there exist five distinct notions of independence in the non-commutative setting assuming certain axioms to hold: see [19] and the following developments in this section. The best way to approach this phenomenon in the context of the present article is through the notion of universal product – devised, among others, to create independent copies of random variables in the various non-commutative probability theories. We shall recall it below, and briefly explain how it gives rise to five distinct group structures that will be studied later on in the article. Note that we omit the discussion of the so-called tensor case which relies on constructions similar to the commutative case.

Let us write Alg_n for the category of non-unital associative algebras (of course, an algebra in Alg_n can have a unit, as \mathcal{A} does, but its existence is not required for the constructions presented here). Recall first the construction of free products.

Lemma 7. *The coproduct or free product of algebras, denoted by the symbol \circledast , in the category Alg_n is obtained as follows: let $\mathcal{A}_1, \mathcal{A}_2$ be two associative algebras, then:*

$$\mathcal{A}_1 \circledast \mathcal{A}_2 := \bigoplus_{\varepsilon \in Alt} \mathcal{A}_{\varepsilon_1} \otimes \cdots \otimes \mathcal{A}_{\varepsilon_n},$$

where Alt is the set of sequences $(\varepsilon_1, \dots, \varepsilon_n) \in \{1, 2\}^n$ such that $\varepsilon_{i+1} \neq \varepsilon_i$ for $i < n$. The canonical embeddings of \mathcal{A}_1 and \mathcal{A}_2 in $\mathcal{A}_1 \circledast \mathcal{A}_2$ are denoted respectively ι_1 and ι_2 . An element in a component $\mathcal{A}_{\varepsilon_1} \otimes \cdots \otimes \mathcal{A}_{\varepsilon_n}$ will be called an alternating tensor.

The product of two tensors $h_1 \otimes \cdots \otimes h_n$ and $h'_1 \otimes \cdots \otimes h'_m$ in $\mathcal{A}_1 \circledast \mathcal{A}_2$ is defined as the concatenation product $h_1 \otimes \cdots \otimes h_n \otimes h'_1 \otimes \cdots \otimes h'_m$, if h_n and h'_1 respectively belong to \mathcal{A}_1 and \mathcal{A}_2 or \mathcal{A}_2 and \mathcal{A}_1 and otherwise as: $h_1 \otimes \cdots \otimes (h_n \cdot_{\mathcal{A}_i} h'_1) \otimes \cdots \otimes h'_m$, where $h_n, h'_1 \in \mathcal{A}_i$, $i = 1, 2$. We will use from now on the word notation for tensor products in $\mathcal{A}_1 \circledast \mathcal{A}_2$: $h_1 \cdots h_n := h_1 \otimes \cdots \otimes h_n$.

The problem addressed by the theory of universal products is the one of extending linear forms ϕ_1 and ϕ_2 defined respectively on \mathcal{A}_1 and on \mathcal{A}_2 to a linear form $\phi_1 \phi_2$ on the free product $\mathcal{A}_1 \circledast \mathcal{A}_2$. One requires the construction to be associative, natural (that is, functorial in \mathcal{A}_1 and \mathcal{A}_2) and to satisfy the normalization conditions

$$\begin{aligned} (\phi_1 \phi_2) \circ \iota_1 &= \phi_1 \\ (\phi_1 \phi_2) \circ \iota_2 &= \phi_2 \\ (\phi_1 \phi_2)(\iota_1(a)\iota_2(b)) &= (\phi_1 \phi_2)(\iota_2(b)\iota_1(a)) = \phi_1(a)\phi_2(b), \end{aligned}$$

for $a \in \mathcal{A}_1$ and $b \in \mathcal{A}_2$. These requirements are respectively axioms U_2, U_3, U_4 of Muraki [19].

Solutions were classified by Speicher, Schürmann, and Muraki. They take the following five forms [19, Thm. 2.2]. In the subsequent definitions, ϕ_1 and ϕ_2 are linear forms on \mathcal{A}_1 respectively \mathcal{A}_2 .

Definition 8 (Universal Boolean product). *For an alternating tensor $a_1 \cdots a_n$,*

$$\phi_1 \bullet \phi_2(a_1 \cdots a_n) := \left(\prod_{a_i \in \mathcal{A}_1} \phi_1(a_i) \right) \left(\prod_{a_j \in \mathcal{A}_2} \phi_2(a_j) \right).$$

Definition 9 (Universal free product). *For an alternating tensor $a_1 \cdots a_n$,*

$$(6) \quad \phi_1 *_{\mathcal{F}} \phi_2(a_1 \cdots a_n) := \sum_{I \subseteq [n]} (-1)^{n-|I|+1} \phi_1 *_{\mathcal{F}} \phi_2 \left(\prod_{i \in I} a_i \right) \left(\prod_{\substack{j \notin I \\ a_j \in \mathcal{A}_1}} \phi_1(a_j) \right) \left(\prod_{\substack{k \notin I \\ a_k \in \mathcal{A}_2}} \phi_2(a_k) \right).$$

In this definition (by induction), it is understood that $\phi_1 *_{\mathcal{F}} \phi_2(\prod_{i \in \emptyset} a_i) := 1$ and that products $\prod_{i \in I} a_i$ are taken in increasing order of the indices (e.g., $\prod_{i \in \{3,5,7\}} a_i = a_3 a_5 a_7$). This convention for handling products will remain consistent throughout the entire article.

Remark 10. *It is difficult to get an explicit closed formula for the universal free product (6), and we are not aware of such a formula in the literature. However, the following can be said. Recall first that a noncrossing partition of a totally ordered set can always be obtained (although not uniquely) by recursively extracting intervals from the totally ordered set. For example, the noncrossing partition $\{1,7\} \amalg \{2,4,6\} \amalg \{3\} \amalg \{5\}$ can be obtained by first extracting $\{3\}$ (the remaining ordered set is $\{1,2,4,5,6,7\}$), then $\{5\}$ (the remaining ordered set is $\{1,2,4,6,7\}$), then $\{2,4,6\}$ (which is an interval in $\{1,2,4,6,7\}$) to be left with the last block, $\{1,7\}$. Conversely, any partition obtained by the process of recursively extracting intervals is noncrossing.*

Definition 9 then implies (by an elementary inductive argument based on this characterisation of noncrossing partitions) that there exist universal coefficients α_{π} such that

$$\phi_1 *_{\mathcal{F}} \phi_2(a_1 \cdots a_n) = \sum_{\substack{\pi = (\pi_1, \dots, \pi_k) \in NC_n \\ \pi \leq \{S, S^c\}}} \alpha_{\pi} \prod_{i=1}^k \phi_{1,2} \left(\prod_{j \in \pi_i} a_j \right),$$

where NC_n is the set of noncrossing partitions of $[n]$, $S = \{1, 3, 5, \dots\}$ is the set of odd integers in $[n]$, and where $\phi_{1,2}$ is ϕ_1 or ϕ_2 depending on whether the elements a_j for $j \in \pi_i$ belong to \mathcal{A}_1 or \mathcal{A}_2 . Recall that $\pi \leq \{S, S^c\}$ means that π is finer than $\{S, S^c\}$; remark that this condition implies that, if j and j' belong to the same block π_i , then a_j and $a_{j'}$ belong to the same algebra \mathcal{A}_1 or \mathcal{A}_2 .


Let now $b_1 \cdots b_n$ be a sequence of random variables belonging to \mathcal{A}_1 and \mathcal{A}_2 . We do not require anymore the sequence to be alternating, that is: if $b_i \in \mathcal{A}_1$ (resp. \mathcal{A}_2), then it is not necessarily the case that $b_{i+1} \in \mathcal{A}_2$ (resp. \mathcal{A}_1). We denote by S the set of indices i such that $b_i \in \mathcal{A}_1$. We say that a noncrossing partition $\pi \in NC_n$ of $[n]$ is *adapted to S* if and only if

$$\pi_{\text{Int}}(S) \leq \pi \leq \{S, [n] - S\},$$

i.e., there exists a splitting $\pi = \pi^1 \cup \pi^2$ of the partition into two subsets of blocks such that

- π^1 is a partition of S (and therefore π^2 a partition of $[n] - S$),
- given i , $1 \leq i < n$, in a block π_j of π^p (where $p \in \{1, 2\}$), then either $i+1 \in \pi_j$ or $i+1 \in \pi_l$ with $\pi_l \in \pi^q$ with $q \neq p$.

The set of noncrossing partitions of $[n]$ adapted to S is denoted ANC_n^S and if such a splitting exists for a given $\pi \in NC_n$, then it is unique and we write $(\pi^1, \pi^2) \in ANC_n^S$. In words,

FIGURE 1. Adapted noncrossing partition $\pi_1 \cup \pi_2$.FIGURE 2. Adapted noncrossing partition $\sigma_1 \cup \sigma_2$.FIGURE 3. Adapted noncrossing partition $\tau_1 \cup \tau_2$.

a noncrossing partition π is adapted to S if and only if 1) elements in each of its blocks correspond to lower indices of random variables in the same algebra \mathcal{A}_1 or \mathcal{A}_2 , and 2) when a_i and a_{i+1} belong to the same algebra, i and $i+1$ belong to the same block. As an example, we consider elements $b_1, b_2, b_3 \in \mathcal{A}_1$, $a_1, a_2, a_3, a_4 \in \mathcal{A}_2$, and the word $w = b_1 b_2 a_1 a_2 a_3 b_3 a_4$. Then $S = \{1, 2, 6\}$ and the pairs $\pi_1 = \{\{1, 2\}, \{6\}\}$, $\pi_2 = \{\{3, 4, 5\}, \{7\}\}$ (see Fig. 1) as well as $\sigma_1 = \{\{1, 2, 6\}\}$, $\sigma_2 = \{\{3, 4, 5\}, \{7\}\}$ (see Fig. 2), and $\tau_1 = \{\{1, 2\}, \{6\}\}$, $\tau_2 = \{\{3, 4, 5, 7\}\}$ (see Fig. 3) would be noncrossing partitions adapted to S .

We get:

Lemma 11. *There exist (universal) coefficients α_{π^1, π^2} such that for any sequence $b_1 \cdots b_n$ of random variables in \mathcal{A}_1 and \mathcal{A}_2*

$$(7) \quad \phi_1 *_f \phi_2(b_1 \cdots b_n) = \sum_{(\pi^1, \pi^2) \in \text{ANCS}_n^S} \alpha_{\pi^1, \pi^2} \left(\prod_{\pi_j \in \pi^1} \phi_1(b_{\pi_j}) \right) \left(\prod_{\pi_k \in \pi^2} \phi_2(b_{\pi_k}) \right).$$

The lemma follows by grouping in the word $b_1 \cdots b_n$ the consecutive letters that belong to the same algebra to form an alternating tensor to which Remark 10 can be applied. Notice that, given any $(\pi^1, \pi^2) \in \text{ANCS}_n^S$, one can recover the index set as $S = \bigcup_{\pi_j \in \pi^1} \pi_j$. Such properties of universal free products and reasoning on them can be found already in Speicher [25]. Details of the proof are left to the reader.

Let us point out for later use that when π^1 and π^2 contain each a single block, the coefficient α_{π^1, π^2} is obtained directly from the universal product formula in Definition 9 and is equal to one. More precisely, π^1, π^2 can only both contain a single block in the cases that $b_1 \cdots b_n = c_1 \cdots c_i d_{i+1} \cdots d_n$ with $c_k \in \mathcal{A}_p$ and $d_k \in \mathcal{A}_q$ (which corresponds to $n = 2$, $a_1 = c_1 \cdots c_i$, $a_2 = d_{i+1} \cdots d_n$ in Definition 9) or $b_1 \cdots b_n = c_1 \cdots c_i d_{i+1} \cdots d_j e_{j+1} \cdots e_n$ with $c_k, e_k \in \mathcal{A}_p$ and $d_k \in \mathcal{A}_q$ (which corresponds to $n = 3$, $a_1 = c_1 \cdots c_i$, $a_2 = d_{i+1} \cdots d_j$, $a_3 = e_{j+1} \cdots e_n$ in

Definition 9), where $p \neq q$. We apply Definition 9 in the first case with $I = \emptyset, \{1\}, \{2\}$ and in the second case with $I = \{1, 3\}$.

Definition 12 (Universal monotone product). *For an alternating tensor $a_1 \cdots a_n$,*

$$\phi_1 \blacktriangleright \phi_2(a_1 \cdots a_n) := \phi_1\left(\prod_{a_i \in \mathcal{A}_1} a_i\right) \prod_{a_j \in \mathcal{A}_2} \phi_2(a_j).$$

Definition 13 (Universal anti-monotone product). *For an alternating tensor $a_1 \cdots a_n$,*

$$\phi_1 \blacktriangleleft \phi_2(a_1 \cdots a_n) := \left(\prod_{a_i \in \mathcal{A}_1} \phi_1(a_i)\right) \phi_2\left(\prod_{a_j \in \mathcal{A}_2} a_j\right).$$

As already mentioned, we omit the tensor case, which would be given by the product

$$\phi_1\left(\prod_{a_i \in \mathcal{A}_1} a_i\right) \phi_2\left(\prod_{a_j \in \mathcal{A}_2} a_j\right).$$

In each of the above cases, if X, Y are two random variables in (\mathcal{A}, φ) , we let $(\mathcal{A}_1, \varphi_1)$ and $(\mathcal{A}_2, \varphi_2)$ be two copies of (\mathcal{A}, φ) embedded as usual in $\mathcal{A}_1 \otimes \mathcal{A}_2$ and X_1, Y_2 copies of X and Y respectively in \mathcal{A}_1 and \mathcal{A}_2 . We set $X' := \iota_1(X_1) \in \mathcal{A}_1 \otimes \mathcal{A}_2$ and $Y'' := \iota_2(Y_2) \in \mathcal{A}_1 \otimes \mathcal{A}_2$. The two variables X' and Y'' are independent for the five notions of independence and the respective products of states. The key difference between the five cases is the way the state map φ is extended to $\mathcal{A}_1 \otimes \mathcal{A}_2$. For example, in the monotone case, X' and Y'' are monotone independent for the state map $\phi_1 \blacktriangleright \phi_2$.

For each of the mentioned probability theories, the (additive) convolution of two distributions of variables X, Y belonging to (\mathcal{A}, φ) is then defined as the distribution of the sum $X' + Y''$. For example, the moment of order n associated to the Boolean, respectively monotone convolution of the distributions of X and Y is obtained by computing $\phi_1 \bullet \phi_2((X' + Y'')^n)$, respectively $\phi_1 \blacktriangleright \phi_2((X' + Y'')^n)$. In the first case (using notation from Section 2) we get:

$$\phi_1 \bullet \phi_2((X' + Y'')^n) = \sum_{S \subset [n]} \phi(X^{|S_1|}) \cdots \phi(X^{|S_l|}) \phi(Y^{|J_1^S|}) \cdots \phi(Y^{|J_k^S|}).$$

This expression is symmetric (and equal to $\sum_{S \subset [n]} \phi(Y^{|S_1|}) \cdots \phi(Y^{|S_l|}) \phi(X^{|J_1^S|}) \cdots \phi(X^{|J_k^S|})$): Boolean convolution is commutative.

In the second case:

$$\phi_1 \blacktriangleright \phi_2((X' + Y'')^n) = \sum_{S \subset [n]} \phi(X^{|S|}) \phi(Y^{|J_1^S|}) \cdots \phi(Y^{|J_k^S|}).$$

Notice that, since the monotone and anti-monotone products are non-symmetric, the order matters and the convolution of the distributions of X and Y is different in general from the convolution of the distributions of Y and X .

We are now in the position to develop the three group structures corresponding to free, Boolean and monotone probabilities, by analogy with the commutative case. As the anti-monotone product is simply obtained by exchanging the roles of \mathcal{A}_1 and \mathcal{A}_2 in the free product of algebras, its treatment is omitted here. We recall first the framework that permits to obtain these products as the products arising from a group of characters on a (not necessarily commutative or cocommutative) Hopf algebra. In the following sections, this approach is linked to the one derived from universal products.

5. PRODUCTS OF STATES

Hereafter, as was done in the classical case, we extend the notion of *state* (which we define here as a linear unital map on \mathcal{A}), to the notion of *generalized measure*, i.e., an arbitrary unital linear form on the tensor algebra $T(\mathcal{A})$. Remark that our notion of state is sometimes called a *formal state* (with the term state reserved for positive linear unital maps, which requires the assumption that \mathcal{A} is a $*$ -algebra that we do not assume here).

The generalized measure associated to a state φ is the unital linear form, denoted $\phi : T(\mathcal{A}) \rightarrow \mathbb{K}$:

$$a_1 \cdots a_n \longmapsto \phi(a_1 \cdots a_n) = \varphi(a_1 \underset{\mathcal{A}}{\cdot} \cdots \underset{\mathcal{A}}{\cdot} a_n).$$

A generalized measure ψ extends uniquely to a character Ψ on the double tensor algebra $T(T^+(\mathcal{A}))$ by requiring that

$$\Psi([w_1 | \cdots | w_n]) := \psi(w_1) \cdots \psi(w_n).$$

As ψ is the restriction of Ψ to $T^+(\mathcal{A})$, this correspondence defines a bijection between generalized measures and characters on $T(T^+(\mathcal{A}))$. We denote $\mathcal{G}(\mathcal{A})$ and $\mathcal{G}_T(\mathcal{A})$ the set of generalized measures on \mathcal{A} respectively the set of characters on $T(T^+(\mathcal{A}))$ and write

$$\Psi =: \text{char}(\psi) \quad \text{and} \quad \psi =: \text{gm}(\Psi)$$

for the bijection from $\mathcal{G}(\mathcal{A})$ to $\mathcal{G}_T(\mathcal{A})$ and its inverse.

Recall now the definition of a particular Hopf algebra structure on the double tensor algebra $T(T^+(\mathcal{A}))$.

Definition 14. *The map $\Delta : T^+(\mathcal{A}) \rightarrow (\mathbb{K} \oplus T^+(\mathcal{A})) \otimes T(T^+(\mathcal{A}))$ is defined by*

$$(8) \quad \Delta(a_1 \cdots a_n) := \sum_{S \subseteq [n]} a_S \otimes [a_{J_1^S} | \cdots | a_{J_k^S}] = \sum_{S \subseteq [n]} a_S \otimes [a_{J_{[n]}^S}],$$

with $\Delta(\mathbf{1}) := \mathbf{1} \otimes \mathbf{1}$ and $[a_{J_{[n]}^S}] := [a_{J_1^S} | \cdots | a_{J_k^S}]$. It is then extended multiplicatively to a coproduct on $T(T^+(\mathcal{A}))$

$$\Delta([w_1 | \cdots | w_m]) := \Delta(w_1) \cdots \Delta(w_m).$$

Theorem 15 ([10]). *The graded algebra $T(T^+(\mathcal{A}))$ equipped with the coproduct (8) is a graded connected non-commutative and non-cocommutative Hopf algebra.*

A crucial observation is that the coproduct (8) can be split into two parts as follows. On $T^+(\mathcal{A})$ define the *left half-coproduct* by

$$\Delta_{\prec}^+(a_1 \cdots a_n) := \sum_{1 \in S \subseteq [n]} a_S \otimes [a_{J_{[n]}^S}],$$

and

$$\Delta_{\prec}(a_1 \cdots a_n) := \Delta_{\prec}^+(a_1 \cdots a_n) - a_1 \cdots a_n \otimes \mathbf{1}.$$

The *right half-coproduct* is defined by

$$\Delta_{\succ}^+(a_1 \cdots a_n) := \sum_{1 \notin S \subseteq [n]} a_S \otimes [a_{J_{[n]}^S}],$$

and

$$\Delta_{\succ}(a_1 \cdots a_n) := \Delta_{\succ}^+(a_1 \cdots a_n) - \mathbf{1} \otimes a_1 \cdots a_n.$$

Which yields $\Delta = \Delta_{\prec}^+ + \Delta_{\succ}^+$, and $\Delta(w) = \Delta_{\prec}(w) + \Delta_{\succ}(w) + w \otimes \mathbf{1} + \mathbf{1} \otimes w$. This is extended to $T^+(T^+(\mathcal{A}))$ by defining

$$\begin{aligned}\Delta_{\prec}^+([w_1 | \cdots | w_m]) &:= \Delta_{\prec}^+(w_1) \Delta(w_2) \cdots \Delta(w_m) \\ \Delta_{\succ}^+([w_1 | \cdots | w_m]) &:= \Delta_{\succ}^+(w_1) \Delta(w_2) \cdots \Delta(w_m).\end{aligned}$$

By duality, the dual vector space $T(T^+(\mathcal{A}))^*$ of linear forms on $T(T^+(\mathcal{A}))$ is an algebra with respect to the convolution product defined for $f, g \in T(T^+(\mathcal{A}))^*$ by

$$(9) \quad f * g := (f \otimes g) \circ \Delta,$$

where we implicitly identify $\mathbb{K} \otimes \mathbb{K}$ with \mathbb{K} . The unit for the product (9) is the canonical projection, denoted ν hereafter, from $T(T^+(\mathcal{A}))$ to \mathbb{K} (sending both non-empty words $a_1 \cdots a_n$ and elements $[w_1 | \cdots | w_k]$, $k > 1$, in $T(T^+(\mathcal{A}))$ to zero). We summarize the group-theoretical implications of these ideas in the

Proposition 16. *Characters in $\mathcal{G}_T(\mathcal{A})$ form a group under convolution (9). The corresponding Lie algebra $g_T(\mathcal{A})$ of infinitesimal characters is characterised by $\alpha(\mathbf{1}) = 0$ and $\alpha([w_1 | \cdots | w_n]) = 0$, for $\alpha \in g_T(\mathcal{A})$ and $w_1, \dots, w_n \in T^+(\mathcal{A})$, $n \geq 2$. $\mathcal{G}_T(\mathcal{A})$ and $g_T(\mathcal{A})$ are related bijectively by the convolution exponential and logarithm, \exp^* respectively \log^* .*

The left and right half-convolution or half-shuffle products on $T^+(T^+(\mathcal{A}))^*$ are obtained by dualizing the half-coproducts:

$$(10) \quad f \prec g := (f \otimes g) \circ \Delta_{\prec} \quad \text{and} \quad f \succ g := (f \otimes g) \circ \Delta_{\succ},$$

which split the associative convolution product (9)

$$f * g = f \succ g + f \prec g.$$

The two operations in (10) are not associative. Indeed, they satisfy the so-called *half-shuffle identities* defining a shuffle algebra:

$$(11) \quad (f \prec g) \prec h = f \prec (g * h)$$

$$(12) \quad (f \succ g) \prec h = f \succ (g \prec h)$$

$$(13) \quad f \succ (g \succ h) = (f * g) \succ h.$$

Relations (11)-(13) are extended using Schützenberger's trick, that is, by setting $\nu \succ f := f$, $f \prec \nu := f$, $\nu \prec f := 0$ and $f \succ \nu := 0$ for $f \in T^+(T^+(\mathcal{A}))^*$ ($\nu \prec \nu$ and $\nu \succ \nu$ are left undefined, whereas $\nu * \nu = \nu$).

Recall finally (e.g. from [12]) the following proposition. A (left) pre-Lie algebra [8, Chap. 6] is a vector space V equipped with a bilinear product \triangleright such that the (left) pre-Lie identity

$$(x \triangleright y) \triangleright z - x \triangleright (y \triangleright z) = (y \triangleright x) \triangleright z - y \triangleright (x \triangleright z)$$

is satisfied. The notion of right pre-Lie algebra is defined analogously. The bracket $[x, y] := x \triangleright y - y \triangleright x$ satisfies the Jacobi identity and defines a Lie algebra structure on V . The next statement is a consequence of the fact that any shuffle algebra carries a (left) pre-Lie product defined by antisymmetrization of the half-shuffle products.

Proposition 17. *The space $T^+(T^+(\mathcal{A}))^*$ is equipped with a (left) pre-Lie algebra structure defined by the product*

$$f \triangleright g := f \succ g - g \prec f$$

*and the associated Lie bracket $[f, g] = f \triangleright g - g \triangleright f$ is equal to $f * g - g * f$.*

For $\alpha \in g_T(\mathcal{A})$ the *left* and *right half-shuffle*, or “time-ordered”, exponentials are defined by (see [10] for details)

$$\begin{aligned}\mathcal{E}_{\prec}(\alpha) &:= \exp^{\prec}(\alpha) := \nu + \sum_{n>0} \alpha^{\prec n} \\ \mathcal{E}_{\succ}(\alpha) &:= \exp^{\succ}(\alpha) := \nu + \sum_{n>0} \alpha^{\succ n},\end{aligned}$$

where $\alpha^{\prec n} := \alpha \prec (\alpha^{\prec n-1})$, $\alpha^{\prec 0} := \nu$ and $\alpha^{\succ n} := (\alpha^{\succ n-1}) \succ \alpha$, $\alpha^{\succ 0} := \nu$. They satisfy by definition the fixed point equations

$$(14) \quad \mathcal{E}_{\prec}(\alpha) = \nu + \alpha \prec \mathcal{E}_{\prec}(\alpha) \quad \mathcal{E}_{\succ}(\alpha) = \nu + \mathcal{E}_{\succ}(\alpha) \succ \alpha,$$

and provide bijections between the Lie algebra $g_T(\mathcal{A})$ and the group of characters $\mathcal{G}_T(\mathcal{A})$. Its inverses, known as *left* and *right half-shuffle* logarithms, \mathcal{L}_{\prec} respectively \mathcal{L}_{\succ} , can be deduced from the fixed point equations (14) using the half-shuffle identities (11)-(13):

$$(15) \quad \mathcal{L}_{\prec}(\Phi) = (\Phi - \nu) \prec \Phi^{-1} \quad \text{and} \quad \mathcal{L}_{\succ}(\Phi) = \Phi^{-1} \succ (\Phi - \nu).$$

With this structure in place, the shuffle algebra approach [13] permits to express moment-cumulant relations in non-commutative probability in terms of the convolution or shuffle exponential and logarithm, \exp^* respectively \log^* , and the half-shuffle exponentials and logarithms defined in equations (14) and (15). In addition, the unified setting covers relations between the different families of cumulants [2] in terms of a single fundamental identity in the group $\mathcal{G}_T(\mathcal{A})$

$$\Phi = \exp^*(\rho) = \mathcal{E}_{\prec}(\kappa) = \mathcal{E}_{\succ}(\beta),$$

and the corresponding infinitesimal relation in $g_T(\mathcal{A})$ [9, 12]. Here $\Phi \in \mathcal{G}_T(\mathcal{A})$ and ρ , κ , and β are infinitesimal characters in $g_T(\mathcal{A})$, which correspond respectively to monotone, free and boolean cumulants.

6. THE MONOTONE GROUP OF STATES

The previous two sections lead to the following definitions. We refer the reader to reference [10] for details (in particular the proof that the following constructions define groups – the next one is actually a rephrasing in terms of generalized measures of definitions given in [10] and other articles following the same approach).

Recall from Section 5 that $\mathcal{G}(\mathcal{A})$ denotes the set of generalized measures on \mathcal{A} and that there is a bijection $gm: \mathcal{G}_T(\mathcal{A}) \rightarrow \mathcal{G}(\mathcal{A})$, where $\mathcal{G}_T(\mathcal{A})$ is the set of characters on $T(T^+(\mathcal{A}))$.

Definition 18 (Monotone group of states, algebraic definition). *The set $\mathcal{G}(\mathcal{A})$ equipped with the product*

$$\phi * \psi := gm(char(\phi) * char(\psi))$$

is a group called the monotone group of states.

In the next proposition, the notation of Section 4 is used. We write \mathcal{B}_1 and \mathcal{B}_2 for two copies of the tensor algebra $T(\mathcal{A})$ and ϕ_1, ψ_2 for two copies of elements of $\phi, \psi \in \mathcal{G}(\mathcal{A})$ acting respectively on \mathcal{B}_1 and \mathcal{B}_2 . Given an element a in \mathcal{A} , we write a' and a'' for copies of a in \mathcal{B}_1 respectively \mathcal{B}_2 , and use the canonical embeddings ι_1 and ι_2 to view them as elements of the product $\mathcal{B}_1 \otimes \mathcal{B}_2$.

Proposition 19 (Monotone group of states, probabilistic definition). *The set $\mathcal{G}(\mathcal{A})$ of generalized measures equipped with the product*

$$(16) \quad \phi * \psi(a_1 \cdots a_n) := \phi_1 \blacktriangleright \psi_2((a'_1 + a''_1) \cdots (a'_n + a''_n)),$$

is a group that identifies with the monotone group of states of Definition 18.

Proof. The identity between the two definitions follows from the definition of the coproduct Δ on $T(T^+(\mathcal{A}))$ and the universal monotone product \blacktriangleright given in Definition 12. The definition of Δ indeed immediately leads, for the convolution product defined by Equation (9) to the identity

$$\phi * \psi(a_1 \cdots a_n) = \sum_{S \subseteq [n]} \phi(a_S) \psi(a_{J_1^S}) \cdots \psi(a_{J_k^S}).$$

In the second case (identity (16)), the same identity follows by an elementary enumerative combinatorics argument that we omit – it amounts to parametrizing by subsets of $[n]$ the products of a'_i s and a''_j s appearing in the expansion of $(a'_1 + a''_1) \cdots (a'_n + a''_n)$. \square

Let us redo now the analysis of the classical case in a non-commutative setting. Assume that \mathcal{A} has a basis $\mathcal{B} = (b_i)_{i \in \mathbb{N}}$. Regarding this assumption, recall that a wide class of non-commutative probability spaces are subalgebras of algebras of operators on a Hilbert space and have a dense subspace spanned by a Hilbert space basis.

Given a sequence $I = (i_1, \dots, i_k) \in \mathbb{N}^k$, we set $b_I := b_{i_1} \cdots b_{i_k}$. Notice that b_I is now a word in the b_i (and not anymore a commutative monomial as in the classical case). The set \mathcal{Y} of words b_I defines then a (complete) system of coordinates on $\mathcal{G}(\mathcal{A})$, that is, for $\phi \in \mathcal{G}(\mathcal{A})$,

$$\langle \phi | b_I \rangle := \phi(b_I).$$

The definition of the group law on $\mathcal{G}(\mathcal{A})$ implies that the b_I are representative functions on the group [8, 2.7 Representative functions]. We get the alternative presentation of $\mathcal{G}(\mathcal{A})$ as the group of characters of its Hopf algebra of representative functions generated by the elements of \mathcal{Y} :

Corollary 20. *The monotone group of states is canonically isomorphic to the group, denoted $\mathcal{G}_S(\mathcal{A})$ of characters of the Hopf algebra $S(T^+(\mathcal{A}))$ equipped with the product of polynomials over $T^+(\mathcal{A})$ and the coproduct whose restriction to the generators of the polynomial algebra $S(T^+(\mathcal{A}))$ is given by*

$$(17) \quad \begin{aligned} \Delta_m: T^+(\mathcal{A}) &\rightarrow (\mathbb{K} \oplus T^+(\mathcal{A})) \otimes S(T^+(\mathcal{A})) \\ \Delta_m(a_1 \cdots a_n) &:= \sum_{S \subseteq [n]} a_S \otimes (a_{J_1^S} | \cdots | a_{J_k^S}). \end{aligned}$$

We call Δ_m the monotone coproduct.

For example, let $a_1 a_2 a_3 \in T^+(\mathcal{A})$ and let us calculate the reduced monotone coproduct (that is, $\bar{\Delta}_m(x) := \Delta_m(x) - x \otimes \mathbf{1} - \mathbf{1} \otimes x$):

$$\begin{aligned} \bar{\Delta}_m(a_1 a_2 a_3) &= a_2 a_3 \otimes a_1 + a_1 a_2 \otimes a_3 + a_1 a_3 \otimes a_2 \\ &\quad + a_1 \otimes a_2 a_3 + a_2 \otimes (a_1 | a_3) + a_3 \otimes a_1 a_2. \end{aligned}$$

Notice the fact that $S(T^+(\mathcal{A}))$ is indeed a Hopf algebra (that is, the coproduct is coassociative as well as a map of associative unital algebras and $S(T^+(\mathcal{A}))$ has an antipode) can be

checked easily by hand – the proof is similar to the one for $T(T^+(\mathcal{A}))$, as defined above. However, it also follows automatically from the group structure of $\mathcal{G}(\mathcal{A})$; for general properties of (pro-affine) algebraic groups, and the bijection between the group and linear forms on $T^+(\mathcal{A})$ (or, equivalently, functions on the chosen set of coordinates \mathcal{Y}), see e.g. [8, Sect. 3.5 and 3.6].

The coproduct map (17) has a particular property: it is left-linear, that is, it maps the vector space $T^+(\mathcal{A})$ of the Hopf algebra generators of $S(T^+(\mathcal{A}))$ as a polynomial algebra to $(\mathbf{1} \oplus T^+(\mathcal{A})) \otimes S(T^+(\mathcal{A}))$. This property ensures that the following proposition holds (see e.g. [8, 6.4 Left-Linear Groups and Faà di Bruno]).

Let the map, called later on the linearised coproduct map,

$$(18) \quad \delta_m: T^+(\mathcal{A}) \rightarrow T^+(\mathcal{A}) \otimes T^+(\mathcal{A})$$

be defined in terms of the monotone coproduct (17),

$$\delta_m := (p \otimes p)\Delta_m,$$

where the map $p: S(T^+(\mathcal{A})) \rightarrow T^+(\mathcal{A})$ is the projector that is zero on $\mathbf{1}$ as well as on products in $S^+(T^+(\mathcal{A}))$, i.e., elements $(w|v)$ with $w \in T^+(\mathcal{A})$ and $v \in S^+(T^+(\mathcal{A}))$ are mapped to zero, and it is the identity on $T^+(\mathcal{A})$. The linearised coproduct can be described explicitly. Indeed, for any word $w = a_1 \cdots a_n \in T^+(\mathcal{A})$ we have

$$\delta_m(w) = \sum_{\emptyset \neq S_c \subsetneq [n]} a_{[n]-S_c} \otimes a_{S_c},$$

where $\emptyset \neq S_c \subsetneq [n]$ denotes a non-empty subset which has a single connected component, that is, a non-empty interval in $[n]$. For instance

$$\begin{aligned} \delta_m(a_1 a_2 a_3) &= a_2 a_3 \otimes a_1 + a_1 a_2 \otimes a_3 + a_1 a_3 \otimes a_2 + a_1 \otimes a_2 a_3 + a_3 \otimes a_1 a_2 \\ \delta_m(a_1 a_2 a_3 a_4) &= a_2 a_3 a_4 \otimes a_1 + a_1 a_3 a_4 \otimes a_2 + a_1 a_2 a_4 \otimes a_3 + a_1 a_2 a_3 \otimes a_4 + a_1 \otimes a_2 a_3 a_4 \\ &\quad + a_4 \otimes a_1 a_2 a_3 + a_1 a_2 \otimes a_3 a_4 + a_3 a_4 \otimes a_1 a_2 + a_1 a_4 \otimes a_2 a_3. \end{aligned}$$

Observe that δ_m is neither coassociative nor cocommutative. Instead, we find the following.

Proposition 21. *The linearised coproduct (18) satisfies the left pre-Lie coalgebra identity*

$$(19) \quad \mathfrak{a}_m = (\tau \otimes \text{id})\mathfrak{a}_m,$$

where $\mathfrak{a}_m := (\delta_m \otimes \text{id})\delta_m - (\text{id} \otimes \delta_m)\delta_m$. The map τ is defined by switching tensor products, $\tau(x \otimes y) := y \otimes x$.

For completeness, let us conclude the analysis of the monotone group of states with its Lie theoretic part. Once again, the following results are a consequence of the fact that $S(T^+(\mathcal{A}))$, as a polynomial Hopf algebra, is graded and connected (its degree zero component is the ground field) and left linear (this last property being relevant only for the assertions on the connections between the Lie and pre-Lie structures). These results are strictly similar to those holding for the group of characters on $T(T^+(\mathcal{A}))$ and its pre-Lie and Lie algebra of infinitesimal characters, and follow for the same reasons (see e.g. [12]). They are worth being stated, but we omit proofs.

A linear form $\kappa \in S(T^+(\mathcal{A}))^*$ is called an infinitesimal character, if $\kappa(\mathbf{1}) = 0$ (recall that $\mathbf{1}$ is the unit of the product of $S(T^+(\mathcal{A}))$), and if $\kappa(w_1 | \cdots | w_n) = 0$ for all $w_1, \dots, w_n \in T^+(\mathcal{A})$, $n \geq 2$. Let us write $g(\mathcal{A})$ for the set of linear forms μ on $T^+(\mathcal{A})$ such that $\mu(\mathbf{1}) = 0$ and $g_S(\mathcal{A})$ for the set of infinitesimal characters on $S(T^+(\mathcal{A}))$. There is an obvious bijection between the

two sets: the restriction map from $S(T^+(\mathcal{A}))^*$ to $T^+(\mathcal{A})^*$ defines for example the bijection from $g_S(\mathcal{A})$ to $g(\mathcal{A})$.

Infinitesimal characters are closed under commutator brackets defined on $S(T^+(\mathcal{A}))^*$ in terms of the convolution product \star_m on $S(T^+(\mathcal{A}))^*$ dual to the coproduct Δ_m on $S(T^+(\mathcal{A}))$

$$(20) \quad [\alpha, \beta] := \alpha \star_m \beta - \beta \star_m \alpha,$$

for $\alpha, \beta \in g_S(\mathcal{A})$. Hence, $g_S(\mathcal{A})$ is a Lie algebra. We call \star_m the monotone convolution product (on $S(T^+(\mathcal{A}))^*$). Since $\alpha(\mathbf{1}) = 0$ for $\alpha \in g_S(\mathcal{A})$, the exponential defined by its power series with respect to monotone convolution, $\exp^{\star_m}(\alpha)(w) := \nu + \sum_{j>0} \frac{1}{j!} \alpha^{\star_m j}(w)$, is a finite sum terminating at $j = \deg(w)$, and defines a bijection from $g_S(\mathcal{A})$ onto $\mathcal{G}_S(\mathcal{A})$. The compositional inverse of \exp^{\star_m} is given by the logarithm, $\log^{\star_m}(\nu + \gamma)(w) = \sum_{k \geq 1} \frac{(-1)^{k-1}}{k} \gamma^{\star_m k}(w)$, where $\gamma \in g_S(\mathcal{A})$ and ν is the canonical augmentation map from $S(T^+(\mathcal{A}))$ to the ground field. Again the sum terminates at $k = \deg(w)$ for any $w \in H$ as $\gamma(\mathbf{1}) = 0$. This is a general phenomenon [8, Sect. 3.5].

The exponential \exp^{\star_m} can be computed explicitly:

Theorem 22. *Let (\mathcal{A}, φ) be a non-commutative probability space with unital linear map $\varphi: \mathcal{A} \rightarrow \mathbb{K}$. Let Φ denote its extension to $S(T^+(\mathcal{A}))$ as a character. Let ρ be the infinitesimal character defined as $\rho := \log^{\star_m}(\Phi)$. For a word $w = a_1 \cdots a_n \in T^+(\mathcal{A})$, we have the relation*

$$(21) \quad \Phi(w) = \exp^{\star_m}(\rho)(w) = \sum_{j=1}^n \frac{\rho^{\star_m j}(w)}{j!} = \sum_{\pi \in NC_n} \frac{1}{t(\pi)!} h_\pi(a_1, \dots, a_n).$$

The tree factorial $t(\pi)!$ corresponds to the forest $t(\pi)$ of rooted trees encoding the nesting structure of the non-crossing partition $\pi \in NC_n$, see [2] for definitions, and $h_\pi(a_1, \dots, a_n) := \prod_{\pi_i \in \pi} \rho(a_{\pi_i})$.

Proof. We omit the proof of the theorem. Indeed, the calculation is the same as the one proving the identity when \star_m is replaced by the convolution product $*$ of linear forms on $T(T^+(\mathcal{A}))$ (see [12]). This follows from the fact that the coproducts (8) and (17) differ only by the fact that the terms on the right hand side belong respectively to $T(T^+(\mathcal{A}))$ and $S(T^+(\mathcal{A}))$. \square

Corollary 23. *Equation (21) reproduces the monotone moment-cumulant relation of [16]*

$$\Phi(w) = \sum_{\pi \in NC_n} \frac{1}{t(\pi)!} h_\pi(a_1, \dots, a_n).$$

This implies that $h_n(a_1, \dots, a_n) := \rho(a_1 \cdots a_n)$ identifies with the n th multivariate monotone cumulant maps from (4).

Corollary 24. *The equation also implies that the bijection \exp^{\star_m} between $g_S(\mathcal{A})$ and $\mathcal{G}_S(\mathcal{A})$ and its inverse \log^{\star_m} identify (up to canonical isomorphisms) with the bijection \exp^* , respectively its inverse \log^* between $g_T(\mathcal{A})$ and $\mathcal{G}_T(\mathcal{A})$, the Lie algebra of infinitesimal characters, respectively group of characters in $T(T^+(\mathcal{A}))^*$ (see [12]).*

Going back to Proposition 21, we can dualise the pre-Lie coproduct (18) and obtain a left pre-Lie product on $g(\mathcal{A})$

$$(22) \quad \alpha \triangleright \beta := (\alpha \otimes \beta) \circ \delta_m,$$

which satisfies the dual of (19), i.e., the left pre-Lie relation

$$(\alpha \triangleright \beta) \triangleright \gamma - \alpha \triangleright (\beta \triangleright \gamma) = (\beta \triangleright \alpha) \triangleright \gamma - \beta \triangleright (\alpha \triangleright \gamma).$$

By antisymmetrization, we have another Lie bracket on $g(\mathcal{A})$

$$(23) \quad \llbracket \alpha, \beta \rrbracket := \alpha \triangleright \beta - \beta \triangleright \alpha.$$

Proposition 25. *The space $g(\mathcal{A})$ is closed under the pre-Lie product (22), i.e., $(g(\mathcal{A}), \triangleright)$ is a pre-Lie algebra. Furthermore, on the space $g(\mathcal{A}) \cong g_S(\mathcal{A})$ the two Lie brackets (20) and (23) coincide, that is, $\llbracket \alpha, \beta \rrbracket = [\alpha, \beta]$, for $\alpha, \beta \in g(\mathcal{A})$.*

The proposition should be related to Proposition 17. Here, an important remark is in order: the pre-Lie products defined in these two propositions are related but different. However, they give rise to the same Lie algebra. This reflects the fact that some tools that are available when working with $T(T^+(\mathcal{A}))$, such as the half-shuffle products, are not available when working in what we call here the “probabilistic approach”, i.e., in the context of $S(T^+(\mathcal{A}))$.

7. THE BOOLEAN GROUP OF STATES

Seen from the Hopf algebraic point of view, the construction of a group of states in the Boolean and free probability cases results from a different approach than the one used in the monotone case.

Let us be more precise. We first introduce another notation. Given a linear map $\alpha \in T^+(\mathcal{A})^*$, we extend it to an infinitesimal character denoted $ichar(\alpha)$ on $T(T^+(\mathcal{A}))$. For $k \neq 1$, it maps by definition the components $T^+(\mathcal{A})^{\otimes k}$ of $T(T^+(\mathcal{A}))$ to zero. As such the extension is well defined. Conversely, we write igm for the inverse map from infinitesimal characters to elements of $T(\mathcal{A})^*$. The following definition and its motivation can be found in reference [13]. Recall the definition of the two half-shuffle exponentials and logarithms in Section 5.

Definition 26 (Boolean group of states, algebraic definition). *The set $\mathcal{G}(\mathcal{A})$ equipped with the product*

$$\phi \otimes \psi := gm(\mathcal{E}_{\succ}(\mathcal{L}_{\succ}(char(\phi)) + \mathcal{L}_{\succ}(char(\psi))))$$

is a group called the Boolean group of states.

Recall also the Boolean moment-cumulant relation (3), which we rewrite as

$$(24) \quad \phi(a_1 \cdots a_n) = \sum_{\pi \in \text{Int}_n} b_{\pi}(a_1 \cdots a_n),$$

where as usual $b_{\pi}(a_1 \cdots a_n) := \prod_{\pi_k \in \pi} b_{|\pi_k|}(a_{\pi_k})$. Recall that Int_n stands for the set of Boolean (i.e. interval) partitions of $[n]$. Given a non-commutative probability space (\mathcal{A}, φ) with the usual extension of φ to $\phi \in T^+(\mathcal{A})^*$, relations (24) define implicitly the Boolean cumulants $\{b_n\}_{n>0}$ in terms of ϕ respectively the state map φ . By Möbius inversion,

$$(25) \quad b_n(a_1 \cdots a_n) = \sum_{\pi \in \text{Int}_n} (-1)^{|\pi|-1} \phi_{\pi}(a_1 \cdots a_n).$$

Again, $\phi_{\pi}(a_1 \cdots a_n) := \prod_{\pi_k \in \pi} \phi(a_{\pi_k})$. Notice for later use that Equation (25) shows that $b_n(a_1 \cdots a_n) = \phi(a_1 \cdots a_n) + \text{l.o.t.}$, where l.o.t. refers to a sum of products of evaluations of ϕ on words of length strictly less than n . Recall also from [12] the following statement:

Proposition 27. *With the same notation as above, we have*

$$(26) \quad \begin{aligned} \phi &= gm(\mathcal{E}_{\succ}(i\text{char}(\mathbf{b}))), \\ \mathbf{b} &= igm(\mathcal{L}_{\succ}(\text{char}(\phi))). \end{aligned}$$

Let us translate these results in the language of algebraic groups. We fix a basis $\mathcal{B} = (b_i)_{i \in \mathbb{N}}$ of \mathcal{A} . As previously, given a sequence $I = (i_1, \dots, i_k) \in \mathbb{N}^k$, we set $b_I := b_{i_1} \cdots b_{i_k} \in T^+(\mathcal{A})$. The set \mathcal{Y} of words b_I defines then a (complete) system of coordinates on $\mathcal{G}(\mathcal{A})$. The definition of the Boolean group law on $\mathcal{G}(\mathcal{A})$ implies that $\phi \otimes \psi(b_I)$ can be expanded as a linear combination of products of terms $\phi(b_J)$ and $\psi(b_K)$, where J and K stand for subsets of I . In other words, the b_I are representative functions on the group and $\mathcal{G}(\mathcal{A})$ is the group of characters of its Hopf algebra of representative functions generated by the elements of \mathcal{Y} (see again [8] for details). From (25) and the remark that follows the equation, the monomials

$$K_{b,I} := \sum_{\pi \in \text{Int}_n} (-1)^{|\pi|-1} \prod_{\pi_i \in \pi} b_{\pi_i} \in S(T^+(\mathcal{A}))$$

form another system of coordinates of the group that we call the *Boolean coordinates* and denote by $\mathcal{B}ool$.¹ From the group law \otimes in Definition 26 together with (26), we obtain

$$\mathcal{L}_{\succ}(\text{char}(\phi \otimes \psi)) = \mathcal{L}_{\succ}(\text{char}(\phi)) + \mathcal{L}_{\succ}(\text{char}(\psi))$$

and

$$\phi \otimes \psi(K_{b,I}) = \phi(K_{b,I}) + \psi(K_{b,I}).$$

Hence, in the system of Boolean coordinates, the group law is additive and we get the alternative presentation of the group $(\mathcal{G}(\mathcal{A}), \otimes)$:

Corollary 28. *The Boolean group of states is, up to a canonical isomorphism, the group of characters of the Hopf algebra $S(\mathcal{B}ool) \cong S(T^+(\mathcal{A}))$ equipped with the product of polynomials over $\mathcal{B}ool$ and the coproduct*

$$\Delta_B(K_{b,I}) := K_{b,I} \otimes \mathbf{1} + \mathbf{1} \otimes K_{b,I}.$$

In the following proposition, we use again freely the notation of Section 4. We write \mathcal{B}_1 and \mathcal{B}_2 for two copies of the tensor algebra $T(\mathcal{A})$ and ϕ_1, ψ_2 for two copies of elements $\phi, \psi \in \mathcal{G}(\mathcal{A})$ acting respectively on \mathcal{B}_1 and \mathcal{B}_2 . Given an element a in \mathcal{A} , we write a' and a'' for copies of a in \mathcal{B}_1 respectively \mathcal{B}_2 , both naturally embedded into the product $\mathcal{B}_1 \otimes \mathcal{B}_2$

Proposition 29 (Boolean group of states, probabilistic definition). *The set $\mathcal{G}(\mathcal{A})$ equipped with the product*

$$\phi \otimes \psi(a_1 \cdots a_n) := \phi_1 \bullet \psi_2((a'_1 + a''_1) \cdots (a'_n + a''_n)),$$

is a group that identifies with the Boolean group of states defined previously.

We will decompose the proof of the proposition into several steps. In the following arguments, we will use only the fact that the probabilistic definition of the product implies its associativity (the existence of a group structure will follow later from the identification of the two definitions of the product). Let us first compute explicitly the product law of $\mathcal{G}(\mathcal{A})$ as defined in Proposition 29 in the coordinates \mathcal{Y} , that is, the corresponding coproduct that

¹As pointed out to us by the referee, they can be understood as Boolean analog of (classical) generalized k -statistics, advocated for example by T. P. Speed [24]. The same remark applies in the free case presented in the next section.

we denote temporarily Δ_b . From the definition of the product \bullet of states, we obtain the cocommutative coproduct map:

$$\begin{aligned} \Delta_b: T^+(\mathcal{A}) &\rightarrow S(T^+(\mathcal{A})) \otimes S(T^+(\mathcal{A})), \\ (27) \quad \Delta_b(a_1 \cdots a_n) &:= \sum_{S \subseteq [n]} (a_{S_1} | \cdots | a_{S_l}) \otimes (a_{J_1^S} | \cdots | a_{J_k^S}). \end{aligned}$$

It is extended multiplicatively to the polynomial algebra $S(T^+(\mathcal{A}))$ by $\Delta_b(\mathbf{1}) := \mathbf{1} \otimes \mathbf{1}$ and

$$\Delta_b(w_1 | \cdots | w_n) := \Delta_b(w_1) \cdots \Delta_b(w_n).$$

For example, the reduced coproduct action on the word $a_1 a_2 a_3$ yields:

$$\begin{aligned} \overline{\Delta}_b(a_1 a_2 a_3) &= a_2 a_3 \otimes a_1 + a_1 a_2 \otimes a_3 + (a_1 | a_3) \otimes a_2 \\ &\quad + a_1 \otimes a_2 a_3 + a_2 \otimes (a_1 | a_3) + a_3 \otimes a_1 a_2. \end{aligned}$$

We write H_b for the Hopf algebra $S(T^+(\mathcal{A}))$ equipped with the product of polynomials and with the coproduct defined in (27). The corresponding convolution product of linear forms on H_b is denoted \star_b .

Lemma 30. *Let $\alpha_1, \dots, \alpha_k$ be infinitesimal characters on H_b and $w = a_1 \cdots a_n \in T(\mathcal{A})$ a word. Then, we have:*

$$\langle \alpha_1 \star_b \cdots \star_b \alpha_k, w \rangle = \langle \alpha_1 \otimes \cdots \otimes \alpha_k, \sum_{\pi_1, \dots, \pi_k} a_{\pi_1} \otimes \cdots \otimes a_{\pi_k} \rangle,$$

where the sum is over all disjoint subsets π_1, \dots, π_k of $[n]$ such that $\pi_1 \cup \cdots \cup \pi_k = [n]$ and each π_i is an interval (that is, the sum is over ordered Boolean partitions: Boolean partitions equipped with an arbitrary total ordering of their blocks).

Proof. From the definition of Δ_b , only Boolean partitions can occur in the expansion of $\langle \alpha_1 \star_b \cdots \star_b \alpha_k, w \rangle$ that is obtained recursively from the associativity of \star_b . We get that this expansion has necessarily the form

$$\langle \alpha_1 \otimes \cdots \otimes \alpha_k, \sum_{\pi_1, \dots, \pi_k} \mu_\pi \cdot a_{\pi_1} \otimes \cdots \otimes a_{\pi_k} \rangle,$$

for certain scalar coefficients μ_π , where the sum is over ordered Boolean partitions $\pi = \{\pi_1, \dots, \pi_k\}$. The statement in the lemma amounts therefore to the assertion that $\mu_\pi = 1$, for any π . However, as the \star_b product is commutative (or, equivalently, as the coproduct Δ_b is cocommutative), $\mu_\pi = \mu_{\sigma\pi}$, for any permutation σ of $[k]$ acting on π by permutation of the blocks. We can therefore obtain a proof of the statement in the lemma by showing that $\mu_\pi = 1$ in the particular case where the blocks of π (which are intervals) are in natural order.

Let us prove this last property by induction on the number k of blocks. The base case $k = 1$ is obvious. Let us fix a canonically ordered Boolean partition π on $[n]$ with k blocks and assume that $\pi_k = \{p+1, \dots, n\}$. The formula for the \star_b product shows that, for an arbitrary linear form β on H_b , we have that (as α_k is infinitesimal):

$$\begin{aligned} \langle \beta \star_b \alpha_k, w \rangle &= \sum_{r \leq n} \langle \beta \otimes \alpha_k, a_1 \cdots a_r \otimes a_{r+1} \cdots a_n + a_{r+1} \cdots a_n \otimes a_1 \cdots a_r \rangle \\ &\quad + \sum_{1 \leq r < l < n} \langle \beta \otimes \alpha_k, (a_1 \cdots a_{r-1} | a_{l+1} \cdots a_n) \otimes a_r \cdots a_l \rangle. \end{aligned}$$

Only the component $\langle \beta \otimes \alpha_k, a_1 \cdots a_p \otimes a_{p+1} \cdots a_n + a_{p+1} \cdots a_n \otimes a_1 \cdots a_p \rangle$ is relevant for our computation. Replacing β by $\alpha_1 \star_b \cdots \star_b \alpha_{k-1}$, we get that μ_π is the coefficient of

$$\langle \alpha_1 \otimes \cdots \otimes \alpha_{k-1}, a_{\pi_1} \otimes \cdots \otimes a_{\pi_{k-1}} \rangle,$$

in the expansion of $\langle \alpha_1 \star_b \cdots \star_b \alpha_{k-1}, a_1 \cdots a_p \rangle$, which is equal to 1, by the induction hypothesis. This completes the proof. \square

Theorem 31. *Let (\mathcal{A}, φ) be a non-commutative probability space and Φ the extension of φ to H_b as a character. Let α be the infinitesimal character*

$$\alpha := \log^{\star_b}(\Phi).$$

For the word $w = a_1 \cdots a_n \in T^+(\mathcal{A})$ we obtain

$$\Phi(w) = \exp^{\star_b}(\alpha)(w) = \sum_{j=1}^n \frac{\alpha^{\star_b j}}{j!}(w) = \sum_{I \in \text{Int}_n} \alpha_I(a_1, \dots, a_n).$$

In particular, $\text{igm}(\alpha) = b$, the Boolean cumulant map.

Proof. Indeed, by Lemma 30:

$$\alpha^{\star_b k}(w) = k! \sum_{\pi} \prod_{\pi_i} \alpha(a_{\pi_i}),$$

where the sum is over Boolean partitions with k blocks π_1, \dots, π_k . The factor $k!$ appears since there are $k!$ ways to order the blocks π_1, \dots, π_k to form an ordered Boolean partition. The conclusion follows by summation over k . \square

Proof of Proposition 29. We are now in the position to conclude the proof that the product in Proposition 29 identifies with the one in Definition 26. For notational clarity, we write temporarily \otimes' for the product defined in Proposition 29.

By Corollary 28, it is enough to prove that, given two elements ϕ and ψ in $\mathcal{G}(\mathcal{A})$, we have

$$\phi \otimes' \psi(K_{b,I}) = \phi(K_{b,I}) + \psi(K_{b,I})$$

for I arbitrary. For notational simplicity and without loss of generality, we treat the generic case where $I = (1, \dots, n)$.

By Theorem 31, $\log^{\star_b}(\Phi)(b_1 \cdots b_n) = b(b_1 \cdots b_n) = \phi(K_{b,I})$. Therefore, as Δ_b encodes the definition of the product \otimes' ,

$$\phi \otimes' \psi(K_{b,I}) = \Phi \star_b \Psi(K_{b,I}) = \log^{\star_b}(\Phi \star_b \Psi)(b_1 \cdots b_n),$$

where we write Φ and Ψ for the extensions of ϕ respectively ψ to characters on H_b . However, as \star_b is commutative, $\log^{\star_b}(\Phi \star_b \Psi) = \log^{\star_b}(\Phi) + \log^{\star_b}(\Psi)$, and we get finally

$$\phi \otimes' \psi(K_{b,I}) = (\log^{\star_b}(\Phi) + \log^{\star_b}(\Psi))(b_1 \cdots b_n) = \phi(K_{b,I}) + \psi(K_{b,I}),$$

which yields $\phi \otimes' \psi = \phi \otimes \psi$ and concludes the proof. \square

8. THE FREE GROUP OF STATES

We employ freely the notation used in the previous section. Although the arguments are essentially the same as in the Boolean case, we repeat them for the sake of completeness; the present article aims at being a reference on the topic of algebraic groups in non-commutative probability. The following definition and its motivations can be found in reference [13]. Recall the definition of the half-shuffle exponentials and logarithms in Section 5.

Definition 32 (Free probability group of states, algebraic definition). *The set $\mathcal{G}(\mathcal{A})$ equipped with the product*

$$\phi \otimes \psi := gm(\mathcal{E}_{\prec}(\mathcal{L}_{\prec}(\text{char}(\phi)) + \mathcal{L}_{\prec}(\text{char}(\psi))))$$

is a group called the free convolution group of states.

Recall also the moment-cumulant relations in free probability (2), which we rewrite as

$$(28) \quad \phi(a_1 \cdots a_n) = \sum_{\pi \in NC_n} k_{\pi}(a_1 \cdots a_n),$$

where as usual $k_{\pi}(a_1 \cdots a_n) := \prod_{\pi_k \in \pi} k_{|\pi_k|}(a_{\pi_k})$ and NC_n stands for the set of noncrossing partitions of $[n]$. Given (\mathcal{A}, φ) , let $\phi \in T^+(\mathcal{A})^*$ be the usual extension of the state map φ . Relations (28) define implicitly the family of free cumulants $\{k_n\}_{n>0}$ in terms of the linear form ϕ respectively φ . By Möbius inversion, we can express cumulants in terms of moments

$$(29) \quad k_n(a_1 \cdots a_n) = \sum_{\pi \in NC_n} \mu_{NC}(\pi, 1_n) \phi_{\pi}(a_1 \cdots a_n),$$

where $\mu_{NC}(\pi, 1_n)$ denotes the Möbius function on the lattice of noncrossing partitions [20]. Notice for later use that this equation shows that $k_n(a_1 \cdots a_n) = \phi(a_1 \cdots a_n) + \text{l.o.t.}$, where l.o.t. refers again to a sum of products of evaluations of φ on words of length strictly less than n . Recall also from [12] the following statement:

Proposition 33. *With the same notation as above, we have*

$$(30) \quad \begin{aligned} \phi &= gm(\mathcal{E}_{\prec}(\text{ichar}(\mathbf{k}))), \\ \mathbf{k} &= igm(\mathcal{L}_{\prec}(\text{char}(\phi))). \end{aligned}$$

Let us translate these results into the language of algebraic groups. Recall our notation: as before, we fix a basis $\mathcal{B} = (b_i)_{i \in \mathbb{N}}$ of \mathcal{A} with $b_I := b_{i_1} \cdots b_{i_k}$, for $I = (i_1, \dots, i_k) \in \mathbb{N}^k$. The set \mathcal{Y} of words b_I defines then a (complete) system of coordinates on $\mathcal{G}(\mathcal{A})$. The definition of the free probability group law on $\mathcal{G}(\mathcal{A})$ implies that the terms in $\phi \otimes \psi(b_I)$ rewrite as linear combinations of products of terms $\phi(b_J)$ and $\psi(b_K)$, where J and K stand for subsets of I . In other words, the b_I are again representative functions on the group and $\mathcal{G}(\mathcal{A})$ is the group of characters of its Hopf algebra of representative functions generated by the elements of \mathcal{Y} (see again [8] for details). From (29) and the remark that follows the equation, the monomials

$$K_{f,I} := \sum_{\pi \in NC_n} \mu_{NC}(\pi, 1_n) \prod_{\pi_i \in \pi} b_{\pi_i}$$

form another system \mathcal{F} of coordinates of the group that we call the *free coordinates*. The group law \otimes in Definition 32 and (30) yield

$$\mathcal{L}_{\prec}(\text{char}(\phi \otimes \psi)) = \mathcal{L}_{\prec}(\text{char}(\phi)) + \mathcal{L}_{\prec}(\text{char}(\psi))$$

and

$$\phi \otimes \psi(K_{f,I}) = \phi(K_{f,I}) + \psi(K_{f,I}).$$

Hence, in the system of free coordinates, the group law is additive and we get the alternative presentation of the group $(\mathcal{G}(\mathcal{A}), \otimes)$:

Corollary 34. *The free probability group of states is, up to a canonical isomorphism, the group of characters of the Hopf algebra $S(\mathcal{F}) \cong S(T^+(\mathcal{A}))$ equipped with the product of polynomials over \mathcal{F} and the coproduct*

$$\Delta_f(K_{f,I}) := K_{f,I} \otimes \mathbf{1} + \mathbf{1} \otimes K_{f,I}.$$

In the following proposition, we use again freely the notation of Section 4. We write \mathcal{F}_1 and \mathcal{F}_2 for two copies of the tensor algebra $T(\mathcal{A})$ and ϕ_1, ψ_2 for two copies of elements $\phi, \psi \in \mathcal{G}(\mathcal{A})$ acting respectively on \mathcal{F}_1 and \mathcal{F}_2 . Given an element a in \mathcal{A} , we write a' and a'' for copies of a in \mathcal{F}_1 respectively \mathcal{F}_2 .

Proposition 35 (Free probability group of states, probabilistic definition). *The set $\mathcal{G}(\mathcal{A})$ equipped with the product*

$$\phi \otimes \psi(a_1 \cdots a_n) := \phi_1 *_f \psi_2((a'_1 + a''_1) \cdots (a'_n + a''_n)),$$

is a group that identifies with the free probability group of states defined previously.

We will decompose the proof that the two definitions of the product agree into several steps. In the following arguments, we will use only the fact that the probabilistic definition of the product implies its associativity (the existence of a group structure will follow later from the identification of the two definitions of the product). Let us first compute explicitly the product law of $\mathcal{G}(\mathcal{A})$ as defined in Proposition 35 in the coordinates \mathcal{Y} , that is, the corresponding coproduct that we denote temporarily Δ_f . We get from the definition of the product $*_f$ of states and Equation (7) the coproduct map:

$$\Delta_f: T^+(\mathcal{A}) \rightarrow S(T^+(\mathcal{A})) \otimes S(T^+(\mathcal{A})),$$

$$(31) \quad \Delta_f(a_1 \cdots a_n) := \sum_{S \subseteq [n]} \sum_{(\pi^1, \pi^2) \in \text{ANC}_n^S} \alpha_{\pi^1, \pi^2} \left(\prod_{\pi_j \in \pi^1} a_{\pi_j} \right) \otimes \left(\prod_{\pi_k \in \pi^2} a_{\pi_k} \right).$$

It is extended multiplicatively to the polynomial algebra $S(T^+(\mathcal{A}))$ by $\Delta_f(\mathbf{1}) := \mathbf{1} \otimes \mathbf{1}$ and

$$\Delta_f(w_1 | \cdots | w_n) := \Delta_f(w_1) \cdots \Delta_f(w_n).$$

As observed above, its coassociativity follows by duality from the associativity of \otimes . We can calculate the coproduct at low orders from the inductive formula for universal free products. Let $a_1 a_2 a_3 a_4 \in T^+(\mathcal{A})$. We have

$$\begin{aligned} \Delta_f(a_1 a_2) &= a_1 a_2 \otimes \mathbf{1} + a_1 \otimes a_2 + a_2 \otimes a_1 + \mathbf{1} \otimes a_1 a_2. \\ \Delta_f(a_1 a_2 a_3) &= a_1 a_2 a_3 \otimes \mathbf{1} + a_1 \otimes a_2 a_3 + a_1 a_2 \otimes a_3 + a_1 a_3 \otimes a_2 \\ &\quad + a_2 \otimes a_1 a_3 + a_3 \otimes a_1 a_2 + a_2 a_3 \otimes a_1 + \mathbf{1} \otimes a_1 a_2 a_3 \\ \Delta_f(a_1 a_2 a_3 a_4) &= (\text{id} + \tau)(a_1 a_2 a_3 a_4 \otimes \mathbf{1} + a_1 \otimes a_2 a_3 a_4 + a_1 a_2 \otimes a_3 a_4 + a_1 a_4 \otimes a_2 a_3 \\ &\quad + a_1 a_2 a_3 \otimes a_4 + a_1 a_2 a_4 \otimes a_3 + a_1 a_3 a_4 \otimes a_2 \\ &\quad + (a_1 | a_3) \otimes a_2 a_4 + a_1 a_3 \otimes (a_2 | a_4) - (a_1 | a_3) \otimes (a_2 | a_4)). \end{aligned}$$

Here, as before, the map τ switches tensor products, i.e., $\tau(x \otimes y) := y \otimes x$.

We write H_f for the Hopf algebra $S(T^+(\mathcal{A}))$ equipped with the product of polynomials and with the cocommutative coproduct (31). The corresponding convolution product of linear forms on H_f is denoted \star_f .

Lemma 36. *Let $\kappa_1, \dots, \kappa_k$ be infinitesimal characters on H_f and $w = a_1 \cdots a_n \in T(\mathcal{A})$ a word. Then, we have:*

$$\langle \kappa_1 \star_f \cdots \star_f \kappa_k, w \rangle = \langle \kappa_1 \otimes \cdots \otimes \kappa_k, \sum_{\pi_1, \dots, \pi_k} a_{\pi_1} \otimes \cdots \otimes a_{\pi_k} \rangle,$$

where the sum is over all ordered noncrossing partitions of $[n]$ (that is noncrossing partitions equipped with an arbitrary total ordering of their blocks).

Proof. From the definition of Δ_f , only noncrossing partitions can occur in the expansion of $\langle \kappa_1 \star_f \cdots \star_f \kappa_k, w \rangle$ that is obtained recursively from the associativity of \star_f . This expansion has necessarily the form

$$\langle \kappa_1 \otimes \cdots \otimes \kappa_k, \sum_{\pi_1, \dots, \pi_k} \mu_\pi \cdot a_{\pi_1} \otimes \cdots \otimes a_{\pi_k} \rangle,$$

for certain scalar coefficients μ_π , where the sum is over ordered noncrossing partitions. The lemma amounts therefore to the statement that $\mu_\pi = 1$, for any π . However, as Δ_f is cocommutative, the \star_f product is commutative, i.e., $\mu_\pi = \mu_{\sigma\pi}$, for any permutation σ of $[k]$ acting on π by permutation of the blocks. We can therefore obtain a proof of the statement by showing that $\mu_\pi = 1$ in the particular case where the blocks of π are in a total order compatible with the usual tree ordering of blocks of noncrossing partition (that is, the one defined by $\pi_i > \pi_j$ if and only if $\forall k \in \pi_i, \exists l, m \in \pi_j$ such that $l < k < m$). This choice implies in particular that the maximal block π_k for the total order is an interval $\pi_k = \{p+1, \dots, m\}$. We require furthermore that it is the rightmost interval of π . This last condition implies (by general elementary properties of noncrossing partitions) that $\pi - \{\pi_k\}$ cannot be decomposed as the union of a partition of $[p]$ and of $\{m+1, \dots, n\}$.

Let us prove now the property by induction on k . The base case $k = 1$ is obvious. Let us fix an ordered noncrossing partition π of $[n]$ with k blocks as above: in particular we assume that $\pi_k = \{p+1, \dots, m\}$ is the rightmost interval in π . The formula for the \star_f product shows that, for an arbitrary linear form β on H_f , we have that (as κ_k is infinitesimal):

$$\begin{aligned} \langle \beta \star_f \kappa_k, w \rangle &= \sum_{1 \leq r < l \leq n} \langle \beta \otimes \kappa_k, a_1 \cdots a_{r-1} a_{l+1} \cdots a_n \otimes a_r \cdots a_l \rangle \\ &+ \sum_{1 < r < l < n} \langle \beta \otimes \kappa_k, \alpha_{\{\{1, \dots, r-1\}, \{l+1, \dots, n\}\}, \{r, \dots, l\}} (a_1 \cdots a_{r-1} | a_{l+1} \cdots a_n) \otimes a_r \cdots a_l \rangle. \end{aligned}$$

As $\pi - \{\pi_k\}$ cannot be decomposed as the union of a partition of $[p]$ and of $\{m+1, \dots, n\}$, only the component $\langle \beta \otimes \kappa_k, a_1 \cdots a_p a_{m+1} \cdots a_n \otimes a_{p+1} \cdots a_m \rangle$ is relevant for our computation. Replacing β by $\kappa_1 \star_f \cdots \star_f \kappa_{k-1}$, we get that μ_π is the coefficient of

$$\langle \kappa_1 \otimes \cdots \otimes \kappa_{k-1}, a_{\pi_1} \otimes \cdots \otimes a_{\pi_{k-1}} \rangle,$$

in the expansion of $\langle \kappa_1 \star_f \cdots \star_f \kappa_{k-1}, a_1 \cdots a_p a_{m+1} \cdots a_n \rangle$, that is equal to one, by the induction hypothesis. This completes the proof. \square

Theorem 37. *Let (\mathcal{A}, φ) be a non-commutative probability space and Φ the extension of φ to H_f as a character. Let κ be the infinitesimal character*

$$\kappa := \log^{\star_f}(\Phi).$$

For the word $w = a_1 \cdots a_n \in T^+(\mathcal{A})$ we get:

$$\Phi(w) = \exp^{\star_f}(\kappa)(w) = \sum_{j=1}^n \frac{\kappa^{\star_f j}}{j!}(w) = \sum_{\pi \in NC_n} \kappa_\pi(a_1, \dots, a_n).$$

In particular, $\text{igm}(\kappa) = \mathbf{k}$, the free cumulant map.

Proof. Indeed, by Lemma 36:

$$\kappa^{\star_f k}(w) = k! \sum_{\pi} \prod_{\pi_i} \kappa(a_{\pi_i}),$$

where the sum is over all $\pi \in NC_n$ with k blocks π_1, \dots, π_k . The factor $k!$ appears since there are $k!$ ways to order the blocks. The conclusion follows by summation over k . \square

Proof of Proposition 35. We are now in the position to conclude the proof that the product in Proposition 35 identifies with the one in Definition 32. For notational clarity, we write temporarily \otimes' for the product defined in Proposition 35.

By Corollary 34, it is enough to prove that, given two elements ϕ and ψ in $\mathcal{G}(\mathcal{A})$,

$$\phi \otimes' \psi(K_{f,I}) = \phi(K_{f,I}) + \psi(K_{f,I}),$$

for I arbitrary. For notational simplicity and without loss of generality, we treat the generic case where $I = (1, \dots, n)$.

By Theorem 37, we have $\log^{\star_f}(\Phi)(b_1 \cdots b_n) = \mathbf{k}(b_1 \cdots b_n) = \phi(K_{f,I})$. Therefore, as Δ_f encodes the definition of the product \otimes' ,

$$\phi \otimes' \psi(K_{f,I}) = \Phi \star_f \Psi(K_{f,I}) = \log^{\star_f}(\Phi \star_f \Psi)(b_1 \cdots b_n),$$

where we write Φ and Ψ for the extension of ϕ and ψ to characters on H_f . However, as \star_f is commutative, $\log^{\star_f}(\Phi \star_f \Psi) = \log^{\star_f}(\Phi) + \log^{\star_f}(\Psi)$, and we get finally

$$\phi \otimes' \psi(K_{f,I}) = (\log^{\star_f}(\Phi) + \log^{\star_f}(\Psi))(b_1 \cdots b_n) = \phi(K_{f,I}) + \psi(K_{f,I}),$$

which yields $\phi \otimes' \psi = \phi \otimes \psi$ and concludes the proof. \square

REFERENCES

- [1] N. I. Akhiezer, *The classical moment problem and some related questions in analysis*, Hafner Publishing Co., New York, 1965.
- [2] O. Arizmendi, T. Hasebe, F. Lehner, C. Vargas, *Relations between cumulants in non-commutative probability*, *Advances in Mathematics* **282**, (2015) 56-92.
- [3] A. Ben Ghorbal, M. Schürmann, *On the algebraic foundations of non-commutative probability theory*, Preprintreihe Mathematik **8/99**, Universität Greifswald (1999).
- [4] A. Ben Ghorbal, M. Schürmann, *Non-commutative notions of stochastic independence*, *Mathematical Proceedings of the Cambridge Philosophical Society* **133**, (2002) 531-561.
- [5] A. Ben Ghorbal, M. Schürmann, *Quantum Lévy processes on dual groups*, *Mathematische Zeitschrift* **251**, (2005) 147-165.

- [6] P. Biane, *Free probability and combinatorics*, Proceedings of the International Congress of Mathematicians, Vol. II (Beijing, 2002), 765-774, Higher Ed. Press, Beijing, 2002.
- [7] T. S. Blyth, *Lattices and ordered algebraic structures*. Universitext. Springer-Verlag London, Ltd., London, 2005.
- [8] P. Cartier, F. Patras, *Classical Hopf algebras and their Applications*, Springer, 2021.
- [9] A. Celestino, K. Ebrahimi-Fard, F. Patras, D. Perales, *Cumulant-Cumulant Relations in Free Probability Theory from Magnus' Expansion*, Foundations of Computational Mathematics **22**, (2022) 733-755.
- [10] K. Ebrahimi-Fard, F. Patras, *Cumulants, free cumulants and half-shuffles*, Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences **471**, (2015) 2176.
- [11] K. Ebrahimi-Fard, F. Patras, *The splitting process in free probability theory* International Mathematics Research Notices **2016** (9), 2647-2676.
- [12] K. Ebrahimi-Fard, F. Patras, *Monotone, free, and boolean cumulants: a shuffle algebra approach*, Advances in Mathematics **328**, (2018) 112-132.
- [13] K. Ebrahimi-Fard, F. Patras, *Shuffle group laws: applications in free probability*, Proceedings of the London Mathematical Society **119**, (2019) 814-840.
- [14] K. Ebrahimi-Fard, F. Patras, *A group-theoretical approach to conditionally free cumulants*, IRMA Lectures in Mathematics and Theoretical Physics **31**, (2020) 67-90.
- [15] K. Ebrahimi-Fard, F. Patras, N. Tapia, L. Zambotti, *Hopf-algebraic Deformations of Products and Wick Polynomials*, International Mathematics Research Notices, **2020**, No. 24, (2020) 10064-10099.
- [16] T. Hasebe, H. Saigo, *The monotone cumulants*, Annales de l'Institut Henri Poincaré - Probabilités et Statistiques **47**, No. 4, (2011) 1160-1170.
- [17] S. Manzel, M. Schürmann, *Non-Commutative Stochastic Independence and Cumulants*, Infinite Dimensional Analysis, Quantum Probability and Related Topics **20**, Issue 02, (2017) 1750010 (38 p.).
- [18] J. A. Mingo, R. Speicher, *Free Probability and Random Matrices*, Fields Institute Monographs **1**, Springer-Verlag New York, 2017.
- [19] N. Muraki, *The five independences as natural products*, Infinite Dimensional Analysis, Quantum Probability and Related Topics **6**, No. 3, (2003) 337-371.
- [20] A. Nica, R. Speicher, *Lectures on the combinatorics of free probability*, London Mathematical Society Lecture Note Series, **335** Cambridge University Press (2006).
- [21] G. Peccati, M. S. Taqqu, *Wiener chaos: moments, cumulants and diagrams. A survey with computer implementation*. Bocconi & Springer Series, 1. Springer, Milan; Bocconi University Press, Milan, 2011.
- [22] M. Schürmann, *White Noise on Bialgebras*, Lecture Notes in Mathematics **1544**, Springer-Verlag, Berlin, 1993.
- [23] M. P. Schützenberger, *Contribution aux applications statistiques de la théorie de l'information*, Publications de l'Institut de statistique de l'Université de Paris **3**, no. 1-2, (1954) 3.
- [24] T. P. Speed, *Cumulants and partition lattices. II. Generalised k-statistics*, Journal of the Australian Mathematical Society **40**, no. 1, (1986) 34-53.
- [25] R. Speicher, *A New Example of 'Independence' and 'White Noise'*, Probability Theory and Related Fields **84**, (1990) 141-159.
- [26] R. Speicher, *Multiplicative functions on the lattice of non-crossing partitions and free convolution*, Mathematische Annalen **298**(1), (1994) 611-628.
- [27] R. Speicher, R. Woroudi, *Boolean convolution*, In: Voiculescu, D. V. (ed.) Free Probability Theory. Proceedings, Toronto, Canada 1995, Fields Inst. Commun. **12**, Providence, RI: Amer. Math. Soc., (1997) 267-279.

- [28] R. Speicher, *On universal products*, In: Voiculescu, D. V. (ed.) Free Probability Theory. Proceedings, Toronto, Canada 1995, Fields Institute Communications **12**, Providence, RI: Amer. Math. Soc., (1997) 257-266.
- [29] D. Voiculescu, K. J. Dykema, A. Nica, Free random variables, CRM Monograph Series, vol. 1, American Mathematical Society, Providence, RI, 1992. A non-commutative probability approach to free products with applications to random matrices, operator algebras and harmonic analysis on free groups.
- [30] D. Voiculescu, *Free Probability Theory: Random Matrices and von Neumann Algebras*, Proceedings of the International Congress of Mathematicians, Zürich, Switzerland 1994. Birkhäuser Verlag, Basel, Switzerland (1995).
- [31] W. von Waldenfels, *Proof of an algebraic central limit theorem by moment generating functions*, in: Stochastic Processes - Mathematics and Physics II, Lecture Notes in Math. **1250**, 342-347, Springer-Verlag, Berlin, 1985.

SCHOOL OF MATHEMATICS, THE UNIVERSITY OF EDINBURGH, AND THE MAXWELL INSTITUTE FOR THE MATHEMATICAL SCIENCES, JAMES CLERK MAXWELL BUILDING, THE KING'S BUILDINGS, EDINBURGH, EH9 3FD, UNITED KINGDOM

Email address: ichevyrev@gmail.com

URL: <https://ilyachevyrev.wordpress.com/>

DEPARTMENT OF MATHEMATICAL SCIENCES, NORWEGIAN UNIVERSITY OF SCIENCE AND TECHNOLOGY, NO 7491 TRONDHEIM, NORWAY.

Email address: kurusch.ebrahimi-fard@ntnu.no

URL: <https://folk.ntnu.no/kurusche/>

UNIV. CÔTE D'AZUR, CNRS, UMR 7351, PARC VALROSE, 06108 NICE CEDEX 02, FRANCE.

Email address: patras@unice.fr

URL: www-math.unice.fr/~patras