

PAPER • OPEN ACCESS

# Form factors of the tricritical three-state Potts model in its scaling limit

To cite this article: Giuseppe Mussardo *et al* *J. Stat. Mech.* (2024) 033103

View the [article online](#) for updates and enhancements.

You may also like

- [High-precision percolation thresholds and Potts-model critical manifolds from graph polynomials](#)  
Jesper Lykke Jacobsen
- [Generalized susceptibilities along the phase boundary of the three-dimensional three-state Potts model](#)  
Xue Pan, Mingmei Xu and Yuanfang Wu
- [Parafermionic conformal field theory on the lattice](#)  
Roger S K Mong, David J Clarke, Jason Alicea et al.

PAPER: Quantum statistical physics, condensed matter, integrable systems

## Form factors of the tricritical three-state Potts model in its scaling limit

Giuseppe Mussardo<sup>1</sup>, Marco Panero<sup>2</sup>  
and Andrea Stampiggi<sup>1,\*</sup>

<sup>1</sup> SISSA and INFN, Sezione di Trieste, via Bonomea 265, I-34136 Trieste, Italy

<sup>2</sup> Physics Department, University of Turin and INFN, Sezione di Torino, Via Pietro Giuria 1, I-10125 Turin, Italy

E-mail: [astampig@sissa.it](mailto:astampig@sissa.it), [mussardo@sissa.it](mailto:mussardo@sissa.it) and [marco.panero@unito.it](mailto:marco.panero@unito.it)

Received 13 November 2023

Accepted for publication 30 January 2024

Published 5 March 2024



Online at [stacks.iop.org/JSTAT/2024/033103](https://stacks.iop.org/JSTAT/2024/033103)  
<https://doi.org/10.1088/1742-5468/ad2926>

**Abstract.** We compute the form factors of the order and disorder operators, together with those of the stress–energy tensor, of a two-dimensional three-state Potts model with vacancies along its thermal deformation at the critical point. At criticality, the model is described by the non-diagonal partition function of the unitary minimal model  $\mathcal{M}_{6,7}$  of conformal field theories and is accompanied by an internal  $S_3$  symmetry. The off-critical thermal deformation is an integrable massive theory that is still invariant under  $S_3$ . The presence of infinitely many conserved quantities, whose spin spectrum is related to the exceptional Lie algebra  $E_6$ , allows us to determine the analytic  $S$ -matrix, the exact mass spectrum and the matrix elements of local operators of this model in an exact non-perturbative way. We use the spectral representation series of the correlators and the fast convergence of these series to compute several universal ratios of the renormalization group.

**Keywords:** correlation functions, form factors, integrable quantum field theory

\* Author to whom any correspondence should be addressed.



Original Content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](https://creativecommons.org/licenses/by/4.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

**Contents**

<b>1. Introduction</b> .....	<b>2</b>
<b>2. Critical TPM: the CFT approach</b> .....	<b>4</b>
<b>3. Thermal deformation of the TPM</b> .....	<b>8</b>
3.1. Scattering theory .....	10
<b>4. Generalities on FFs</b> .....	<b>11</b>
<b>5. FFs of the stress–energy tensor</b> .....	<b>14</b>
<b>6. FFs of order and disorder leading and subleading operators</b> .....	<b>17</b>
6.1. Leading disorder operator $\mu$ .....	17
6.2. Subleading disorder operator $\zeta$ .....	20
6.3. Leading order operator $\sigma$ and subleading order operator $Z$ .....	21
<b>7. Universal ratios of the renormalization group</b> .....	<b>23</b>
<b>8. Comparison with a Monte Carlo study</b> .....	<b>26</b>
<b>9. Discussion and conclusions</b> .....	<b>33</b>
<b>Acknowledgment</b> .....	<b>33</b>
<b>Appendix. Analytic properties of FFs</b> .....	<b>33</b>
<b>References</b> .....	<b>36</b>

**1. Introduction**

An interesting problem in theoretical physics consists of the study of two-dimensional statistical models which exhibit a second-order phase transition for certain values of their coupling constants. As is well known, the proper language to frame the scaling limit of these microscopical critical models is the one of the conformal field theories (CFTs) [1], which, in two dimensions, have the peculiar feature of the underlying infinite-dimensional Virasoro algebra and the possibility of defining consistent models employing only a finite number of families of irreducible representations of this algebra (the so-called degenerate fields). These ‘minimal models’ capture the universal scaling behavior of the two-dimensional critical models and are exactly solvable, in the sense that it is possible to obtain—at least in principle—closed expressions for the multi-point correlation functions of scaling operators by exploiting properties of the Virasoro algebra and null-vector conditions of the corresponding degenerate fields [1, 2].

However, no general framework has yet been formulated for generic off-critical models. Nonetheless, powerful tools of analysis can be employed if the deformed theory turns

out to be integrable, i.e. if it has infinitely many commuting conserved local quantities. When this happens, these conserved quantities impose severe constraints on the structure of the scattering amplitudes, which are non-zero only for the elastic processes and can be computed exactly [3–9]. In practice, it is sufficient to find all possible two-particle amplitudes, since all other scattering amplitudes involving  $n > 2$  particles can be expressed in terms of their  $n(n-1)/2$  two-particle scatterings.

Notable examples of these two-dimensional integrable field theories are the ones described by the Ising model in a magnetic field, the thermally deformed tricritical Ising model and the thermally deformed three-state tricritical Potts model (TPM). These models possess conserved charges whose spins match the Coxeter exponents of three exceptional algebras ( $E_8$ ,  $E_7$  and  $E_6$ , respectively [4–11]).

Once the  $S$ -matrix of an integrable theory is known, it is possible to study off-critical two-point correlation functions through the technique of form factors (FFs) [9, 11, 12]. These quantities are defined as matrix elements of (semi-)local operators between the vacuum of the theory and multi-particle states. The novel contribution of this paper is the exact computation of FFs of the leading and sub-leading order and disorder operators in the thermal deformation of the TPM. We also compute some universal renormalization group ratios (see [13] and references therein), which can be useful for numerical and experimental studies.

The  $q$ -state Potts model has been studied in great detail in the literature—see e.g. [14] for a review. A generalization of this model, called ‘dilute’ due to the presence of vacancies, has been studied in [15]. Its lattice realization has a Hamiltonian, invariant under the permutation group  $S_3$ , of the form

$$\mathcal{H} = - \sum_{\langle i,j \rangle} t_i t_j (\mathcal{K} + \mathcal{F} \delta_{s_i, s_j}) + \mathcal{D} \sum_i t_i , \quad (1)$$

where each site  $i$  is associated with a Potts ‘spin’  $s_i = \{0, 1, 2\}$  and a vacancy variable  $t_i = \{0, 1\}$ . In the equation,  $\mathcal{K}$  and  $\mathcal{F}$  are interaction parameters associated with vacancies and Potts spins, respectively, while  $\mathcal{D}$  plays the role of chemical potential associated to vacancies. The usual Potts model is retrieved when the chemical potential  $\mathcal{D}$  is negative and large in absolute value. Since the TPM possesses only two relevant  $S_3$  invariant operators [2], the energy  $\epsilon$  and the vacancy  $t$ , it is retrieved from equation (1) in the limit of vanishing  $\mathcal{K}$ . The TPM is also exactly solvable on a honeycomb lattice and proven to be self-dual, i.e. there is a mapping between its disordered (high-temperature) and ordered (low-temperature) phases [16].

The content of this paper is organized as follows. In section 2, we recall a few basic properties of minimal models of the CFT, useful to set most of the notation, followed by a thorough discussion of the TPM, in particular, the operator content and the fusion rules of this model. Section 3 is dedicated to the thermal deformation of the TPM, regarded as an integrable field theory; the scattering theory and the mass spectrum are also discussed. The FFs are introduced in section 4 (their analytic properties are gathered in the appendix). In section 5, we review the FFs of the stress–energy tensor originally obtained in [10]. The FFs of the order and disorder operators of the thermal deformation of the TPM are presented in section 6. The two-point correlation functions and the universal ratios of the renormalization group are discussed in section 7. Some

preliminary results from the Monte Carlo simulations of the model are presented in section 8. Our conclusions are finally summarized in section 9.

## 2. Critical TPM: the CFT approach

It is well known that minimal models  $\mathcal{M}_{p,q}$  of the CFT are characterized by the following properties [1, 2]:

- (i) Their central charge  $c$  is always less than one and can be parameterized by two coprime integers  $p, q$  (with  $q > p$ )

$$c(p, q) = 1 - 6 \frac{(p - q)^2}{pq}. \tag{2}$$

- (ii) There are a finite number of families of degenerate fields, both in the analytic and anti-analytic sectors of the CFT. In the analytic sector, a degenerate primary field  $\phi_{r,s}(z)$  has conformal dimension

$$\Delta_{r,s} = \frac{(pr - qs)^2 - (q - p)^2}{4pq}, \tag{3}$$

with  $1 \leq r \leq q - 1$  and  $1 \leq s \leq p - 1$ , coincident to that of its anti-analytic counterpart  $\bar{\phi}_{r,s}(\bar{z})$ :  $\bar{\Delta}_{r,s} = \Delta_{r,s}$ . Unitary minimal models are retrieved by setting in the previous formulas  $m = q = p + 1$  with  $m \geq 3$ .

- (iii) Correlation functions of all conformal fields satisfy a set of linear differential equations that can be exactly solved, yielding in turn the structure constants of the operator algebra.
- (iv) The modular-invariant partition function can be determined exactly [17–19].

Another important feature of minimal models is that the operator product expansion (OPE) algebra of degenerate fields, in the analytic and anti-analytic sectors separately, is ‘closed’: the OPE of two degenerate fields can be written in terms of degenerate fields and their descendants. In particular, the operators appearing in the OPE are obtained by iterative multiplication of the fields  $\phi_{2,1}$  and  $\phi_{1,2}$ . The general formula, which does not give information about the structure constants, is

$$\phi_{r_1, s_1} \times \phi_{r_2, s_2} = \sum_{r_3 = \{|r_2 - r_1| + 1, |r_2 - r_1| + 3, \dots\}}^{\min(r_1 + r_2 - 1, 2q - 1 - r_1 - r_2)} \sum_{s_3 = \{|s_2 - s_1| + 1, |s_2 - s_1| + 3, \dots\}}^{\min(s_1 + s_2 - 1, 2p - 1 - s_1 - s_2)} \phi_{r_3, s_3}. \tag{4}$$

Finding the modular-invariant partition functions of a given universality class associated with a minimal model is equivalent to determining the operator content of the model. The problem of the modular-invariant partition functions for unitary theories was first addressed by Cardy [18, 19] and the solution for minimal models with periodic boundary conditions was found soon after by Cappelli *et al.* [17].

**Table 1.** Kač table of the universality class  $\mathcal{M}_{6,7}$ .

5	5	$\frac{22}{7}$	$\frac{12}{7}$	$\frac{5}{7}$	$\frac{1}{7}$	0
4	$\frac{23}{8}$	$\frac{85}{56}$	$\frac{33}{56}$	$\frac{5}{56}$	$\frac{1}{56}$	$\frac{3}{8}$
3	$\frac{4}{3}$	$\frac{10}{21}$	$\frac{1}{21}$	$\frac{1}{21}$	$\frac{10}{21}$	$\frac{4}{3}$
2	$\frac{3}{8}$	$\frac{1}{56}$	$\frac{5}{56}$	$\frac{33}{56}$	$\frac{85}{56}$	$\frac{23}{8}$
1	0	$\frac{1}{7}$	$\frac{5}{7}$	$\frac{12}{7}$	$\frac{22}{7}$	5
s $\uparrow$ , r $\rightarrow$	1	2	3	4	5	6

The conformal dimensions of the fields in  $\mathcal{M}_{6,7}$  are summarized in table 1. The universality class  $\mathcal{M}_{6,7}$  contains two algebras of fields which close under the OPE: the pentacritical  $\varphi^{10}$  Landau–Ginzburg field theory and the TPM.

Corresponding to the minimal model  $\mathcal{M}_{6,7}$  there are then two modular-invariant partition functions: a diagonal one, which corresponds to the  $\varphi^{10}$  Landau–Ginzburg theory

$$Z^{(1)} = |\chi_0|^2 + |\chi_{\frac{1}{7}}|^2 + |\chi_{\frac{5}{7}}|^2 + |\chi_{\frac{12}{7}}|^2 + |\chi_{\frac{22}{7}}|^2 + |\chi_5|^2 + |\chi_{\frac{3}{8}}|^2 + |\chi_{\frac{1}{56}}|^2 + |\chi_{\frac{5}{56}}|^2 + |\chi_{\frac{33}{56}}|^2 + |\chi_{\frac{85}{56}}|^2 + |\chi_{\frac{23}{8}}|^2 + |\chi_{\frac{4}{3}}|^2 + |\chi_{\frac{10}{21}}|^2 + |\chi_{\frac{1}{21}}|^2 \quad (5)$$

and an off-diagonal one relative to the TPM

$$Z^{(2)} = |\chi_0 + \chi_5|^2 + |\chi_{\frac{1}{7}} + \chi_{\frac{22}{7}}|^2 + |\chi_{\frac{5}{7}} + \chi_{\frac{12}{7}}|^2 + 2|\chi_{\frac{4}{3}}|^2 + 2|\chi_{\frac{10}{21}}|^2 + 2|\chi_{\frac{1}{21}}|^2. \quad (6)$$

Here,  $\chi_\Delta$  is the character of the irreducible representation of the Virasoro algebra of weight  $\Delta$ , whose explicit formulas can be found in [20].

Using the notation of table 2 for the various fields, it is possible to show that the fusion rules of table 3 are compatible with an  $S_3$  internal symmetry, so that operators can be organized according to irreducible representations of this discrete group, i.e. either in doublets (**s**, **Z** and **A**) or in the one-dimensional neutral representation (**1**,  $\epsilon$ ,  $t$ , **X**, **Y** and **B**). In the model, there are no operators that transform according to the alternating representation.

Doublet operators such as **s** and **Z** are charged under the generator  $\Omega$  of the subgroup  $Z_3$ . Let the components of the doublet corresponding to the leading operator **s** be, respectively,  $\sigma$  and  $\tilde{\sigma}$ . Similarly, one defines the components of the subleading order operator **Z** to be  $Z$  and  $\tilde{Z}$ . As it happens in the case of the Ising model [21], the tricritical Ising model [22] and the three-state Potts model [23, 24], in view of the self-duality of the lattice model, we can introduce a leading disorder operator  $(\mu, \tilde{\mu})$  associated with the leading-order operator  $(\sigma, \tilde{\sigma})$ . In complete analogy to its leading counterpart,  $(\zeta, \tilde{\zeta})$  is the subleading disorder operator associated with the subleading order operator  $(Z, \tilde{Z})$ . The conformal dimensions of these operators are:

**Table 2.** Operator content of the TPM at criticality.

$\phi_{r,s}$	$(\Delta, \bar{\Delta})$	Operator	Interpretation	$S_3$ representation
$\phi_{1,1} \times \bar{\phi}_{1,1}$	$(0, 0)$	$\mathbb{1}$	Identity	1
$\phi_{3,3} \times \bar{\phi}_{3,3}$	$(\frac{1}{21}, \frac{1}{21})$	$\mathbf{s}$	Leading order	2
$\phi_{2,1} \times \bar{\phi}_{2,1}$	$(\frac{1}{7}, \frac{1}{7})$	$\epsilon$	Energy	1
$\phi_{2,3} \times \bar{\phi}_{2,3}$	$(\frac{10}{21}, \frac{10}{21})$	$\mathbf{Z}$	Subleading order	2
$\phi_{3,1} \times \bar{\phi}_{3,1}$	$(\frac{5}{7}, \frac{5}{7})$	$t$	Vacancy	1
$\phi_{1,3} \times \bar{\phi}_{1,3}$	$(\frac{4}{3}, \frac{4}{3})$	$\mathbf{A}$	Irrelevant	2
$\phi_{4,1} \times \bar{\phi}_{4,1}$	$(\frac{12}{7}, \frac{12}{7})$	$\mathbf{X}$	Irrelevant	1
$\phi_{5,1} \times \bar{\phi}_{5,1}$	$(\frac{22}{7}, \frac{22}{7})$	$\mathbf{B}$	Irrelevant	1
$\phi_{6,1} \times \bar{\phi}_{6,1}$	$(5, 5)$	$\mathbf{Y}$	Irrelevant	1
$\phi_{4,1} \times \bar{\phi}_{3,1}$	$(\frac{12}{7}, \frac{5}{7})$	$\mathcal{F}$	Irrelevant, spin 1	1
$\phi_{3,1} \times \bar{\phi}_{4,1}$	$(\frac{5}{7}, \frac{12}{7})$	$\bar{\mathcal{F}}$	Irrelevant, spin $-1$	1
$\phi_{4,1} \times \bar{\phi}_{2,1}$	$(\frac{12}{7}, \frac{1}{7})$	$\mathcal{U}$	Irrelevant, spin 3	1
$\phi_{2,1} \times \bar{\phi}_{4,1}$	$(\frac{1}{7}, \frac{12}{7})$	$\bar{\mathcal{U}}$	Irrelevant, spin $-3$	1
$\phi_{6,1} \times \bar{\phi}_{1,1}$	$(5, 0)$	$\mathcal{W}$	Irrelevant, spin 5	1
$\phi_{1,1} \times \bar{\phi}_{6,1}$	$(0, 5)$	$\bar{\mathcal{W}}$	Irrelevant, spin $-5$	1

$$\begin{aligned}
 (\Delta_\sigma, \bar{\Delta}_\sigma) &= (\Delta_{\tilde{\sigma}}, \bar{\Delta}_{\tilde{\sigma}}) = (\Delta_\mu, \bar{\Delta}_\mu) = (\Delta_{\tilde{\mu}}, \bar{\Delta}_{\tilde{\mu}}) = \left(\frac{1}{21}, \frac{1}{21}\right), \\
 (\Delta_Z, \bar{\Delta}_Z) &= (\Delta_{\tilde{Z}}, \bar{\Delta}_{\tilde{Z}}) = (\Delta_\zeta, \bar{\Delta}_\zeta) = (\Delta_{\tilde{\zeta}}, \bar{\Delta}_{\tilde{\zeta}}) = \left(\frac{10}{21}, \frac{10}{21}\right).
 \end{aligned}
 \tag{7}$$

The transformation properties under  $S_3$  of the order and disorder operators are as follows. Under the generator  $\Omega$  of  $Z_3$  the disorder operators are invariant and the order operators acquire a phase:

$$\begin{aligned}
 \Omega\sigma &= \omega\sigma, & \Omega\tilde{\sigma} &= \omega^*\tilde{\sigma}, & \Omega Z &= \omega^2 Z, & \Omega\tilde{Z} &= (\omega^*)^2 \tilde{Z}, \\
 \Omega\mu &= \mu, & \Omega\tilde{\mu} &= \tilde{\mu}, & \Omega\zeta &= \zeta, & \Omega\tilde{\zeta} &= \tilde{\zeta},
 \end{aligned}
 \tag{8}$$

where  $\omega = \exp(2i\pi/3)$ . On the other hand, the generator  $\mathcal{C}$  of  $Z_2$  acts like a charge conjugation operator:

$$\begin{aligned}
 \mathcal{C}\sigma &= \tilde{\sigma}, & \mathcal{C}Z &= \tilde{Z}, \\
 \mathcal{C}\mu &= \tilde{\mu}, & \mathcal{C}\zeta &= \tilde{\zeta}.
 \end{aligned}
 \tag{9}$$

The assumption that only order operators are charged with respect to  $Z_3$  is consistent with an alternative definition of the TPM in terms of parafermionic algebra of

**Table 3.** Fusion rules of the TPM at criticality.

$\mathbf{s} \times \mathbf{s} = \mathbb{1} + \epsilon + t + \mathbf{X} + \mathbf{B} + \mathbf{Y} + \mathbf{s} + \mathbf{Z} + \mathbf{A}$	$\epsilon \times \epsilon = \mathbb{1} + t$
$\mathbf{Z} \times \mathbf{Z} = \mathbb{1} + \epsilon + t + \mathbf{X} + \mathbf{Y} + \mathbf{s} + \mathbf{A}$	$t \times t = \mathbb{1} + t + \mathbf{B}$
$\mathbf{A} \times \mathbf{A} = \mathbb{1} + \mathbf{Y} + \mathbf{A}$	$\mathbf{X} \times \mathbf{X} = \mathbb{1} + t + \mathbf{B}$
$\mathbf{B} \times \mathbf{B} = \mathbb{1} + t$	$\mathbf{Y} \times \mathbf{Y} = \mathbb{1}$
$\mathbf{s} \times \epsilon = \mathbf{s} + \mathbf{Z}$	$\mathbf{s} \times t = \mathbf{s} + \mathbf{Z} + \mathbf{A}$
$\mathbf{s} \times \mathbf{Z} = \epsilon + t + \mathbf{X} + \mathbf{B} + \mathbf{s} + \mathbf{Z}$	$\mathbf{s} \times \mathbf{A} = t + \mathbf{X} + \mathbf{s}$
$\mathbf{s} \times \mathbf{X} = \mathbf{s} + \mathbf{Z} + \mathbf{A}$	$\mathbf{s} \times \mathbf{B} = \mathbf{s} + \mathbf{Z}$
$\mathbf{s} \times \mathbf{Y} = \mathbf{s}$	
$\epsilon \times \mathbf{Z} = \mathbf{s} + \mathbf{A}$	$\epsilon \times t = \epsilon + \mathbf{X}$
$\epsilon \times \mathbf{A} = \mathbf{Z}$	$\epsilon \times \mathbf{X} = t + \mathbf{Y}$
$\epsilon \times \mathbf{B} = \mathbf{X} + \mathbf{Y}$	$\epsilon \times \mathbf{Y} = \mathbf{B}$
$\mathbf{Z} \times t = \mathbf{s} + \mathbf{Z}$	$\mathbf{Z} \times \mathbf{A} = \epsilon + \mathbf{B} + \mathbf{Z}$
$\mathbf{Z} \times \mathbf{X} = \mathbf{s} + \mathbf{Z}$	$\mathbf{Z} \times \mathbf{B} = \mathbf{s} + \mathbf{A}$
$\mathbf{Z} \times \mathbf{Y} = \mathbf{Z}$	
$t \times \mathbf{A} = \mathbf{s}$	$t \times \mathbf{X} = \epsilon + \mathbf{X} + \mathbf{Y}$
$t \times \mathbf{B} = t + \mathbf{B}$	$t \times \mathbf{Y} = \mathbf{X}$
$\mathbf{A} \times \mathbf{X} = \mathbf{s}$	$\mathbf{A} \times \mathbf{B} = \mathbf{Z}$
$\mathbf{A} \times \mathbf{Y} = \mathbf{A}$	
$\mathbf{X} \times \mathbf{B} = \epsilon + \mathbf{X}$	$\mathbf{X} \times \mathbf{Y} = t$
$\mathbf{B} \times \mathbf{Y} = \epsilon$	

central charge  $6/7$ —see equation (A.8) of [25]. The parafermionic theory is defined to be invariant under the product group  $Z_3 \times \tilde{Z}_3$ . The first  $Z_3$  group acts non-trivially only on order operators, while the second only on disorder ones. However, the symmetry group of the TPM is  $S_3$ , which contains  $Z_3$  and  $Z_2$  as subgroups. While  $Z_3$  is the group of symmetry of the parafermionic theory, the second group  $Z_2$  acts as a charge conjugation operator.

There are two important features of parafermionic theories. The first is that they are self-dual by definition: order and disorder operators in the ordered phase can be mapped into disorder and order operators of the disordered phase, respectively. As a consequence, we will only focus on the FFs in the disordered phase. The second is that the order and disorder operators are not mutually local, in general. In particular, for the TPM the non-locality factors are

$$\begin{aligned} \gamma_{\sigma,\mu} = \gamma_{Z,\zeta} = -1/3, \quad \gamma_{\sigma,\zeta} = \gamma_{Z,\mu} = +1/3, \\ \gamma_{\tilde{\sigma},\mu} = \gamma_{\tilde{Z},\zeta} = 1/3, \quad \gamma_{\tilde{\sigma},\zeta} = \gamma_{\tilde{Z},\mu} = -1/3. \end{aligned} \tag{10}$$

Identical results apply if we charge-conjugate both the first and second entries in the  $\gamma$ s.



### 3. Thermal deformation of the TPM

Consider the fixed point action  $\mathcal{A}_0$  of the TPM and define the action  $\mathcal{A}$  of its thermal deformation as

$$\mathcal{A} = \mathcal{A}_0 + \lambda \int d^2x \epsilon(x), \quad (11)$$

where  $\epsilon$  is the energy operator and  $\lambda$  is its coupling constant. In the scaling limit of the lattice version of the model, i.e. the limit in which the temperature  $T$  approaches its critical value  $T_c$  and the chemical potential is fixed at its critical value, we have  $\lambda \propto (T - T_c)$ . The action  $\mathcal{A}$  corresponds to a massive theory and there is a non-vanishing trace of the stress–energy tensor  $\Theta(z, \bar{z})$  given by

$$\Theta(z, \bar{z}) = 2\pi \lambda (2 - 2\Delta_\epsilon) \epsilon(z, \bar{z}). \quad (12)$$


In the following, we will often denote the quantum field theory associated with the thermal deformation of the TPM as  $\epsilon$ -TPM. In addition to the stress–energy tensor, in the  $\epsilon$ -TPM there are infinitely many local conserved quantities which commute with each other and therefore the theory is integrable [3, 7]. The spins of conserved quantities for this model are given by an infinite sequence of residues

$$\{1, 4, 5, 7, 8, 11\} \pmod{h = 12}. \quad (13)$$

In addition, they coincide, modulo the Coxeter number  $h = 12$ , with the Coxeter exponents of the exceptional algebra  $E_6$ .

The hidden  $E_6$  structure of the model can be understood employing a different CFT description of the TPM, i.e. the so-called coset construction [26], which in this case consists of  $[(E_6)_1 \times (E_6)_1]/(E_6)_2$ . Let us discuss this point in more detail. The affine algebra  $(E_6)_k$  (figure 1) has different integrable highest weights depending on the level  $k$ . Calling  $w_i$  ( $i = 0, 1, \dots, 7$ ) the fundamental weights of  $(E_6)_k$ , which are the basis of the weight space in the natural basis  $w_i = \hat{e}_{i+1}$  (e.g.  $w_1 = (0, 1, 0, 0, 0, 0)$ ), the integrable highest weights at level  $k=1$  are  $w_0$ ,  $w_1$  and  $w_5$ , while at level  $k=2$  we have  $2w_0$ ,  $2w_1$ ,  $2w_5$ ,  $(w_0 + w_1)$ ,  $(w_0 + w_5)$ ,  $(w_1 + w_5)$ ,  $w_2$  and  $w_6$ . Through the coset fields, we can recover the operator content of the TPM. As discussed for instance in [26], the procedure consists in finding the allowed branchings, the fractional dimensions of the integrable representations, conjugate representations (related to the symmetry group of the non-affine Dynkin diagram) and coset fields, i.e. classes of equivalent branchings related by automorphisms ( $Z_3$  in the case of affine  $E_6$ ). The results are summarized in table 4.

The coset construction of the TPM only holds for critical theory. Away from criticality there is, however, a correspondence between the thermal deformation of the TPM and the exceptional algebra  $E_6$  and this involves Toda field theories. These theories are built starting from a Lagrangian associated to a Lie algebra of finite rank (see, e.g. [9, 27] and references therein). In [28] it was proven that the deformation of the Toda field theory of  $E_6$  perturbed with the adjoint representation representation is integrable and has conserved charges whose spins coincide with those of the thermal deformation of



**Figure 1.** Dynkin diagram of the affine  $E_6$  algebra. The first term in parentheses labels the root, while the second its comark.

**Table 4.** Identification of coset fields of  $[(E_6)_1 \times (E_6)_1]/(E_6)_2$  with the operator content of the TPM. An element of the coset  $[\lambda, \mu, \nu]$  is such that it appears in the branching  $\lambda + \mu = \sum_{\nu} b'_{\lambda, \mu} \nu$ .

Cosets	Operators
$\{[w_0, w_0, 2w_0] \sim [w_1, w_1, 2w_5] \sim [w_5, w_5, 2w_5]\}$	$\mathbf{1}, \mathbf{Y}$
$\{[w_0, w_1, w_0 + w_1] \sim [w_5, w_0, w_0 + w_5] \sim [w_1, w_5, w_1 + w_5]\}$	$\sigma$
$\{[w_1, w_0, w_0 + w_1] \sim [w_0, w_5, w_0 + w_5] \sim [w_5, w_1, w_1 + w_5]\}$	$\tilde{\sigma}$
$\{[w_0, w_0, w_6] \sim [w_1, w_1, w_2]\}$	$\epsilon, \mathbf{B}$
$\{[w_1, w_5, w_6] \sim [w_5, w_0, w_2]\}$	$Z$
$\{[w_5, w_0, w_2] \sim [w_5, w_1, w_6]\}$	$\tilde{Z}$
$\{[w_0, w_0, w_1 + w_5] \sim [w_1, w_1, w_0 + w_5] \sim [w_5, w_5, w_0 + w_1]\}$	$t, \mathbf{X}$
$\{[w_0, w_1, 2w_5] \sim [w_5, w_0, 2w_1] \sim [w_1, w_5, 2w_0]\}$	$A$
$\{[w_1, w_0, 2w_5] \sim [w_0, w_5, 2w_1] \sim [w_5, w_1, 2w_0]\}$	$\tilde{A}$

the TPM. Thus, the scattering matrix of this perturbed Toda field theory has the same minimal scattering matrix as our model: its mass spectrum contains six particles, two particle–antiparticle doublets,  $(l, \bar{l})$  and  $(h, \bar{h})$ , and two self-conjugated particles,  $L$  and  $H$ , and the masses of these particles can be expressed as

$$\begin{aligned}
 m_l &= m_{\bar{l}} = M(\lambda), & m_L &= 2 \cos(\pi/4) m_l, \\
 m_h &= m_{\bar{h}} = 2 \cos(\pi/12) m_l, & m_H &= 4 [\cos(\pi/4)]^2 m_l,
 \end{aligned}
 \tag{14}$$

where  $M(\lambda)$  is the mass of the lightest kink of the theory and depends on the coupling constant  $\lambda$  in equation (11) as [29]



$$M(\lambda) = C_{\epsilon} |\lambda|^{\frac{7}{12}}, \tag{15}$$

where

$$C_{\epsilon} = \left[ 4\pi^2 \gamma\left(\frac{4}{7}\right) \gamma\left(\frac{9}{14}\right) \gamma\left(\frac{5}{7}\right) \gamma\left(\frac{11}{14}\right) \right]^{\frac{7}{24}} \frac{2\Gamma\left(\frac{1}{4}\right) \Gamma\left(\frac{13}{12}\right)}{\Gamma\left(\frac{1}{2}\right) \Gamma\left(\frac{2}{3}\right) \Gamma\left(\frac{7}{6}\right)} \approx 3.74656. \tag{16}$$

In the latter equation,  $\gamma(z)$  is a shorthand for  $\Gamma(z)/\Gamma(1-z)$ .

**Table 5.** Conventions for incoming and outgoing states, where  $\theta_1 > \theta_2 > \theta_3$ .

Picture	State	Convention
	Incoming	$ A_a(\theta_1)A_b(\theta_2)A_c(\theta_3)\dots\rangle_{\text{IN}}$
	Outgoing	$ \dots A_c(\theta_3)A_b(\theta_2)A_a(\theta_1)\rangle_{\text{OUT}}$

For the thermal deformation of the TPM, the excitations are ordinary particles only in the disordered phase  $\lambda > 0$ , while some of them become kinks in the ordered phase  $\lambda < 0$ . In a quantum field theory with localized interactions, particles are associated with asymptotic states which transform according to irreducible representations of the symmetry group of the theory. For the thermal deformation of the TPM, we have two  $S_3$ -doublets and two singlets. Since the particles are in the doublets  $(l, \bar{l})$  and  $(h, \bar{h})$  are generated by order (magnetization-like) operators, we define the non-locality properties of the particles with respect to disorder operators to be

$$\gamma_{l,\mu} = \gamma_{l,\zeta} = 1/3, \quad \gamma_{\bar{l},\mu} = \gamma_{\bar{l},\zeta} = -1/3. \tag{17}$$

Identical relations hold if  $l$  is replaced by  $h$ . This choice of non-local factors is consistent with the scattering theory of the  $\epsilon$ -TPM, which will now be outlined.

### 3.1. Scattering theory

In  $(1 + 1)$ -dimensional theories, all particles scatter along a straight line. It is useful to use the rapidity variable  $\theta$  which parameterizes the on-shell momentum of the particle of type  $a$  and mass  $m_a$  as

$$p_a(\theta_a) = m_a(\cosh \theta_a, \sinh \theta_a) \tag{18}$$

and denote the one-particle state as  $|A_a(\theta)\rangle$ . For the multiparticle states we assume, as a convention, that all incoming particles can be written from left to right, with decreasing rapidity. On the other hand, outgoing states are written in the converse: particles with the highest rapidity on the right and particles with lowest rapidity on the left. Examples are provided in table 5.

Integrable models have commuting conserved quantities  $Q_s$  that can be diagonalized simultaneously with the Hamiltonian. This condition imposes strong constraints on the scattering matrix. Apart from being unitary and satisfying crossing symmetry, the  $S$ -matrix is elastic and all  $n$ -particle processes can be factorized in terms of two-particle scatterings [7, 9]. These two properties, along with bootstrap assumptions, i.e. treating bound states on the same footing as asymptotic particles, allow for the determination of the analytic two-body scattering amplitudes. In the case of the  $\epsilon$ -TPM, the  $S$ -matrix [3, 6] is diagonal, i.e. reflection amplitudes vanish, and the various amplitudes are reported in table 6. They can be expressed in terms of the building block functions

**Table 6.**  $S$ -matrix of the  $\epsilon$ -TPM [3, 6]. Here  $[x]$  is a shorthand for  $s_{x/12}(\theta)$  (see equation (19)). The superscript refers to the associated bound state. Notice that 12 is the Coxeter number of the exceptional algebra  $E_6$ .

$a$	$b$	$S_{ab}$	$a$	$b$	$S_{ab}$
$l$	$l$	$\bar{l} \quad \bar{h}$ [8][6][2]	$L$	$L$	$-[10] \begin{smallmatrix} L \\ \bar{l} \end{smallmatrix} [8] \begin{smallmatrix} H \\ \bar{h} \end{smallmatrix} [6]^2 [4] [2]$
$\bar{l}$	$\bar{l}$	$\begin{smallmatrix} l \\ \bar{l} \end{smallmatrix} \quad \begin{smallmatrix} h \\ \bar{h} \end{smallmatrix}$ [8][6][2]	$L$	$h$	$\begin{smallmatrix} l \\ \bar{l} \end{smallmatrix} [10] [8]^2 [6]^2 [4]^2 [2]$
$l$	$\bar{l}$	$\begin{smallmatrix} L \\ \bar{l} \end{smallmatrix}$ $-[10] [6] [4]$	$L$	$\bar{h}$	$\begin{smallmatrix} \bar{l} \\ \bar{h} \end{smallmatrix} [10] [8]^2 [6]^2 [4]^2 [2]$
$l$	$L$	$\begin{smallmatrix} l \\ \bar{l} \end{smallmatrix} \quad \begin{smallmatrix} h \\ \bar{h} \end{smallmatrix}$ [9][7][5][3]	$L$	$H$	$\begin{smallmatrix} L \\ \bar{l} \end{smallmatrix} [11] [9]^2 [7]^3 [5]^3 [3]^2 [1]$
$\bar{l}$	$L$	$\begin{smallmatrix} \bar{l} \\ \bar{h} \end{smallmatrix} \quad \begin{smallmatrix} \bar{h} \\ \bar{l} \end{smallmatrix}$ [9][7][5][3]	$h$	$h$	$\begin{smallmatrix} \bar{l} \\ \bar{h} \end{smallmatrix} [10] [8]^3 [6]^3 [4]^2 [2]^2$
$l$	$h$	$\begin{smallmatrix} \bar{h} \\ \bar{l} \end{smallmatrix} [9] [7] [5]^2 [3] [11]$	$\bar{h}$	$\bar{h}$	$\begin{smallmatrix} l \\ \bar{h} \end{smallmatrix} [10] [8]^3 [6]^3 [4]^2 [2]^2$
$\bar{l}$	$\bar{h}$	$\begin{smallmatrix} h \\ \bar{l} \end{smallmatrix} [9] [7] [5]^2 [3] [11]$	$h$	$\bar{h}$	$-[10]^2 [8]^2 [6]^3 [4]^3 [2]$
$l$	$\bar{h}$	$\begin{smallmatrix} L \\ \bar{l} \end{smallmatrix} [9] [7]^2 [5] [3] [1]$	$h$	$H$	$\begin{smallmatrix} l \\ \bar{h} \end{smallmatrix} [11] [9]^3 [7]^4 [5]^4 [3]^3 [1]$
$\bar{l}$	$h$	$\begin{smallmatrix} L \\ \bar{l} \end{smallmatrix} [9] [7]^2 [5] [3] [1]$	$\bar{h}$	$H$	$\begin{smallmatrix} \bar{l} \\ \bar{h} \end{smallmatrix} [11] [9]^3 [7]^4 [5]^4 [3]^3 [1]$
$l$	$H$	$\begin{smallmatrix} h \\ \bar{h} \end{smallmatrix} [10] [8]^2 [6]^2 [4]^2 [2]$	$H$	$H$	$-[10]^3 [8]^5 [6]^6 [4]^5 [2]^3$
$\bar{l}$	$H$	$\begin{smallmatrix} \bar{h} \\ \bar{l} \end{smallmatrix} [10] [8]^2 [6]^2 [4]^2 [2]$			


$$[x] \equiv s_{x/12}(\theta) = \frac{\sinh[(\theta + i\pi x/12)/2]}{\sinh[(\theta - i\pi x/12)/2]} \tag{19}$$

By looking at bound states with positive residue of two-particle scattering amplitudes, it is possible to extract the mass spectrum of the model, which also coincides with the one of the  $E_6$  Toda field theory (equation (14)).

In the  $\epsilon$ -TPM, the degeneracy on the particle–antiparticle states  $(l, \bar{l})$  and  $(h, \bar{h})$  is lifted if we require that the particle singlets and doublets transform according to the singlet and doublet irreducible representations of  $S_3$ , respectively. In particular, we suppose that particles  $l$  and  $h$  have  $Z_3$ -charge  $\omega$ ,  $\bar{l}$  and  $\bar{h}$  have  $Z_3$ -charge  $\omega^*$ , while  $L$  and  $H$  do not transform under the generator  $\Omega$  of  $S_3$ . On the other hand,  $Z_2$  charge conjugation maps particles to their respective antiparticles. In the specific case of the two singlets,  $L$  and  $H$ , the particle and antiparticle coincide.

#### 4. Generalities on FFs

The aim of this section is to outline the main properties of FFs; in particular, the ones useful to address one- and two-particle states. A further discussion of the analytic properties of such objects can be found in the [appendix](#).



**Figure 2.** Form factor of the operator  $\mathcal{O}$ .

For a relativistic theory, a generic matrix element of a Hermitian operator  $\mathcal{O}$  between an incoming and outgoing state is given by

$$\begin{aligned} & \text{OUT}\langle A_{b_1}(q_1) \dots A_{b_m}(q_m) | \mathcal{O}(x) | A_{a_1}(p_1) \dots A_{a_n}(p_n) \rangle_{\text{IN}} \\ &= \exp \left\{ i \left( \sum_{j=1}^m (q_j)_\mu - \sum_{i=1}^n (p_i)_\mu \right) x^\mu \right\} \times \text{OUT}\langle A_{b_1}(q_1) \dots A_{b_m}(q_m) | \mathcal{O}(0) | A_{a_1}(p_1) \dots A_{a_n}(p_n) \rangle_{\text{IN}}. \end{aligned} \tag{20}$$

FFs are nothing more than matrix elements between the vacuum of the theory and incoming multi-particle states, as pictorially described in figure 2. They are related to the previous equation by crossing, are analytic functions of the rapidity difference  $\theta_{ij} = \theta_i - \theta_j$  and are defined as

$$F_{a_1 \dots a_n}^{\mathcal{O}}(\theta_1, \dots, \theta_n) = \langle 0 | \mathcal{O}(0) | A_{a_1}(\theta_1) \dots A_{a_n}(\theta_n) \rangle. \tag{21}$$

For a scalar operator, knowledge of the FFs allows us to write the spectral series for the correlation functions which, for the two-point correlators, read [30]

$$\langle \mathcal{O}(x) \mathcal{O}(0) \rangle = \sum_{n=0}^{\infty} \int \frac{d\theta_1 \dots d\theta_n}{(2\pi)^n n!} |F_{a_1 \dots a_n}^{\mathcal{O}}(\theta_1, \dots, \theta_n)|^2 \exp \left\{ -|x| \sum_{k=1}^n m_k \cosh \theta_k \right\}. \tag{22}$$

An important feature of these spectral series is their rapid convergence [9, 31]. Practically, lowest mass particle FFs contribute more than higher mass particle ones, as it will be evidenced later by applications of the  $\Delta$ -theorem [32], which is a sum rule for the conformal weight of an operator  $\Phi$

$$\Delta_\Phi = -\frac{1}{4\pi \langle \Phi \rangle} \int d^2x \langle \Theta(x) \Phi(0) \rangle. \tag{23}$$

This sum rule is similar to Zamolodchikov’s  $c$ -theorem sum rule for the UV central charge [33]

$$c = \frac{3}{2} \int dr r^3 \langle \Theta(r) \Theta(0) \rangle. \tag{24}$$

For integrable theories, FFs of scalar operators can be parameterized as

$$F_{a_1 \dots a_n}^{\mathcal{O}}(\theta_1, \dots, \theta_n) = \frac{Q_{a_1 \dots a_n}^{\mathcal{O}}(\theta_1, \dots, \theta_n)}{D_{a_1 \dots a_n}(\theta_1, \dots, \theta_n)} \prod_{i < j} F_{a_i a_j}^{\min}(\theta_{ij}), \quad (25)$$

where  $D_{a_1 \dots a_n}(\theta_1, \dots, \theta_n)$  is a  $2\pi i$ -periodic even function of the rapidity differences,  $Q_{a_1 \dots a_n}^{\mathcal{O}}(\theta_1, \dots, \theta_n)$  encodes the non-locality properties of  $\mathcal{O}$  with respect to the operator generating particles, and  $F_{a_i a_j}^{\min}(\theta_{ij})$  are the so-called ‘minimal’ FFs, which satisfy

$$\begin{aligned} F_{ij}^{\min}(\theta) &= S_{ij}(\theta) F_{ji}^{\min}(-\theta), \\ F_{ij}^{\min}(\theta + 2\pi i) &= F_{ji}^{\min}(-\theta). \end{aligned} \quad (26)$$

Being relevant to section 6, the non-locality equation satisfied by FFs in the two-particle case reads

$$F_{a_1 a_2}^{\mathcal{O}}(\theta) = S_{a_1 a_2}(\theta) e^{i2\pi \gamma_{a_2, \mathcal{O}}} F_{a_1 a_2}^{\mathcal{O}}(\theta + 2\pi i), \quad (27)$$

with  $\theta = \theta_1 - \theta_2$ .

The FFs can be found by solving a set of linear equations in correspondence with kinematic and dynamical poles, for a suitable parametrization of  $Q_{a_1 \dots a_n}^{\mathcal{O}}(\theta_1, \dots, \theta_n)$ . Concerning bound-state poles, e.g.  $|A_i(\theta)A_j(\theta + iu_{ij}^k)\rangle \rightarrow |A_k(iu_{ij}^k)\rangle$ , a two-particle FF obeys

$$-i \lim_{\theta \rightarrow iu_{ij}^k} (\theta - iu_{ij}^k) F_{a_i, a_j}^{\mathcal{O}}(\theta) = \Gamma_{ij}^k F_{a_k}^{\mathcal{O}}, \quad (28)$$

where  $\Gamma_{ij}^k$  is the real three-point vertex

$$\Gamma_{ij}^k = \left[ -i \lim_{\theta \rightarrow iu_{ij}^k} (\theta - iu_{ij}^k) S_{ij}(\theta) \right]^{1/2}. \quad (29)$$

There are also residual equations coming from double poles of the  $S$ -matrix. The equation for a pole located at  $i\phi$  reads

$$-i \lim_{\theta \rightarrow i\phi} (\theta - i\phi) F_{ij}(\theta) = \Gamma_{id}^c \Gamma_{\bar{d}j}^e F_{ce}(i\gamma), \quad (30)$$

where  $\gamma = \pi - u_{cd}^{\bar{i}} - u_{d\bar{e}}^{\bar{j}}$  ( $\bar{j}$  denoting the antiparticle of  $j$ ). The interested reader can find a thorough discussion of these equations in references [10, 34]. The kinematic poles are due to particle–antiparticle annihilation and in these cases the two-particle FFs obey [35]

$$-i \lim_{\theta \rightarrow i\pi} (\theta - i\pi) F_{a_i, a_{\bar{i}}}^{\mathcal{O}}(\theta) = [1 - e^{i2\pi \gamma_{a_i, \mathcal{O}}}] \langle \mathcal{O} \rangle, \quad (31)$$

where  $\langle \mathcal{O} \rangle$  is the vacuum expectation value (VEV) of the operator  $\mathcal{O}$ .

In the two-particle case,  $Q_{a_i a_j}^{\mathcal{O}}(\theta)$  is a polynomial in  $e^\theta$ , which satisfies an upper bound on its maximal degree [34]. Define  $y_{\mathcal{O}}$  to be the coefficient multiplying  $\theta_i$  as it approaches infinity, i.e.

$$F_{a_i a_j}^{\mathcal{O}}(\theta_i, \theta_j) \underset{|\theta_i| \rightarrow \infty}{\sim} \exp\{y_{\mathcal{O}}|\theta_i|\}. \quad (32)$$

The upper bound on the degree of the FF reads

$$y_{\mathcal{O}} \leq \Delta_{\mathcal{O}}, \quad (33)$$

where  $\Delta_{\mathcal{O}}$  is the conformal weight of the operator  $\mathcal{O}$  in the UV CFT.

The latter bound and the aforementioned linear equations are in most cases sufficient to fix the FFs as functions of the VEV of some operator. However, it may happen that they are not enough. By computing the FFs of order and disorder, leading and subleading, operators of the thermally deformed TPM, we also made use of second-order equations coming from the asymptotic factorization of the FFs [32].

## 5. FFs of the stress–energy tensor

This section outlines the computation of the FFs of the stress–energy tensor  $\Theta(x)$ , as originally obtained in [10].

Since the FFs of the stress–energy tensor corresponding to the same particle states all have the same pole structure, the two-particle FFs of the trace  $\Theta$  can be parametrized by the following polynomial:

$$Q_{ab}^{\Theta}(\theta) = \left( \cosh \theta + \frac{m_a^2 + m_b^2}{2m_a m_b} \right)^{1 - \delta_{m_a, m_b}} \sum_{i=0}^{\deg Q_{ab}^{\Theta}} c_{ab}^i (\cosh \theta)^i, \quad (34)$$

where the degree is determined by the asymptotic behavior equation (33). Notice that when particles have the same mass, i.e. in the case of a particle–antiparticle doublet or a singlet, the prefactor in the latter equation is completely absorbed by the unknown coefficients  $c_{ab}^{(i)}$ . Being a conserved quantity of the theory, the FFs of  $\Theta$  do not obey the usual residue equations for particle–antiparticle annihilation, rather:

$$F_{a\bar{a}}^{\Theta}(i\pi) = 2\pi m_a^2. \quad (35)$$

Apart from these properties,  $\Theta$  is a  $Z_3$ -invariant operator, like  $\mu$  and  $\zeta$ . The problem of finding the one- and two-particle FFs is therefore similar for these three operators. In the following, we will present the parametrization and the equations satisfied by the one-particle and first two-particle FFs (table 7).

**Table 7.** Multi-particle states with lowest center of mass energy and non-trivial FFs with respect to the  $Z_3$ -invariant operators.

state	$E_{CM}/m_l$
$L$	1.41421...
$\bar{l}\bar{l}, \bar{l}l$	$\geq 2$
$H$	2.73205...
$LL$	$\geq 2.82843...$
$\bar{l}h, \bar{l}h, \bar{h}l, h\bar{l}$	$\geq 2.93185...$
$ll, \bar{l}\bar{l}$	$\geq 3$

The first two-particle FFs of  $\Theta$  are parametrized as follows

$$\begin{aligned}
 F_{\bar{l}\bar{l}}^\Theta(\theta) &= c_{\bar{l}\bar{l}}^0 \frac{F_{\bar{l}\bar{l}}^{\min}(\theta)}{D_{\bar{l}\bar{l}}(\theta)}, \\
 F_{LL}^\Theta(\theta) &= (c_{LL}^0 + c_{LL}^1 \cosh \theta) \frac{F_{LL}^{\min}(\theta)}{D_{LL}(\theta)}, \\
 F_{\bar{l}h}^\Theta(\theta) &= c_{\bar{l}h}^0 \left( \cosh \theta + \frac{m_l^2 + m_h^2}{2m_l m_h} \right) \frac{F_{\bar{l}h}^{\min}(\theta)}{D_{\bar{l}h}(\theta)},
 \end{aligned} \tag{36}$$

and satisfy the equations

$$F_{\bar{l}\bar{l}}^\Theta(i\pi) = 2\pi m_l^2, \tag{37}$$

$$-i \lim_{\theta \rightarrow iu_{\bar{l}\bar{l}}^L} (\theta - iu_{\bar{l}\bar{l}}^L) F_{\bar{l}\bar{l}}^\Theta(\theta) = \Gamma_{\bar{l}\bar{l}}^L F_L^\Theta, \tag{38}$$

$$F_{LL}^\Theta(i\pi) = 2\pi m_L^2, \tag{39}$$

$$-i \lim_{\theta \rightarrow iu_{LL}^L} (\theta - iu_{LL}^L) F_{LL}^\Theta(\theta) = \Gamma_{LL}^L F_L^\Theta, \tag{40}$$

$$-i \lim_{\theta \rightarrow iu_{LL}^H} (\theta - iu_{LL}^H) F_{LL}^\Theta(\theta) = \Gamma_{LL}^H F_H^\Theta, \tag{41}$$

$$-i \lim_{\theta \rightarrow i\pi/2} (\theta - i\pi/2) F_{LL}^\Theta(\theta) = \Gamma_{LL}^l \Gamma_{L\bar{l}}^{\bar{l}} F_{\bar{l}\bar{l}}^\Theta(0), \tag{42}$$

$$-i \lim_{\theta \rightarrow iu_{\bar{l}h}^L} (\theta - iu_{\bar{l}h}^L) F_{\bar{l}h}^\Theta(\theta) = \Gamma_{\bar{l}h}^L F_L^\Theta, \tag{43}$$

$$-i \lim_{\theta \rightarrow iu_{\bar{l}h}^H} (\theta - iu_{\bar{l}h}^H) F_{\bar{l}h}^\Theta(\theta) = \Gamma_{\bar{l}h}^H F_H^\Theta, \tag{44}$$

$$-i \lim_{\theta \rightarrow i\pi 7/12} (\theta - i\pi 7/12) F_{\bar{l}h}^\Theta(\theta) = \Gamma_{\bar{l}\bar{l}}^{\bar{l}} \Gamma_{\bar{h}l}^l F_{\bar{l}\bar{l}}^\Theta(i\pi 2/12). \tag{45}$$

Solutions of this system of equations are provided in table 8 and reproduce the ones of [10]. As a further check of consistency, we computed the contributions to the  $c$ -theorem equation (24) (table 9).



**Table 8.** Parameters of the first FFs of the trace of the stress–energy tensor  $\Theta$  and the energy operator  $\epsilon$ .

Param./ $m_l^2$		Param./ $\langle\epsilon\rangle$	
$F_L^\Theta$	1.2613539473...	$F_L^\epsilon$	-0.9499625864...
$c_{\bar{l}}^0$	6.2831853071...	$d_{\bar{l}}^0$	-4.7320508076...
$F_H^\Theta$	0.2920374047...	$F_H^\epsilon$	-0.2199419194...
$c_{LL}^0$	21.765592370...	$d_{LL}^0$	-16.392304845...
$c_{LL}^1$	9.1992217564...	$d_{LL}^1$	-6.9282032303...
$c_{\bar{l}\bar{h}}^0$	25.226482640...	$d_{\bar{l}\bar{h}}^0$	-18.998802632...

**Table 9.**  $c$ -theorem contributions of the first five particle states ( $c = 6/7 \approx 0.85714$ ).

State	$c$ -series
$A_L$	0.7596528
$A_{\bar{l}}$	0.0853913
$A_H$	0.0029236
$A_{LL}$	0.0024957
$A_{\bar{l}\bar{h}}$	0.0049010
Total	0.85 536...
Relative error	0.207%

Since the fundamental mass  $m_l$  obeys equation (35), it is possible to compute  $\langle\Theta\rangle$  from the cluster equation [36]

$$\lim_{\theta \rightarrow \infty} F_{LL}^\Theta(\theta) = \frac{w_{LL}}{\langle\Theta\rangle} F_L^\Theta F_L^\Theta, \tag{46}$$

where  $w_{LL}$  is a phase, which can be either 1 or  $-1$ . In order to make the  $\Delta$ -theorem contributions for the energy operator  $\epsilon$  positive, we choose the phase  $w_{LL}$  such that  $\langle\Theta\rangle$  is negative, yielding

$$\langle\Theta\rangle = -1.3277932893 \dots m_l^2. \tag{47}$$

Combining equations (12) and (15) we get

$$\langle\epsilon\rangle = -1.7303450451 \dots \lambda^{1/6}, \tag{48}$$

which coincides with the VEV of the energy operator in the thermal perturbation of the whole minimal model  $\mathcal{M}_{6,7}$  in the conformal normalization [37].

**Table 10.**  $\Delta$ -theorem contributions for the energy operator  $\epsilon$  of the first five particle states ( $\Delta_\epsilon = 1/7 \approx 0.14286$ ).

State	$\Delta_\epsilon$ -series
$A_L$	0.0953527
$A_{\bar{l}}$	0.0313356
$A_H$	0.0013696
$A_{LL}$	0.0026516
$A_{\bar{h}}$	0.0046187
Total	0.135 328
Relative error	5.27%

The FFs of  $\epsilon$  retain the same pole structure and polynomial dependence as the ones of  $\Theta$ , with modified coefficients; in particular, the FFs are parametrized as follows

$$\begin{aligned}
 F_{\bar{l}}^\epsilon(\theta) &= d_{\bar{l}}^0 \frac{F_{\bar{l}}^{\min}(\theta)}{D_{\bar{l}}(\theta)}, \\
 F_{LL}^\epsilon(\theta) &= (d_{LL}^0 + d_{LL}^1 \cosh \theta) \frac{F_{LL}^{\min}(\theta)}{D_{LL}(\theta)}, \\
 F_{\bar{h}}^\epsilon(\theta) &= d_{\bar{h}}^0 \left( \cosh \theta + \frac{m_l^2 + m_h^2}{2m_l m_h} \right) \frac{F_{\bar{h}}^{\min}(\theta)}{D_{\bar{h}}(\theta)},
 \end{aligned} \tag{49}$$

with the unknown coefficients satisfying

$$d_{ab}^i = \frac{c_{ab}^i}{\langle \Theta \rangle} \langle \epsilon \rangle. \tag{50}$$

For the sake of completeness, their values are summarized in table 8 and the contributions of the respective FFs in the  $\Delta$  sum rule are shown in table 10.

## 6. FFs of order and disorder leading and subleading operators

This section is dedicated to the derivation of the first FFs of magnetic operators in the thermally deformed TPM. Let us start with the leading disorder operator.

### 6.1. Leading disorder operator $\mu$

Because of  $\mathcal{C}$ -invariance,  $\langle \tilde{\mu} \rangle$  and  $\langle \mu \rangle$  coincide. In table 7, we present the first states with lowest mass and non-trivial FFs. Since the computation of these FFs involves non-locality factors, the derivation will be outlined step-by-step.

Let us start from  $F_{\bar{l}}^\mu(\theta)$ . Since in the end we are concerned with computing the off-critical two-point correlation functions of the operators, it is important to notice that

the only contribution that needs to be evaluated numerically in equation (22) is that of  $F_{\bar{l}l}^\mu(\theta)$ . Indeed, because of  $\mathcal{C}$ -invariance, the FFs of the operator  $\tilde{\mu}$  are read from that of  $\mu$ :

$$F_{\bar{l}l}^{\tilde{\mu}}(\theta) = F_{\bar{l}l}^\mu(\theta), \quad F_{\bar{l}l}^{\tilde{\mu}}(\theta) = F_{\bar{l}l}^\mu(\theta) = S_{\bar{l}l}(\theta) F_{\bar{l}l}^\mu(-\theta), \quad (51)$$

where the last equality is a consequence of equation (26).

Since the disorder operator has non-zero non-locality factor with respect to particles  $l$  and  $\bar{l}$ , the FF will also possess an additional term which takes into account particle–antiparticle annihilation. Due to the asymptotic behaviors of  $F_{\bar{l}l}^{\min}(\theta)$  and  $D_{\bar{l}l}(\theta)$ ,  $Q_{\bar{l}l}^\mu(\theta)$  satisfies the following inequality on its degree:

$$(\deg Q_{\bar{l}l}^\mu) \theta - \frac{5}{6}|\theta| \leq \frac{1}{21}|\theta|. \quad (52)$$

Notice the absence of the absolute value in the first term on the l.h.s. The polynomial  $Q_{\bar{l}l}^\mu(\theta)$  is not symmetric in the exchange  $\theta \rightarrow -\theta$ , since the non-locality factors of  $\mu$  with respect to particles  $l$  and  $\bar{l}$  are non-zero. The non-locality equation that must be satisfied is

$$F_{\bar{l}l}^\mu(\theta) = S_{\bar{l}l}(\theta) e^{i2\pi\gamma_{\bar{l},\mu}} F_{\bar{l}l}^\mu(\theta + 2\pi i), \quad (53)$$

with  $\gamma_{\bar{l},\mu} = -1/3$ . Solutions are sought of the form  $\exp(\alpha\theta)$ , provided that they satisfy

$$e^{\alpha\theta} = -e^{-i2\pi/3} e^{\alpha(\theta+2\pi i)}. \quad (54)$$

Therefore the parametrization for these FFs is

$$F_{\bar{l}l}^\mu(\theta) = \left( a_{\bar{l}l}^0 e^{\frac{5}{6}\theta} + a_{\bar{l}l}^1 e^{-\frac{1}{6}\theta} \right) \frac{F_{\bar{l}l}^{\min}(\theta)}{\cosh\left(\frac{\theta}{2}\right) D_{\bar{l}l}(\theta)}. \quad (55)$$

Pole equations

$$\begin{aligned} -i \lim_{\theta \rightarrow i\pi} (\theta - i\pi) F_{\bar{l}l}^\mu(\theta) &= \left[ -i \lim_{\theta \rightarrow i\pi} (\theta - i\pi) F_{\bar{l}l}^\mu(\theta) \right]^* = \left( 1 - e^{i2\pi/3} \right) \langle \mu \rangle, \\ -i \lim_{\theta \rightarrow iu_{\bar{l}l}^L} (\theta - iu_{\bar{l}l}^L) F_{\bar{l}l}^\mu(\theta) &= -i \lim_{\theta \rightarrow iu_{\bar{l}l}^L} (\theta - iu_{\bar{l}l}^L) F_{\bar{l}l}^\mu(\theta) = \Gamma_{\bar{l}l}^L F_L^\mu, \end{aligned} \quad (56)$$

allow us to fix all three unknowns— $a_{\bar{l}l}^0$ ,  $a_{\bar{l}l}^1$  and  $F_L^\mu$ —as functions of the VEV of the leading disorder operator  $\mu$  (table 11).

As far as  $F_{LL}^\mu(\theta)$  is concerned, the absence of non-locality phases simplifies the parametrization of the FF:

$$F_{LL}^\mu(\theta) = \left( a_{LL}^0 + a_{LL}^1 \cosh \theta \right) \frac{F_{LL}^{\min}(\theta)}{D_{LL}(\theta)}. \quad (57)$$

**Table 11.** Parameters of the first FFs of the leading disorder operator  $\mu$  and sub-leading disorder operator  $\zeta$ .

Param./ $\langle\mu\rangle$		Param./ $\langle\zeta\rangle$	
$F_L^\mu$	-0.3477104392...	$F_L^\zeta$	-2.2476356112...
$a_{l\bar{l}}^0$	-0.3169872981...	$b_{l\bar{l}}^0$	-4.4150635077...
$a_{l\bar{l}}^1$	-1.1830127018...	$b_{l\bar{l}}^1$	-7.6471143143...
		$b_{l\bar{l}}^2$	-2.3660254027...
$F_H^\mu$	-0.0589332596...	$F_H^\zeta$	-0.8408344177...
$a_{LL}^0$	-5.1961524220...	$b_{LL}^0$	-49.9807620886...
$a_{LL}^1$	-0.9282032290...	$b_{LL}^1$	-38.7646096627...
$a_{l\bar{h}}^0$	-0.6587868452...	$b_{l\bar{h}}^0$	-22.6098992318...
$a_{l\bar{h}}^1$	-6.0223791136...	$b_{l\bar{h}}^1$	-74.3815188798...
$a_{l\bar{h}}^2$	-4.2584651101...	$b_{l\bar{h}}^2$	-59.3127674978...
		$b_{l\bar{h}}^3$	-9.4994013120...

However, the corresponding  $S$ -matrix has a second-order pole; therefore, the set of pole equations is

$$\begin{aligned}
 -i \lim_{\theta \rightarrow iu_{LL}^L} (\theta - iu_{LL}^L) F_{LL}^\mu(\theta) &= \Gamma_{LL}^L F_L^\mu, \\
 -i \lim_{\theta \rightarrow iu_{LL}^H} (\theta - iu_{LL}^H) F_{LL}^\mu(\theta) &= \Gamma_{LL}^H F_H^\mu, \\
 -i \lim_{\theta \rightarrow i\pi/2} (\theta - i\pi/2) F_{LL}^\mu(\theta) &= \Gamma_{L\bar{l}}^{\bar{l}} \Gamma_{Ll}^l F_{l\bar{l}}^\mu(0) = \Gamma_{Ll}^l \Gamma_{L\bar{l}}^{\bar{l}} F_{l\bar{l}}^\mu(0).
 \end{aligned}
 \tag{58}$$

Again, all three unknowns ( $F_H^\mu$ ,  $a_{LL}^0$  and  $a_{LL}^1$ ) can be fixed by simply solving these linear equations (table 11). The first non-trivial check of the consistency of these solutions is given by the second-order cluster equation

$$\lim_{\theta \rightarrow \infty} F_{LL}^\mu(\theta) = \frac{w_{LL}}{\langle\mu\rangle} F_L^\mu F_L^\mu,
 \tag{59}$$

satisfied when the phase  $w_{LL} = 1$ .

The FF  $F_{l\bar{h}}^\mu(\theta)$  is obtained along the lines of  $F_{l\bar{l}}^\mu(\theta)$ . In particular, its parametrization reads

$$F_{l\bar{h}}^\mu(\theta) = \left( a_{l\bar{h}}^0 e^{\frac{4}{3}\theta} + a_{l\bar{h}}^1 e^{\frac{1}{3}\theta} + a_{l\bar{h}}^2 e^{-\frac{2}{3}\theta} \right) \frac{F_{l\bar{h}}^{\min}(\theta)}{D_{l\bar{h}}(\theta)},
 \tag{60}$$

and satisfies the pole equations

$$\begin{aligned}
 -i \lim_{\theta \rightarrow iu_{l\bar{h}}^L} (\theta - iu_{l\bar{h}}^L) F_{l\bar{h}}^\mu(\theta) &= \Gamma_{l\bar{h}}^L F_L^\mu, \\
 -i \lim_{\theta \rightarrow iu_{l\bar{h}}^H} (\theta - iu_{l\bar{h}}^H) F_{h\bar{l}}^\mu(\theta) &= \Gamma_{l\bar{h}}^H F_H^\mu, \\
 -i \lim_{\theta \rightarrow i\pi 7/12} (\theta - i\pi 7/12) F_{l\bar{h}}^\mu(\theta) &= \Gamma_{ll}^{\bar{l}} \Gamma_{l\bar{l}}^l F_{l\bar{l}}^\mu(i\pi 2/12), \\
 -i \lim_{\theta \rightarrow i\pi 7/12} (\theta - i\pi 7/12) F_{h\bar{l}}^\mu(\theta) &= \Gamma_{ll}^{\bar{l}} \Gamma_{l\bar{l}}^l F_{l\bar{l}}^\mu(i\pi 2/12).
 \end{aligned} \tag{61}$$

Notice that the latter two pole equations have a peculiar dependence on the two-particle FFs  $F_{l\bar{l}}^\mu(\theta)$  and  $F_{l\bar{l}}^\mu(\theta)$ . Solutions are reported in table 11. We also compute the first contributions to the  $\Delta$ -theorem (table 12).

### 6.2. Subleading disorder operator $\zeta$

The computation of the FFs for the subleading disorder operator  $\zeta$  presents few differences if compared to the case of  $\mu$ . First of all, since the  $\zeta$  has the same transformation properties as  $\mu$ , the non-vanishing FFs of  $\zeta$  are the same as the ones of  $\mu$  (table 7). The only difference in computing the FFs between the leading and subleading disorder operators lies in their different anomalous dimensions, which in turn affects the upper bound on the FF degree. On the other hand, non-locality coefficients remain unchanged in passing from the leading to the subleading case. These considerations are accounted for in the parametrization of some of the  $Q$  polynomials:

$$\begin{aligned}
 F_{l\bar{l}}^\zeta(\theta) &= \left( b_{l\bar{l}}^0 e^{\frac{5}{6}\theta} + b_{l\bar{l}}^1 e^{-\frac{1}{6}\theta} + b_{l\bar{l}}^2 e^{-\frac{7}{6}\theta} \right) \frac{F_{l\bar{l}}^{\min}(\theta)}{\cosh\left(\frac{\theta}{2}\right) D_{l\bar{l}}(\theta)}, \\
 F_{LL}^\zeta(\theta) &= (b_{LL}^0 + b_{LL}^1 \cosh \theta) \frac{F_{LL}^{\min}(\theta)}{D_{LL}(\theta)}, \\
 F_{l\bar{h}}^\zeta(\theta) &= \left( b_{l\bar{h}}^0 e^{\frac{4}{3}\theta} + b_{l\bar{h}}^1 e^{\frac{1}{3}\theta} + b_{l\bar{h}}^2 e^{-\frac{2}{3}\theta} + b_{l\bar{h}}^3 e^{-\frac{5}{3}\theta} \right) \frac{F_{l\bar{h}}^{\min}(\theta)}{D_{l\bar{h}}(\theta)}.
 \end{aligned} \tag{62}$$

The set of linear equations for  $\zeta$  remains the same as the ones for  $\mu$ , except for a change in label  $\mu \rightarrow \zeta$  in all of them. However, it turns out that there is one linearly dependent equation, which does not allow fixing one of the unknowns, e.g.  $F_L^\zeta$ . Studying the cluster equation [36]

$$\lim_{\theta \rightarrow \infty} F_{LL}^\zeta(\theta) = \frac{w_{LL}}{\langle \zeta \rangle} F_L^\zeta F_L^\zeta, \tag{63}$$

provides a solution to the problem. If  $w_{LL} = 1$ , we have two non-physical solutions

$$F_L^\zeta = (-0.2725122015\dots, 2.8678582521\dots) \langle \zeta \rangle,$$

**Table 12.**  $\Delta$ -theorem contributions for the leading disorder operator  $\mu$  ( $\Delta_\mu = 1/21 \approx 0.047619$ ) and the subleading disorder operator  $\zeta$  of the first five particle states ( $\Delta_\zeta = 10/21 \approx 0.47619$ ).

State	$\Delta_\mu$ -series	$\Delta_\zeta$ -series
$A_L$	0.0349015	0.2256069
$A_{\bar{l}}$	0.0095534	0.1266482
$A_H$	0.0003670	0.0051113
$A_{LL}$	0.0005463	0.0121824
$A_{\bar{h}}$	0.0010622	0.0257981
Total	0.0464304	0.395347
Relative error	2.50%	16.98%

since the first value gives a contribution to the  $\Delta$ -theorem which is positive but too small, while the second gives a negative contribution. If  $w_{LL} = -1$ , one finds

$$F_L^\zeta = (-2.2476356112\dots, -0.3477104393\dots) \langle \zeta \rangle.$$

While the second solution is associated with the disorder operator  $\mu$ , the first one is acceptable: its contribution to the  $\Delta$ -theorem is given in table 12 and it is compatible with the conformal weight of the operator. Notice that convergence is slower than in the leading case, as it generally happens when one considers less relevant operators—see also [22]. Solving the system of equations, together with equation (63), yields the results of table 11.

### 6.3. Leading order operator $\sigma$ and subleading order operator $Z$

The FFs of the leading and subleading order operators are simpler to compute if compared to their disordered counterparts, because there are no non-locality factors involved. The first non-trivial FFs are summarized in table 13. It turns out that the solution of their pole equations requires the knowledge of the FFs of disorder operators.

The parametrizations of the first FFs of  $\sigma$  are

$$\begin{aligned}
 F_{ll}^\sigma(\theta) &= a_{ll}^0 \frac{F_{ll}^{\min}(\theta)}{D_{ll}(\theta)}, \\
 F_{lL}^\sigma(\theta) &= (a_{lL}^0 + a_{lL}^1 \cosh \theta) \frac{F_{lL}^{\min}(\theta)}{D_{lL}(\theta)}, \\
 F_{lh}^\sigma(\theta) &= (a_{lh}^0 + a_{lh}^1 \cosh \theta) \frac{F_{lh}^{\min}(\theta)}{D_{lh}(\theta)}, \\
 F_{hL}^\sigma(\theta) &= \left( a_{hL}^0 + a_{hL}^1 \cosh \theta + a_{hL}^2 (\cosh \theta)^2 \right) \frac{F_{hL}^{\min}(\theta)}{D_{hL}(\theta)}
 \end{aligned} \tag{64}$$

**Table 13.** Multi-particle states with lowest center of mass energy and non-trivial FF with respect to the order operator  $\sigma$ . The ones of the subleading order operator  $Z$  are retrieved by charge-conjugating all particles.

State	$E_{CM}/m_l$
$\bar{l}$	1
$\bar{h}$	1.93185...
$ll$	$\geq 2$
$\bar{l}L$	$\geq 2.41421...$
$lh$	$\geq 2.93185...$
$\bar{l}\bar{l}$	$\geq 3$
$\bar{h}L$	$\geq 3.34606...$

and the set of pole equations is

$$-i \lim_{\theta \rightarrow iu_{ll}^{\bar{l}}} (\theta - iu_{ll}^{\bar{l}}) F_{ll}^{\sigma}(\theta) = \Gamma_{ll}^{\bar{l}} F_{\bar{l}}^{\sigma}, \tag{65}$$

$$-i \lim_{\theta \rightarrow iu_{ll}^{\bar{h}}} (\theta - iu_{ll}^{\bar{h}}) F_{ll}^{\sigma}(\theta) = \Gamma_{ll}^{\bar{h}} F_{\bar{h}}^{\sigma}, \tag{66}$$

$$-i \lim_{\theta \rightarrow iu_{lL}^{\bar{l}}} (\theta - iu_{lL}^{\bar{l}}) F_{lL}^{\sigma}(\theta) = \Gamma_{lL}^{\bar{l}} F_{\bar{l}}^{\sigma}, \tag{67}$$

$$-i \lim_{\theta \rightarrow iu_{lL}^{\bar{h}}} (\theta - iu_{lL}^{\bar{h}}) F_{lL}^{\sigma}(\theta) = \Gamma_{lL}^{\bar{h}} F_{\bar{h}}^{\sigma}, \tag{68}$$

$$-i \lim_{\theta \rightarrow iu_{lh}^{\bar{l}}} (\theta - iu_{lh}^{\bar{l}}) F_{lh}^{\sigma}(\theta) = \Gamma_{lh}^{\bar{l}} F_{\bar{l}}^{\sigma}, \tag{69}$$

$$-i \lim_{\theta \rightarrow iu_{lh}^{\bar{h}}} (\theta - iu_{lh}^{\bar{h}}) F_{lh}^{\sigma}(\theta) = \Gamma_{lh}^{\bar{h}} F_{\bar{h}}^{\sigma}, \tag{70}$$

$$-i \lim_{\theta \rightarrow i\pi 5/12} (\theta - i\pi 5/12) F_{lh}^{\sigma}(\theta) = \Gamma_{lh}^{\bar{l}} \Gamma_{ll}^L F_{lL}^{\sigma}(i\pi/12), \tag{71}$$

$$-i \lim_{\theta \rightarrow iu_{hL}^{\bar{l}}} (\theta - iu_{hL}^{\bar{l}}) F_{hL}^{\sigma}(\theta) = \Gamma_{hL}^{\bar{l}} F_{\bar{l}}^{\sigma}, \tag{72}$$

$$-i \lim_{\theta \rightarrow i\pi 8/12} (\theta - i\pi 8/12) F_{hL}^{\sigma}(\theta) = \Gamma_{lh}^l \Gamma_{lL}^l F_{ll}^{\sigma}(i\pi 4/12), \tag{73}$$

$$-i \lim_{\theta \rightarrow i\pi 4/12} (\theta - i\pi 4/12) F_{hL}^{\sigma}(\theta) = \Gamma_{lh}^l \Gamma_{lL}^h F_{lh}^{\sigma}(i\pi/12), \tag{74}$$

$$-i \lim_{\theta \rightarrow i\pi 6/12} (\theta - i\pi 6/12) F_{hL}^{\sigma}(\theta) = \Gamma_{hL}^{\bar{l}} \Gamma_{ll}^L F_{lL}^{\sigma}(i\pi/12). \tag{75}$$

These equations are linearly dependent and  $F_{\bar{l}}^{\sigma}$  cannot be fixed in this way. This value, however, can be obtained employing the cluster equation

$$\lim_{\theta \rightarrow +\infty} F_{ll}^{\mu}(\theta) = \frac{1}{\langle \mu \rangle} F_{\bar{l}}^{\bar{\sigma}} F_{\bar{l}}^{\sigma} = \frac{1}{\langle \mu \rangle} (F_{\bar{l}}^{\sigma})^2. \tag{76}$$

This equation allows us to fix  $F_l^\sigma$  up to a sign, which, however, is not relevant to the computation of two-point correlation functions. The parameters are summarized in table 14.

For the subleading order operator, the parametrization of the FFs is

$$\begin{aligned}
 F_{l\bar{l}}^Z(\theta) &= (b_{l\bar{l}}^0 + b_{l\bar{l}}^1 \cosh \theta) \frac{F_{l\bar{l}}^{\min}(\theta)}{D_{l\bar{l}}(\theta)}, \\
 F_{lL}^Z(\theta) &= (b_{lL}^0 + b_{lL}^1 \cosh \theta) \frac{F_{lL}^{\min}(\theta)}{D_{lL}(\theta)}, \\
 F_{l\bar{h}}^Z(\theta) &= \left( b_{l\bar{h}}^0 + b_{l\bar{h}}^1 \cosh \theta + b_{l\bar{h}}^2 (\cosh \theta)^2 \right) \frac{F_{l\bar{h}}^{\min}(\theta)}{D_{l\bar{h}}(\theta)}, \\
 F_{hL}^Z(\theta) &= \left( b_{hL}^0 + b_{hL}^1 \cosh \theta + b_{hL}^2 (\cosh \theta)^2 \right) \frac{F_{hL}^{\min}(\theta)}{D_{hL}(\theta)}
 \end{aligned} \tag{77}$$

and the set of pole equations is the same as the one of  $\sigma$ , provided that we substitute  $\sigma \rightarrow Z$  and charge-conjugate all particles, since  $\sigma$  and  $Z$  have complex conjugate  $Z_3$  phases. It turns out that, in solving the set of linear equations,  $F_l^Z$  and  $F_h^Z$  remain unknown. Conjointly solving

$$\begin{aligned}
 \lim_{\theta \rightarrow \infty} F_{lL}^Z(\theta) &= \frac{w_{lL}}{\langle \zeta \rangle} F_l^Z F_L^\zeta, \\
 \lim_{\theta \rightarrow \infty} F_{hL}^Z(\theta) &= \frac{w_{hL}}{\langle \zeta \rangle} F_h^Z F_L^\zeta, \\
 \lim_{\theta \rightarrow +\infty} F_{l\bar{l}}^\zeta(\theta) &= \frac{w_{l\bar{l}}}{\langle \zeta \rangle} (F_l^Z)^2,
 \end{aligned} \tag{78}$$

allows us finding  $F_l^Z$  and  $F_h^Z$  without any phase ambiguity between them— $F_l^Z$  is fixed up to an overall sign, still. The parameters of the FFs of  $Z$  are summarized in table 14.

## 7. Universal ratios of the renormalization group

In the following we are going to consider some universal ratios of the renormalization group associated with the thermal deformation of TPM, i.e. with the action equation (11). Consider an operator  $\phi_j$  of the theory which has at criticality the anomalous dimension  $\mathcal{X}_j = 2\Delta_j$  and therefore renormalization group eigenvalue  $y_j = 2 - 2\Delta_j$ . The coupling  $\lambda$  of the theory is a dimensional quantity related to the lightest excitation of the theory  $m_l$  as in equation (15). Denoting by  $K_j$  the metric factor associated to the operator  $\phi_j$ , on general grounds the scaling form of the VEV  $\langle \phi_j \rangle$  can be written as (see [13] for a more detailed discussion)

$$\langle \phi_j \rangle = B_{j\epsilon} \lambda^{\mathcal{X}_j/y_\epsilon}, \tag{79}$$



**Table 14.** Parameters of the first FFs of the leading order operator  $\sigma$  and subleading order operator  $Z$ .

Param./ $\langle\mu\rangle$		Param./ $\langle\zeta\rangle$	
$F_l^\sigma$	0.6143373207...	$F_l^Z$	2.2927380937...
$F_h^\sigma$	-0.1788394265...	$F_h^Z$	-1.6446346242...
$a_{ll}^0$	3.2187030369...	$b_{ll}^0$	18.4497693307...
$a_{ll}^1$	5.8427632193...	$b_{ll}^1$	12.8748121373...
$a_{lL}^0$	1.7461593992...	$b_{lL}^0$	46.9842992716...
$a_{lL}^1$	66.6327984972...	$b_{lL}^1$	42.1249702982...
$a_{lh}^0$	61.9658050968...	$b_{lh}^0$	695.4260997639...
$a_{lh}^1$	45.6244273941...	$b_{lh}^1$	960.8141050048...
$a_{hL}^0$	45.6244273770...	$b_{hL}^0$	276.4554994639...
$a_{hL}^1$	5.4770385951...	$b_{hL}^1$	531.5286291647...
$a_{hL}^2$		$b_{hL}^2$	851.3434042817...
		$b_{hL}^3$	325.5818137048...

with  $B_{j\epsilon} := K_j K_\epsilon^{\mathcal{X}_j/y_\epsilon}$ . Generalized susceptibilities

$$\hat{\Gamma}_{jk}^\epsilon = \Gamma_{jk}^\epsilon \lambda_\epsilon^{(2-y_j-y_k)/y_\epsilon}, \tag{80}$$

with  $\Gamma_{jk}^i := K_j K_k K_\epsilon^{(2-y_j-y_k)/y_\epsilon}$ , are related to two-point connected off-critical correlators through the fluctuation-dissipation theorem

$$\hat{\Gamma}_{jk}^\epsilon = \int d^2x \langle \phi_j(x) \phi_k(0) \rangle_\epsilon^C. \tag{81}$$

Universal ratios are defined as ratios involving the correlation length  $\xi$ , the expectation values  $\langle \phi_j \rangle$  and generalized susceptibilities  $\hat{\Gamma}_{jk}^\epsilon$  such that the dependence on the metric factors  $K_j$  cancels out [13]. These quantities are associated with the universality class and therefore are the same for different microscopic realizations of the model. The first trivial universal ratio is the ratio between the correlation lengths in the disordered and ordered phase:

$$\frac{\xi_+}{\xi_-} = \frac{m_L}{m_l} = 2 \cos(\pi/4) = \sqrt{2} \approx 1.41421. \tag{82}$$

In order to find non-trivial universal ratios starting from the FFs computed in previous sections, equation (81) must be rewritten as to make the dependence on  $\lambda$  manifest. Notice, however, that some susceptibilities are identically zero, because of  $Z_3$ -symmetry

**Table 15.** First contributions for the generalized susceptibilities of  $Z_3$ -invariant operators.

State	$\hat{\Gamma}_{\mu\mu}^\epsilon \frac{m_l^2}{\langle\mu\rangle^2}$	$\hat{\Gamma}_{\epsilon\epsilon}^\epsilon \frac{m_l^2}{\langle\epsilon\rangle^2}$	$\hat{\Gamma}_{\zeta\zeta}^\epsilon \frac{m_l^2}{\langle\zeta\rangle^2}$	$\hat{\Gamma}_{\mu\epsilon}^\epsilon \frac{m_l^2}{\langle\mu\rangle\langle\epsilon\rangle}$	$\hat{\Gamma}_{\mu\zeta}^\epsilon \frac{m_l^2}{\langle\mu\rangle\langle\zeta\rangle}$	$\hat{\Gamma}_{\epsilon\zeta}^\epsilon \frac{m_l^2}{\langle\epsilon\rangle\langle\zeta\rangle}$
$A_L$	0.12090	0.90243	5.05186	0.33031	0.78153	2.13517
$A_{\bar{l}}$	0.02860	0.29656	6.90261	0.09041	0.35195	1.19861
$A_H$	0.00093	0.01296	0.18053	0.00347	0.01296	0.04837
$A_{LL}$	0.00110	0.02509	0.53618	0.02329	0.02329	0.11530
$A_{\bar{l}h}$	0.00237	0.04371	1.90493	0.01005	0.05368	0.24416
Sum	0.15390	1.28076	14.5761	0.45754	1.22341	3.74160

**Table 16.** First contributions for the generalized susceptibilities of magnetization operators.

State	$\hat{\Gamma}_{\sigma\sigma}^\epsilon \frac{m_l^2}{\langle\mu\rangle^2}$	$\hat{\Gamma}_{Z\bar{Z}}^\epsilon \frac{m_l^2}{\langle\zeta\rangle^2}$	$\hat{\Gamma}_{\sigma\bar{Z}}^\epsilon \frac{m_l^2}{\langle\mu\rangle\langle\zeta\rangle}$
$A_{\bar{l}}$	0.75482	10.5133	2.81702
$A_{\bar{h}}$	0.01714	1.44951	0.15762
$A_{ll}$	0.00838	4.49913	0.13430
$A_{\bar{l}L}$	0.00919	2.17994	0.13840
$A_{lh}$	0.00120	1.33517	0.02814
$A_{\bar{h}L}$	0.00041	0.33581	0.01113
Sum	0.79113	20.3128	3.28662

as, for instance,  $\hat{\Gamma}_{\sigma\mu}^\epsilon$ ,  $\hat{\Gamma}_{\sigma Z}^\epsilon$  and  $\hat{\Gamma}_{\sigma\bar{\sigma}}^\epsilon$ . Charge conjugation also reduces the number of quantities to compute, e.g.  $\hat{\Gamma}_{\bar{\mu}\bar{\mu}}^\epsilon = \hat{\Gamma}_{\mu\mu}^\epsilon$ . In table 15 are presented the generalized susceptibilities involving  $Z_3$ -invariant operators ( $\mu$ ,  $\epsilon$  and  $\zeta$ ), while table 16 is dedicated to magnetization ones ( $\sigma$  and  $Z$ ).

In general, FFs depend naturally on the VEV of some operator

$$F_{a_1, \dots, a_n}^{\phi_j}(\theta_1, \dots, \theta_n) = f_{a_1, \dots, a_n}^{\phi_j}(\theta_1, \dots, \theta_n) \langle \phi'_j \rangle_i, \tag{83}$$

where  $\phi'_j$  is often coincident with  $\phi_j$ , like in the case of  $\mu$ ,  $\epsilon$  and  $\zeta$ , but not always, as it happens for  $\sigma$  and  $Z$ . Manipulations of equation (22) yield

$$\Gamma_{jk}^\epsilon = 2\pi C_\epsilon^{-2} (B'_{j\epsilon})^* B'_{k\epsilon} \mathcal{F}_{jk}^\epsilon, \tag{84}$$

where

$$\begin{aligned} \mathcal{F}_{jk}^i = & \int_0^\infty d\tau \tau \left\{ \sum_{\{n\}} \int_{-\infty}^\infty \frac{d\theta_1 \dots d\theta_n}{n! (2\pi)^n} \exp\left(-\tau \sum \beta_{a_i} \cosh \theta_i\right) \right. \\ & \left. \times \left[ f_{a_1, \dots, a_n}^{\phi_j}(\theta_1, \dots, \theta_n) \right]^* f_{a_1, \dots, a_n}^{\phi_k}(\theta_1, \dots, \theta_n) \right\}. \end{aligned} \tag{85}$$

Here,  $\beta_{a_j} = m_{a_j}/m_l$ . By approximating the two-point correlation function with the sum of the contributions associated to the one- and two-particle states with lowest center-of-mass energy, we present hereafter some universal ratios of the thermal deformation of the TPM:

$$(R_c)_{\sigma\sigma}^\epsilon = \frac{\Gamma_{\epsilon\epsilon}^\epsilon \Gamma_{\sigma\sigma}^\epsilon}{B_{\mu\epsilon} B_{\mu\epsilon}} \approx 0.0153975135, \tag{86}$$

$$(R_c)_{\sigma Z}^\epsilon = \frac{\Gamma_{\epsilon\epsilon}^\epsilon \Gamma_{\sigma Z}^\epsilon}{B_{\mu\epsilon} B_{\zeta\epsilon}} \approx 0.0639662952, \tag{87}$$

$$(R_c)_{\mu\mu}^\epsilon = \frac{\Gamma_{\epsilon\epsilon}^\epsilon \Gamma_{\mu\mu}^\epsilon}{B_{\mu\epsilon} B_{\mu\epsilon}} \approx 0.0029953833, \tag{88}$$

$$(R_c)_{\mu\epsilon}^\epsilon = \frac{\Gamma_{\epsilon\epsilon}^\epsilon \Gamma_{\mu\epsilon}^\epsilon}{B_{\mu\epsilon} B_{\epsilon\epsilon}} \approx 0.0089049910, \tag{89}$$

$$(R_c)_{\mu\zeta}^\epsilon = \frac{\Gamma_{\epsilon\epsilon}^\epsilon \Gamma_{\mu\zeta}^\epsilon}{B_{\mu\epsilon} B_{\zeta\epsilon}} \approx 0.0238108180, \tag{90}$$

$$(R_c)_{\epsilon\epsilon}^\epsilon = \frac{\Gamma_{\epsilon\epsilon}^\epsilon \Gamma_{\epsilon\epsilon}^\epsilon}{B_{\epsilon\epsilon} B_{\epsilon\epsilon}} \approx 0.0249269560, \tag{91}$$

$$(R_c)_{\zeta\epsilon}^\epsilon = \frac{\Gamma_{\epsilon\epsilon}^\epsilon \Gamma_{\zeta\epsilon}^\epsilon}{B_{\zeta\epsilon} B_{\epsilon\epsilon}} \approx 0.0728215391, \tag{92}$$

$$(R_c)_{ZZ}^\epsilon = \frac{\Gamma_{\epsilon\epsilon}^\epsilon \Gamma_{ZZ}^\epsilon}{B_{\zeta\epsilon} B_{\zeta\epsilon}} \approx 0.3953414813, \tag{93}$$

$$(R_c)_{\zeta\zeta}^\epsilon = \frac{\Gamma_{\epsilon\epsilon}^\epsilon \Gamma_{\zeta\zeta}^\epsilon}{B_{\zeta\epsilon} B_{\zeta\epsilon}} \approx 0.2836900244. \tag{94}$$

It should be noted that the universal ratios involving the subleading order and disorder operators are subject to more uncertainty than those containing more relevant operators. Indeed for less relevant operators, the spectral series converges more slowly, as evidenced by the  $\Delta$ -theorem contributions.

### 8. Comparison with a Monte Carlo study

In this section we present some results coming from numerical simulations of the thermal deformation of the tricritical three-state Potts model on isotropic square lattices of spacing  $a$ . Hereafter we denote the sizes of the lattice in the two directions as  $L_t = aN_t$  and  $L_x = aN_x$ ; for all of the simulations described here, we used  $L_t = L_x \equiv L$ . Periodic boundary conditions are imposed in both directions; this choice, which preserves translational symmetry under translations by integer multiples of the lattice spacing along the  $t$ - and  $x$ -axes, minimizes the impact of artifacts due to the finiteness of the system. The reduced Hamiltonian of the lattice model is defined by equation (1), which here we re-write in the following form,

$$\frac{H}{k_B T} = -K \sum_{\langle i,j \rangle} \delta_{s_i, s_j} (1 - \delta_{s_i, 0}) - V \sum_{\langle i,j \rangle} \delta_{s_i, 0} \delta_{s_j, 0} - D \sum_i \delta_{s_i, 0}, \tag{95}$$

to account for the implementation used in our simulation code, where the spin and vacancy variables are simultaneously encoded into a single variable  $s_i \in \{0, 1, 2, 3\}$  defined on each lattice site  $i$ , with the value 0 corresponding to a vacancy, while the other three values correspond to the site being occupied by a spin. As in equation (1), the  $\langle i, j \rangle$  notation denotes pairs of nearest-neighbor lattice sites. To simplify our numerical study, we restricted our attention to simulations for the  $V=0$  case only—that is, we neglected the possibility of a vacancy–vacancy coupling—so that the dynamics of the model depends only on the two parameters  $K$  (the coupling between nearest-neighbor spins) and  $D$  (which acts as a chemical potential for the vacancies).

For our Monte Carlo study, we created a dedicated C++ simulation algorithm, which generates the configurations of  $s_i$  variables combining Swendsen–Wang cluster updates [38] (for the spins) with local Metropolis updates [39] (for both the spins and the vacancies). One complete update consists of five Metropolis updates of the entire lattice alternated with five cluster updates of the whole system. While the non-local cluster updates dramatically reduce the critical slowing down close to the critical points, this numerical problem is not completely solved, due to the persistence of non-negligible autocorrelations for the vacancies, whose evolution in Monte Carlo time is affected only indirectly by the cluster updates.

We studied the model for a range of  $K$  values in the proximity of the tricritical value  $K_{tc} = 1.649903(5)$ , holding  $D$  fixed to its tricritical value  $D_{tc} = 3.152152(2)$ . These values for both  $K_{tc}$  and  $D_{tc}$  coincide with those reported in [40] where they were estimated by means of a different method, i.e. the transfer-matrix technique. We performed simulations at 81 equally spaced values of  $K$  in the  $[K_{tc} - 0.04, K_{tc} + 0.04]$  interval; note that values of  $K$  larger than  $K_{tc}$  correspond to thermal deformations driving the system into a broken-symmetry phase, in which the magnetization is expected to be non-zero in the thermodynamic limit. The square lattices that we simulated had sizes  $(L/a)^2 = 64^2$ ,  $128^2$ ,  $256^2$  and  $512^2$ . For each value of  $K$  and each lattice size, after at least  $10^4$  complete updates for reaching Monte Carlo thermalization, we collected  $8 \times 10^5$  configurations for lattices of sizes  $(L/a)^2 = 64^2$  and  $128^2$ ,  $4 \times 10^5$  configurations for lattices of size  $(L/a)^2 = 256^2$ , and a number of configurations between 387610 and 751375 for the lattices of size  $(L/a)^2 = 512^2$ . We used these ensembles of configurations to compute the expectation values of all observables described below, with the exception of the spin–spin correlation function, which was estimated from ensembles consisting of  $4 \times 10^5$  configurations for each value of  $K$  on lattices of sizes  $(L/a)^2 = 64^2$ ,  $128^2$  and  $(L/a)^2 = 256^2$ , and a number of configurations between 201835 and 386155 for each  $K$  value on lattices of size  $(L/a)^2 = 512^2$ .

The observables that we evaluated in our Monte Carlo simulations include:

- The density of vacancies, which is the fraction of sites in the  $s_i = 0$  state

$$\rho = \frac{1}{(L/a)^2} \sum_i \delta_{s_i, 0}. \quad (96)$$

- The contribution to the energy (in units of  $k_B T$ ) coming from spin–spin interactions, divided by the number of lattice sites, which is denoted as

$$\epsilon = \frac{K}{(L/a)^2 k_B T} \sum_{\langle i,j \rangle} \delta_{s_i, s_j} (1 - \delta_{s_i, 0}). \quad (97)$$

In the following (with a slight abuse of terminology), we will also refer to this quantity simply as ‘energy’; in addition, we also considered the associated susceptibility.

- The magnetization in each configuration, which is defined as

$$\sigma = (1 - \rho) \frac{q f_{\max} - 1}{q - 1}, \quad (98)$$

where  $f_{\max}$  denotes the fraction of spins in the majority state in the given configuration,

$$f_{\max} = \max_{k \in \{1,2,3\}} \left( \frac{1}{(L/a)^2} \sum_i \delta_{s_i, k} \right). \quad (99)$$

- Finally, the two-point spin–spin correlation function, which is defined as

$$G_{ss}(r) = \frac{1}{2(L/a)^2} \sum_i \sum_{k=0}^1 (1 - \delta_{s_i, 0}) \delta_{s_i, s_{i+r\hat{u}_k}}, \quad (100)$$

where  $\hat{u}_0$  and  $\hat{u}_1$  respectively denote the positively oriented unit vectors in the  $t$  and  $x$  directions of the lattice.


For all of these variables, we computed the integrated autocorrelation time (for a discussion of the numerical estimate of this quantity in Monte Carlo simulations, see, e.g. [41]), and grouped the whole set of ‘measurements’ of each observable obtained in each simulation into bins of length equal to the integrated autocorrelation time. We verified that, as expected, the average over each bin produced data with negligible residual autocorrelation.

Figure 3 shows how the distribution of vacancies extracted from our simulations varies as a function of  $K_{\text{tc}} - K$ . We first observe that the results from simulations on lattices of different sizes that we considered exhibit a remarkable collapse of data, indicating that finite-volume corrections are under control. Furthermore, the data are consistent with an inflection point at  $K = K_{\text{tc}}$  and the value of  $\rho$  at that point is in full agreement with the result (denoted by the black cross) obtained through the transfer-matrix technique in [40].


Next, we consider the average value of  $\epsilon$ , which is plotted in figure 4, together with the results for the associated susceptibility (displayed in the inset plot, with a logarithmic scale on the vertical axis). As for the vacancy density, also for  $\langle \epsilon \rangle$  we observe a clear collapse of the data obtained from simulations on lattices of different areas, and an inflection point at a location consistent with  $K = K_{\text{tc}}$ . For this observable, the Monte Carlo results are expected to be modeled by a function of the form:

$$\langle \epsilon \rangle = e_0 (K_{\text{tc}} - K)^{1/6} + e_1. \quad (101)$$

Form factors of the tricritical three-state Potts model in its scaling limit



**Figure 3.** Average density of vacancies, displayed as a function of  $K_{tc} - K$ , obtained from our simulations on square lattices of different sizes (denoted by symbols of different shapes and colors). The plot also shows the average density of vacancies at  $K_{tc}$  (shown by the black cross) that was computed in [40] using a different numerical technique.



**Figure 4.** Average contribution to the energy density from spin–spin interactions, plotted against  $K_{tc} - K$ , from Monte Carlo calculations on square lattices of linear size ranging from  $L/a = 64$  to  $L/a = 512$ ; the curve obtained by fitting the data for  $0.005 \leq K_{tc} - K \leq 0.04$  to equation (101) is also shown. The inset plot shows the results for the susceptibility associated with  $\epsilon$  (rescaled by the number of lattice sites), which are displayed with a logarithmic scale for the vertical axis.

Fitting the data in the range  $0.005 \leq K_{\text{tc}} - K \leq 0.04$  to equation (101), one obtains  $e_0 = 2.036(4)$  and  $e_1 = -1.6124(23)$ . The curve obtained from the fit is shown by the black line in figure 4. The good quality of this fit, in which the exponent of  $(K_{\text{tc}} - K)$  is fixed to the value predicted analytically, confirms the validity of the CFT description of the results obtained from Monte Carlo simulations, at least for the range of  $K$  values and for the lattice sizes under consideration in this study.

Our Monte Carlo results for the average magnetization are displayed in figure 5; note that, in the broken-symmetry phase, the results obtained from simulations on lattices of different area are consistent with each other. On the other hand, in the symmetric phase, the results obtained from smaller lattices are systematically larger than those from larger lattices: this is an artifact of the definition of the magnetization in equation (98), which is based on the fraction of spins in the majority state in each configuration: by definition, for a system with a finite number of degrees of freedom, in the symmetric phase  $f_{\text{max}}$  is always larger than  $1/3$ —except in the exceedingly rare configurations in which the total number of variables in the spin states 1, 2 and 3 are exactly equal to each other. Nevertheless, our Monte Carlo results show that the magnetization in the symmetric phase vanishes when it is extrapolated to the thermodynamic limit.

In the broken-symmetry phase, fitting the results for the magnetization in the range  $-0.04 \leq K_{\text{tc}} - K \leq -0.005$  to

$$\langle \sigma \rangle = d_0 |K_{\text{tc}} - K|^{1/18} + d_1, \quad (102)$$

one obtains  $d_0 = 0.9526(15)$  and  $d_1 = 0.1185(13)$ . The result of this fit is shown by the black line in figure 5. As for  $\langle \epsilon \rangle$ , the fact that one obtains a good modeling of Monte Carlo results using this simple functional form, with the exponent of  $|K_{\text{tc}} - K|$  fixed to the value predicted by the CFT, confirms the validity of this description of Monte Carlo results.


Next, we show examples of our results for the two-point spin–spin correlation function in figure 6. In particular, it is interesting to study the universal ratio between the correlation lengths in the symmetric and in the symmetry-breaking phases. Sufficiently close to the tricritical point, it is expected that the ratio of correlation lengths obtained at values of  $K$  that are symmetric with respect to  $K_{\text{tc}}$  should become consistent with the theoretical prediction  $\xi_+/\xi_- = \sqrt{2} = 1.41421\dots$

As an example to illustrate this type of study, the data displayed in figure 6 are obtained from two test ensembles of Monte Carlo simulations on a lattice of size  $(L/a)^2 = 256^2$  at  $K = K_{\text{tc}} \pm 0.01$ , i.e. close to the tricritical point<sup>3</sup>. Expressing all distances in units of the lattice spacing, the correlators can be fitted to

$$G_{ss}(r) = a_0 \frac{\exp(-r/\xi)}{r^{4/21}} + a_2, \quad (103)$$

<sup>3</sup> Each of these two additional ensembles consisted of  $4 \times 10^5$  configurations, that were produced after  $10^4$  complete updates for thermalization, like the other simulations on the lattices of size  $(L/a)^2 = 256^2$ . Note that, in order to (slightly) reduce the computing times, for these two test ensembles the normalization used in the Monte Carlo calculation of the correlators explicitly dismissed an additive contribution to  $G_{ss}(r)$ ; this additive contribution does not depend on  $r$ , and, hence, has no effect on the evaluation of the correlation lengths.

Form factors of the tricritical three-state Potts model in its scaling limit



**Figure 5.** Average magnetization, plotted against  $K_{tc} - K$ , from our Monte Carlo simulations on lattices with the same linear sizes as in figures 3 and 4. The curve obtained fitting our data in the broken-symmetry phase to equation (102) is also shown (black line).

yielding  $\xi = 2.672(82)$  for  $K = 1.639903$  and  $\xi = 1.800(45)$  for  $K = 1.659903$ , whose ratio

$$\frac{\xi_{K=1.639903}}{\xi_{K=1.659903}} = 1.484(59) \tag{104}$$


is within less than 1.2 standard deviations from the analytically predicted value.

It should be noted that the extraction of the correlation length from the Monte Carlo results for the correlators is non-trivial, since it is based on a three-parameter, non-linear fit of cross-correlated data, and the fitted parameters are also quite strongly correlated with each other. This, in turn, leads to a likely overestimate of the uncertainties affecting the fitted parameters, including, in particular, the correlation length. However, we followed a very conservative approach to the error budget, without attempting to reduce these uncertainties. Figure 7 shows the correlation lengths computed from Monte Carlo simulations on lattices of different sizes and at different  $K$  values. In particular, the main plot (in which a logarithmic scale is used for both axes) displays the data in the symmetric phase and their fit to the power-law dependence on  $|K_{tc} - K|$  predicted from the theory:


$$\xi = a_0 |K_{tc} - K|^{-7/12}. \tag{105}$$

The plot makes manifest that the simulation results precisely follow the power-law behavior predicted by the CFT. The only free parameter of the fit to equation (105) is the overall amplitude, for which we obtain  $a_0 = 0.356(1)$ . Analogous results for the broken-symmetry phase are shown in the inset plot (where a linear scale is used for both axes).





**Figure 6.** Two-point spin correlation functions computed from Monte Carlo simulations on a lattice of size  $(L/a)^2 = 256^2$  at two different  $K$  values, shown as a function of the spatial separation  $r$  in units of the lattice spacing  $a$ . The dashed lines are the curves obtained from the fits to equation (103).



**Figure 7.** Correlation lengths computed from Monte Carlo simulations at different  $K$  values and different lattice sizes, and their fit to the power-law behavior predicted from conformal field theory (black line), given by equation (105). The main plot refers to simulations at  $K < K_{tc}$ , whereas the inset plot shows analogous results in the broken-symmetry phase.

## 9. Discussion and conclusions

In this work we have addressed the computation of the FFs of the relevant operators of the TPM once this model is perturbed away from its critical point by varying the temperature. The corresponding massive theory is integrable and has a distinguished fingerprint associated with the  $E_6$  exceptional Lie algebra. We have determined the FFs associated to the leading and subleading order and disorder operators and we have checked their correct identification using the  $\Delta$  sum rule. We have shown how to theoretically compute several universal ratios of the renormalization group associated with the universality class of the model in any of its microscopic realizations. We also presented some results coming from a Monte Carlo study of the lattice model, which provided clear evidence that, in the range of parameters that we considered, the results are in agreement with the expectations from the CFT, and that, in addition, the universal ratio of the masses in the high and low-temperature phases of the model is fully consistent with our theoretical prediction.

### Acknowledgment

G M acknowledges the grants PNRR MUR Project PE0000023-NQSTI and PRO3 Quantum Pathfinder. The work of M P has been partially supported by the Spoke 1 ‘FutureHPC & BigData’ of the Italian Research Center on High-Performance Computing, Big Data and Quantum Computing (ICSC) funded by MUR (M4C2-19) – Next Generation EU (NGEU), by the Italian PRIN ‘Progetti di Ricerca di Rilevante Interesse Nazionale—Bando 2022’, prot. 2022TJFCYB, and by the ‘Simons Collaboration on Confinement and QCD Strings’ funded by the Simons Foundation. The simulations were run on CINECA computers. We acknowledge support from the SFT Scientific Initiative of the Italian Nuclear Physics Institute (INFN).


### Appendix. Analytic properties of FFs

In this appendix, we will briefly touch upon some properties of FFs, complementing the discussion of section 4. Let us start with the connection between the matrix element in the r.h.s. of equations (20) and (21), given by crossing

$$F_{a_1, \dots, a_m, \bar{a}_{m+1}, \dots, \bar{a}_n}^{\mathcal{O}}(\theta_1, \dots, \theta_m, \theta_{m+1} - i\pi, \dots, \theta_n - i\pi) = F_{a_1, \dots, a_n}^{\mathcal{O}}(\theta_{ij}; \theta_{Ak}; \theta_{BC}), \quad (\text{A1})$$

where  $1 \leq i < j \leq m$ ,  $m+1 \leq k \leq m < A \leq n$  and  $m+1 \leq B < C \leq n$ .

In analogy to the  $S$ -matrix, we expect the FFs to be analytic functions in the strip  $0 < \text{Im}\theta_{ij} < 2\pi$  and to have poles in correspondence of bound states. To understand the analytic properties satisfied by FFs, it is useful to



**Figure 8.** The analytic continuation that defines non-local fields.

define the Faddeev–Zamolodchikov (FZ) algebra, which follows from the definition of the scattering matrix. The FZ algebra is an associative algebra of vertex operators which satisfies the following commutation relations involving the  $S$ -matrix:

$$\begin{aligned} A_{a_i}(\theta_i) A_{a_j}(\theta_j) &= S_{ij}(\theta) A_{a_j}(\theta_j) A_{a_i}(\theta_i), \\ A_{a_i}^\dagger(\theta_i) A_{a_j}^\dagger(\theta_j) &= S_{ij}(\theta) A_{a_j}^\dagger(\theta_j) A_{a_i}^\dagger(\theta_i), \\ A_{a_i}(\theta_i) A_{a_j}^\dagger(\theta_j) &= S_{ij}(\theta) A_{a_j}^\dagger(\theta_j) A_{a_i}(\theta_i) + 2\pi \delta_{a_i a_j} \delta(\theta), \end{aligned} \tag{A2}$$

with  $\theta = \theta_i - \theta_j$ . Here  $A_{a_i}(\theta_i)$  is the destruction operator and  $A_{a_i}^\dagger(\theta_i)$  its creation counterpart. Also  $a_i$  is a shorthand for the set of quantum numbers that distinguishes a particle from the others in the spectrum.

It is straightforward to rephrase the definition of FFs equation (21) in the vertex operator formalism. Suppose that the vertex operator  $A_{a_i}^\dagger$  is non-local with respect to the operator  $\mathcal{O}$  [25], which means that after a counterclockwise rotation of  $A_{a_i}^\dagger(\theta')$  around  $\mathcal{O}(\theta)$ , the OPE  $A_{a_i}^\dagger(\theta')\mathcal{O}(\theta)$  acquires a phase  $\exp(i2\pi\gamma_{a_i,\mathcal{O}})$ . If  $\gamma_{a_i,\mathcal{O}}$  is zero, then we have ‘mutual locality’. A graphical representation of this process is presented in figure 8.

Watson equations are direct consequences of the commutation relations equation (A2) and the non-locality properties of the vertex operators with respect to  $\mathcal{O}$ :

$$F_{\dots,a_m,a_{m+1},\dots}^\mathcal{O}(\dots,\theta_m,\theta_{m+1},\dots) = S_{a_m,a_{m+1}}(\theta_m - \theta_{m+1}) F_{\dots,a_{m+1},a_m,\dots}^\mathcal{O}(\dots,\theta_{m+1},\theta_m,\dots), \tag{A3}$$

$$F_{a_1\dots a_n}^\mathcal{O}(\theta_1 + 2\pi i, \theta_2, \dots, \theta_n) = e^{i2\pi\gamma_{a_1,\mathcal{O}}} F_{a_2\dots a_n,a_1}^\mathcal{O}(\theta_2, \dots, \theta_n, \theta_1). \tag{A4}$$

From these equations, it is clear that minimal FFs introduced in section 4 are nothing more than building-block solutions for the two-particle Watson equation, if non-locality is not accounted for. For degenerate mass spectra with reflectionless, i.e. diagonal,  $S$ -matrix, the general structure of a two-particle amplitude is

$$S_{ij}(\theta) = (-1)^{\delta_{i\bar{j}}} \prod_{x \in \mathcal{A}_{ij}} s_x(\theta), \tag{A5}$$

where  $s_x(\theta)$  is the building-block function equation (19) and  $\mathcal{A}_{ij}$  is the set of poles associated with the scattering amplitude—see e.g. table 6. Minimal FFs can be parametrized as

$$F_{ij}^{\min}(\theta) = \left[ -i \sinh\left(\frac{\theta}{2}\right) \right]^{\delta_{ij}} \prod_{x \in \mathcal{A}_{ij}} [h_x(\theta)]^{p_x}, \tag{A6}$$

where  $p_x$  is the multiplicity associated with  $x \in \mathcal{A}_{ij}$ , which coincides with the order of the pole in the corresponding  $S$ -matrix element. The building-block functions  $h_x(\theta)$  satisfy

$$h_x(\theta) = -s_x(\theta) h_x(-\theta), \quad h_x(\theta + 2\pi i) = h_x(-\theta), \tag{A7}$$

and their minimal solution is

$$h_x(\theta) = \exp \left\{ 2 \int_0^\infty \frac{dt \sinh((1-x)t)}{t \sinh^2(t)} \sinh^2\left(\frac{\hat{\theta}t}{2i\pi}\right) \right\}, \tag{A8}$$

with  $\hat{\theta} = i\pi - \theta$ . It should be noted that  $h_x(\theta)$  has poles on the imaginary axis of  $\theta$ , as can be clearly seen after performing an analytic continuation. There are two equivalent ways, the first is

$$h_x(\theta) = \prod_{k=0}^\infty \frac{[\Gamma(k + \frac{1}{2} + \frac{x}{2})]^2 \Gamma(k + 1 - \frac{x}{2} - \frac{i\theta}{2\pi}) \Gamma(k + 2 - \frac{x}{2} + \frac{i\theta}{2\pi})}{[\Gamma(k + \frac{3}{2} - \frac{x}{2})]^2 \Gamma(k + \frac{x}{2} - \frac{i\theta}{2\pi}) \Gamma(k + 1 + \frac{x}{2} + \frac{i\theta}{2\pi})}, \tag{A9}$$

while the second reads

$$h_x(\theta) = \prod_{k=0}^\infty \left[ \frac{1 + \left(\frac{\hat{\theta}/2\pi}{k + \frac{1}{2} + \frac{x}{2}}\right)^2}{1 + \left(\frac{\hat{\theta}/2\pi}{k + \frac{3}{2} - \frac{x}{2}}\right)^2} \right]^{k+1}. \tag{A10}$$

It is possible to find analytic continuations with a finite number of factors, useful in numerical evaluations

$$h_x(\theta) = \prod_{k=0}^{N-1} \frac{[\Gamma(k + \frac{1}{2} + \frac{x}{2})]^2 \Gamma(k + 1 - \frac{x}{2} - \frac{i\theta}{2\pi}) \Gamma(k + 2 - \frac{x}{2} + \frac{i\theta}{2\pi})}{[\Gamma(k + \frac{3}{2} - \frac{x}{2})]^2 \Gamma(k + \frac{x}{2} - \frac{i\theta}{2\pi}) \Gamma(k + 1 + \frac{x}{2} + \frac{i\theta}{2\pi})} \times \exp \left\{ 2 \int_0^\infty \frac{dt \sinh((1-x)t)}{t \sinh^2(t)} e^{-2Nt} \sinh^2 \left( \frac{\hat{\theta}t}{2i\pi} \right) \right\}, \quad (\text{A11})$$

$$h_x(\theta) = \prod_{k=0}^{N+1} \left[ \frac{1 + \left( \frac{\hat{\theta}/2\pi}{k + \frac{1}{2} + \frac{x}{2}} \right)^2}{1 + \left( \frac{\hat{\theta}/2\pi}{k + \frac{3}{2} - \frac{x}{2}} \right)^2} \right]^{k+1} \times \exp \left\{ 2 \int_0^\infty \frac{dt \sinh((1-x)t)}{t \sinh^2(t)} (1 + N - Ne^{-2t}) e^{-2Nt} \sinh^2 \left( \frac{\hat{\theta}t}{2i\pi} \right) \right\}. \quad (\text{A12})$$

The last property of  $h_x(\theta)$  we mention is its asymptotic behavior [10], which finds a natural application in the asymptotic factorization of FFs:

$$h_x(\theta) \underset{|\theta| \rightarrow \infty}{\sim} \exp \left\{ \int_0^\infty \frac{dt}{t} \left[ \frac{\sinh((1-x)t)}{\sinh^2(t)} - \frac{1-x}{t} \right] \right\} \exp \left\{ \frac{1-x}{2} (|\theta| - i\pi) \right\}. \quad (\text{A13})$$

Concerning the parametrization of FFs equation (25), kinematic poles can be implemented by adding a term  $\cosh(\theta_{i\bar{i}}/2)$  at the denominator of the FF. While for the poles coming from the  $S$ -matrix one can use:

$$D_{ij}(\theta) = \prod_{x \in \mathcal{A}_{ij}} [P_x(\theta)]^{n_x}, \quad (\text{A14})$$

with

$$P_x(\theta) = \frac{\cos(\pi x) - \cosh \theta}{2 \cos^2(\pi x/2)} \quad (\text{A15})$$

and

$$n_x = \begin{cases} m + 1, & \text{if } p_x = 2m + 1 \text{ and the pole is in the s-channel;} \\ m, & \text{if } p_x = 2m + 1 \text{ and the pole is in the t-channel;} \\ m, & \text{if } p_x = 2m. \end{cases} \quad (\text{A16})$$

## References

- [1] Belavin A A, Polyakov A M and Zamolodchikov A B 1984 Infinite conformal symmetry in two-dimensional quantum field theory *Nucl. Phys. B* **241** 333–80
- [2] Friedan D, Qiu Z and Shenker S 1984 Conformal invariance, unitarity and critical exponents in two dimensions *Phys. Rev. Lett.* **52** 1575–8
- [3] Fateev V A and Zamolodchikov A B 1990 *Conformal Field Theory and Purely Elastic S-Matrices* (World Scientific) pp 245–70

- [4] Christe P and Mussardo G 1990 Integrable systems away from criticality: the Toda field theory and  $S$ -matrix of the tricritical Ising model *Nucl. Phys. B* **330** 465–87
- [5] Christe P and Mussardo G 1990 Elastic  $S$ -matrices in  $(1+1)$ -dimensions and Toda field theories *Int. J. Mod. Phys. A* **5** 4581–628
- [6] Sotkov G and Zhu C 1989 Bootstrap fusions and tricritical Potts model away from criticality *Phys. Lett. B* **229** 391–7
- [7] Zamolodchikov A B 1989 Integrable field theory from conformal field theory *Integrable Sys Quantum Field Theory* (Elsevier) pp 641–74
- [8] Mussardo G 1992 Off-critical statistical models: factorized scattering theories and bootstrap program *Phys. Rep.* **218** 215–379
- [9] Mussardo G 2020 *Statistical Field Theory: An Introduction to Exactly Solved Models in Statistical Physics (Ogt:C)* 2nd edn (Oxford University Press)
- [10] Acerbi C, Valleriani A and Mussardo G 1996 Form factors and correlation functions of the stress-energy tensor in massive deformation of the minimal models  $((E_n)_1 \otimes (E_n)_1)/(E_n)_2$  *Int. J. Mod. Phys. A* **11** 5327–64
- [11] Smirnov F A 1992 *Form-Factors in Completely Integrable Models of Quantum Field Theory (Advanced Series in Mathematical Physics vol 14)* (World Scientific) pp 1–208
- [12] Karowski M and Weisz P 1978 Exact form-factors in  $(1+1)$ -dimensional field theoretic models with soliton behavior *Nucl. Phys. B* **139** 455–76
- [13] Fioravanti D, Mussardo G and Simon P 2000 Universal amplitude ratios of the renormalization group: two-dimensional tricritical Ising model *Phys. Rev. E* **63** 016103
- [14] Wu F Y 1982 The Potts model *Rev. Mod. Phys.* **54** 235–68
- [15] Nienhuis B, Berker A N, Riedel E K and Schick M 1979 First- and second-order phase transitions in Potts models: renormalization-group solution *Phys. Rev. Lett.* **43** 737–40
- [16] Nienhuis B 1991 Locus of the tricritical transition in a two-dimensional  $q$ -state Potts model *Physica A* **177** 109–13
- [17] Cappelli A, Itzykson C and Zuber J-B 1987 Modular invariant partition functions in two dimensions *Nucl. Phys. B* **280** 445–65
- [18] Cardy J L 1986 Effect of boundary conditions on the operator content of two-dimensional conformally invariant theories *Nucl. Phys. B* **275** 200–18
- [19] Cardy J L 1986 Operator content of two-dimensional conformally invariant theories *Nucl. Phys. B* **270** 186–204
- [20] Rocha-Caridi A and Wallach N R 1984 Characters of irreducible representations of the Virasoro algebra *Math. Z.* **185** 1–21
- [21] Fradkin E 2017 Disorder operators and their descendants *J. Stat. Phys.* **167** 427–61
- [22] Cortés Cubero A, Konik R, Lencsés M, Mussardo G and Takács G 2022 Duality and form factors in the thermally deformed two-dimensional tricritical Ising model *SciPost Phys.* **12** 162
- [23] Capizzi L, Horváth D X, Calabrese P and Castro-Alvaredo O A 2022 Entanglement of the 3-state Potts model via form factor bootstrap: total and symmetry resolved entropies *J. High Energy Phys.* **JHEP05(2022)113**
- [24] Caselle M, Delfino G, Grinza P, Jahn O and Magnoli N 2006 Potts correlators and the static three-quark potential *J. Stat. Mech.* **03008–03008**
- [25] Zamolodchikov A B and Fateev V A 1985 Nonlocal (parafermion) currents in two-dimensional conformal quantum field theory and self-dual critical points in  $Z_N$ -symmetric statistical systems *Sov. Phys. JETP* **62** 215–25
- [26] Di Francesco P, Mathieu P and Sénéchal D 1997 *Conformal Field Theory Graduate Texts in Contemporary Physics* (Springer)
- [27] Braden H W, Corrigan E, Dorey P E and Sasaki R 1990 Affine Toda field theory and exact  $S$ -matrices *Nucl. Phys. B* **338** 689–746
- [28] Eguchi T and Yang S 1989 Deformations of conformal field theories and soliton equations *Phys. Lett. B* **224** 373–8
- [29] Fateev V A 1994 The exact relations between the coupling constants and the masses of particles for the integrable perturbed conformal field theories *Phys. Lett. B* **324** 45–51
- [30] Zamolodchikov A B 1991 Two-point correlation function in scaling Lee-Yang model *Nucl. Phys. B* **348** 619–41
- [31] Cardy J L and Mussardo G 1993 Universal properties of self-avoiding walks from two-dimensional field theory *Nucl. Phys. B* **410** 451–93
- [32] Delfino G, Simonetti P and Cardy J L 1996 Asymptotic factorisation of form factors in two-dimensional quantum field theory *Phys. Lett. B* **387** 327–33
- [33] Zamolodchikov A B 1986 Irreversibility of the flux of the renormalization group in a 2-D field theory *JEPT Lett.* **43** 12

- [34] Delfino G and Mussardo G 1995 The spin-spin correlation function in the two-dimensional Ising model in a magnetic field at  $T = T_c$  *Nucl. Phys. B* **455** 724–58
- [35] Yurov V P and Zamolodchikov A B 1991 Correlation functions of integrable 2D models of the relativistic field theory: Ising model *Int. J. Mod. Phys. A* **06** 3419–40
- [36] Koubek A and Mussardo G 1993 On the operator content of the sinh-Gordon model *Phys. Lett. B* **311** 193–201
- [37] Fateev V, Lukyanov S, Zamolodchikov A and Zamolodchikov A 1998 Expectation values of local fields in the Bullough-Dodd model and integrable perturbed conformal field theories *Nucl. Phys. B* **516** 652–74
- [38] Swendsen R H and Wang J 1987 Nonuniversal critical dynamics in Monte Carlo simulations *Phys. Rev. Lett.* **58** 86–88
- [39] Metropolis N, Rosenbluth A W, Rosenbluth M N, Teller A H and Teller E 1953 Equation of state calculations by fast computing machines *J. Chem. Phys.* **21** 1087–92
- [40] Qian X, Deng Y and Blöte H W J 2005 Dilute Potts model in two dimensions *Phys. Rev. E* **72** 056132
- [41] Wolff U 2004 Monte Carlo errors with less errors *Comput. Phys. Commun.* **156** 143–53 Wolff U 2007 *Comput. Phys. Commun.* **176** 383 (erratum)