




Contents lists available at ScienceDirect

Advances in Mathematics

journal homepage: www.elsevier.com/locate/aim

Stability of Sobolev inequalities on Riemannian manifolds with Ricci curvature lower bounds



Francesco Nobili, Ivan Yuri Violo*

University of Jyväskylä, Department of Mathematics and Statistics, P.O. Box 35 (MaD), FI-40014, Finland

ARTICLE INFO

Article history:

Received 18 October 2022
 Received in revised form 9 October 2023
 Accepted 27 January 2024
 Available online 12 February 2024
 Communicated by Gang Tian

Keywords:

Ricci curvature
 Sobolev inequalities
 Concentration compactness
 Stability

ABSTRACT

We study the qualitative stability of two classes of Sobolev inequalities on Riemannian manifolds. In the case of positive Ricci curvature, we prove that an almost extremal function for the sharp Sobolev inequality is close to an extremal function of the round sphere. In the setting of non-negative Ricci curvature and Euclidean volume growth, we show an analogous result in comparison with the extremal functions in the Euclidean Sobolev inequality. As an application, we deduce a stability result for minimizing Yamabe metrics. The arguments rely on a generalized Lions' concentration compactness on varying spaces and on rigidity results of Sobolev inequalities on singular spaces.

© 2024 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Contents

1.	Introduction	2
2.	Preliminaries	9
2.1.	Calculus on metric measure spaces	9
2.2.	RCD-spaces	11
2.3.	Sobolev inequalities	13
2.4.	Convergence and stability under pmGH-convergence	14

* Corresponding author.

E-mail addresses: francesco.nobili@dm.unipi.it (F. Nobili), ivan.violo@sns.it (I.Y. Violo).

3.	Pólya-Szegő inequality	16
3.1.	Non-compact case	16
3.2.	Rigidity	19
4.	Regularity of extremal functions	24
5.	Rigidity of extremal functions in the Sobolev inequality	27
5.1.	Compact case	27
5.2.	Non-compact case	29
6.	Compactness of extremizing sequences	30
6.1.	Density upper bound	30
6.2.	Concentration compactness for Sobolev extremals	32
7.	Radial functions: technical results	36
8.	Proof of the main results	40
8.1.	Stability in the compact case	40
8.2.	Stability in the non-compact case	46
	Acknowledgments	48
	Appendix A. Concentration compactness: non-compact case	48
A.1.	Technical convergence lemmas	48
A.2.	Concentration compactness principles	51
	Appendix B. Technical results	53
	References	55

1. Introduction

The sharp Sobolev inequality on the standard round sphere S^n , $n > 2$, reads as

$$\|u\|_{L^{2^*}}^2 \leq \frac{2^* - 2}{n} \|\nabla u\|_{L^2}^2 + \|u\|_{L^2}^2, \quad \forall u \in W^{1,2}(S^n), \tag{1.1}$$

where $2^* := 2n/(n - 2)$ and the norms are computed with the renormalized volume measure $\frac{\text{Vol}_{S^n}}{\text{Vol}_{S^n}(S^n)}$. This inequality goes back to the work of Aubin [15], who also characterized non-constant extremizers (see also [68, Chapter 5]) having the following expression (denoting by d the distance induced by the metric):

$$u := \frac{a}{(1 - b \cos(d(\cdot, z_0)))^{\frac{n-2}{2}}}, \quad \text{with } a \in \mathbb{R}, b \in (0, 1), z_0 \in S^n. \tag{1.2}$$

We will refer to them as *spherical bubbles*. A natural question is the one of stability:

(Q) Is a function satisfying almost equality in (1.1) close to a spherical bubble?

Up to a change of coordinates via the stereographic projection (see e.g. [79,43,45]), this question is equivalent to the stability of the Euclidean Sobolev inequality

$$\|u\|_{L^{2^*}(\mathbb{R}^n)} \leq \text{Eucl}(n, 2) \|\nabla u\|_{L^2(\mathbb{R}^n)}, \quad \forall u \in \dot{W}^{1,2}(\mathbb{R}^n), \tag{1.3}$$

where $\dot{W}^{1,2}(\mathbb{R}^n) := \{u \in L^{2^*}(\mathbb{R}^n) : |\nabla u| \in L^2(\mathbb{R}^n)\}$ and $\text{Eucl}(n, 2) > 0$ is the sharp constant, computed by Aubin [16] and Talenti [97] (see (2.10) for its precise value).

Extremizers, i.e. functions u for which equality occurs in (1.3), are also in this case completely characterized:

$$u(x) := \frac{a}{(1 + b|x - z_0|^2)^{\frac{n-2}{2}}}, \quad a \in \mathbb{R}, b > 0, z_0 \in \mathbb{R}^n. \tag{1.4}$$

We shall refer to these functions as *Euclidean bubbles* (usually called Talenti or Aubin-Talenti bubbles). The first *quantitative* stability result was obtained by Bianchi and Egnell [25] who showed that

$$\inf \frac{\|\nabla(u - w)\|_{L^2(\mathbb{R}^n)}}{\|\nabla u\|_{L^2(\mathbb{R}^n)}} \leq C_n \left(\frac{\|\nabla u\|_{L^2(\mathbb{R}^n)}}{\|u\|_{L^{2^*}(\mathbb{R}^n)}} - \text{Eucl}(n, 2)^{-1} \right)^{\frac{1}{2}}, \quad \forall u \in \dot{W}^{1,2}(\mathbb{R}^n), \tag{1.5}$$

for a dimensional constant $C_n > 0$ and the infimum taken among all w as in (1.4). This stability is *strong*, in the sense that the L^2 -norm of the difference of gradients is the biggest possible norm that can be controlled, and optimal, as the exponent $1/2$ is sharp. We mention that quantitative stability for the case of the p -Sobolev inequality in \mathbb{R}^n has also been obtained in sharp form (see [40,47,87,48]). The stability of (1.3) in *qualitative* form, meaning that if the right-hand side of (1.5) is small then so is the left-hand side (in a non-quantified sense), can be deduced via concentration compactness [80,81].

In this note, we address the analogous stability of (Q) for Sobolev inequalities on more general Riemannian manifolds.

Let us consider a closed n -dimensional Riemannian manifold (M, g) , $n > 2$, satisfying

$$\text{Ric}_g \geq (n - 1)g.$$

Under these assumptions the same Sobolev inequality (1.1) as in the sphere holds [72]:

$$\|u\|_{L^{2^*}}^2 \leq \frac{2^* - 2}{n} \|\nabla u\|_{L^2}^2 + \|u\|_{L^2}^2, \quad \forall u \in W^{1,2}(M), \tag{1.6}$$

where the norms are with the renormalized volume measure. Proofs of this inequality using different methods are also given in [19,21,50,68,20,44]. We can ask an analogous stability:

(Q') Is a function satisfying almost equality in (1.6) close to a spherical bubble?

Almost equality here means that

$$Q(u) := \frac{\|u\|_{L^{2^*}}^2 - \|u\|_{L^2}^2}{\|\nabla u\|_{L^2}^2} \sim \frac{2^* - 2}{n}.$$

In the previous work [88], we proved that if $|Q(u) - \frac{2^*-2}{n}|$ is small, then M is qualitatively close in the measure Gromov-Hausdorff sense to a spherical suspension, which

roughly said is a possibly-singular generalization of the round sphere. In particular, when $\sup \mathcal{Q}(u) = n^{-1}(2^* - 2)$, rigidity occurs, i.e. M is isometric to \mathbb{S}^n . These facts already suggested an affirmative answer to (Q') and in fact here we will confirm that this is indeed the case. More precisely, for M as above, every $a \in \mathbb{R}$, $b \in [0, 1)$ and $z \in M$, set

$$w_{a,b,z}(\cdot) := \frac{a}{(1 - b \cos(d(\cdot, z_0)))^{\frac{n-2}{2}}}, \quad (1.7)$$

with the convention that $w_{a,0,z} \equiv a$. Our main result is then the following (as before, all the norms are with respect to the renormalized volume measure):

Theorem 1.1. *For every $\varepsilon > 0$ and $n > 2$ there exists $\delta := \delta(\varepsilon, n) > 0$ such that the following holds. Let (M, g) be an n -dimensional Riemannian manifold with $\text{Ric}_g \geq (n-1)g$ and suppose there exists $u \in W^{1,2}(M)$ non-constant satisfying*

$$\mathcal{Q}(u) > \frac{2^* - 2}{n} - \delta. \quad (1.8)$$

Then, there exist $a \in \mathbb{R}$, $b \in [0, 1)$ and $z \in M$ such that

$$\frac{\|\nabla(u - w_{a,b,z})\|_{L^2} + \|u - w_{a,b,z}\|_{L^{2^*}}}{\|u\|_{L^{2^*}}} \leq \varepsilon. \quad (1.9)$$

Moreover, if $w_{a,b,z} \equiv a$ (i.e. $b = 0$), then $a \in \mathbb{R}$ can be chosen so that the remainder

$$R := u - a$$

satisfies

$$\|R \cdot \|R\|_{L^2}^{-1} - \sqrt{N+1} \cos(d(\cdot, p))\|_{L^2} \leq C_n(\varepsilon^\alpha + \delta)^\beta, \quad (1.10)$$

for some $p \in M$ and positive constants α, β, C_n depending only on n .

The above theorem is the first stability result for the Sobolev inequality that covers a wide class of Riemannian manifolds; indeed up to our best knowledge only very special symmetric cases had been studied so far: see [24] for the hyperbolic space and [51] for $\mathbb{S}^1(1/\sqrt{d-2}) \times \mathbb{S}^{n-1}(1)$.

Some comments on the above statement are in order.

- i) The value of δ depends only on n and $\varepsilon > 0$, but not on the manifold M . Moreover, up to scaling, an analogous statement holds assuming $\text{Ric}_g \geq K$ for some $K > 0$, with δ depending also on K .
- ii) Even if Theorem 1.1 is stated completely in the smooth-setting, its proof will require the study of the Sobolev inequality also in singular spaces (see below the strategy for more details).

iii) The result (1.9) actually holds under a slightly weaker assumption than (1.8), namely:

$$\|u\|_{L^{2^*}(\text{Vol}_g)}^2 \geq A\|\nabla u\|_{L^2(\text{Vol}_g)}^2 + B\|u\|_{L^2(\text{Vol}_g)}^2, \tag{1.11}$$

with $|A - \frac{2^*-2}{n}| + |B - 1| \leq \delta$ (see Remark 8.2).

- iv) The first part of Theorem 1.1 holds also restricting to the class of non constant spherical bubbles, that is $w_{a,b,z}$ with $b \neq 0$.
- v) The second part of Theorem 1.1 should be read as follows: if the almost extremal function u is close to a constant, then (up to changing the constant) the remainder is close in L^2 -sense to a cosine of the distance. Thus, since

$$1 + \varepsilon \cos(d) \sim \frac{1}{(1 + \varepsilon \cos(d))^{\frac{n-2}{2}}},$$

this means that u still retains, at a ‘second-order’ approximation, the shape of a spherical bubble. This extra information essentially comes from the fact that the linearization of the Sobolev inequality is the Poincaré inequality, which means that plugging in (1.6) functions of the type $1 + \varepsilon f$ and sending $\varepsilon \rightarrow 0$ gives the sharp Poincaré inequality for f (see e.g. [88, Lemma 6.7]). Therefore if $1 + \varepsilon f$ satisfies almost equality in (1.6), then f almost satisfies equality in the sharp Poincaré inequality and thus should be close to a cosine of the distance (see [38]).

- vi) When M is not the round sphere, the existence of an extremizer, that is a function which maximizes $\mathcal{Q}(u)$, is unknown in general. This question is contained in [68, Question 4B, Pag. 120] as part of the so-called AB -program around Sobolev inequalities on general Riemannian manifolds. In this direction, we mention the Sobolev-alternative statement proved in [88, Theorem 6.8].

Nevertheless, thanks to the above theorem, we are able to say something about the shape of functions for which this ratio is large, i.e. satisfying (1.8).

Remark 1.2. Note that above we deal only with $p = 2$. The reason is that the inequality

$$\|u\|_{L^{p^*}}^p \leq A\|\nabla u\|_{L^p}^p + \|u\|_{L^p}^p, \quad \forall u \in W^{1,p}(M), \tag{1.12}$$

is false for any $p > 2$, $A > 0$ and any (M, g) closed manifold (see [68, Prop. 4.1]). ■

As an application of Theorem 1.1, we prove a stability-type result for minimizing Yamabe metrics. Recall that a solution to the Yamabe problem on a Riemannian manifold (M, g) is a smooth positive function u such that the metric $u^{\frac{4}{n-2}}g$ has constant scalar curvature (see [99] and also the surveys [78,29]). After the works [98,15,92] it is known that a solution exists on every closed Riemannian manifold and that can be found as a minimizer of

$$Y(M, g) := \inf_{\substack{u \in W^{1,2}(M) \\ u \neq 0}} \mathcal{E}(u) := \inf_{\substack{u \in W^{1,2}(M) \\ u \neq 0}} n(n-1) \frac{\int \frac{2^*-2}{n} |\nabla u|^2 + \frac{\text{Scal}_g}{n(n-1)} u^2 d\text{Vol}_g}{\left(\int |u|^{2^*} d\text{Vol}_g\right)^{2/2^*}}, \quad (1.13)$$

where Scal_g is the scalar curvature of g and Vol_g is the (non-renormalized) volume measure. $Y(M, g)$ is called Yamabe constant of (M, g) and it is a conformal invariant. Note that in the case of \mathbb{S}^n , the minimizers of $\mathcal{E}(u)$ are precisely the spherical bubbles in (1.2).

Corollary 1.3. *For every $n > 2$ and $\varepsilon > 0$ there exists $\delta := \delta(\varepsilon, n) > 0$ such that the following holds. Let (M, g) be an n -dimensional Riemannian manifold with $\text{Ric}_g \geq (n-1)g$ and $u \in W^{1,2}(M)$ non-zero such that*

$$d_{GH}(M, \mathbb{S}^n) \leq \delta, \quad |\mathcal{E}(u) - Y(M, g)| \leq \delta. \quad (1.14)$$

Then, there exist $a \in \mathbb{R}, b \in (0, 1)$ and $z_0 \in M$ satisfying

$$\frac{\|u - w_{a,b,z}\|_{W^{1,2}}}{\|u\|_{W^{1,2}}} \leq \varepsilon,$$

where $w_{a,b,z}$ is as in (1.7).

Here d_{GH} denotes the Gromov-Hausdorff distance. A similar stability for almost minimizers of $\mathcal{E}(\cdot)$ has been recently proved in [45] in quantitative form and under no assumptions on the metric. The novelty here is that we have a comparison with an explicit class of functions, while in [45] no information is known about the shape of the minimizers.

We discuss now a second stability result on non-compact Riemannian manifolds. Our motivations come from the fact that, to prove Theorem 1.1, non-compact setting will naturally arise in our investigation (see below the main strategy of proof).

Let us consider an n -dimensional Riemannian manifolds (M, g) , $n > 2$, satisfying

$$\text{Ric}_g \geq 0, \quad \text{AVR}(M) := \lim_{R \rightarrow \infty} \frac{\text{Vol}(B_R(x))}{\omega_n R^n} > 0, \quad (1.15)$$

for $x \in M$. The latter condition is called *Euclidean volume growth* property and $\text{AVR}(M)$ is the asymptotic volume ratio. Notice that the limit exists and is independent of x , by the Bishop-Gromov inequality.

In [22], the following sharp Euclidean-type Sobolev inequality was derived under the assumptions (1.15):

$$\|u\|_{L^{2^*}} \leq \text{AVR}(M)^{-\frac{1}{n}} \text{Eucl}(n, 2) \|\nabla u\|_{L^2}, \quad \forall u \in \dot{W}^{1,2}(M). \quad (1.16)$$

Moreover, they proved that equality occurs in (1.16) for some non-zero function $u \in \dot{W}^{1,2}(M)$, then M is isometric to \mathbb{R}^n and u is in particular an Euclidean bubble. Actually

in [22] this rigidity requires also $u \in C^n(M)$ and $u \geq 0$, however these additional assumptions can be removed after the results in [13] and [33] (see also Theorem 5.3).

The natural stability question is what happens if a function satisfies almost equality in (1.16). Clearly, differently from (1.6), we cannot deduce anything about the geometry of M . Indeed the inequality is sharp on every M as in (1.15), which means that we can always find functions so that $\frac{\|u\|_{L^{2^*}}}{\|\nabla u\|_{L^2}}$ is arbitrary close to $\text{AVR}(M)^{-\frac{1}{n}} \text{Eucl}(n, 2)$. We can prove however that a function for which almost equality occurs in (1.16) is close to a Euclidean bubble. Set

$$v_{a,b,z} := \frac{a}{(1 + b d(\cdot, z)^2)^{\frac{n-2}{2}}}, \quad \text{for } a \in \mathbb{R}, b > 0, z \in M.$$

Theorem 1.4. *For every $\varepsilon > 0, V \in (0, 1)$ and $n > 2$, there exists $\delta := \delta(\varepsilon, n, V) > 0$ such that the following holds. Let (M, g) be an n -dimensional Riemannian manifold as in (1.15) with $\text{AVR}(M) \geq V$ and assume there exists $u \in \dot{W}^{1,2}(M)$ non-zero satisfying*

$$\frac{\|u\|_{L^{2^*}}}{\|\nabla u\|_{L^2}} > \text{AVR}(M)^{-\frac{1}{n}} \text{Eucl}(n, 2) - \delta.$$

Then, there exist $a \in \mathbb{R}, b > 0$, and $z \in M$ so that

$$\frac{\|\nabla(u - v_{a,b,z})\|_{L^2}}{\|\nabla u\|_{L^2}} \leq \varepsilon.$$

Notice that the stability is strong in the sense that we control the gradient norm as in the Euclidean case (1.5).

A direct consequence of the above theorem is:

Corollary 1.5. *Let (M, g) be an n -dimensional Riemannian manifold as in (1.15). Then*

$$\text{AVR}(M)^{\frac{1}{n}} \text{Eucl}^{-1}(n, 2) = \inf_{a \in \mathbb{R}, b > 0, z \in M} \frac{\|\nabla v_{a,b,z}\|_{L^2}}{\|v_{a,b,z}\|_{L^{2^*}}}.$$

Remark 1.6. Our main results in Theorem 1.1 and Theorem 1.4, even if stated on smooth Riemannian manifolds, actually hold also in the context of weighted Riemannian manifolds and more generally in the singular setting of metric measure spaces with a synthetic Ricci curvature lower bound. The generalized version of these statements can be found in Theorem 8.1 and Theorem 8.4. ■

Strategy of proof and non-smooth setting. We outline the argument for Theorem 1.1 (Theorem 1.4 is simpler and follows by the same strategy). The underlying idea is classical, that is to argue by contradiction and concentration compactness. However, the novelty is that the space is not homogeneous and also not fixed, since we need to deal with a whole class of Riemannian manifolds. Moreover, singular and non-compact limit

spaces must also be considered. In particular, the whole analysis will be carried out in the more general setting of RCD spaces, which are metric measure spaces with a synthetic notion of Ricci curvature bounded below (see Section 2 for details and references).

Suppose that Theorem 1.1 is false. Then, there exist $\varepsilon > 0$, a sequence $\{M_k\}_{k \in \mathbb{N}}$ of n -dimensional Riemannian manifolds with $\text{Ric}_k \geq n - 1$ and non-constant functions $u_k : M_k \rightarrow \mathbb{R}$, $\|u_k\|_{L^{2^*}} = 1$, which satisfy (1.8) for some $\delta_k \downarrow 0$, but so that for any $k \in \mathbb{N}$

$$\inf \|u_k - w\|_{L^{2^*}} + \|\nabla(u_k - w)\|_{L^2} > \varepsilon, \tag{1.17}$$

where the inf runs among all spherical bubbles $w = a(1 - b \cos(\mathbf{d}_k(\cdot, z)))^{\frac{2-n}{2}}$ (\mathbf{d}_k being the distance on M_k). Similarly to the classical concentration compactness [80,81] in \mathbb{R}^n , we choose points $y_k \in M_k$ and constants $\sigma_k > 0$ so that, defining

$$(Y_k, \rho_k, \mu_k) := (M_k, \sigma_k \mathbf{d}_k, \text{Vol}_k(M_k)^{-1} \sigma_k^n \text{Vol}_k), \quad u_{\sigma_k} = \sigma_k^{-n/2^*} u_k, \tag{1.18}$$

we have

$$\int_{B_1^{Y_k}(y_k)} |u_{\sigma_k}|^{2^*} d\mu_k = \frac{1}{2},$$

(in the actual proof we choose a suitable constant close to 1). The spaces (Y_k, ρ_k, μ_k) are in particular metric measure spaces which are rescalings of the original manifolds M_k . Note that it can happen that $\sigma_k \uparrow \infty$, which corresponds to a concentrating behavior of the sequence u_k . In this case, the diameter of Y_k goes to infinity and we are in a sense performing a blow-up along M_k .

Thanks to Gromov’s precompactness theorem [64] it is possible to show that, up to a subsequence, $(Y_k, \rho_k, \mu_k, y_k)$ converges in the pointed-measure-Gromov-Hausdorff sense to a limit RCD space (Y, ρ, μ, \bar{y}) (which might be non-smooth). Using a generalized version of Lions’ concentration compactness for a sequence of RCD spaces (see Section 6), we show that up to a further subsequence, u_{σ_k} converges L^{2^*} -strongly (on varying spaces, see Definition 2.9 below) to some $u \in L^{2^*}(\mu)$. It also follows that u is extremal for a ‘limit Sobolev inequality’ on Y , that might be both as in (1.6) or of Euclidean-type as in (1.16), depending if there is concentration or not along the original sequence u_k . The key point is proving:

- Concentration $\Rightarrow Y$ is a metric-cone and u is a Euclidean bubble
- Non-concentration $\Rightarrow Y$ is a spherical suspension and u is a spherical bubble

We will show these two facts by proving suitable rigidity theorems for the Sobolev inequalities on RCD spaces (see Section 5). The proof will be then completed by carefully bringing back this information from u to the sequence u_k to find a contradiction with (1.17). It is worth noticing that, in case of concentration, the scaled functions u_{σ_k} tend

to a Euclidean bubble but, to reach a contradiction, the original sequence u_k must be close to the family of spherical bubbles. This turns out to be true because a concentrated spherical bubble looks locally, around the point where it is concentrated, like a Euclidean bubble (see Lemma 7.3).

We conclude this introduction by mentioning that generalized concentration compactness techniques on varying spaces, in a similar spirit to the present work, have been recently developed in [11,12] and applied to study the problem of existence of isoperimetric regions on non-compact Riemannian manifolds [10].

2. Preliminaries

2.1. Calculus on metric measure spaces

A metric measure space is a triple (X, d, \mathbf{m}) , where (X, d) is a complete and separable metric space and $\mathbf{m} \neq 0$ is a non-negative and boundedly finite Borel measure. Two metric measure spaces are *isomorphic*, provided there exists a measure preserving isometry between them. To avoid technicalities, we will always assume $\text{supp}(\mathbf{m}) = X$. We will denote by $\text{LIP}(X)$ and $\text{LIP}_{bs}(X)$ respectively the space of Lipschitz functions and Lipschitz functions with bounded support in (X, d) . We recall the notion of local Lipschitz constant of a Lipschitz function $f \in \text{LIP}(X)$:

$$\text{lip } f(x) := \overline{\lim}_{y \rightarrow x} \frac{|f(y) - f(x)|}{d(x, y)},$$

set to $+\infty$ if x is isolated. The Sobolev space on a metric measure space was introduced in [39] and [93] (inspired by the notion of upper gradient [69,70]). Here we follow the axiomatization of [5] (equivalent to that of [93,39]).

Let (X, d, \mathbf{m}) be a metric measure space and define the Cheeger energy $\text{Ch}: L^2(\mathbf{m}) \rightarrow [0, \infty]$

$$\text{Ch}(f) := \inf \left\{ \varliminf_{n \rightarrow \infty} \int \text{lip}^2 f_n \, d\mathbf{m} : (f_n) \subset L^2(\mathbf{m}) \cap \text{LIP}(X), f_n \rightarrow f \text{ in } L^2(\mathbf{m}) \right\}.$$

The Sobolev space is defined as $W^{1,2}(X) := \{f \in L^2(\mathbf{m}) : \text{Ch}(f) < \infty\}$ and equipped with the norm $\|f\|_{W^{1,2}(X)}^2 := \|f\|_{L^2(\mathbf{m})}^2 + \text{Ch}(f)$ turning it into a Banach space. We recall also (see e.g. [5]) that for every $f \in W^{1,2}(X)$ there exists a minimal \mathbf{m} -a.e. object $|\nabla f| \in L^2(\mathbf{m})$ called minimal weak upper gradient so that

$$\text{Ch}(f) = \int |\nabla f|^2 \, d\mathbf{m}.$$

To lighten the notation, we will often write $\|\nabla f\|_{L^2(\mathbf{m})}$ in place of $\| |\nabla f| \|_{L^2(\mathbf{m})}$. We shall often use the *locality* of minimal weak upper gradients:

$$|\nabla f| = |\nabla g|, \quad \mathbf{m}\text{-a.e. in } \{f = g\},$$

for every $f, g \in W^{1,2}(X)$. For $\Omega \subset X$ open we say that $f \in W_{loc}^{1,2}(\Omega)$, provided $\eta f \in W^{1,2}(X)$ for every $\eta \in \text{LIP}_{bs}(X)$ with $d(\text{supp}(\eta), X \setminus \Omega) > 0$. By locality, the object

$$|\nabla f| := |\nabla(\eta f)|, \quad \mathbf{m}\text{-a.e. on } \{\eta = 1\},$$

is well defined as an $L^2_{loc}(\Omega)$ -function and will be called again minimal weak upper gradient. It can be easily checked that if $f \in W_{loc}^{1,2}(X)$ with $f, |\nabla f| \in L^2(\mathbf{m})$, then $f \in W^{1,2}(X)$.

We shall need also the following semicontinuity result:

$$\begin{aligned} f_n \in W_{loc}^{1,2}(X), f_n \rightarrow f \text{ m-a.e.} &\Rightarrow f \in W_{loc}^{1,2}(X), |\nabla f| \in L^2(\mathbf{m}) \\ \underline{\lim}_n \|\nabla f_n\|_{L^2(\mathbf{m})} < \infty &\Rightarrow \|\nabla f\|_{L^2(\mathbf{m})} \leq \underline{\lim}_n \|\nabla f_n\|_{L^2(\mathbf{m})} \end{aligned} \tag{2.1}$$

The $W_{loc}^{1,2}$ regularity can be directly proved by appealing to the semicontinuity (see, e.g., [59, Prop 2.1.13]) in the space $W^{1,2}(X)$ and a cut-off argument. The fact that $|\nabla f| \in L^2(\mathbf{m})$ follows by noticing that, for any ball $B \subset X$, $\int_B |\nabla f|^2 \, d\mathbf{m} \leq \int_B |\nabla(\eta f)|^2 \, d\mathbf{m} \leq \underline{\lim}_n \|\nabla f_n\|_{L^2(\mathbf{m})}^2$, where $\eta \in \text{LIP}_c(X)^+$ with $\eta \equiv 1$ on B , having used twice the locality of the minimal weak upper gradient and again [59, Prop 2.1.13]. This proves (2.1) by arbitrariness of B .

For $\Omega \subseteq X$ open, we define the Sobolev space of functions vanishing at the boundary $W_0^{1,2}(\Omega) \subset W^{1,2}(X)$ as the closure of $\text{LIP}_c(\Omega)$ with respect to the $W^{1,2}$ norm.

A metric measure space is called infinitesimally Hilbertian [54] provided

$$|\nabla(f + g)|^2 + |\nabla(f - g)|^2 = 2|\nabla f|^2 + 2|\nabla g|^2, \quad \mathbf{m}\text{-a.e., } \forall f, g \in W^{1,2}(X),$$

or equivalently if $W^{1,2}(X)$ is Hilbert. This allows defining a formal scalar product between gradients of Sobolev functions by polarization

$$\langle \nabla f, \nabla g \rangle := |\nabla f|^2 + |\nabla g|^2 - |\nabla(f - g)|^2 \in L^1(\mathbf{m}), \quad \forall f, g \in W^{1,2}(X), \tag{2.2}$$

that is bilinear on its entries. By locality, it is possible to consider also a scalar product for functions in $W_{loc}^{1,2}(\Omega)$.

We recall next the *measure-valued Laplacian* as in [54], in the case of X proper and infinitesimally Hilbertian. We say that $f \in W_{loc}^{1,2}(\Omega)$ has a measure-valued Laplacian on Ω , and we write $f \in D(\Delta, \Omega)$, provided there exists a (signed) Radon measure μ such that

$$\int g \, d\mu = - \int \langle \nabla f, \nabla g \rangle \, d\mathbf{m}, \quad \forall g \in \text{LIP}_c(\Omega).$$

Here signed Radon measure means difference of two positive Radon measures (see also [37] for a related discussion). The unique measure μ satisfying the above is denoted

by Δf and depends linearly on f . If $\Omega = X$ we simply write $f \in D(\Delta)$. Moreover, if $\Delta f \ll \mathfrak{m}$, we write $\Delta f := \frac{d\Delta f}{d\mathfrak{m}} \in L^1_{loc}(\Omega)$.

Next, we introduce the sets of finite perimeter following [3,84]. For $E \subset X$ Borel and $A \subset X$ open, define

$$\text{Per}(E, A) := \inf \left\{ \liminf_{n \rightarrow \infty} \int_A \text{lip } f_n \, d\mathfrak{m} : f_n \subset \text{LIP}_{loc}(A), f_n \rightarrow \chi_E \text{ in } L^1_{loc}(A) \right\}.$$

If $\text{Per}(E, X) < \infty$ we say that E has finite perimeter. In this case, the map $A \mapsto \text{Per}(E, A)$ is the restriction to open sets of a non-negative finite Borel measure called the perimeter measure of E (see [3] and also [84]). As a convention, when $A = X$ we simply write $\text{Per}(E)$ instead of $\text{Per}(E, X)$.

2.2. RCD-spaces

In this note, we shall work with spaces that encode Ricci lower bounds in a synthetic sense as introduced first and independently in [82] and [95,96]. For $K \in \mathbb{R}, N \in [1, \infty)$, the Curvature Dimension condition $\text{CD}(K, N)$ for a metric measure space is a weak notion of Ricci curvature bounded below by K and dimension bounded above by N . We will actually consider here the subclass of spaces satisfying the so-called Riemannian Curvature Dimension condition. The RCD-condition has been defined first in the infinite dimensional setting [6] and later in [54] in finite dimension. We also recall [18,7,4,9,46,35] for key contributions on this theory and for the study of the equivalence of different definitions and approaches. We refer to [2] for more details and references.

Definition 2.1. A metric measure space (X, d, \mathfrak{m}) satisfies the $\text{RCD}(K, N)$ condition for some $K \in \mathbb{R}, N \in (1, \infty)$, if it is infinitesimally Hilbertian and satisfies the $\text{CD}(K, N)$ condition.

To keep the exposition shorter will not recall the definition of the $\text{CD}(K, N)$ condition and instead focus on recalling the key properties of RCD spaces used in this note.

We start recalling that $\text{RCD}(K, N)$ spaces satisfy the Bishop-Gromov inequality [95, 96]:

$$\frac{\mathfrak{m}(B_R(x))}{v_{K,N}(R)} \leq \frac{\mathfrak{m}(B_r(x))}{v_{K,N}(r)}, \quad \text{for any } 0 < r < R \leq \pi \sqrt{\frac{N-1}{K^+}} \text{ and } x \in X, \tag{2.3}$$

where K^+ is the positive part of K and $v_{K,N}(r)$ is the volume of a ball of radius r in the (K, N) -model space, see [95,96] for the precise definition. We only recall the particular case $v_{0,N}(r) = \omega_N r^N$. In particular $\text{RCD}(K, N)$ spaces are uniformly locally doubling and, since they support a weak local Poincaré inequality [91], by the work [39] we have:

$$|\nabla f| = \text{lip } f, \quad \mathfrak{m}\text{-a.e., } \forall f \in \text{Lip}_{bs}(X). \tag{2.4}$$

Since $\text{RCD}(K, N)$ spaces are geodesic and uniformly locally doubling, they admit a reverse doubling inequality. We omit the standard argument (see e.g. [63, Prop. 3.3]).

Lemma 2.2. *Let (X, d, m) be an $\text{RCD}(K, N)$ space for some $N \in (1, \infty), K \in \mathbb{R}$. Then there exists $\gamma = \gamma(N) > 0$ and $R_{K^-, N} > 0$ (with $R_{0, N} = +\infty$) such that for every ball $B_R(x) \subsetneq X$ with $R \leq R_{K^-, N}$, it holds*

$$\frac{m(B_r(x))}{m(B_R(x))} \leq \left(\frac{r}{R}\right)^\gamma, \quad \forall r \in (0, R/2). \tag{2.5}$$

We recall also the following version of the *coarea formula* from [84, Proposition 4.2] adapted to RCD -setting after [57].

Theorem 2.3 (Coarea formula). *Let (X, d, m) be an $\text{RCD}(K, N)$ space, $N < +\infty, \Omega \subset X$ open and $f \in \text{LIP}_{loc}(\Omega)$. Then given any Borel function $g : X \rightarrow [0, \infty)$, it holds that*

$$\int_{\{s < f < t\}} g |\nabla f| dm = \iint_s^t g \, d\text{Per}(\{f > r\}, \cdot) \, dr, \quad \forall s, t \in [0, \infty), s < t, \{f > s\} \subset\subset \Omega. \tag{2.6}$$

Proof. Fix s, t as in (2.6) and $U \subset\subset \Omega$ open and containing $\{f > s\}$. We can suppose that $s > 0$. Let $\eta \in \text{LIP}_c(\Omega)$ with $\eta = 1$ in $U, 0 \leq \eta \leq 1$ and set $\tilde{f} := \eta f \in \text{LIP}_c(X)$. Then by [84, Remark 4.3] and the results in [57] about the identification of total variation and minimal weak upper gradient, (2.6) holds for s, t , any g and with \tilde{f} in place of f . To pass to f simply use the locality of the weak upper gradient and note that by construction $\{\tilde{f} > r\} = \{f > r\}$ for every $r > s$. \square

We also report a regularity result from [73].

Theorem 2.4. *Let (X, d, m) be an $\text{RCD}(K, N)$ space for some $K \in \mathbb{R}, N \in (1, \infty)$ and let $u \in D(\Delta)$ with $\Delta u = gu$ for some $g \in L^\infty(m), \|g\|_{L^\infty(m)} \leq M$. Then for every $x_0 \in X$ and every $R > 0$ it holds*

$$\| |\nabla u| \|_{L^\infty(B_R(x_0))} \leq C(K, N, R, M) \int_{B_{2R}(x_0)} |u| dm.$$

In particular $u \in \text{LIP}_{loc}(X)$.

We say that an $\text{RCD}(0, N)$ space (X, d, m) has Euclidean volume growth, if

$$\text{AVR}(X) := \lim_{R \rightarrow \infty} \frac{m(B_R(x))}{\omega_N R^N} > 0, \tag{2.7}$$

for one (and thus, any) $x \in X$. In this setting, a sharp isoperimetric inequality was proved in [22] (previous versions in the smooth-setting already appeared in [28,1,49,74]). A slightly weaker inequality holds also in the MCP setting ([34]).

Theorem 2.5. *Let (X, d, \mathbf{m}) be an $\text{RCD}(0, N)$ space with $N \in (1, \infty)$, $\text{AVR}(X) > 0$. Then*

$$\text{Per}(E) \geq N(\text{AVR}(X)\omega_N)^{1/N} \mathbf{m}(E)^{\frac{N-1}{N}}, \quad \forall E \subset X \text{ Borel, } \mathbf{m}(E) < +\infty. \quad (2.8)$$

Here $\omega_N := \pi^{N/2}\Gamma^{-1}(N/2 + 1)$, where $\Gamma(\cdot)$ is the Gamma-function. We shall need also the rigidity of (2.8) in the RCD-setting. This has been proved in [13] under the noncollapsed assumption, which was recently removed (with a different argument) in [33].

Theorem 2.6. *Let (X, d, \mathbf{m}) be an $\text{RCD}(0, N)$ space with $N \in (1, \infty)$, $\text{AVR}(X) > 0$. Equality holds in (2.8) for some $E \subset X$ Borel with $\mathbf{m}(E) < +\infty$ if and only if X is a N -Euclidean metric measure cone and E is (up to \mathbf{m} -negligible sets) a metric ball centered at one of the tips of X .*

Theorem 2.6 is stated in [33] with the extra assumption that E is bounded, however this assumption can be dropped thanks to the recent [14].

Recall that for $N \in [1, \infty)$, the N -Euclidean cone over a metric measure space (Z, \mathbf{m}_Z, d_Z) is defined to be the space $Z \times [0, \infty)/(Z \times \{0\})$ endowed with the following distance and measure

$$\begin{aligned} d((t, z), (s, z')) &:= \sqrt{t^2 + s^2 - 2st \cos(d_Z(z, z') \wedge \pi)}, \\ \mathbf{m} &:= t^{N-1} dt \otimes \mathbf{m}_Z. \end{aligned}$$

The point $Z \times \{0\}$ is called tip of the cone.

2.3. Sobolev inequalities

We next report the main Sobolev inequalities of this note starting in the compact setting. On an $\text{RCD}(N - 1, N)$ space (X, d, \mathbf{m}) for some $N \in (2, \infty)$ with $\mathbf{m}(X) = 1$, we recall the following Sobolev inequality ([90,36])

$$\|u\|_{L^{2^*}(\mathbf{m})}^2 \leq \frac{2^* - 2}{N} \|\nabla u\|_{L^2(\mathbf{m})}^2 + \|u\|_{L^2(\mathbf{m})}^2, \quad \forall u \in W^{1,2}(X), \quad (2.9)$$

where $2^* = 2N/(N - 2)$.

Moving to the non-compact setting, we start recalling a classical one-dimensional inequality by Bliss [27] (see also [17,97,40]). To state it we introduce some notations. For all $N \in (2, \infty)$, we define $\sigma_{N-1} := N\omega_N$ and recall the sharp Euclidean Sobolev constant

$$\text{Eucl}(N, 2) := \left(\frac{4}{N(N-2)\sigma_N^{2/N}} \right)^{\frac{1}{2}}. \tag{2.10}$$

Lemma 2.7 (*Bliss inequality*). *Let $u: [0, \infty) \rightarrow \mathbb{R}$ be locally absolutely continuous, $N \in (2, \infty)$ and define $2^* := 2N/(N-2)$. Then*

$$\left(\sigma_{N-1} \int_0^\infty |u|^{2^*}(t) t^{N-1} dt \right)^{\frac{1}{2^*}} \leq \text{Eucl}(N, 2) \left(\sigma_{N-1} \int_0^\infty |u'|^2(t) t^{N-1} dt \right)^{\frac{1}{2}}, \tag{2.11}$$

whenever one side is finite. Moreover, equality holds if and only if u is of the type:

$$v_{a,b}(r) := a(1 + br^2)^{\frac{2-N}{2}}, \quad a \in \mathbb{R}, b > 0. \tag{2.12}$$

We recall the sharp Sobolev Euclidean-type inequality [88] (first appeared in [22] for manifolds).

Theorem 2.8. *Let (X, d, m) be an $\text{RCD}(0, N)$ space, $N \in (2, \infty)$, with Euclidean volume growth. Then, for every $u \in W_{loc}^{1,2}(X)$ with $m(\{|u| > t\}) < +\infty$ for all $t > 0$, it holds*

$$\|u\|_{L^{2^*}(m)} \leq \text{Eucl}(N, 2) \text{AVR}(X)^{-\frac{1}{N}} \|\nabla u\|_{L^2(m)}. \tag{2.13}$$

Moreover, (2.13) is sharp.

Proof. Combine [88, Theorem 1.13] and Lemma B.1. \square

For convenience in the rest of this note, we adopt the following notation.

Convention: We say that an $\text{RCD}(K, N)$ space (X, d, m) , with $N \in (2, \infty)$, supports a Sobolev inequality with constants $A > 0, B \geq 0$, if, setting $2^* := 2N/(N-2)$,

$$\|u\|_{L^{2^*}(m)}^2 \leq A \|\nabla u\|_{L^2(m)}^2 + B \|u\|_{L^2(m)}^2, \quad \forall u \in W^{1,2}(X). \tag{S}$$

Inequality (S), if true, actually holds for all $u \in W_{loc}^{1,2}(X)$ satisfying $m(\{|u| > t\}) < +\infty$ for all $t > 0$ (recall Lemma B.1).

2.4. Convergence and stability under pmGH-convergence

We start recalling the notion of *pointed-measure Gromov Hausdorff convergence* (pmGH convergence for short) following [58]. This presentation is not standard (see e.g. [32,64]), but it is equivalent in the case of a sequence of uniformly locally doubling metric measure spaces ([58]).

Set $\bar{\mathbb{N}} := \mathbb{N} \cup \{\infty\}$ and consider a sequence of pointed metric measure spaces (X_n, d_n, m_n, x_n) , with $x_n \in X_n$. We say that X_n pmGH-converge to X_∞ if there exist isometric embeddings $\iota_n : X_n \rightarrow (Z, d)$, $n \in \bar{\mathbb{N}}$, into a common metric space (Z, d) such that

$(\iota_n)_\# \mathbf{m}_n \rightarrow (\iota_\infty)_\# \mathbf{m}_\infty$ in duality with $C_{bs}(Z)$ and $\iota_n(x_n) \rightarrow \iota_\infty(x_\infty)$ in Z .

In the case of a sequence of uniformly locally doubling spaces (as in the case of $RCD(K, N)$ -spaces for fixed $K \in \mathbb{R}, N < \infty$) we can also take (Z, d) to be proper.

It will be also convenient to adopt the so-called *extrinsic approach* and identify X_n with their isomorphic copies in (Z, d) . This allows writing $\mathbf{m}_n \rightarrow \mathbf{m}_\infty$ in duality with $C_{bs}(Z)$. A choice of space (Z, d) together with isomorphic copies of the spaces X_n will be often called a *realization of the convergence*.

For the scope of this note, it is important to recall the notion of convergence of functions along pmGH-convergence [71,58,8] and their properties. We fix in what follows a pmGH-convergent sequence of pointed metric measure spaces as discussed above.

Definition 2.9. Let $p \in (1, \infty)$ and fix a realization of the convergence in (Z, d) . We say:

- i) $f_n \in L^p(\mathbf{m}_n)$ converges L^p -weak to $f_\infty \in L^p(\mathbf{m}_\infty)$, provided $\sup_{n \in \mathbb{N}} \|f_n\|_{L^p(\mathbf{m}_n)} < \infty$ and $f_n \mathbf{m}_n \rightarrow f_\infty \mathbf{m}_\infty$ in $C_{bs}(Z)$;
- ii) $f_n \in L^p(\mathbf{m}_n)$ converges L^p -strong to $f_\infty \in L^p(\mathbf{m}_\infty)$, provided it converges L^p -weak and $\overline{\lim}_n \|f_n\|_{L^p(\mathbf{m}_n)} \leq \|f_\infty\|_{L^p(\mathbf{m}_\infty)}$;
- iii) $f_n \in W^{1,2}(X_n)$ converges $W^{1,2}$ -weak to $f_\infty \in W^{1,2}(X)$ provided it converges L^2 -weak and $\sup_{n \in \mathbb{N}} \|\nabla f_n\|_{L^2(\mathbf{m}_n)} < \infty$;
- iv) $f_n \in W^{1,2}(X_n)$ converges $W^{1,2}$ -strong to $f_\infty \in W^{1,2}(X)$ provided it converges L^2 -strong and $\|\nabla f_n\|_{L^2(\mathbf{m}_n)} \rightarrow \|\nabla f_\infty\|_{L^2(\mathbf{m}_\infty)}$;
- v) $f_n \in L^p(\mathbf{m}_n)$ converges L^p_{loc} -strong to $f_\infty \in L^p(\mathbf{m}_\infty)$, provided ηf_n converges L^p -strong to ηf_∞ for every $\eta \in C_{bs}(Z)$.

Recall from [71,58,8] the linearity of convergence: if f_n, g_n converge L^p -strong to f_∞, g_∞ , respectively, then

$$f_n + g_n \text{ converges } L^p\text{-strong to } f_\infty + g_\infty. \tag{2.14}$$

We point out the following simple fact: for any $p \in (1, \infty)$ it holds

$$f_n \text{ } L^p\text{-weak converges to } f_\infty \quad \Rightarrow \quad \|f_\infty\|_{L^2(\mathbf{m}_\infty)} \leq \varliminf_{n \rightarrow \infty} \|f_n\|_{L^2(\mathbf{m}_n)}. \tag{2.15}$$

Indeed, if the above liminf above is $+\infty$, then there is nothing to prove. So let us assume it to be finite and also to be a limit, hence f_n is L^2 -bounded. Then there exists an L^2 -weak convergent subsequence (see [58]) to some $h \in L^2(\mathbf{m}_\infty)$ and in particular $\|h\|_{L^2(\mathbf{m}_\infty)} \leq \varliminf_n \|f_n\|_{L^2(\mathbf{m}_n)}$. By uniqueness of limits we have $h = f_\infty$, which shows (2.15).

After the works in [95,96,82,52,6,58] and thanks to Gromov’s precompactness theorem [64] we have the following precompactness result.

Theorem 2.10. *Let $(X_n, d_n, \mathbf{m}_n, x_n)$ be a sequence of pointed $RCD(K_n, N_n)$ spaces, $n \in \mathbb{N}$, with $\mathbf{m}_n(B_1(x_n)) \in [v^{-1}, v]$, for $v > 1$ and $K_n \rightarrow K \in \mathbb{R}, N_n \rightarrow N \in [1, \infty)$.*

Then, there exists a subsequence $(X_{n_k}, \mathbf{d}_{n_k}, \mathbf{m}_{n_k}, x_{n_k})$ pmGH-converging to a pointed RCD(K, N) space $(X_\infty, \mathbf{d}_\infty, \mathbf{m}_\infty, x_\infty)$.

We report from [58] the Mosco-convergence of the Cheeger energies for pmGH-converging RCD-spaces: if f_n is L^2 -weak convergent to f_∞ , then

$$\text{Ch}(f_\infty) \leq \varliminf_{n \rightarrow \infty} \text{Ch}(f_n). \tag{2.16}$$

Moreover, for any $f_\infty \in L^2(\mathbf{m}_\infty)$, there exists $f_n \in L^2(\mathbf{m}_n)$ converging L^2 -strong to f_∞ and

$$\overline{\lim}_{n \rightarrow \infty} \text{Ch}(f_n) \leq \text{Ch}(f_\infty).$$

In particular, the above is a limit.

3. Pólya-Szegő inequality

3.1. Non-compact case

In this part we extend to the non-compact case the Pólya-Szegő inequality of Euclidean-type obtained in [88].

We need first to recall basic notations and facts about monotone decreasing rearrangements for functions in a m.m.s. $(X, \mathbf{d}, \mathbf{m})$ (for more details we refer to [86]). Let $\Omega \subseteq X$ be an open set (possibly unbounded) and $u : \Omega \rightarrow [0, +\infty)$ be a Borel function such that $\mathbf{m}(\{u > t\}) < \infty$ for any $t > 0$. We define $\mu : [0, +\infty) \rightarrow [0, \infty)$, the distribution function of u as $\mu(t) := \mathbf{m}(\{u > t\})$. For u and μ as above, let us consider the generalized inverse $u^\#$ of μ :

$$u^\#(s) := \begin{cases} \text{ess sup } u & \text{if } s = 0, \\ \inf \{t : \mu(t) < s\} & \text{if } s > 0. \end{cases}$$

Note that $u^\#$ is non-increasing. In this note, we will perform rearrangements into the *Euclidean model space* $I_N := ([0, \infty), |\cdot|, \mathbf{m}_N)$, equipped with the standard Euclidean distance and weighted measure $\mathbf{m}_N := \sigma_{N-1} t^{N-1} \mathcal{L}^1$, for $N \in (1, \infty)$. For any open set $\Omega \subset X$ we set $\Omega^* := [0, r]$ with $\mathbf{m}_N([0, r]) = \mathbf{m}(\Omega)$ (i.e. $r^N = \omega_N^{-1} \mathbf{m}(\Omega)$), with the convention $\Omega^* = [0, \infty)$ if $\mathbf{m}(\Omega) = +\infty$. The Euclidean monotone rearrangement $u_N^* : \Omega^* \rightarrow \mathbb{R}^+$ is then defined by

$$u_N^*(x) := u^\#(\mathbf{m}_N([0, x])) = u^\#(\omega_N x^N), \quad \forall x \in \Omega^*.$$

Note that u_N^* is always a non-increasing function, since so is $u^\#$. To lighten the notation, we shall often drop the subscript and just write u^* . We collect basic facts about rear-

rangements, that can be proved by standard arguments as in the Euclidean case (see, e.g. [76]):

$$u \leq v \Rightarrow u^* \leq v^*, \tag{3.1}$$

$$(\varphi(u))^* = \varphi(u^*), \quad \forall \varphi : [0, \infty) \rightarrow [0, \infty) \text{ non-decreasing.} \tag{3.2}$$

$$\|u\|_{L^p(\mathbf{m})} = \|u^*\|_{L^p(\mathbf{m}_N)}, \quad \forall u \in L^p(\Omega). \tag{3.3}$$

Lemma 3.1. *Let (X, d, \mathbf{m}) be a metric measure space and $N \in (1, \infty)$. Let $(u_n) : X \rightarrow \mathbb{R}^+$ be an non-decreasing sequence of Borel functions. Denote $u := \sup_n u_n$ and suppose that $\mathbf{m}(\{u > t\}) < +\infty$ for every $t > 0$. Then, $u_n^* : I_N \rightarrow \mathbb{R}^+$ (which exists by the assumptions) is a monotone non-decreasing sequence and $\lim_n u_n^* = u^*$ a.e. in $[0, \infty)$.*

Proof. The fact that (u_n^*) is monotone non-decreasing follows by the order preserving property of the rearrangement (3.1). Set $g := \sup_n u_n^* = \lim_n u_n^*$ pointwise on $[0, \infty)$. In particular $\{u_n^* > t\} \uparrow \{g > t\}$ and $\{u_n > t\} \uparrow \{u > t\}$ for any $t > 0$. Therefore

$$\mathbf{m}_N(\{g > t\}) = \lim_n \mathbf{m}_N(\{u_n^* > t\}) = \lim_n \mathbf{m}(\{u_n > t\}) = \mathbf{m}(\{u > t\}) = \mathbf{m}_N(\{u^* > t\}).$$

So $g, u^* : [0, \infty) \rightarrow [0, +\infty]$ are equimeasurable and non-increasing (indeed g is the supremum of non-increasing functions), therefore they coincide a.e. (see e.g. the proof [76, Prop. 1.1.4]). \square

We will need the following approximation result to pass from the bounded to the unbounded case in the Euclidean Pólya-Szegő inequality. It will be needed also in other parts of this note.

Lemma 3.2. *Let (X, d, \mathbf{m}) be a metric measure space and $u \in W_{loc}^{1,2}(X)$ such that $\mathbf{m}(\{|u| > t\}) < +\infty$ for all $t > 0$ and $|\nabla u| \in L^2(\mathbf{m})$. Then there exists a sequence $u_n \in W^{1,2}(X)$ of functions with bounded support, such that $u_n \rightarrow u$ \mathbf{m} -a.e. and $|\nabla(u_n - u)| \rightarrow 0$ in $L^2(\mathbf{m})$.*

Moreover if $u \geq 0$ (resp. $u \in L^p(\mathbf{m})$, $p \in [1, \infty)$) we can take (u_n) non-decreasing (resp. so that $u_n \rightarrow u$ in $L^p(\mathbf{m})$).

Proof. We first deal with the case $u \geq 0$ and $u \in L^\infty(\mathbf{m})$ with $\mathbf{m}(\text{supp}(u)) < +\infty$. Fix $x \in X$ and consider the sequence $(\eta_n) \subset \text{LIP}(X)$ given by $\eta_n(\cdot) := (2 - \frac{d(\cdot, x)}{n})^+ \wedge 1$. Note that (η_n) is non-decreasing with $\text{LIP}(\eta_n) \leq n^{-1}$, $\eta_n = 1$ in $B_n(x)$ and $\text{supp}(\eta_n) \subset B_{2n}(x)$. Take $u_n := u\eta_n \in W^{1,2}(X)$ with bounded support. Clearly $u_n \uparrow u$ pointwise and if $u \in L^p(\mathbf{m})$ also $u_n \rightarrow u$ in $L^p(\mathbf{m})$ by dominated convergence. Moreover, by locality

$$\int |\nabla(u - u\eta_n)|^2 d\mathbf{m} \leq 2 \int_{B_n^c(x)} (|\nabla u|^2 + |\nabla(\eta_n u)|^2) d\mathbf{m},$$

and by the Leibniz rule

$$\begin{aligned} \|\nabla(\eta_n u)\|_{L^2(B_n(x)^c)} &\leq 2n^{-1}\|u\|_{L^\infty(\mathfrak{m})}\mathfrak{m}(\text{supp}(u))^{\frac{1}{2}} + \|\eta_n|\nabla u|\|_{L^2(B_n^c(x))} \\ &\leq 2n^{-1}\|u\|_{L^\infty(\mathfrak{m})}\mathfrak{m}(\text{supp}(u))^{\frac{1}{2}} + \|\nabla u\|_{L^2(B_n^c(x))} \rightarrow 0. \end{aligned}$$

This proves that $|\nabla u - \nabla(u_n \eta_n)| \rightarrow 0$ in $L^2(\mathfrak{m})$.

If $u \geq 0$, take $u_k := ((u - 1/k)^+) \wedge k$, $k \in \mathbb{N}$, which is a non-decreasing sequence of functions. Clearly

$$\int |\nabla(u - u_k)|^2 \mathfrak{m} \leq \int_{\{0 < u < 1/k\}} |\nabla u|^2 \mathfrak{m} \rightarrow 0,$$

by dominated convergence. Moreover, since $u_k \in L^\infty(\mathfrak{m})$ and $\mathfrak{m}(\text{supp}(u_k)) < +\infty$, the conclusion in this case follows from the previous one and a diagonal argument (multiplying by the functions η_n). Monotonicity of the sequence is preserved because $\eta_n f \leq \eta_{\bar{n}} g$ \mathfrak{m} -a.e. for every $\bar{n} > n$ and assuming $0 \leq f \leq g$ \mathfrak{m} -a.e. The pointwise \mathfrak{m} -a.e. convergence is also kept, since it remains true on every ball, recalling that $\eta_n = 1$ in $B_n(x)$.

Finally for a general u we approximate first u^+ and then u^- by functions u_n and v_n respectively as we did in the above steps. Clearly if $u \in L^p(\mathfrak{m})$ then $u_n - v_n \rightarrow u$ in $L^p(\mathfrak{m})$. Moreover by construction we have that $u_n - v_n = \chi_{\{u > 0\}} u_n - \chi_{\{u < 0\}} v_n$. Therefore $|\nabla(u - (u_n - v_n))| = |\nabla(u^+ - u_n)| + |\nabla(u^- - v_n)| \rightarrow 0$ in $L^2(\mathfrak{m})$. This concludes the proof also in this case. \square

We can now prove the Pólya-Szegő inequality in the non compact case.

Proposition 3.3. *Let (X, d, \mathfrak{m}) be an $\text{RCD}(0, N)$ space for some $N \in (1, \infty)$ with $\text{AVR}(X) > 0$. Let $u \in W_{loc}^{1,2}(X)$ be non-negative and such that $\mathfrak{m}(\{u > t\}) < \infty$ for any $t > 0$. Then,*

$$\int |\nabla u|^2 \mathfrak{m} \geq \text{AVR}(X)^{2/N} \int_0^\infty |\nabla u^*|^2 \mathfrak{m}_N, \tag{3.4}$$

meaning that, if the left hand side is finite, then $u^* \in W_{loc}^{1,2}(I_N)$ and (3.4) holds.

Proof. First, if $\|\nabla u\|_{L^2(\mathfrak{m})} = \infty$, there is nothing to prove. So, suppose $|\nabla u| \in L^2(\mathfrak{m})$. By Lemma 3.2 there exists a non-decreasing sequence $u_n \in W^{1,2}(X)$ of functions with bounded support, such $u_n \rightarrow u$ \mathfrak{m} -a.e. and $\|\nabla u_n\|_{L^2(\mathfrak{m})} \rightarrow \|\nabla u\|_{L^2(\mathfrak{m})}$. Applying the Pólya-Szegő inequality for bounded domains in [88, Theorem 3.6], we have $u_n^* \in W^{1,2}(I_N)$ and

$$\int |\nabla u_n|^2 \mathfrak{m} \geq \text{AVR}(X)^{2/N} \int |\nabla u_n^*|^2 \mathfrak{m}_N.$$

Moreover by Lemma 3.1 the sequence u_n^* is non-increasing and $\sup_n u_n^* = u^*$ pointwise. The proof is now concluded since we have that $u^* \in W_{loc}^{1,2}(I_N)$ and $\liminf_n \int |\nabla u_n^*|^2 \mathfrak{m}_N \geq \int |\nabla u^*|^2 \mathfrak{m}_N$ by semicontinuity (recall (2.1)). \square

3.2. Rigidity

In this section, we prove the rigidity in the Pólya-Szegő inequality of Proposition 3.3. The idea is that if equality in (3.4) is attained, the superlevel sets are isoperimetric sets, so Theorem 2.6 implies that the space is a cone. This line of thoughts follows classical arguments that date back to the work of [89] in Euclidean contexts and [23] for manifolds with Ricci curvature lower bounds.

Moreover, under additional regularity, the function can also be proven to be radial. A similar rigidity result was proved in [86] in the compact case for a different Pólya-Szegő inequality.

Theorem 3.4 (*Rigidity of the Euclidean Pólya-Szegő inequality*). *Let (X, d, m) be an $RCD(0, N)$ space for some $N \in (1, \infty)$ with $AVR(X) > 0$. Suppose equality holds in (3.4) (with both sides finite) for $u \in \text{LIP}_{loc}(X)$ non-negative satisfying $u(x) \rightarrow 0$ as $d(x, z) \rightarrow \infty$, for $z \in X$ and with $(u^*)' \neq 0$ a.e. in $\{u^* > 0\}$. Then, X is isomorphic to an N -Euclidean metric measure cone.*

Moreover, if $|\nabla u| \neq 0$ m-a.e. on $\{u > 0\}$, then u is radial, i.e.

$$u(x) = u^* \circ AVR(X)^{\frac{1}{N}} d(x, x_0)$$

for a suitable tip x_0 of X .

Proof. We divide the proof into different steps.

Step 1. We establish an improved version of (3.4) for a function u as in the statement. Fix such u . By Theorem 2.8 we know that $u \in L^{2^*}(\mathbf{m})$. For every $n \in \mathbb{N}$ set $v_n := (u - 1/n)^+$ and notice that they are supported in the open set $\Omega_n := \{u > 1/(2n)\}$, which is bounded. Therefore $v_n \in \text{LIP}_c(X)$. In particular by the Lipschitz-to-Lipschitz property of the rearrangement in the compact case (see [88, Prop. 3.4]) we have $v_n^* \in \text{LIP}_c([0, R_n])$ for suitable $R_n > 0$. From (3.2) we also have $v_n^* = (u^* - 1/n)^+$, which is non-increasing and $(v_n^*)' \neq 0$ a.e. in $\{v_n^* > 0\}$. In particular $u^* \in \text{LIP}_{loc}(0, \infty)$.

Define the functions $\varphi_n, \psi_n, \mu_n : [0, \sup v_n] \rightarrow [0, +\infty)$ as

$$\varphi_n(t) := \int_{\{v_n > t\}} |\nabla v_n|^2 \, dm, \quad \psi_n(t) := \int_{\{v_n > t\}} |\nabla v_n| \, dm, \quad \mu_n(t) := m(\{v_n > t\})$$

and analogously $\varphi, \psi, \mu : [0, \sup u] \rightarrow [0, +\infty]$ replacing everywhere v_n with u . Note that, thanks to the locality of the gradient, $\varphi(t) = \varphi_n(t - 1/n)$ for all $t \in (1/n, \infty)$ and the same holds for ψ and μ . We claim that

a) μ_n is absolutely continuous with

$$-\mu_n'(t) = \frac{\text{Per}(\{v_n^* > t\})}{|(v_n^*)'|((v_n^*)^{-1}(t))}, \quad \text{a.e. } t \in (0, \sup v_n). \tag{3.5}$$

If moreover $|\nabla u| \neq 0$ m-a.e. in $\{u > 0\}$ then also

$$-\mu'_n(t) = \int |\nabla v_n|^{-1} d\text{Per}(\{v_n > t\}) \quad \text{a.e. } t \in (0, \sup v_n); \tag{3.6}$$

b) φ_n, ψ_n are absolutely continuous with

$$\varphi'_n(t) = - \int |\nabla v_n| d\text{Per}(\{v_n > t\}), \quad \psi'_n(t) = -\text{Per}(\{v_n > t\}), \quad \text{for a.e. } t \in (0, \sup v_n). \tag{3.7}$$

Claim (3.5) in a) follows from [86, Lemma 3.10-3.11], since $\mu_n(t) = \mathbf{m}_N(\{v_n^* > t\})$ and $\text{Per}(\{v_n^* > t\})$ is concentrated on the point $(v_n^*)^{-1}(t)$. Claim b) is instead just a direct verification using the coarea formula (see (2.6)), since $v_n \in \text{LIP}_c(X)$. Under the assumption $|\nabla u| \neq 0$ m-a.e. in $\{u > 0\}$, by the Hölder inequality (using (3.6)) we have

$$-\varphi'_n(t) \geq -\psi'_n(t)^2(-\mu'_n(t))^{-1}, \tag{3.8}$$

at a.e. $t \in (0, \sup v_n)$ which is a differentiability point for μ_n, ψ_n, φ_n . If instead we only know that $(u^*)' \neq 0$ a.e. in $\{u^* > 0\}$, we can still deduce (3.8) applying first Hölder inequality and then differentiating (see the argument in [86, Prop. 3.12]). Integrating the above inequality, recalling that $\text{Per}(\{v_n^* > t\}) = N\omega_N^{\frac{1}{N}}\mu_n(t)^{\frac{N-1}{N}}$, we get for every $r, s \in [0, \sup v_n]$ with $s < r$:

$$\int_{\{s < v_n \leq r\}} |\nabla v_n|^2 dm \geq \int_s^r \left(\frac{\text{Per}(\{v_n > t\})}{N\omega_N^{\frac{1}{N}}\mu_n(t)^{\frac{N-1}{N}}} \right)^2 \int |\nabla v_n^*| d\text{Per}(\{v_n^* > t\}) dt. \tag{3.9}$$

Hence, the isoperimetric inequality (2.8) gives directly

$$\int_{\{s < v_n \leq r\}} |\nabla v_n|^2 dm \geq \text{AVR}(X)^{2/N} \int_{\{s < v_n^* \leq r\}} |\nabla v_n^*|^2 dm_N, \quad \forall 0 \leq s < r \leq \sup v_n, \tag{3.10}$$

having also used coarea formula for the function v_n^* since it is $\text{LIP}([0, R_n])$ as recalled before.

Since $v_n = (u - 1/n)^+$ and $v_n^* = (u^* - 1/n)^+$, from the locality of the gradient we can rewrite (3.10) (after a change of variable) as

$$\int_{\{s+1/n < u \leq r+1/n\}} |\nabla u|^2 dm \geq \text{AVR}(X)^{2/N} \int_{\{s+1/n < u^* \leq r+1/n\}} |\nabla u^*|^2 dm_N, \tag{3.11}$$

for every $s < r$ with $s, r \in (0, \sup u - 1/n]$. Taking the limit as $n \rightarrow +\infty$ we obtain

$$\int_{\{s < u \leq r\}} |\nabla u|^2 dm \geq AVR(X)^{2/N} \int_{\{s < u^* \leq r\}} |\nabla u^*|^2 dm_N, \quad \forall 0 \leq s < r \leq \sup u. \quad (3.12)$$

Step 2. We pass to the proof that X is a cone. We claim that if equality occurs in (3.10) for some $n \in \mathbb{N}$ and some $r, s \in [0, \sup v_n]$ with $r < s$, then

- i) $\text{Per}(\{v_n > t\}) = N(\omega_N AVR(X))^{1/N} \mu_n(t)^{\frac{N-1}{N}}$, for a.e. $t \in (s, r)$.
- ii) If $|\nabla u| \neq 0$ m-a.e. in $\{u > 0\}$, then $|\nabla v_n|$ is constant $\text{Per}(\{v_n > t\})$ -a.e. for a.e. $t \in (s, r)$.

Claim i) follows directly from the way we deduced (3.10) from (3.9) using the isoperimetric inequality (2.8). Claim ii) instead follows by the equality case in the Hölder inequality (3.8).

We now suppose, as in the hypotheses, that u attains equality in (3.4), which means that equality holds in (3.12) with $(s, r) = (0, \sup u)$. We claim that equality must hold in (3.12) also for all $s < r$ with $s, r \in (0, \sup u)$. Suppose it fails for some $s < r$. Then, calling $L(s', r')$ and $R(s', r')$ respectively the left and right hand sides of (3.12), we have

$$L(0, \sup u) = L(0, s) + L(s, r) + L(r, \sup u) > R(0, s) + R(s, r) + R(r, \sup u) \geq R(0, \sup u),$$

which contradicts the equality for $(0, \sup u)$. This proves the claim. Thus, equality holds in (3.11) for every $s < r$, with $s, r \in (0, \sup u - 1/n]$ which is equivalent to equality in (3.10) for every $s < r$ with $r, s \in [0, \sup v_n]$. Therefore i) holds and, provided $|\nabla u| \neq 0$ at m-a.e. point in $\{u > 0\}$, also ii) holds for every $s < r$ with $r, s \in [0, \sup v_n]$ and $n \in \mathbb{N}$. Putting these together and by arbitrariness of n , implies that

$$\text{Per}(\{u > t\}) = N(AVR(X)\omega_N)^{1/N} \mathbf{m}(\mu(t))^{\frac{N-1}{N}}, \quad \text{a.e. } t \in (0, \sup(u)), \quad (3.13)$$

and, if $|\nabla u| \neq 0$ m-a.e. in $\{u > 0\}$, we get

$$|\nabla u| \equiv c_t \quad \text{Per}(\{u > t\})\text{-a.e. for some constant } c_t \geq 0 \quad (3.14)$$

for a.e. $t \in (0, \sup u)$. Therefore, there exists t with $\mu(t) > 0$ so that equality occurs in (3.13), and recalling the rigidity in Theorem 2.6, we get that X is isomorphic to an N -Euclidean metric measure cone.

Step 3. Here we prove the functional rigidity of u , i.e. we prove that u is radial under the additional assumption: $|\nabla u| \neq 0$ m-a.e. on $\{u > 0\}$.

We first claim that (3.13) actually holds for every $t \in (0, \sup u)$. Let $t \in (0, \sup u)$ and consider a sequence $t_n \downarrow t$ for which (3.13) holds in every t_n . Then, by lower-semicontinuity of the perimeter (see, e.g., [84, Proposition 3.6]) and continuity of μ , we get

$$\text{Per}(\{u > t\}) \leq \liminf_{n \rightarrow \infty} \text{Per}(\{u > t_n\}) \stackrel{(3.13)}{=} N(AVR(X)\omega_N)^{1/N} \mu(t)^{\frac{N-1}{N}}.$$

Being the converse inequality always true (from (2.8)), the claim follows. Since $\{u > t\}$ are bounded (recall that u tends to zero at infinity), we can apply the rigidity Theorem 2.6 to deduce that for every $t \in (0, \sup u)$ there exists a radius $R_t > 0$ and $x_t \in X$ a tip for X (recall that X is a cone from Step 2) so that $\mathbf{m}(\{u > t\} \Delta B_{R_t}(x_t)) = 0$, where Δ denotes the symmetric difference. However $\{u > t\}$ is open. Thus

$$\{u > t\} = B_{R_t}(x_t). \tag{3.15}$$

We stress that the notation x_t is chosen because the cone structure may depend *a priori* on the isoperimetric superlevel set $\{u > t\}$. From here, the rest of the proof is devoted to show that x_t is in fact independent of t and u is radial. To do so we will follow the lines of the argument used in [86, Theorem 5.1], for the compact case.

Using (3.14) and (3.5) (recall that $\mu(t) = \mu_n(t - 1/n)$) we get

$$Nc_t^{-1}(\text{AVR}(X)\omega_N)^{\frac{1}{N}}\mu(t)^{\frac{N}{N-1}} = \int |\nabla u|^{-1} d\text{Per}(\{u > t\}) = -\mu'(t) = \frac{N\omega_N^{\frac{1}{N}}\mu(t)^{\frac{N}{N-1}}}{|(u^*)'((u^*)^{-1}(t))|},$$

for a.e. $t \in (0, \sup u)$. In particular,

$$|\nabla u| = \text{AVR}(X)^{\frac{1}{N}} |(u^*)'((u^*)^{-1}(t))| \quad \text{Per}(\{u > t\})\text{-a.e. and a.e. } t \in (0, \sup u). \tag{3.16}$$

Let $M := \|u\|_{L^\infty(\mathbf{m})} \in [0, +\infty)$. From the hypotheses u^* is non-negative, strictly decreasing and locally absolutely continuous (in fact locally Lipschitz) in $\{u^* > 0\} = [0, A)$ for some $A \in (0, +\infty]$ (in fact $A = \mathbf{m}(\{u > 0\})$). Hence it admits a strictly decreasing continuous inverse $(u^*)^{-1} : (0, M] \rightarrow [0, A)$, locally absolutely continuous in $(0, M)$. Since $(u^*)^{-1}(M) = 0$, we can extend it by zero in $[M, \infty)$ and call $H : (0, \infty) \rightarrow [0, A)$ this extension. In particular $H \in \text{AC}_{loc}(0, \infty)$. Observe that H might blow up at zero. Note also that, since u^* is locally Lipschitz in $(0, A)$, it preserves \mathcal{L}^1 -null sets. Hence pre-images of \mathcal{L}^1 -null subsets of $(0, M)$ via $H = (u^*)^{-1}$ are also \mathcal{L}^1 -null. Therefore for a.e. $t \in (0, A)$ the function u^* is differentiable at $(u^*)^{-1}(t)$, the function H is differentiable at t and

$$(u^*)'((u^*)^{-1}(t))H'(t) = (u^*((u^*)^{-1}(t)))' = 1. \tag{3.17}$$

To conclude the proof, we need to show that $f := \text{AVR}(X)^{-\frac{1}{N}} H \circ u : \{u > 0\} \rightarrow [0, \infty)$ satisfies

$$f(\cdot) = \mathbf{d}(x_0, \cdot), \tag{3.18}$$

for some point $x_0 \in \{u > 0\}$. Observe that f is continuous. We start proving that:

$$f \in \text{LIP}_{loc}(\{u > 0\}) \text{ and } |\nabla f| = 1 \text{ m-a.e. in } \{u > 0\}. \tag{3.19}$$

To show this we will use the chain rule in Lemma B.3 with $u, \Omega := \{u > 0\}, \varphi := H$ and $I := (0, \infty)$. To check the hypotheses we observe that by continuity $u(\Omega') \subset\subset (0, \infty)$ for all $\Omega' \subset\subset \Omega$. Moreover by (3.16) and (3.17) we have that for a.e. $t \in (0, M)$ it holds

$$|H'(u)||\nabla u| = |H'(t)||((u^*)'((u^*)^{-1}(t)))|AVR(X)^{\frac{1}{N}} = AVR(X)^{\frac{1}{N}}, \quad \text{Per}(\{u > t\})\text{-a.e.}$$

Therefore by coarea (recall (2.6)) and the fact that $\mathbf{m}(\{|\nabla u| = 0\} \cap \Omega) = 0$, we easily deduce that $|H'(u)||\nabla u| = AVR(X)^{\frac{1}{N}}$ \mathbf{m} -a.e. in Ω . In particular $|H'(u)||\nabla u| \in L^2_{loc}(\mathbf{m})$ and we can apply Lemma B.3 to deduce that $f \in W^{1,2}_{loc}(\{u > 0\})$ with $|\nabla f| = 1$, \mathbf{m} -a.e. in $\{u > 0\}$. Moreover from the local Sobolev-to-Lipschitz property (see [62, Prop. 1.10]) we deduce that $f \in \text{LIP}_{loc}(\{u > 0\})$ and

$$|f(x) - f(y)| \leq \mathbf{d}(x, y), \quad \forall x, y \in \{u > 0\}, \quad \text{with } \mathbf{d}(x, y) \leq \mathbf{d}(x, \{u = 0\}). \quad (3.20)$$

This proves (3.19). Next, we claim that

$$\{f < t\} = B_t(x_t), \quad \forall t \in (0, A), \quad (3.21)$$

with $x_t \in \{u > 0\}$. We already know by (3.15) and since H is strictly decreasing, that for every $t \in (0, A)$ the set $\{f < t\}$ is a ball $B_{r_t}(x_t)$ for some $r_t \geq 0$ and x_t tip of X . In particular $\mathbf{m}(\{f < t\}) = \omega_N \theta (r_t)^N$ and $\text{Per}(\{f < t\}) = (\omega_N \theta)^{\frac{1}{N}} N \theta (r_t)^{N-1}$, where $\theta := AVR(X)$. Moreover by coarea formula (2.6) applied to $-f$ and using (3.19)

$$\omega_N \theta [(r_t)^N - (r_s)^N] = \mathbf{m}(\{f < t\}) - \mathbf{m}(\{f < s\}) = \int_{\{s \leq f < t\}} |\nabla f| \, \mathbf{d}\mathbf{m} = \int_s^t \text{Per}(\{f < r\}) \, dr.$$

Therefore the function $(r_t)^N$ is absolutely continuous with

$$\frac{d}{dt}(r_t)^N = (\omega_N \theta)^{-1} \text{Per}(\{f < t\}) = N(r_t)^{N-1}, \quad \text{a.e. } t \in (0, A),$$

from which follows that $r_t = a + t$, for all $t \in (0, A)$, for some constant $a \geq 0$. We claim that $a = 0$. Indeed by continuity and Bishop-Gromov inequality we have

$$a^N \omega_N AVR(X) \leq \mathbf{m}(\cap_{t>0} B_{a+t}(x_t)) = \mathbf{m}(\cap_{t>0} \{f < t\}) = \mathbf{m}(\{f = 0\}) = \mathbf{m}(\{u = M\}) = 0,$$

where in the last equality we used that $|\nabla u| \neq 0$ \mathbf{m} -a.e. in $\{u > 0\}$. This proves (3.21).

It remains to prove that $x_t \equiv x_0$ for all $t \in (0, A)$. This would show (3.18) and conclude the proof. We argue by contradiction and suppose that $x_t \neq x_{\bar{t}}$ for some $\bar{t} < t < A$. Set $\delta := \mathbf{d}(x_t, x_{\bar{t}}) > 0$. Recall that x_t is a tip of X , hence there is a ray emanating from it and containing \bar{x}_t , i.e. an isometry $\gamma : [0, \infty) \rightarrow X$ with $\gamma_0 = x_t$ and $\gamma_\delta = x_{\bar{t}}$. Consider the points $x := \gamma_t \in \partial B_t(x_t) = \{f = t\}$ and $y := \gamma_{\delta+\bar{t}} \in \partial B_{\bar{t}}(x_{\bar{t}}) = \{f = \bar{t}\}$.

Since $\gamma_{\delta+\bar{t}} \in B_t(x_t)$ and γ is an isometry, $\delta + \bar{t} < t$. Therefore applying (3.20), since $d(y, \{u = 0\}) \geq d(y, \partial B_t(x_t)) = d(x, y)$, we finally find a contradiction:

$$t - \bar{t} = f(x) - f(y) \leq d(x, y) = t - (\bar{t} + \delta). \quad \square$$

From Step 1 of the above proof, we deduce the following that has its own interest.

Proposition 3.5 (Improved Pólya-Szegő inequality). *Let (X, d, m) be an $RCD(0, N)$ space with $N \in (1, \infty)$ and $AVR(X) > 0$. Then for every $u \in \text{LIP}_{loc}(X)$, non-negative, $u(x) \rightarrow 0$ as $d(x, z) \rightarrow +\infty$ for some $z \in X$, and with $(u^*)' \neq 0$ -a.e. in $\{u^* > 0\}$, it holds*

$$\int_{\{s < u < r\}} |\nabla u|^2 dm \geq \int_s^r \left(\frac{\text{Per}(\{u > t\})}{N \omega_{\frac{1}{N}} \mu(t)^{\frac{N-1}{N}}} \right)^2 \int |\nabla u_N^*| d\text{Per}(\{u_N^* > t\}) dt, \quad \forall 0 \leq s < r \leq \sup u. \tag{3.22}$$

Remark 3.6. Even if we shall not need it, we observe that Proposition 3.3, Proposition 3.5 and Theorem 3.4 hold replacing $p = 2$ with any $p \in (1, \infty)$, the proof is the same.

We point out that the improved rearrangement inequality (3.22) appeared also in [13, Eq. (3.46)] for non-collapsed spaces and for functions defined on open sets (with finite volume) and with zero-Dirichlet boundary conditions. ■

Remark 3.7 (On the necessity of $(u^*)' \neq 0$ and $|\nabla u| \neq 0$). We point out that, the hypothesis $(u^*)' \neq 0$ in Theorem 3.4 is necessary to prove that u is radial. This is well-known, see e.g. [31, Example 4.6] for an easy counterexample (in \mathbb{R}^n) of a Lipschitz function saturating the Pólya-Szegő inequality with $(u^*)' = 0$ occurring on a set of positive measure.

In Theorem 3.4 we also assumed $|\nabla u| \neq 0$ at m -a.e. point of $\{u > 0\}$. This was needed to carry out key computations by differentiating the distribution functions (see, e.g., (3.6) above), as also done in [86]. It is not clear to us at the moment if this assumption can be removed. ■

4. Regularity of extremal functions

We discuss here the general regularity properties of extremal functions for the Sobolev inequalities (S) considered in this note.

Theorem 4.1 (Regularity of extremal functions). *Fix $N \in (2, \infty)$ and set $2^* := 2N/(N - 2)$. Let (X, d, m) be an $RCD(K, N)$ space, for some $K \in \mathbb{R}, N \in (2, \infty)$ supporting a Sobolev inequality (S) with constant $A > 0, B \geq 0$. Suppose that equality occurs in (S) for some $u \in W_{loc}^{1,2}(X)$ satisfying $\|u\|_{L^{2^*}(m)} = 1$. Then $u \in D(\Delta)$ and*

$$-A\Delta u = (|u|^{2^*-2}u - Bu). \tag{4.1}$$

Moreover if $u \in L^\infty(\mathfrak{m})$, then $u \in \text{LIP}_{loc}(X)$, $|u| > 0$ on X and if $B = 0$ then $|\nabla u| \neq 0$ \mathfrak{m} -a.e.

For the proof, we need two additional results.

Proposition 4.2 (Hopf strong maximum principle). *Let (X, d, \mathfrak{m}) be an $\text{RCD}(K, N)$ space for $K \in \mathbb{R}, N < \infty$. Let $\Omega \subset X$ be open and connected and $u \in D(\Delta, \Omega) \cap C(\Omega)$ satisfying $\Delta u - cum \geq 0$ for some constant $c \geq 0$ and $u(x_0) = \sup_\Omega u \geq 0$, with $x_0 \in \Omega$. Then u is constant.*

Proof. We first prove the following weaker maximum principle:

let $U \subset X$ be open and bounded, and suppose that $v \in D(\Delta, U) \cap C(\bar{U})$ satisfies $\Delta v - cv\mathfrak{m} \geq \delta\mathfrak{m}$ with $\delta > 0$, and $m := \max_{\bar{U}} v \geq 0$, then

$$\max_U v \leq \sup_{\partial U} v. \tag{4.2}$$

Let v and U be as above. Set $C := \{x \in \bar{U} : v(x) = m\}$. If $C \cap \partial U \neq \emptyset$ we are done, hence we can assume that $C \subset U$. Since C is closed $\emptyset \neq \partial C \subset C \subset U$. Let $z_0 \in \partial C$. By continuity there exists r small enough so that $B_r(z_0) \subset U$ and $v \geq -\delta/(2c)$ in $B_r(z_0)$. Then $\Delta v \geq cv\mathfrak{m} + \delta\mathfrak{m} \geq \delta/2\mathfrak{m}$ in $B_r(z_0)$ and in particular v is subharmonic. Then from the strong maximum principle for subharmonic functions [61] (see also [26]) (recall that balls in X are connected) we deduce that $v \equiv m$ in $B_r(z_0)$, which contradicts the fact that $z_0 \in \partial C \subset U$.

We now go back to the proof. The argument is essentially the same in [61], only that we will use the above weak maximum principle instead of the weak maximum principle for subharmonic functions.

Define the set $C := \{u = u(x_0)\} \subset \Omega$. If $C = \Omega$ we are done. Otherwise there exists $x \in \Omega \setminus C$ such that exists a unique $y \in C$ satisfying $r := d(x, y) = d(x, C) < d(x, \Omega^c)$ (see [61]). Define the function $h(z) := e^{-Ad(x,z)^2} - e^{-Ar^2}$, with $A \gg 1$ to be chosen. Let $r' < r/2$ be such that $B_{r'}(y) \subset \Omega$. To finish the proof it is sufficient to show that

$$u(y) = u(y) + \varepsilon h(y) \leq \sup_{\partial B_{r'}(y)} u + \varepsilon h, \quad \forall \varepsilon > 0, \tag{4.3}$$

indeed the conclusion then follows arguing exactly as at the end of [61].

By Laplacian comparison [54] (with computations similar to [61]) we can show that, provided A is chosen large enough depending on r and c , $\Delta|_{B_{r/2}(y)} h \geq 2ce^{-Ad(x,\cdot)^2}\mathfrak{m}|_{B_{r/2}(y)}$. Therefore

$$(\Delta h - ch\mathfrak{m})|_{B_{r/2}(y)} \geq ce^{-Ad(x,\cdot)^2}\mathfrak{m}|_{B_{r/2}(y)} \geq ce^{-4Ar^2}\mathfrak{m}|_{B_{r/2}(y)}.$$

In particular for every $\varepsilon > 0$

$$(\Delta(u + \varepsilon h) - c(u + \varepsilon h)\mathbf{m})|_{B_{r'}(y)} \geq \varepsilon c e^{-4Ar^2} \mathbf{m}|_{B_{r'}(y)},$$

from which (4.3) follows from (4.2) with $v := u + \varepsilon h$, $U := B_{r'}(y)$, noticing that $\sup_{B_{r'}(y)} v \geq u(y) + \varepsilon h(y) = u(x_0) \geq 0$. \square

Proposition 4.3. *Let (X, d, \mathbf{m}) be an $\text{RCD}(K, N)$ space for some $K \in \mathbb{R}$, $N < +\infty$. Consider $\Omega \subset X$ open and $u \in D(\Delta, \Omega)$ with $\Delta u \in L^2_{loc}(\Omega)$. Then*

$$\Delta u = 0 \quad \mathbf{m}\text{-a.e. in } \{|\nabla u| = 0\}. \tag{4.4}$$

Proof. We adapt an argument present in [83] in the Euclidean setting.

It is enough to consider $\Omega = X$ and $\Delta u \in L^2(\mathbf{m})$ with $u \in W^{1,2}(X)$, the general case follows multiplying by Lipschitz cut-off functions with bounded Laplacian (see [85]). We have $|\nabla u| \in W^{1,2}(X)$ (see e.g. [42, Lemma 3.5]) and in particular for every $\varepsilon > 0$, $\frac{|\nabla u|}{|\nabla u| + \varepsilon} \in W^{1,2}(X)$ with

$$\nabla \left(\frac{|\nabla u|}{|\nabla u| + \varepsilon} \right) = \nabla |\nabla u| \frac{\varepsilon}{(|\nabla u| + \varepsilon)^2}$$

(see [55] for the notion of gradient of a Sobolev function). Fix $\varphi \in \text{LIP}(X)$ with $\text{supp}(\varphi) \subset \Omega$. Then integrating by parts

$$\int \varphi \Delta u \frac{|\nabla u|}{|\nabla u| + \varepsilon} \, d\mathbf{m} = - \int \langle \nabla \varphi, \nabla u \rangle \frac{|\nabla u|}{|\nabla u| + \varepsilon} \, d\mathbf{m} + \int \varphi \langle \nabla |\nabla u|, \nabla u \rangle \frac{\varepsilon}{(|\nabla u| + \varepsilon)^2} \, d\mathbf{m}.$$

Since $\left| \frac{\varepsilon |\nabla u|}{(|\nabla u| + \varepsilon)^2} \right| \leq 1$, sending $\varepsilon \rightarrow 0^+$ and applying dominated convergence we obtain

$$\int_{\{|\nabla u| \neq 0\}} \varphi \Delta u \, d\mathbf{m} = - \int \langle \nabla \varphi, \nabla u \rangle \, d\mathbf{m} = \int \Delta u \varphi \, d\mathbf{m}.$$

From the arbitrariness of φ the conclusion follows. \square

Remark 4.4. Even if not needed here, we observe that Proposition 4.3 actually holds in the more general setting of $\text{RCD}(K, \infty)$ spaces (with the same proof). \blacksquare

We can now prove the regularity result for Sobolev extremals.

Proof of Theorem 4.1. The fact that $u \in D(\Delta)$ and that (4.1) holds follows from a straight-forward computation exploiting the fact that u is a minimizer of

$$\inf \frac{\|\nabla v\|_{L^2(\mathbf{m})}^2 + B/A \|v\|_{L^2(\mathbf{m})}^2}{\|v\|_{L^{2^*}(\mathbf{m})}^2} = \frac{1}{A},$$

where the infimum is among all $v \in W_{loc}^{1,2}(X)$ such that $\mathfrak{m}(\{|v| > t\}) < +\infty$ for every $t > 0$ and taking variations of the form $u + \varepsilon v$, $v \in \text{LIP}_c(X)$ as $\varepsilon \rightarrow 0$. See e.g. [88, Prop. 8.3] for the details in the compact case.

We pass to the second part, assuming that u is in $L^\infty(\mathfrak{m})$. From (4.1) we have that $\Delta u \in L^\infty(\mathfrak{m})$, therefore Theorem 2.4 shows that $u \in \text{LIP}_{loc}(X)$.

From now on we will identify u with its continuous representative. We need to show that $|u| > 0$ in X . Suppose this is not the case, i.e. $|u|(x_0) = 0$ for some $x_0 \in X$. Note that $|u|$ also satisfies the hypotheses of the theorem, hence $-\Delta|u| = |u|A^{-1}(|u|^{2^*-2} - B)$. Consider the function $v := -|u| \leq 0$. Then, since $u \in L^\infty(\mathfrak{m})$,

$$\Delta v - Cv = |u|(A^{-1}|u|^{2^*-2} - A^{-1}B + C) \geq 0,$$

provided we choose the constant $C > 0$ big enough. In particular, v satisfies the assumption of the maximum principle of Proposition 4.2 with $v(x_0) = 0 = \max v$. Hence $v \equiv 0$ in X , which is a contradiction because u is assumed non-zero. Finally, if $B = 0$, since u never vanishes, we have that also Δu never vanishes, hence $|\nabla u| \neq 0$ \mathfrak{m} -a.e. thanks to (4.4). \square

5. Rigidity of extremal functions in the Sobolev inequality

5.1. Compact case

We study here the equality case for the Sobolev inequality as in (2.9).

As a technical tool we will need the following result that is a standard application of the Moser iteration scheme (see e.g. [67, Theorem 4.4]). This is known to be still valid in our setting, relying only on the Sobolev inequality (see also the discussion after [56, Theorem 5.7]).

Lemma 5.1. *Let (X, d, \mathfrak{m}) be a compact $\text{RCD}(K, N)$ space, $N < +\infty$, and $u \in D(\Delta)$ satisfying for some $g \in L^{N/2}(\mathfrak{m})$*

$$\Delta u = gum.$$

Then $u \in L^q(\mathfrak{m})$ for every $q < +\infty$.

We can now state and prove the main result of this section. Note that the fact that X is spherical suspension already follows from [88, Theorem 1.9]. Here, we are mainly interested in the explicit expression of extremal functions.

Theorem 5.2. *Let (X, d, \mathfrak{m}) be an $\text{RCD}(N - 1, N)$ space, $\mathfrak{m}(X) = 1$, $N \in (2, \infty)$ and set $2^* = 2N/(N - 2)$. Let $u \in W^{1,2}(X)$ be non-constant with $\|u\|_{L^{2^*}} = 1$ satisfying*

$$\|u\|_{L^{2^*}(\mathfrak{m})}^2 = \frac{2^* - 2}{N} \|\nabla u\|_{L^2(\mathfrak{m})}^2 + \|u\|_{L^2(\mathfrak{m})}^2.$$

Then, X is isomorphic to a spherical suspension and, for some $a \in \mathbb{R}, b \in (0, 1)$ and $z_0 \in X$:

$$u = a(1 - b \cos d(\cdot, z_0))^{\frac{2-N}{2}}.$$

Proof. The argument is inspired by the computations in [44, Section 2.1].

First, we need to deduce some regularity on the extremal function u . From Theorem 4.1 we know that $u \in D(\Delta)$ and that

$$\frac{2^* - 2}{N} \Delta u = u - |u|^{2^* - 2} u. \tag{5.1}$$

Since $u^{2^* - 2} \in L^{N/2}(\mathfrak{m})$, by Lemma 5.1 below we deduce that $u \in L^q(\mathfrak{m})$ for all $q < +\infty$. In particular $\Delta u \in L^q(\mathfrak{m})$ for all $q < +\infty$. Therefore by [75, Corollary 6] we have $u \in \text{LIP}(X)$ and so $u \in L^\infty(\mathfrak{m})$ (alternatively we could have showed $u \in L^\infty(\mathfrak{m})$ applying [90, Lemma 4.1] and then deduced the Lipschitzianity from Theorem 4.1). Then we can apply the second part of Theorem 4.1 to deduce that either $u > 0$ or $u < 0$ in X . Note also that $\Delta u \in W^{1,2}(X)$.

Without loss of generality, we can assume that $u > 0$. Set $v := u^{\frac{-2}{N-2}}$. By the chain rule for the Laplacian (see e.g. [59, Prop. 5.2.3]) $v \in D(\Delta)$ with

$$\Delta v = u^{\frac{-2}{N-2}} \left(\frac{-2}{N-2} u^{-1} \Delta u + \frac{2N}{(N-2)^2} u^{-2} |\nabla u|^2 \right) \in W^{1,2}(X) \cap L^\infty(\mathfrak{m}),$$

indeed $|\nabla u|^2 \in W^{1,2}(X)$ by [55, Prop. 3.1.3]. Noting that $|\nabla v|^2 = \frac{4}{(N-2)^2} |\nabla u|^2 u^{-2} u^{\frac{-4}{N-2}}$, an easy computation using (5.1) shows

$$v \Delta v = -\frac{N}{2} (v^2 - 1) + \frac{N}{2} |\nabla v|^2. \tag{5.2}$$

Since v is bounded above and away from zero, by the chain rule for the Laplacian we also have that $v^{1-N} \in D(\Delta)$ with $\Delta v^{1-N} \in L^\infty(\mathfrak{m})$. We can then multiply (5.2) by Δv^{1-N} and integrate

$$-\frac{N}{2} \int \Delta v^{1-N} v^2 \, d\mathfrak{m} = \int \Delta v^{1-N} \left(v \Delta v - \frac{N}{2} |\nabla v|^2 \right) \, d\mathfrak{m}.$$

We now proceed to integrate by parts. To do this note that $v \Delta v \in W^{1,2}(X)$ and $|\nabla v|^2 \in D(\Delta)$ (see [55, Prop. 3.1.3]). Moreover by the Leibniz rule for the divergence $\text{div}(\nabla v \Delta v) = \langle \nabla v, \nabla \Delta v \rangle + (\Delta v)^2 \in L^1(\mathfrak{m})$ by the Leibniz rule (see [55,60] for the notion of divergence and e.g. [62, Prop. 3.2] for a version of the Leibniz rule that applies here). Hence

$$N(1 - N) \int |\nabla v|^2 v^{1-N} \, d\mathfrak{m} = -\frac{N}{2} \int \Delta v^{1-N} |\nabla v|^2 - \int \langle \nabla v^{1-N}, \nabla v \Delta v + v \nabla \Delta v \rangle \, d\mathfrak{m}$$

$$\begin{aligned}
 &= -\frac{N}{2} \int v^{1-N} \Delta |\nabla v|^2 \\
 &\quad + \int v^{1-N} (\Delta v)^2 + N \langle \nabla v, \nabla \Delta v \rangle v^{1-N} \, \mathbf{m}.
 \end{aligned}$$

Combining the above with the dimensional Bochner inequality ([46,66]) and with $v > 0$, we get

$$\frac{1}{2} \Delta |\nabla v|^2 - \langle \nabla \Delta v, \nabla v \rangle \mathbf{m} = \frac{(\Delta v)^2}{N} \mathbf{m} + (N - 1) |\nabla v|^2 \mathbf{m}.$$

Integrating and using that $\int d\Delta |\nabla v|^2 = 0$ gives

$$\int (\Delta v)^2 \, \mathbf{m} = - \int \langle \nabla \Delta v, \nabla v \rangle = \int \frac{(\Delta v)^2}{N} \, \mathbf{m} + (N - 1) \int |\nabla v|^2 \, \mathbf{m},$$

from which $\int (\Delta v)^2 \, \mathbf{m} = N \int |\nabla v|^2 \, \mathbf{m}$. In particular $\int \tilde{v}^2 \, \mathbf{m} = N \int |\nabla \tilde{v}|^2$, where $\tilde{v} := (v - \int v \, \mathbf{m})$. Then by [77] we deduce that X is a spherical suspension and

$$\tilde{v}(x) = c \cos d(x, z_0) = -c \cos d(x, \bar{z}_0), \quad \forall x \in X,$$

for some constant $c > 0$ and $z_0, \bar{z}_0 \in X$ tips of the spherical suspension with $d(z, \bar{z}_0) = \pi$. Recalling that $v = u^{\frac{2}{2-N}}$ concludes the proof. \square

5.2. Non-compact case

Here we investigate the equality case in the Euclidean-type Sobolev inequality (2.13).

Theorem 5.3. *Let (X, d, \mathbf{m}) be an $\text{RCD}(0, N)$ space with $N \in (2, \infty)$, $\text{AVR}(X) > 0$ and set $2^* = 2N/(N - 2)$. Suppose that for some non-zero $u \in W_{loc}^{1,2}(X)$ with $\mathbf{m}(\{|u| > t\}) < \infty$ for all $t > 0$, it holds*

$$\|u\|_{L^{2^*}(\mathbf{m})} = \text{Eucl}(N, 2) \text{AVR}(X)^{-\frac{1}{N}} \|\nabla u\|_{L^2(\mathbf{m})} \tag{5.3}$$

(both being finite). Then, X is isomorphic to a N -Euclidean metric measure cone and

$$u = a(1 + b d^2(\cdot, z_0))^{\frac{2-N}{2}}, \tag{5.4}$$

for some $a \in \mathbb{R}, b > 0$ and z_0 one of the tips of X .

Proof. We will apply Theorem 3.4. First we need to prove the required regularity of u .

Notice that we can equivalently suppose that $\|u\|_{L^{2^*}(\mathbf{m})} = 1$, by scaling invariance. Moreover also $|u|$ satisfies the equality in (5.3). By assumptions, it is possible to perform a Euclidean rearrangement $|u|^*$ of $|u|$. By the Pólya-Szegő inequality and the one-dimensional Bliss inequality we get

$$\begin{aligned} \|u\|_{L^{2^*}(\mathfrak{m})} &= \text{Eucl}(N, 2)\text{AVR}(X)^{-\frac{1}{N}} \|\nabla u\|_{L^2(\mathfrak{m})} \\ &\stackrel{(3.4)}{\geq} \text{Eucl}(N, 2)\|\nabla|u|^*\|_{L^2(\mathfrak{m}_N)} \stackrel{(2.11)}{\geq} \|u^*\|_{L^{2^*}(\mathfrak{m}_N)}. \end{aligned}$$

Note that we can apply (2.11) since by Proposition 3.3, $u^* \in W_{loc}^{1,2}(I_N)$ and thus u is locally absolutely continuous in $(0, \infty)$ (see e.g. [88, Section 2.2]). By (3.3) we see that the inequalities in the above are all equalities, and therefore equality holds in the Bliss inequality. Therefore $|u|^*(t) = a(1 + bt^2)^{\frac{2-N}{2}}$ for some $a \in \mathbb{R}, b > 0$. In particular, since $\|u\|_{L^\infty} = \|u^*\|_{L^\infty} < \infty$ by equimeasurability, we have $u \in W_{loc}^{1,2}(X) \cap L^\infty(\mathfrak{m})$ and we can invoke Theorem 4.1 (with $B = 0$) to deduce $u \in \text{Lip}_{loc}(X) \cap D(\Delta)$, $\mathfrak{m}(\{|\nabla u| = 0\}) = 0$, $u > 0$ or $u < 0$, and (assuming $u > 0$):

$$\text{Eucl}^2(N, 2)\text{AVR}(X)^{-\frac{2}{N}} \Delta u = -u^{2^*-1}.$$

Recalling Theorem 2.4, since $u \in L^\infty(\mathfrak{m})$, we get that $|\nabla u| \in L^\infty(\mathfrak{m})$. By the Sobolev-to-Lipschitz property (see [53,6]), u has a Lipschitz representative, still denoted by u in what follows. It remains to show that $u(x) \rightarrow 0$ as $\mathfrak{d}(z, x) \rightarrow \infty$, for $z \in X$. Suppose, by contradiction, that there is a sequence $(x_n) \subset X$ satisfying $\mathfrak{d}(x_n, z) \rightarrow \infty$ as $n \uparrow \infty$ and with the property that $u(x_n) \geq c > 0$ for all $n \in \mathbb{N}$. Since $u \in \text{LIP}(X)$, denoting $L := \text{Lip}(f)$, we see that for any $x \in B_{c/(2L)}(x_n)$ we have $u(x) \geq u(x_n) - L\mathfrak{d}(x, x_n) \geq c/2$ and therefore

$$\int_{B_{c/(2L)}(x_n)} |u|^{2^*} \, \mathfrak{d}\mathfrak{m} \geq (c/2)^{2^*} \mathfrak{m}(B_{c/(2L)}(x_n)) \geq \omega_N c^{2^*} \text{AVR}(X)(c/(2L))^N > 0.$$

However this contradicts $u \in L^{2^*}(\mathfrak{m})$.

We deduced all the regularity required to invoke Theorem 3.4, so we know that X is an N -Euclidean metric measure cone with tip z_0 and u is radial, i.e. $u(x) = u^* \circ \text{AVR}(X)^{\frac{1}{N}} \mathfrak{d}(x, z_0)$. The conclusion follows since $u^*(t) = |u|^*(t) = a(1 + bt^2)^{\frac{2-N}{2}}$ for some $a \in \mathbb{R}, b > 0$. \square

6. Compactness of extremizing sequences

A classical result using concentration compactness is that a sequence extremizing functions for the Sobolev inequality in \mathbb{R}^n , up to a rescaling, dilation and translation, converges up to a subsequence to an extremal function. In this part, we generalize this method to an extremizing sequence of functions defined on a sequence of $\text{RCD}(0, N)$ spaces (Theorem 6.2).

6.1. Density upper bound

We first address a technical density bound that will be needed in the proof of Theorem 6.2 to get pre-compactness in the pmGH-topology. This part is needed only for

collapsed RCD-spaces: a reader interested in the case of smooth manifolds can skip this subsection.

Lemma 6.1 (Density bound from reverse Sobolev). *For every $N \in (2, \infty)$, $K \in \mathbb{R}$, there are constants $\lambda_{N,K} \in (0, 1)$, $r_{K^-,N} > 0$ (with $r_{0,N} = +\infty$), $C_{N,K} > 0$ such that the following holds. Let (X, d, \mathbf{m}) be an $\text{RCD}(K, N)$ space and $u \in W_{loc}^{1,2}(X) \cap L^{2^*}(\mathbf{m})$, non-constant satisfying*

$$\|u\|_{L^{2^*}(\mathbf{m})}^2 \geq A \|\nabla u\|_{L^2(\mathbf{m})}^2, \tag{6.1}$$

for some $A > 0$. Assume also that for some $\eta \in (0, \lambda_{N,K})$, $\rho \in (0, r_{K^-,N} \wedge \frac{\lambda_{N,K}}{8} \text{diam}(X))$ and $x \in X$ it holds

$$\|u\|_{L^{2^*}(B_\rho(x))}^{2^*} \geq (1 - \eta) \|u\|_{L^{2^*}(\mathbf{m})}^{2^*}.$$

Then

$$\frac{\mathbf{m}(B_\rho(x))}{\rho^N} \leq \frac{C_{N,K}}{A^{N/2}}. \tag{6.2}$$

Proof. We fix a constant $\lambda = \lambda_{N,K} \in (0, 1)$ sufficiently small and to be chosen later. We also fix a constant $r_{K^-,N} > 0$, with $r_{0,N} = +\infty$ and with $r_{K^-,N}$ small and to be chosen later in the case $K < 0$ ($r_{K^-,N}$ will be chosen after $\lambda_{N,K}$). Assume $\rho \leq r_{K^-,N}$ and $\eta \leq \lambda_{N,K}$ are as in the hypotheses.

Observe that $B_{4\lambda^{-1}\rho}(x) \subsetneq X$. Up to choosing $r_{K^-,N}$ small enough (when $K < 0$) we can assume that $4\lambda^{-1}\rho \leq \tilde{r}_{K^-,N}$, where $\tilde{r}_{K^-,N} > 0$ is the one given by Lemma B.2. Set $r := 4\lambda^{-1}\rho \geq 4\rho$ and note that $B_r(x) \subsetneq X$.

Fix a cut-off function $\varphi \in \text{LIP}_c(B_{r/2}(x))$ such that $\varphi = 1$ in $B_{r/4}(x)$, $0 \leq \varphi \leq 1$ and $\text{Lip}(\varphi) \leq 10/r$. Then from (B.2), since $r \leq \tilde{r}_{K^-,N}$, we have

$$\begin{aligned} \|u\|_{L^{2^*}(B_\rho(x))} &\leq \|u\varphi\|_{L^{2^*}(\mathbf{m})} \leq \frac{C_{N,K}r}{\mathbf{m}(B_r(x))^{1/N}} \|\nabla u\|_{L^2(\mathbf{m})} + \frac{10C_{N,K}}{\mathbf{m}(B_r(x))^{1/N}} \|u\|_{L^2(B_r(x))} \\ &\leq \frac{C_{N,K}r}{\mathbf{m}(B_r(x))^{1/N}} \|\nabla u\|_{L^2(\mathbf{m})} + \frac{10C_{N,K}}{\mathbf{m}(B_r(x))^{1/N}} (\|u\|_{L^2(B_\rho(x))} + \|u\|_{L^2(B_r(x) \setminus B_\rho(x))}) \\ &\leq \frac{C_{N,K}r}{\mathbf{m}(B_r(x))^{1/N}} \|\nabla u\|_{L^2(\mathbf{m})} + \frac{10C_{N,K} \|u\|_{L^{2^*}(\mathbf{m})}}{\mathbf{m}(B_r(x))^{1/N}} (\mathbf{m}(B_\rho(x))^{1/N} + \lambda^{1/2^*} \mathbf{m}(B_r(x))^{1/N}) \end{aligned}$$

Substituting (6.1), applying (2.5) (up to choosing $r_{K^-,N}$ small enough so that $r \leq R_{K^-,N}$), using that $1 - \lambda < 1 - \eta$ and simplifying $\|u\|_{L^{2^*}(\mathbf{m})}$, we reach

$$(1 - \lambda)^{1/2^*} \leq \frac{C_{N,K}r}{\sqrt{A} \mathbf{m}(B_r(x))^{1/N}} + 10C_{N,K}((\lambda/4)^\gamma + \lambda^{1/2^*}),$$

where $\gamma > 0$ is a constant depending only on N . Choosing λ small enough with respect to N and K gives

$$\frac{m(B_\rho(x))}{r^N} \leq \frac{m(B_r(x))}{r^N} \leq \frac{C_{N,K}}{A^{N/2}}. \tag{6.3}$$

Recalling that $r = 4\lambda^{-1}\rho$ proves (6.2). \square

6.2. Concentration compactness for Sobolev extremals

In the following theorem we show that a sequence of extremizing functions defined on a sequence of $\text{RCD}(0, N)$ spaces, after a suitable rescaling of both the function and the space, admits a subsequence converging to a limit extremal function on some limit $\text{RCD}(0, N)$ space. The idea is similar to the classical Lions’ concentration-compactness principle ([80,81]). The first step is a characterization of the failure of compactness in the critical Sobolev embedding by specific concentration and splitting of the mass phenomena (see Appendix A.2). The second step is observing that the extra information that the sequence is extremizing for the Sobolev inequality will prevent these pathological phenomena and ensure compactness. A crucial point will be to exploit the strict concavity property of the Sobolev inequality, and in particular of the function $t \mapsto t^{2/2^*}$, to deduce that splitting the mass is not convenient in an extremizing sequence.

Theorem 6.2. *For every $N \in (2, \infty)$, exists $\eta_N \in (0, 1/2)$ such that the following holds. Let $(Y_n, \rho_n, \mu_n, y_n)$ be a sequence of pointed $\text{RCD}(0, N)$ spaces supporting a Sobolev inequality (S) with $A_n \rightarrow A > 0$ and $B_n \rightarrow B \in [0, \infty)$ and also satisfying either $\sup_n \mu_n(B_1(y_n)) < +\infty$ or $\text{diam}(Y_n) > \eta_N^{-1}$.*

Suppose there exist non-constant functions $u_n \in W^{1,2}(Y_n)$ with $\|u_n\|_{L^{2^}(\mu_n)} = 1$ and*

$$\sup_{y \in Y_n} \int_{B_1(y)} |u_n|^{2^*} d\mu_n = \int_{B_1(y_n)} |u_n|^{2^*} d\mu_n = 1 - \eta, \tag{6.4}$$

$$\|u_n\|_{L^{2^*}(\mu_n)}^2 \geq \tilde{A}_n \|\nabla u_n\|_{L^2(\mu_n)}^2 + B_n \|u_n\|_{L^2(\mu_n)}^2, \tag{6.5}$$

for $\tilde{A}_n \rightarrow A$, and some $\eta \in (0, \eta_N)$. Then, up to a subsequence, it holds:

- i) Y_n pmGH-converges to a pointed $\text{RCD}(0, N)$ -space (Y, ρ, μ, y) supporting a Sobolev inequality as in (S) with constants A, B ;
- ii) u_n converges L^{2^*} -strong to some $u \in W_{loc}^{1,2}(Y)$ with $|\nabla u| \in L^2(\mu)$ and

$$\int |\nabla u_n|^2 d\mu_n \rightarrow \int |\nabla u|^2 d\mu, \quad \text{as } n \uparrow \infty.$$

If $B > 0$, then the convergence is also $W^{1,2}$ -strong.

iii) It holds

$$\|u\|_{L^{2^*}(\mu)}^2 = A\|\nabla u\|_{L^2(\mu)}^2 + B\|u\|_{L^2(\mu)}^2.$$

Proof. We subdivide the proof into different steps.

STEP 1. We take $\eta_N := \frac{\lambda_{0,N}}{8} \wedge \frac{1}{3}$, with $\lambda_{0,N}$ as in Lemma 6.1. In light of Theorem 2.10, to extract a subsequence converging pmGH it is sufficient to check that $\mu_n(B_1(y_n)) \in (v^{-1}, v)$ for some $v > 1$. If $\text{diam}(Y_n) > \eta_N^{-1} \geq 8\lambda_{0,N}^{-1}$, thanks to the assumptions (6.4) and (6.5), we can apply Lemma 6.1 to obtain

$$\overline{\lim}_n \mu_n(B_1(y_n)) \leq \overline{\lim}_n \frac{C_N}{(\tilde{A}_n)^{N/2}} = \frac{C_N}{A^{N/2}} < +\infty,$$

otherwise $\sup_n \mu_n(B_1(y_1)) < +\infty$ is directly true by the assumptions. On the other hand, since by assumption the spaces Y_n satisfy a Sobolev inequality with constants A_n, B_n , plugging in functions $\varphi_n \in \text{LIP}(Y_n)$ such that $\varphi_n = 1$ in $B_1(y_n)$ with $\text{supp} \varphi_n \subset B_2(y_n)$, $0 \leq \varphi_n \leq 1$ and $\text{Lip}(\varphi_n) \leq 1$, we get

$$\mu_n(B_1(y_n))^{2/2^*} \leq (A_n + B_n)\mu_n(B_2(y_n)) \leq 2^N(A_n + B_n)\mu_n(B_1(y_n)),$$

where we used the Bishop-Gromov inequality. Since $\lim_n(A_n + B_n) = A + B > 0$ we also obtain $\underline{\lim}_n \mu_n(B_1(y_n)) > 0$. Therefore up to a not relabeled subsequence, the spaces Y_n pmGH converge to a pointed RCD(0, N) space (Y, ρ, μ, y) . Moreover, the stability of the Sobolev inequalities [88, Lemma 4.1] ensures that Y supports a Sobolev inequality as in (S) with constants A, B . This settles point i).

STEP 2. From now on we assume to have fixed a realization of the convergence in a proper metric space (Z, d) (as in Section 2.4). Let $\nu_n := |u_n|^{2^*} \mu_n \in \mathcal{P}(Z)$. Moreover we will denote by $B_r(z)$, $z \in Z$, and by $B_r^n(y)$, $y \in Y_n$, respectively the balls in (Z, d) and in (Y_n, ρ_n) , recalling that we are identifying (Y_n, ρ_n) as a subset of (Z, d) . From Lemma A.6 we have that, up to a subsequence, (exactly) one of cases i), ii), iii) in the statement of Lemma A.6 holds. We claim i) (i.e. compactness) occurs. First, notice that vanishing as in case ii) cannot occur:

$$\overline{\lim}_{n \rightarrow \infty} \sup_{y \in Y_n} \nu_n(B_R(y)) \geq \overline{\lim}_{n \rightarrow \infty} \nu_n(B_1(y_n)) \stackrel{(6.4)}{=} 1 - \eta, \quad \forall R \geq 1.$$

Thus, it remains to exclude the dichotomy case iii). Suppose by contradiction that iii) of Lemma A.6 holds for some $\lambda \in (0, 1)$ (with $\lambda \geq \overline{\lim}_n \sup_z \nu_n(B_R(z))$ for all $R > 0$), sequences $R_n \uparrow \infty$, $(z_n) \subset Z$ and measures ν_n^1, ν_n^2 with $\text{supp}(\nu_n^1) \subset B_{R_n}(z_n)$ and $\text{supp}(\nu_n^2) \subset Z \setminus B_{10R_n}(z_n)$. We claim first that $\text{supp}(\nu_n^1) \subset B_{3R_n}(y_n)$ and $\text{supp}(\nu_n^2) \subset Z \setminus B_{4R_n}(y_n)$. Indeed $\lambda \geq \overline{\lim}_n \nu_n(B_1(y_n)) = 1 - \eta$ and

$$\underline{\lim}_n \nu_n(B_{R_n}(z_n)) \geq \underline{\lim}_n \nu_n^1(B_{R_n}(z_n)) = \lim_n \nu_n^1(Z) = \lambda \geq 1 - \eta.$$

Since $\nu_n(B_1(y_n)) = 1 - \eta$ and $\eta < 1/2$, this implies that for n large enough $B_{R_n}(z_n) \cap B_1(y_n) \neq \emptyset$, which implies the claim, provided $R_n \geq 1$.

Let φ_n be a Lipschitz cut-off so that $0 \leq \varphi_n \leq 1$, $\varphi_n \equiv 1$ on $B_{3R_n}^n(y_n)$, $\text{supp}(\varphi_n) \subset B_{4R_n}^n(y_n)$ and $\text{Lip}(\varphi_n) \leq R_n^{-1}$, for every $n \in \mathbb{N}$. Since

$$1 \geq |\varphi_n|^2 + |(1 - \varphi_n)|^2, \quad \text{in } Z, \tag{6.6}$$

we can estimate by triangular inequality, the Leibniz rule and Young inequality

$$\begin{aligned} \|\nabla u_n\|_{L^2(\mu_n)}^2 &\geq \|\varphi_n|\nabla u_n\|_{L^2(\mu_n)}^2 + \|(1 - \varphi_n)|\nabla u_n\|_{L^2(\mu_n)}^2 \\ &\geq \|\nabla(u_n\varphi_n)\|_{L^2(\mu_n)}^2 + \|\nabla(u_n(1 - \varphi_n))\|_{L^2(\mu_n)}^2 \\ &\quad - \underbrace{2(1 + \delta^{-1})\|u_n|\nabla\varphi_n\|_{L^2(\mu_n)}^2 - 2\delta\|\nabla u_n\|_{L^2(\mu_n)}^2}_{:=R_n(\delta)} \end{aligned} \tag{6.7}$$

for every $\delta > 0$ and every n . Setting $O_n := B_{4R_n}^n(y_n) \setminus B_{3R_n}^n(y_n)$, we have by the Hölder inequality

$$\|u_n|\nabla\varphi_n\|_{L^2(\mu_n)}^2 \leq R_n^{-2}\|u_n\|_{L^{2^*}(O_n)}^2\mu_n(O_n)^{2/N} \leq 16v^{2/N}\|u_n\|_{L^{2^*}(O_n)}^2,$$

having used that $\mu_n(O_n) \leq \mu_n(B_{4R_n}^n(y_n)) \leq (4R_n)^N\mu_n(B_1(y_n)) \leq (4R_n)^Nv$, by the Bishop-Gromov inequality. Notice that we also have

$$\overline{\lim}_{n \rightarrow \infty} \|u_n\|_{L^{2^*}(O_n)} \leq \overline{\lim}_{n \rightarrow \infty} \left|1 - \nu_n^1(Z) - \nu_n^2(Z)\right|^{1/2^*} = 0,$$

from which we get $\lim_n \|u_n|\nabla\varphi_n\|_{L^2(\mu_n)}^2 = 0$. Therefore, recalling that $\|\nabla u_n\|_{L^2(\mu_n)}^2$ is uniformly bounded by (6.5), choosing appropriately $\delta_n \rightarrow 0$, we get

$$R_n(\delta_n) \rightarrow 0. \tag{6.8}$$

Combining (6.7) with (6.8), recalling that $\lim_n A_n = \lim_n \tilde{A}_n$, we get

$$\begin{aligned} 1 &\stackrel{(6.5)}{\geq} \overline{\lim}_{n \rightarrow \infty} A_n \|\nabla(u_n\varphi_n)\|_{L^2(\mu_n)}^2 + A_n \|\nabla(u_n(1 - \varphi_n))\|_{L^2(\mu_n)}^2 + B_n \|u_n\|_{L^2(\mu_n)}^2 \\ &\stackrel{(S)}{\geq} \overline{\lim}_{n \rightarrow \infty} \|u_n\varphi_n\|_{L^{2^*}(\mu_n)}^2 + \|u_n(1 - \varphi_n)\|_{L^{2^*}(\mu_n)}^2 \\ &\quad + B_n \left(\|u_n\|_{L^2(\mu_n)}^2 - \|u_n\varphi_n\|_{L^2(\mu_n)}^2 - \|u_n(1 - \varphi_n)\|_{L^2(\mu_n)}^2\right) \\ &\stackrel{(6.6)}{\geq} \overline{\lim}_{n \rightarrow \infty} (\nu_n^1(Z))^{2/2^*} + (\nu_n^2(Z))^{2/2^*} \\ &\geq \lambda^{2/2^*} + (1 - \lambda)^{2/2^*} > 1, \end{aligned}$$

having used the strict concavity of $t \mapsto t^{2/2^*}$ and the fact that $\lambda \in (0, 1)$. This gives a contradiction, hence dichotomy in iii) cannot happen.

STEP 3. In the previous step, we proved that case i) in Lemma A.6 occurs, i.e. there exists $(z_n) \subset Z$ such that for every $\varepsilon > 0$ there exists $R := R(\varepsilon)$ so that $\int_{B_R^n(z_n)} |u_n|^{2^*} d\mu_n \geq 1 - \varepsilon$ for all $n \in \mathbb{N}$. As soon as $\varepsilon < 1/2$, we have $B_R^n(z_n) \cap B_1^n(y_n) \neq \emptyset$ and

$$\int_{B_{2R+1}^n(y_n)} |u_n|^{2^*} d\mu_n \geq 1 - \varepsilon \quad \forall n \in \mathbb{N}. \tag{6.9}$$

Moreover $y_n \rightarrow y$ in Z , hence the sequence of probabilities $|u_n|^{2^*} \mu_n$ is tight (Z is proper) and, along a not relabeled subsequence, converges in duality with $C_b(Z)$ to some $\nu \in \mathcal{P}(Y)$. Additionally, up to a further subsequence we have that u_n is L^{2^*} -weak convergent to some $u \in L^{2^*}(\mu)$ ([8]) with $\sup_n \|\nabla u_n\|_{L^2(\mu_n)} < \infty$ and also that $|\nabla u_n|^2 d\mu_n \rightharpoonup \omega$ in duality with $C_{bs}(Z)$ for some bounded Borel measure ω . Applying Lemma A.3, up to a further subsequence, we also deduce that u_n converges L^2_{loc} -strong to some $u \in L^2_{loc}(\mu)$, together with the facts $u \in W^{1,2}_{loc}(Y)$ and $|\nabla u| \in L^2(\mu)$. Note that if $B > 0$ then actually $u \in W^{1,2}(Y)$, by (6.5) and the lower semicontinuity of the L^2 -norm (2.15).

We are in position to invoke Lemma A.7 to infer the existence of countably many points $\{x_j\}_{j \in J} \subset Y$ and positive weights $(\nu_j), (\omega_j) \subset \mathbb{R}^+$, so that $\nu = |u|^{2^*} \mu + \sum_{j \in J} \nu_j \delta_{x_j}$ and $\omega \geq |\nabla u|^2 \mu + \sum_{j \in J} \omega_j \delta_{x_j}$, with $A\omega_j \geq \nu_j^{2/2^*}$ and in particular $\sum_j \nu_j^{2/2^*} < \infty$. Moreover up to passing to a subsequence we can, and will, from now on assume that the limits $\lim_n \|\nabla u_n\|_{L^2(\mu_n)}^2$ and $\lim_n B_n \|u\|_{L^2(\mu_n)}^2$ exist. Finally, by the lower semicontinuity of the L^2 -norm (see (2.15)) we have $B \|u\|_{L^2(\mu)}^2 \leq \lim_n B_n \|u\|_{L^2(\mu_n)}^2$, where $B \|u\|_{L^2(\mu)}^2$ is taken to be zero when $B = 0$ and $\|u\|_{L^2(\mu)}^2 = +\infty$. Also $\lim_n \|\nabla u_n\|_{L^2(\mu_n)}^2 \geq \omega(Z)$. Therefore

$$\begin{aligned} 1 &= \lim_{n \rightarrow \infty} \int |u_n|^{2^*} d\mu_n \geq \lim_{n \rightarrow \infty} \tilde{A}_n \|\nabla u_n\|_{L^2(\mu_n)}^2 + \lim_{n \rightarrow \infty} B_n \|u_n\|_{L^2(\mu_n)}^2 \\ &\geq A\omega(Z) + B \|u\|_{L^2(\mu)}^2 \\ &\geq A \int |\nabla u|^2 d\mu + \sum_{j \in J} \nu_j^{2/2^*} + B \|u\|_{L^2(\mu)}^2 \\ &\stackrel{(8)}{\geq} \left(\int |u|^{2^*} d\mu \right)^{2/2^*} + \sum_{j \in J} \nu_j^{2/2^*} \\ &\geq \left(\int |u|^{2^*} d\mu + \sum_{j \in J} \nu_j \right)^{2/2^*} = \nu(Y)^{2/2^*} = 1, \end{aligned}$$

having used, in the last inequality, the concavity of the function $t^{2/2^*}$. In particular, all the inequalities must be equalities and, since $t^{2/2^*}$ is strictly concave, we infer that every term in the sum $\int |u|^{2^*} d\mu + \sum_{j \in J} \nu_j^{2/2^*}$ must vanish except one. By the assumption (6.4) and $|u|^{2^*} \mu_n \rightharpoonup \nu$ in $C_b(Z)$, we have $\nu_j \leq 1 - \eta$ for every $j \in J$. Hence $\nu_j = 0$

and $\|u\|_{L^{2^*}(\mu)} = 1$. This means that u_n converges L^{2^*} -strong to u . Moreover, retracing the equalities in the above we have that $\lim_n \int |\nabla u_n|^2 d\mu_n = \int |\nabla u|^2 d\mu$ and, when $B > 0$, $\lim_n \int |u_n|^2 d\mu_n = \int |u|^2 d\mu$. This proves point ii). Finally, equality in the fourth inequality is precisely part iii) of the statement. The proof is now concluded. \square

7. Radial functions: technical results

In this section, we prove results about convergence and approximation of radial functions.

The first one (Lemma 7.2 below) says that, given a sequence of RCD spaces converging in the pmGH-sense, a radial function on the limit space is the limit of the same radial functions along the sequence.

We will need the following simple fact. We omit the proof, which is an easy consequence of Cavalieri’s formula and Bishop-Gromov inequality. In this section, we denote $d_z(\cdot) := d(z, \cdot)$ the distance function from a point z .

Lemma 7.1. *Let (X, d, m) be an $RCD(0, N)$ space for some $N \in (2, \infty)$. Then for every $\alpha > N$, $z \in X$ and $r > 0$ it holds*

$$\int_{B_r(z)^c} d_z(\cdot)^{-\alpha} dm \leq \frac{m(B_r(z))}{r^N} C_{N,\alpha} r^{N-\alpha}. \tag{7.1}$$

Lemma 7.2. *Let $(Y_n, \rho_n, \mu_n, z_n)$ be a sequence of $RCD(K, N)$ spaces, for some $K \in \mathbb{R}$, $N \in (2, \infty)$, that is pmGH-converging to (Y, ρ, μ, z_0) . Let $p \in (1, \infty)$ and $f \in C(\mathbb{R})$ satisfying $|f(t)|^p \leq C|t|^{-\alpha}$, for some $\alpha > 0$. Suppose also that*

$$\lim_{R \rightarrow +\infty} \sup_n \int_{B_R(z_n)^c} \rho_{z_n}^{-\alpha} d\mu_n = 0, \tag{7.2}$$

where $\rho_{z_n}(\cdot) := \rho_n(\cdot, z_n)$. Then, $f \circ \rho_{z_n}$ converges L^p -strong to $f \circ \rho_{z_0}$. In particular, for any $u_n \in L^p(\mu_n)$ that converges L^p -strong to $f \circ \rho_{z_0}$, it holds

$$\|u_n - f \circ \rho_{z_n}\|_{L^p(\mu_n)} \rightarrow 0. \tag{7.3}$$

Proof. We only need to prove that $f \circ \rho_{z_n}$ converges L^p -strong to $f \circ \rho_{z_0}$, then (7.3) follows from the linearity of the L^p -convergence (2.14).

The assumptions on f imply that f is uniformly continuous and we denote by $\omega : [0, \infty) \rightarrow [0, \infty)$ a global modulus of continuity for f . Observe that f is also bounded. In the sequel, we fix (Z, d) a realization of the convergence and recall that $d|_{Y_n \times Y_n} = \rho_n$. We can estimate

$$\int |f \circ d_{z_0} - f \circ d_{z_n}|^p d\mu_n$$

$$\begin{aligned} &\leq \int_{B_R(z_n)} |f \circ d_{z_0} - f \circ d_{z_n}|^p d\mu_n + 2^p \int_{Z \setminus B_R(z_n)} |f|^p \circ d_{z_0} + |f|^p \circ d_{z_n} d\mu_n \\ &\leq \mu_n(B_R(z_n))\omega(d(z_0, z_n))^p + 2^p C \int_{B_R(z_n)^c} d(z_0, \cdot)^{-\alpha} + d(z_n, \cdot)^{-\alpha} d\mu_n \\ &\leq \mu_n(B_R(z_n))\omega(d(z_0, z_n))^p + 2^p C \cdot 2^\alpha \sup_n \int_{B_R(z_n)^c} d(z_n, \cdot)^{-\alpha} d\mu_n, \end{aligned}$$

where in the last step we assume that n is big enough so that $d(z_n, z_0) < R/2$, which ensures $d^{-1}(z_0, \cdot) \leq 2d^{-1}(z_n, \cdot)$ in $B_R(z_n)^c$. Since $\sup_n \mu_n(B_R(z_0)) < +\infty$ for every $R > 0$, by the pmGH-convergence, we can send first $n \uparrow \infty$ and then $R \uparrow \infty$ to obtain $\|f \circ d_{z_0} - f \circ d_{z_n}\|_{L^p(\mu_n)} \rightarrow 0$. Fix $\varphi \in C_{bs}(Z)$ and $R > 0$ so that $\text{supp}(\varphi) \subset B_R(z_0)$, then

$$\begin{aligned} &\left| \int \varphi f \circ d_{z_0} d\mu - \int \varphi f \circ d_{z_n} d\mu_n \right| \leq \\ &\leq \|\varphi\|_\infty \int_{\text{supp}\varphi} |f \circ d_{z_0} - f \circ d_{z_n}| d\mu_n + \left| \int \varphi f \circ d_{z_0} d\mu_n - \int \varphi f \circ d_{z_0} d\mu \right| \\ &\leq \|\varphi\|_\infty \mu_n(B_R(z_0))^{1-1/p} \|f \circ d_{z_0} - f \circ d_{z_n}\|_{L^p(\mu_n)} + \left| \int \varphi f \circ d_{z_0} d\mu_n - \int \varphi f \circ d_{z_0} d\mu \right|. \end{aligned}$$

Sending $n \uparrow \infty$ we obtain that $f \circ d_{z_n} d\mu_n \rightharpoonup f \circ d_{z_0} \mu$ in duality with $C_{bs}(Z)$. It remains to prove that $\|f \circ \rho_{z_n}\|_{L^p(\mu_n)} \rightarrow \|f \circ \rho_{z_0}\|_{L^p(\mu)}$. Since $\|f \circ d_{z_0} - f \circ d_{z_n}\|_{L^p(\mu_n)} \rightarrow 0$, it is enough to show that $\|f \circ d_{z_0}\|_{L^p(\mu_n)} \rightarrow \|f \circ \rho_{z_0}\|_{L^p(\mu)}$. Clearly $\|f \circ \rho_{z_0}\|_{L^p(\mu)} = \|f \circ d_{z_0}\|_{L^p(\mu)} \leq \underline{\lim}_n \|f \circ d_{z_0}\|_{L^p(\mu_n)}$, hence we only need to show $\|f \circ d_{z_0}\|_{L^p(\mu)} \geq \overline{\lim}_n \|f \circ d_{z_0}\|_{L^p(\mu_n)}$. We can assume n is big enough so that $d(z_0, z_n) \leq 1$. For every $R \geq 4$ fix a cut-off function $\varphi_R \in C_{bs}(Z)$, $0 \leq \varphi_R \leq 1$, such that $\varphi_R \equiv 1$ in $B_R(z_0)$ and with support in $B_{2R}(z_0)$. Then

$$\int |\varphi_R(|f|^p \circ d_{z_0}) - |f|^p \circ d_{z_0}| d\mu_n \leq \int_{B_R(z_0)^c} |f|^p \circ d_{z_0} d\mu_n \leq 2 \cdot 2^\alpha C \sup_n \int_{B_{R/2}(z_n)^c} d_{z_n}^{-\alpha} d\mu_n,$$

where we have used that $B_R(z_0)^c \subset B_{R/2}(z_n)^c$ and $d_{z_0}^{-1} \leq 2d_{z_n}^{-1}$ in $B_{R/2}(z_n)^c$. This shows that

$$\left| \int \varphi_R(|f|^p \circ d_{z_0}) - |f|^p \circ d_{z_0} d\mu_n \right| \leq \varepsilon_R \rightarrow 0, \quad \text{as } R \uparrow \infty,$$

where ε_R is independent of n . Therefore

$$-\varepsilon_R + \overline{\lim}_n \int |f|^p \circ d_{z_0} d\mu_n \leq \overline{\lim}_n \int \varphi_R(|f|^p \circ d_{z_0}) d\mu_n = \int \varphi_R(|f|^p \circ d_{z_0}) d\mu \leq \int |f|^p \circ \rho_{z_0} d\mu.$$

Sending R to infinity, we conclude the proof. \square

The second result of this section is a technical fact that will play a key role in the proof of our main theorem. It states that a Euclidean bubble which is strongly concentrated around a point is close to a spherical bubble.

Lemma 7.3. *For every $N \in (2, \infty)$, there are constants $C_N, \alpha = \alpha(N) > 0$ such that the following holds. Given $\sigma \geq 1$, set $2^* = 2N/(N - 2)$ and*

$$f_{\text{eu}}(t) := \frac{\sigma^{\frac{N-2}{2}}}{(1 + \sigma^2 t^2)^{\frac{N-2}{2}}}, \quad f_{\text{sphere}}(t) := \frac{\sigma^{\frac{N-2}{2}}}{(1 + 2\sigma^2(1 - \cos(t)))^{\frac{N-2}{2}}}, \quad t \in [0, \pi].$$

Let (X, d, \mathbf{m}) be $\text{RCD}(N - 1, N)$, $z \in X$, $d_z(\cdot) := d(z, \cdot)$ and $v := \sigma^N \mathbf{m}(B_{\sigma^{-1}}(z))$. Then

$$\|(f_{\text{eu}} - f_{\text{sphere}})(d_z)\|_{L^{2^*}(\mathbf{m})} + \|\nabla(f_{\text{eu}} - f_{\text{sphere}})(d_z)\|_{L^2(\mathbf{m})} \leq C_N \sigma^{-\alpha} (\sqrt{v} + 1).$$

Proof. We fix $\eta \in (0, 1)$ to be chosen later. Denote $B := B_{\frac{1}{\eta\sigma}}(z)$. In what follows $C_N > 0$ is a constant depending only on N , its value may vary from line to line without notice and without being relabeled. By Bishop-Gromov and the assumptions, we get

$$\mathbf{m}(B) \leq v(\eta\sigma)^{-N}. \tag{7.4}$$

We divide the proof into two steps, one for the L^{2^*} -norm and one for the L^2 -norm of the gradient.

STEP 1. We start estimating

$$\begin{aligned} \|(f_{\text{eu}} - f_{\text{sphere}})(d_z)\|_{L^{2^*}(\mathbf{m})} &\leq \|(f_{\text{eu}} - f_{\text{sphere}})(d_z)\|_{L^{2^*}(B)} \\ &\quad + \|f_{\text{sphere}}(d_z)\|_{L^{2^*}(B^c)} + \|f_{\text{eu}}(d_z)\|_{L^{2^*}(B^c)} =: \text{I} + \text{II} + \text{III}. \end{aligned}$$

We analyze each term separately. We start with I. Recall that

$$|2(1 - \cos(t)) - t^2| \leq ct^4, \quad 1 - \cos(t) \leq ct^2, \quad \forall t \geq 0,$$

for some numerical constant $c > 0$. Using $\| |x|^p - |y|^p \| \leq C_p |x - y| (|x|^{p-1} + |y|^{p-1})$ with $p = (N - 2)/2$ and the above estimates we have for all $t \in [0, (\eta\sigma)^{-1}]$ the following:

$$\begin{aligned} &|f_{\text{eu}} - f_{\text{sphere}}|(t) \\ &\leq \frac{C_N \sigma |2(1 - \cos(t)) - t^2| \left(\left| \frac{1}{\sigma} + 2\sigma(1 - \cos(t)) \right|^{\frac{N-2}{2}-1} + \left| \frac{1}{\sigma} + \sigma t^2 \right|^{\frac{N-2}{2}-1} \right)}{\left(\frac{1}{\sigma} + \sigma t^2 \right)^{\frac{N-2}{2}} \left(\frac{1}{\sigma} + 2\sigma(1 - \cos(t)) \right)^{\frac{N-2}{2}}} \\ &\leq C_N \frac{\sigma \frac{1}{(\eta\sigma)^4} \cdot (\sigma^{-1} + (\eta^2\sigma)^{-1})^{\frac{N-2}{2}-1}}{\sigma^{2-N}} \leq C_N \eta^{-N} \sigma^{\frac{N-2}{2}-2}. \end{aligned}$$

This and (7.4) directly implies that

$$(I)^{2^*} = \int_{B_{\frac{1}{\eta\sigma}}(z_n)} |f_{eu} - f_{sphere}|^{2^*}(\mathbf{d}_z) \, \mathbf{d}\mathbf{m} \leq C_N v \eta^{-N(2^*+1)} \sigma^{-2 \cdot 2^*}.$$

We pass to II. Note that $|f_{sphere}(t)|^{2^*}, |f_{eu}(t)|^{2^*} \leq C_N \sigma^{-N} t^{-2N}$, having used that $1 - \cos(t) \geq ct^2$ in $[0, \pi]$ for some numerical constant $c > 0$. Hence applying Lemma 7.1 and using (7.4)

$$(II)^{2^*} + (III)^{2^*} \leq C_N v \sigma^{-N} (\sigma \eta)^N \leq v C_N \eta^N.$$

STEP 2. From the chain rule for the gradient and the fact that $|\nabla \mathbf{d}(z, \cdot)| = 1$ m-a.e., we have

$$\begin{aligned} \|\nabla(f_{eu} - f_{sphere})(\mathbf{d}_z)\|_{L^2(\mathbf{m})} &\leq \|(f'_{eu} - f'_{sphere})(\mathbf{d}_z)\|_{L^2(B)} \\ &\quad + \|f'_{sphere}(\mathbf{d}_z)\|_{L^2(B^c)} + \|f'_{eu}(\mathbf{d}_z)\|_{L^2(B^c)} =: I' + II' + III'. \end{aligned}$$

We start with I'_n . Reasoning similarly to Step 1, we can estimate for all $t \in [0, (\eta\sigma)^{-1}]$

$$\begin{aligned} |f'_{eu} - f'_{sphere}|(t) &= (N - 2)\sigma \left| \frac{t(\frac{1}{\sigma} + 2\sigma(1 - \cos(t)))^{\frac{N}{2}} - \sin(t)(\frac{1}{\sigma} + \sigma t^2)^{\frac{N}{2}}}{(\frac{1}{\sigma} + \sigma t^2)^{\frac{N}{2}}(\frac{1}{\sigma} + 2\sigma(1 - \cos(t)))^{\frac{N}{2}}} \right| \\ &\leq C_N \sigma^{N+2} t \left| 2(1 - \cos(t)) - t^2 \right| \left(\left(\frac{1}{\sigma} + 2\sigma(1 - \cos(t)) \right)^{\frac{N}{2}-1} \right. \\ &\quad \left. + \left(\frac{1}{\sigma} + \sigma t^2 \right)^{\frac{N}{2}-1} \right) + C_N \sigma^{N+1} |\sin(t) - t| \left(\frac{1}{\sigma} + \sigma t^2 \right)^{\frac{N}{2}} \\ &\leq C_N \sigma^{N+2} t^5 \left(\frac{1}{\sigma} + \frac{1}{\sigma \eta^2} \right)^{\frac{N}{2}-1} + C_N \sigma^{N+1} t^3 \left(\frac{1}{\sigma} + \frac{1}{\sigma \eta^2} \right)^{\frac{N}{2}-1} \\ &\leq C_N \sigma^{\frac{N}{2}-2} \eta^{-N-3}. \end{aligned}$$

Therefore, again using (7.4) we deduce $(II'_n)^2 \leq C v \eta^{-3N-6} \sigma^{-4}$. As above we can directly estimate

$$|f'_{eu}|, |f'_{sphere}|^2 \leq C_N \sigma^{-N+2} t^{2-2N}, \quad t \in [0, \pi],$$

having used $|\sin(t)| \leq ct$ and $1 - \cos(t) \geq ct^2$ in $[0, \pi]$. Hence by Lemma 7.1 and using (7.4)

$$(II'_n)^2 + (III'_n)^2 \leq C_N v \sigma^{-N+2} (\sigma \eta)^{N-2} \leq v C_N \eta^{N-2}.$$

Combining all cases and taking $\eta := \sigma^{-\beta}$ with $\beta > 0$ small enough depending on N we conclude, using also that $v^{1/2^*} + v^{1/2} \leq 2 + 2\sqrt{v}$. \square

8. Proof of the main results

8.1. Stability in the compact case

In this part, we prove the main qualitative stability result of this note. Note that this proves our main Theorem 1.1. We will also provide a proof of Corollary 1.3 at the end.

Given $N > 2$ the family of spherical bubbles in a metric space (X, d) is denoted by

$$\mathcal{M}_{\text{sphere}}(X) := \{a(1 - b \cos d(x, z_0))^{\frac{2-N}{2}} : a \in \mathbb{R}, b \in (0, 1), z_0 \in X\} \cup \{u \equiv a : a \in \mathbb{R}\}.$$

Theorem 8.1. *For every $\varepsilon > 0$ and $N \in (2, \infty)$ there exists $\delta := \delta(\varepsilon, N) > 0$ such that the following holds. Let (X, d, m) be an $\text{RCD}(N - 1, N)$ space for some $N \in (2, \infty)$ with $m(X) = 1$, set $2^* = 2N/(N - 2)$ and suppose that there exists $u \in W^{1,2}(X)$ non-constant satisfying*

$$\frac{\|u\|_{L^{2^*}(m)}^2 - \|u\|_{L^2(m)}^2}{\|\nabla u\|_{L^2(m)}^2} > \frac{2^* - 2}{N} - \delta. \tag{8.1}$$

Then there exists $w \in \mathcal{M}_{\text{sphere}}(X)$ such that

$$\frac{\|\nabla(u - w)\|_{L^2(m)} + \|u - w\|_{L^{2^*}(m)}}{\|u\|_{L^{2^*}(m)}} \leq \varepsilon. \tag{8.2}$$

Moreover if $w \equiv a \in \mathbb{R}$, then $a \in \mathbb{R}$ can be chosen so that the remainder

$$R := u - a$$

satisfies for some $x \in X$

$$\|R \cdot \|R\|_{L^2}^{-1} - \sqrt{N + 1} \cos(d(\cdot, x))\|_{L^2} \leq C_N(\varepsilon^\alpha + \delta)^\beta, \tag{8.3}$$

for some positive constants α, β, C_N depending only on N .

Proof. By scaling invariance, it is not restrictive to assume $\|u\|_{L^{2^*}(m)} = 1$. We only need to prove the first part, as the second follows from Proposition 8.3 below.

We argue by contradiction and suppose that there exist $\varepsilon > 0$, a sequence of $\text{RCD}(N - 1, N)$ spaces (X_n, d_n, m_n) and non-constant functions $u_n \in W^{1,2}(X_n)$ with $\|u_n\|_{L^{2^*}(m_n)} = 1$ so that

$$\|u_n\|_{L^{2^*}(m_n)}^2 \geq \tilde{A}_n \|\nabla u_n\|_{L^2(m_n)} + \|u_n\|_{L^2(m_n)}^2, \tag{8.4}$$

with $\tilde{A}_n \rightarrow \frac{2^*-2}{N}$ and satisfying

$$\inf_{w \in \mathcal{M}_{\text{sphere}}(X_n)} \|\nabla(u_n - w)\|_{L^2(\mathfrak{m}_n)} + \|u_n - w\|_{L^{2^*}(\mathfrak{m}_n)} > \varepsilon, \quad \forall n \in \mathbb{N}. \tag{8.5}$$

Let us fix $\eta < (\eta_N \wedge \frac{1}{3})$, where η_N is as in Theorem 6.2. For every n there exist $y_n \in X_n$ and $t_n < \text{diam}(X_n)$ such that

$$1 - \eta = \int_{B_{t_n}(y_n)} |u_n|^{2^*} \, d\mathfrak{m}_n = \sup_{y \in X_n} \int_{B_{t_n}(y)} |u_n|^{2^*} \, d\mathfrak{m}_n, \quad \forall n \in \mathbb{N}. \tag{8.6}$$

This follows directly by Bishop Gromov inequality and the properness of the space. Define now $\sigma_n := t_n^{-1}$ and consider the sequence $(Y_n, \rho_n, \mu_n, y_n) := (X_{\sigma_n}, \mathfrak{d}_{\sigma_n}, \mathfrak{m}_{\sigma_n}, y_n)$, where $\mathfrak{d}_{\sigma_n} := \sigma_n \mathfrak{d}_n$, $\mathfrak{m}_{\sigma_n} := \sigma_n^N \mathfrak{m}_n$ and $u_{\sigma_n} := \sigma_n^{-N/2^*} u_n \in W^{1,2}(Y_n)$. In particular, by scaling, it holds that

$$1 - \eta = \int_{B_1(y_n)} |u_{\sigma_n}|^{2^*} \, d\mu_n = \sup_{y \in Y_n} \int_{B_1(y)} |u_{\sigma_n}|^{2^*} \, d\mathfrak{m}_n, \tag{8.7}$$

and also

$$1 = \|u_{\sigma_n}\|_{L^{2^*}(\mu_n)}^2 \geq \tilde{A}_n \|\nabla u_{\sigma_n}\|_{L^2(\mu_n)}^2 + \sigma_n^{-2} \|u_{\sigma_n}\|_{L^2(\mu_n)}^2, \tag{8.8}$$

for all $n \in \mathbb{N}$. Moreover, Y_n supports a Sobolev inequality with constants $A_n = (2^* - 2)/N$, $B_n = \sigma_n^{-2}$. Since $t_n \leq \text{diam}(X_n) \leq \pi$, up to a subsequence we have that $\lim_n \sigma_n = \sigma \in [\pi^{-1}, +\infty]$ and, consequently, that $B_n \rightarrow \sigma^{-2} \in [0, \pi^2]$. Thanks to [88, Theorem 1.10] and up to passing to a subsequence, we can assume that $\text{diam}(X_n) \geq \pi/2$ and in particular that $\text{diam}(Y_n) \geq \sigma_n \pi/2$. Moreover $\mu_n(B_1(y_n)) \leq \mu_n(Y_n) = \sigma_n^N$. Hence, up to a subsequence and no matter the value of σ , the hypotheses of Theorem 6.2 are satisfied. Applying Theorem 6.2 we get that, up to a further subsequence, Y_n pmGH-converge to a pointed metric measure space (Y, ρ, μ, \bar{y}) and that u_{σ_n} converges L^{2^*} -strong to some $u \in W^{1,2}(Y)$ satisfying

$$\|u\|_{L^{2^*}(\mu)}^2 = \frac{2^* - 2}{N} \|\nabla u\|_{L^2(\mu)}^2 + \sigma^{-2} \|u\|_{L^2(\mu)}^2,$$

(where it is intended that $\sigma^{-2} \|u\|_{L^2(\mu)}^2 = 0$ if $\sigma = \infty$ and $u \notin L^2(\mu)$). We distinguish now two cases, depending on the value of σ .

CASE 1: $\sigma < \infty$. In this case, $B_n \rightarrow B := \sigma^{-2} > 0$ and Y_n are compact of uniformly bounded diameter. Therefore, Y_n mGH-converges to Y and u_{σ_n} converges also $W^{1,2}$ -strong to $u \in W^{1,2}(Y)$ (recall *ii*) in Theorem 6.2 when $B > 0$). Define $(X_\infty, \mathfrak{d}_\infty, \mathfrak{m}_\infty) := (Y, \rho/\sigma, \mu/\sigma^N)$ so that X_∞ is a RCD($N - 1, N$) space with $\mathfrak{m}_\infty(X_\infty) = 1$. By *iii*) in Theorem 6.2 the function $v := \sigma^{N/2^*} u \in W^{1,2}(X_\infty)$ satisfies

$$\|v\|_{L^{2^*}(\mathfrak{m}_\infty)}^2 = \frac{2^* - 2}{N} \|\nabla v\|_{L^2(\mathfrak{m}_\infty)}^2 + \|v\|_{L^2(\mathfrak{m}_\infty)}^2.$$

Here, we distinguish two situations: v is constant, or not. If v is constant, then $v \equiv 1$ and $u \equiv \sigma^{-N/2^*}$. By linearity of convergence (2.14), $u_{\sigma_n} - \sigma^{-N/2^*}$ converges $W^{1,2}$ -strong and L^{2^*} -strong to zero so, by scaling, we reach

$$\begin{aligned} 0 &= \lim_{n \rightarrow \infty} \|\nabla(u_{\sigma_n} - \sigma^{-N/2^*})\|_{L^2(\mathfrak{m}_{\sigma_n})}^2 + \|u_{\sigma_n} - \sigma^{-N/2^*}\|_{L^{2^*}(\mathfrak{m}_{\sigma_n})}^2 \\ &= \lim_{n \rightarrow \infty} \|\nabla(u_n - (\sigma_n/\sigma)^{N/2^*})\|_{L^2(\mathfrak{m}_n)}^2 + \|u_n - (\sigma_n/\sigma)^{N/2^*}\|_{L^{2^*}(\mathfrak{m}_n)}^2. \end{aligned}$$

This yields a contradiction with (8.5).

If v is not constant, by Theorem 5.2, there exist $a \in \mathbb{R}, b \in (0, 1), z_0 \in X_\infty$ so that

$$v(x) = \frac{a}{(1 - b \cos(\mathfrak{d}_\infty(x, z_0)))^{\frac{N-2}{2}}}, \quad \forall x \in X_\infty.$$

Denoting $f(t) := a(1 - b \cos(t))^{\frac{2-N}{2}}$ for $t \in [0, \pi]$, it is clear that $u = \tilde{f} \circ \rho(\cdot, z_0)$, where $\tilde{f}(s) := \sigma^{-N/2^*} f(\sigma^{-1}s)$, $s \in \mathbb{R}$. Take now a sequence $z_n \rightarrow z_0$ GH-converging and invoke Lemma 7.2 (here (7.2) is trivially satisfied by equi-boundedness of the diameters) to get that $\tilde{f} \circ \mathfrak{d}_{\sigma_n}(\cdot, z_n)$ converges L^{2^*} -strong to u and

$$\lim_n \|u_{\sigma_n} - \tilde{f} \circ \mathfrak{d}_{\sigma_n}(\cdot, z_n)\|_{L^{2^*}(\mathfrak{m}_{\sigma_n})} = 0. \tag{8.9}$$

We want to scale back this information to the original sequence u_n . Simple estimates and triangular inequalities give

$$\begin{aligned} \overline{\lim}_{n \rightarrow \infty} \|u_n - f \circ \mathfrak{d}_n(\cdot, z_n)\|_{L^{2^*}(\mathfrak{m}_n)} &= \overline{\lim}_{n \rightarrow \infty} \|\sigma_n^{-N/2^*} (u_n - f \circ \mathfrak{d}_n(\cdot, z_n))\|_{L^{2^*}(\mathfrak{m}_{\sigma_n})} \\ &\leq \overline{\lim}_{n \rightarrow \infty} \|u_{\sigma_n} - \sigma^{-N/2^*} f \circ \mathfrak{d}_n(\cdot, z_n)\|_{L^{2^*}(\mathfrak{m}_{\sigma_n})} \\ &\quad + |\sigma^{-N/2^*} - \sigma_n^{-N/2^*}| \|f \circ \mathfrak{d}_n(\cdot, z_n)\|_{L^{2^*}(\mathfrak{m}_{\sigma_n})} \\ &\leq \overline{\lim}_{n \rightarrow \infty} \|u_{\sigma_n} - \tilde{f} \circ \mathfrak{d}_{\sigma_n}(\cdot, z_n)\|_{L^{2^*}(\mathfrak{m}_{\sigma_n})} + C_{f,\sigma} \overline{\lim}_{n \rightarrow \infty} |\sigma^{-1} - \sigma_n^{-1}| \stackrel{(8.9)}{=} 0, \end{aligned}$$

using that \tilde{f} is bounded and that σ_n is away from zero. We pass now to the gradient norm. From the chain rule of weak gradients and the fact that $|\nabla \rho(\cdot, z_0)| = 1$ μ -a.e., we have $|\nabla u| = |\tilde{f}'| \circ \rho(\cdot, z_0)$ μ -a.e. and similarly $|\nabla(\tilde{f} \circ \mathfrak{d}_{\sigma_n}(\cdot, z_n))| = |\tilde{f}'| \circ \mathfrak{d}_{\sigma_n}(\cdot, z_n)$ at \mathfrak{m}_{σ_n} -a.e. point. In particular again by Lemma 7.2 we have that $|\tilde{f}'| \circ \mathfrak{d}_{\sigma_n}(\cdot, z_n)$ converges L^2 -strong to $|\nabla u|$. This means that the convergence of $\tilde{f} \circ \mathfrak{d}_{\sigma_n}(\cdot, z_0)$ to u is $W^{1,2}$ -strong. Moreover, as we said above, also u_{σ_n} $W^{1,2}$ -strong converges to u . This together with Lemma A.5 and (2.2) gives

$$\lim_{n \rightarrow \infty} \|\nabla(u_{\sigma_n} - \tilde{f} \circ \mathfrak{d}_{\sigma_n}(\cdot, z_n))\|_{L^2(\mathfrak{m}_{\sigma_n})}^2 = 0. \tag{8.10}$$

Arguing as above for the 2^* -norm we can scale back the above information to obtain

$$\lim_{n \rightarrow \infty} \|\nabla(u_n - f \circ d_n(\cdot, z_n))\|_{L^2(\mathfrak{m}_n)} = 0.$$

We omit the computation since it is analogous. Since $f \circ d_n(\cdot, z_n) \in \mathcal{M}_{\text{sphere}}(X_n)$, we again reached a contradiction with (8.5).

CASE 2: $\sigma = \infty$. Here $B_n \rightarrow B := 0$ and we know that (Y, ρ, μ, \bar{y}) supports a Sobolev inequality (S) with constants $A = (2^* - 2)/N, B = 0$. In particular, $\text{AVR}(Y) > 0$ thanks to [88, Theorem 4.6] and $\sqrt{A} \geq \text{Eucl}(N, 2)\text{AVR}(Y)^{-1/N}$ by sharpness in (2.13). The sequence u_{σ_n} (that we recall is L^{2^*} -strong converging to some $u \in L^{2^*}(\mu)$) is so that $\|\nabla u_{\sigma_n}\|_{L^2(\mathfrak{m}_{\sigma_n})} \rightarrow \|\nabla u\|_{L^2(\mu)}$, hence

$$\text{Eucl}(N, 2)\text{AVR}(Y)^{-1/N}\|\nabla u\|_{L^2(\mu)} \geq \|u\|_{L^{2^*}(\mu)} = \sqrt{A}\|\nabla u\|_{L^2(\mu)}.$$

Therefore $\sqrt{A} = \text{Eucl}(N, 2)\text{AVR}(Y)^{-1/N}$ (recall that u is non-zero) and in particular $\text{AVR}(Y)$ depends only on N . Recalling the rigidity in Theorem 5.3 we get that Y is isomorphic to an N -Euclidean metric measure cone with tip z_0 and u is radial of the following form

$$u(y) = \frac{a}{(1 + b\rho^2(y, z_0))^{\frac{N-2}{2}}}, \quad y \in Y,$$

for some $a \in \mathbb{R}, b > 0$.

Pick now a sequence $z_n \in Y_n$ with $z_n \rightarrow z_0$ in Z . Note that, since $z_n \rightarrow z_0$ and z_0 is a tip of Y , by pmGH convergence we have

$$\lim_n \sigma_n^N \mathfrak{m}_n(B_{1/\sigma_n}(z_n)) = \lim_n \mathfrak{m}_{\sigma_n}(B_1(z_n)) = \mu(B_1(z_0)) = \text{AVR}(Y)\omega_N.$$

Hence up to a subsequence, since $\text{AVR}(Y)$ depends only on N , for every n it holds

$$\mathfrak{m}_{\sigma_n}(B_1(z_n)) = \mathfrak{m}_n(B_{\sigma_n^{-1}}(z_n))\sigma_n^N \leq C_N. \tag{8.11}$$

Denote

$$f(t) := \frac{a}{(1 + bt^2)^{\frac{N-2}{2}}}, \quad t \geq 0.$$

Note that $|f|^{2^*}, |f'|^2 \leq Ct^{-2N+2}$ and for every $R \geq 1$,

$$\int_{B_R(z_n)^c} d_{\sigma_n}(\cdot, z_n)^{-2N+2} d\mathfrak{m}_{\sigma_n} \stackrel{(7.1)}{\leq} C_N \mathfrak{m}_{\sigma_n}(B_1(z_n))R^{-N+2} \stackrel{(8.11)}{\leq} C_N R^{-N+2}. \tag{8.12}$$

Hence assumption (7.2) in Lemma 7.2 is satisfied for Y_n and both f', f and we can apply the result twice to get that $f \circ d_{\sigma_n}(\cdot, z_n)$ converges L^{2^*} -strong to u , that

$$\lim_{n \rightarrow \infty} \|u_{\sigma_n} - f \circ d_{\sigma_n}(\cdot, z_n)\|_{L^{2^*}(\mathfrak{m}_{\sigma_n})} \stackrel{(7.3)}{=} 0, \tag{8.13}$$

and that $|f'| \circ d_{\sigma_n}(\cdot, z_n)$ converges L^2 -strong to $|\nabla u| = |f'| \circ \rho(\cdot, z_0)$. By Lemma A.5 and the convergence of the gradient norms, we immediately get from the parallelogram identity

$$\lim_{n \rightarrow \infty} \int |\nabla(u_{\sigma_n} - f \circ d_{\sigma_n}(\cdot, z_n))|^2 d\mathfrak{m}_{\sigma_n} = 0. \tag{8.14}$$

Scaling all back to X_n we can rewrite the above convergences as

$$\lim_{n \rightarrow \infty} \|u_n - f_n \circ d_n(\cdot, z_n)\|_{L^{2^*}(\mathfrak{m}_n)} + \|\nabla(u_n - f_n \circ d_n(\cdot, z_n))\|_{L^{2^*}(\mathfrak{m}_n)} = 0,$$

where

$$f_n := \frac{ab^{\frac{2-N}{4}}(\sqrt{b}\sigma_n)^{\frac{N-2}{2}}}{(1 + (\sqrt{b}\sigma_n)^2 t^2)^{\frac{N-2}{2}}}.$$

Using (8.11) we deduce

$$\mathfrak{m}_n(B_{(\sqrt{b}\sigma_n)^{-1}}(z_n))(\sqrt{b}\sigma_n)^N \leq C_N(\sqrt{b} \vee 1)^N.$$

This is obvious if $b \geq 1$, while for $b \leq 1$ it follows by the Bishop-Gromov inequality. Having this density bound, we can now apply Lemma 7.3 to get

$$\lim_{n \rightarrow \infty} \|u_n - g_n \circ d_n(\cdot, z_n)\|_{L^{2^*}(\mathfrak{m}_n)} + \|\nabla(u_n - g_n \circ d_n(\cdot, z_n))\|_{L^{2^*}(\mathfrak{m}_n)} = 0,$$

where

$$g_n := \frac{ab^{\frac{2-N}{4}}(\sqrt{b}\sigma_n)^{\frac{N-2}{2}}}{(1 + (\sqrt{b}\sigma_n)^2 - (\sqrt{b}\sigma_n)^2 \cos(t))^{\frac{N-2}{2}}}.$$

Multiplying and dividing by $1 + (\sqrt{b}\sigma_n)^2$ shows that $g_n \circ d_n(\cdot, z_n) \in \mathcal{M}_{\text{sphere}}(X_n)$ and gives a contradiction with (8.5). Having examined all the possible cases, the proof is now concluded. \square

Remark 8.2. It is evident from the proof that (8.2) holds true assuming only that

$$\|u\|_{L^{2^*}(\mathfrak{m})}^2 \geq A\|\nabla u\|_{L^2(\mathfrak{m})}^2 + B\|u\|_{L^2(\mathfrak{m})}^2,$$

with $|A - \frac{2^*-2}{N}| + |B - 1| < \delta$, which is a weaker assumption than (8.1). Indeed, the starting point of the argument is the reverse Sobolev inequality (8.4) (for u_n) and adding here a sequence $B_n \rightarrow 1$ in front of $\|u_n\|_{L^2(\mathfrak{m}_n)}^2$ does not influence the subsequent steps. \blacksquare

Proposition 8.3. *Let (X, d, \mathbf{m}) be an $\text{RCD}(N - 1, N)$ space, $N > 2$, with $\mathbf{m}(X) = 1$ and set $2^* = 2N/(N - 2)$. Let $u \in W^{1,2}(X)$ be non-constant and set*

$$\delta := \frac{2^* - 2}{N} - \frac{\|u\|_{L^{2^*(\mathbf{m})}}^2 - \|u\|_{L^2(\mathbf{m})}^2}{\|\nabla u\|_{L^2(\mathbf{m})}^2} \geq 0.$$

Then setting $g := u - \int u$ we have for some $x \in X$

$$\|g\|_{L^2(\mathbf{m})} \|g\|_{L^{2^*(\mathbf{m})}}^{-1} - \sqrt{N + 1} \cos(d(\cdot, x))\|_{L^2(\mathbf{m})} \leq C_N((\|\nabla u\|_{L^2(\mathbf{m})} \|u\|_{L^{2^*(\mathbf{m})}}^{-1})^\alpha + \delta)^\beta, \tag{8.15}$$

for some positive constants α, β depending only on N .

Proof. We can clearly assume that $\int u = 1$. Moreover we can assume that $\|\nabla u\|_{L^2(\mathbf{m})} \leq \varepsilon_N \|u\|_{L^{2^*(\mathbf{m})}}$ for some small constant $\varepsilon_N > 0$, otherwise the statement is trivial. Analogously we can assume that δ is small with respect to N . By the Sobolev and the Poincaré inequalities, provided ε_N is small enough, we have $\|u\|_{L^{2^*}} \leq 2$. Set $g := u - \int u$. Then by [88, Lemma 6.7] and the Poincaré inequality we have, provided δ and ε_N are small enough,

$$\left| N - \frac{\int |\nabla g|^2 d\mathbf{m}}{\int g^2 d\mathbf{m}} \right| \leq C_N(\|\nabla u\|_{L^2(\mathbf{m})}^\alpha + \delta) \leq \tilde{C}_N((\|\nabla u\|_{L^2(\mathbf{m})} \|u\|_{L^{2^*(\mathbf{m})}}^{-1})^\alpha + \delta),$$

for some $\alpha > 0$ depending only on N . Now (8.15) follows directly from the quantitative Obata theorem in [38] (there, written for Lipschitz functions but by density in $W^{1,2}$, the statement directly extends to Sobolev functions recalling (2.4)). \square

We conclude this part with the proof of the stability result for the Yamabe minimizers in the smooth setting.

Proof of Corollary 1.3. Take as in the hypotheses (M, g) so that $\text{Ric}_g \geq n - 1$ and $d_{GH}(M, \mathbb{S}^n) \leq \delta$. Let $u \in W^{1,2}(M)$ non-zero satisfying $|\mathcal{E}(u) - Y(M, g)| \leq \delta$. Set ν the renormalized volume measure. Since $\text{Scal}_g \geq n(n - 1)$, we have by the Sobolev inequality (1.6) that

$$1 \leq \frac{\frac{2^*-2}{n} \|\nabla u\|_{L^2(\nu)}^2 + \|u\|_{L^2(\nu)}^2}{\|u\|_{L^{2^*(\nu)}}^2} \leq \frac{\mathcal{E}(u)}{n(n - 1)\text{Vol}_g(M)^{2/n}} \leq \frac{(Y(M, g) + \delta)}{n(n - 1)\text{Vol}_g(M)^{2/n}},$$

where the norms are computed using the renormalized volume measure. Recall also that by [41] we have that $\text{Vol}_g(M) \geq (1 - \varepsilon')\text{Vol}(\mathbb{S}^n)$, where $\varepsilon' = \varepsilon'(\delta, n)$ goes to zero as $\delta \rightarrow 0$. This in particular gives that $Y(M, g) \geq c(n) > 0$ if δ is chosen small enough (depending on n). Therefore, combining the above with the inequality (see [16])

$$Y(M, g) \leq Y(\mathbb{S}^n) = n(n - 1)\text{Vol}(\mathbb{S}^n)^{2/n}$$

gives

$$\frac{2^{*-2} \|\nabla u\|_{L^2(\nu)}^2 + \|u\|_{L^2(\nu)}^2}{\|u\|_{L^{2^*}(\nu)}^2} \leq (1 + \delta c(n)^{-1})(1 - \varepsilon')^{-2/n}.$$

The conclusion now follows applying Theorem 1.1 (in the stronger version given by (1.11)). \square

8.2. Stability in the non-compact case

We now prove the qualitative stability result for the sharp Euclidean-type Sobolev inequality. Note that this proves also Theorem 1.4. Given $N > 2$, the family of Euclidean bubbles in a metric space (X, d) is denoted by

$$\mathcal{M}_{\text{eu}}(X) := \{a(1 + bd^2(x, z_0))^{\frac{2-N}{2}} : a \in \mathbb{R}, b > 0, z_0 \in X\}.$$

Theorem 8.4. *For every $\varepsilon > 0, V \in (0, 1)$ and $N \in (2, \infty)$, there exists $\delta := \delta(\varepsilon, N, V) > 0$ such that the following holds. Let (X, d, m) be an $\text{RCD}(0, N)$ space with $\text{AVR}(X) \in (V, V^{-1})$ and, setting $2^* = 2N/(N - 2)$, assume there exists $u \in W_{\text{loc}}^{1,2}(X) \cap L^{2^*}(m)$ non constant with $m(|u| > t) < \infty$ for every $t > 0$ satisfying*

$$\frac{\|u\|_{L^{2^*}(m)}}{\|\nabla u\|_{L^2(m)}} > \text{AVR}(X)^{-\frac{1}{N}} \text{Eucl}(N, 2) - \delta.$$

Then, there exists $v \in \mathcal{M}_{\text{eu}}(X)$ so that

$$\frac{\|\nabla(u - v)\|_{L^2(m)}}{\|\nabla u\|_{L^2(m)}} \leq \varepsilon.$$

Proof. We can clearly assume that $\|u\|_{L^{2^*}(m)} = 1$. Moreover by approximation it is also sufficient to prove the statement for $u \in W^{1,2}(X)$ (see Lemma 3.2).

We proceed by contradiction and suppose that there exist $\varepsilon > 0$, a sequence (X_n, d_n, m_n) of $\text{RCD}(0, N)$ spaces with $\text{AVR}(X_n) \in (V, V^{-1})$ and a sequence $u_n \in W^{1,2}(X_n) \cap L^{2^*}(m_n)$ of non-constant functions satisfying

$$\|u_n\|_{L^{2^*}(m_n)} \geq (A_n - 1/n) \|\nabla u_n\|_{L^2(m_n)}, \tag{8.16}$$

where $A_n := \text{AVR}(X_n)^{-\frac{1}{N}} \text{Eucl}(N, 2)$, and

$$\inf_{v \in \mathcal{M}_{\text{eu}}(X_n)} \frac{\|\nabla(u_n - v)\|_{L^2(m_n)}}{\|\nabla u_n\|_{L^2(m_n)}} > \varepsilon, \quad \forall n \in \mathbb{N}. \tag{8.17}$$

For every $\eta \in (0, 1)$, let $y_n \in X_n$ and $t_n > 0$ so that (arguing as for (8.6))

$$1 - \eta = \int_{B_{t_n}(y_n)} |u_n|^{2^*} \, d\mathbf{m}_n = \sup_{y \in X_n} \int_{B_{t_n}(y)} |u_n|^{2^*} \, d\mathbf{m}_n, \quad n \in \mathbb{N}.$$

Define now $\sigma_n := t_n^{-1}$ and $(Y_n, \rho_n, \mu_n, y_n) := (X_{\sigma_n}, \mathbf{d}_{\sigma_n}, \mathbf{m}_{\sigma_n}, y_n)$, where $\mathbf{d}_{\sigma_n} := \sigma_n \mathbf{d}_n$, $\mathbf{m}_{\sigma_n} := \sigma_n^N \mathbf{m}_n$ and $u_{\sigma_n} := \sigma_n^{-N/2^*} u_n \in W^{1,2}(Y_n)$. In particular, by scaling, for every $n \in \mathbb{N}$ we have

$$1 - \eta = \int_{B_1(y_n)} |u_{\sigma_n}|^{2^*} \, d\mu_n \quad \text{and} \quad \|u_{\sigma_n}\|_{L^{2^*}(\mu_n)} \geq (A_n - 1/n) \|\nabla u_{\sigma_n}\|_{L^2(\mu_n)}.$$

By the assumption, we have the uniform bounds $2V^{\frac{1}{N}} \text{Eucl}(N, 2) \leq A_n \leq 2V^{-\frac{1}{N}} \text{Eucl}(N, 2)$. Thus, up to subsequences, we can clearly suppose that $A_n \rightarrow A$, for some $A > 0$ finite. We can now invoke Theorem 6.2 (the assumptions are satisfied as $\text{diam}(Y_n) = +\infty$) with $\eta := \eta_N/2$ and get that up to a subsequence $(Y_n, \rho_n, \mu_n, y_n)$ pmGH-converges to some $\text{RCD}(0, N)$ space (Y, ρ, μ, \bar{y}) supporting a Sobolev inequality (S) with constant $A > 0, B = 0$. Moreover we have L^{2^*} -strong convergence of u_{σ_n} to a function $u \in W_{loc}^{1,2}(Y)$ attaining equality in this said Sobolev inequality and $\|\nabla u_{\sigma_n}\|_{L^2(\mathbf{m}_{\sigma_n})} \rightarrow \|\nabla u\|_{L^2(\mu)}$. From [88, Theorem 4.6] we have $\text{AVR}(Y) = (\text{Eucl}(N, 2)/A)^N$ and in particular u satisfies the assumptions of Theorem 5.3, which gives that Y is isomorphic to a N -Euclidean metric measure cone with tip z_0 and

$$u(y) = \frac{a}{(1 + b\rho^2(y, z_0))^{\frac{N-2}{2}}}, \quad y \in Y,$$

for suitable $a \in \mathbb{R}, b > 0$.

Take any $z_n \rightarrow z_0$. Then up to subsequence we can assume that $\mathbf{m}_{\sigma_n}(B_1(z_n)) \leq C_N \text{AVR}(Y)$ hold for every n . Writing $f(t) := a(1 + bt^2)^{\frac{2-N}{2}}$ for every $t \in \mathbb{R}^+$, recalling $|f|^{2^*}, |f'|^2 \leq Ct^{-2N+2}$ and arguing as for (8.12), we see that all the hypotheses of Lemma 7.2 are fulfilled both for $f \circ \rho(\cdot, z_0)$ and for $f' \circ \rho(\cdot, z_0)$. We therefore apply Lemma 7.2 twice to get that $f \circ \mathbf{d}_{\sigma_n}(\cdot, z_n)$ converges L^{2^*} -strong to u and that $|f'| \circ \mathbf{d}_{\sigma_n}(\cdot, z_n)$ converges L^2 -strong to $|\nabla u|$. We can thus combine Lemma A.5 with the convergence of the gradient norms to deduce, from the parallelogram identity, that

$$\lim_{n \rightarrow \infty} \|\nabla(u_{\sigma_n} - f \circ \mathbf{d}_{\sigma_n}(\cdot, z_n))\|_{L^2(\mathbf{m}_{\sigma_n})} = 0. \tag{8.18}$$

Scaling back, (8.18) becomes

$$\lim_{n \rightarrow \infty} \|\nabla(u_n - (\sigma_n^{N/2^*} f) \circ (\sigma_n \mathbf{d}_n(\cdot, z_n)))\|_{L^2(\mathbf{m}_n)} = 0.$$

This means that the sequence $v_n := a\sigma_n^{N/2^*} (1 + b\sigma_n^2 \mathbf{d}_n(\cdot, z_n)^2)^{\frac{2-N}{2}} \in \mathcal{M}_{\text{eu}}(X_n)$, satisfies

$$\overline{\lim}_{n \rightarrow \infty} \frac{\|\nabla(u_n - v_n)\|_{L^2(\mathbf{m}_n)}}{\|\nabla u_n\|_{L^2(\mathbf{m}_n)}} = 0,$$

having used that $\|\nabla u_n\|_{L^2(\mathfrak{m}_n)} \geq C_N \text{AVR}(X_n)^{1/N} \|u_n\|_{L^{2^*}} \geq C_N V^{1/N}$. This is a contradiction with (8.17) and concludes the proof. \square

From the above stability, the next corollary directly follows (proving also Corollary 1.5).

Corollary 8.5. *Let (X, d, \mathfrak{m}) be an $\text{RCD}(0, N)$ space with $N \in (2, \infty)$, $\text{AVR}(X) > 0$. Then*

$$\text{AVR}(X)^{\frac{1}{N}} \text{Eucl}^{-1}(N, 2) = \inf_{v \in \mathcal{M}_{\text{eu}}(X)} \frac{\|\nabla v\|_{L^2(\mathfrak{m})}}{\|v\|_{L^{2^*}(\mathfrak{m})}}.$$

Acknowledgments

F.N. is supported by the Academy of Finland Grant No. 314789. I.Y.V. is supported by the Academy of Finland projects *Incidences on Fractals*, Grant No. 321896 and *Singular integrals, harmonic functions, and boundary regularity in Heisenberg groups*, Grant No. 328846.

Appendix A. Concentration compactness: non-compact case

Here we extend the concentration compactness tools for a sequence of converging RCD spaces (developed in [88] in compact setting) to the non-compact case. The main difference is that here mass can also escape to infinity and so we need an additional result (see Lemma A.6). Some additional technical convergence results will be also needed and proved in Section A.1.

A.1. Technical convergence lemmas

Throughout this part we fix a sequence $(X_n, d_n, \mathfrak{m}_n, x_n)$ of pointed $\text{RCD}(K, N)$ spaces, $n \in \mathbb{N} \cup \{\infty\}$, for some $K \in \mathbb{R}, N \in (1, \infty)$ with $X_n \xrightarrow{pmGH} X_\infty$. We also fix a proper metric space (Z, d) realizing the convergence via extrinsic approach [58] (see Section 2.4). We start with a version of the Brezis-Lieb Lemma [30].

Lemma A.1 (*Brezis-Lieb type Lemma*). *Let $q, q' \in (1, \infty)$ and suppose that $u_n \in L^q(\mathfrak{m}_n)$ satisfy $\sup_n \|u_n\|_{L^q(\mathfrak{m}_n)} < +\infty$ and that u_n converges in $L^{q'}$ -strong to some $u_\infty \in L^{q'} \cap L^q(\mathfrak{m}_\infty)$. Then, for any sequence $v_n \in L^q(\mathfrak{m}_n)$ such that $v_n \rightarrow u_\infty$ strongly both in $L^{q'}$ and L^q , it holds*

$$\lim_{n \rightarrow \infty} \int |u_n|^q d\mathfrak{m}_n - \int |u_n - v_n|^q d\mathfrak{m}_n = \int |u_\infty|^q d\mathfrak{m}_\infty. \tag{A.1}$$

Proof. The proof is the same as in [88, Prop. 6.2]. Even if the argument there is done assuming finite reference measure, it is used only at the end when applying the Hölder

inequality. In that step here is enough to multiply by an arbitrary $\varphi \in C_{bs}(Z)$ and argue in the same way. (Note also that the assumptions $q \in [2, \infty)$ and $q' \in (1, q)$, even if present in the statement of [88, Prop. 6.2] are actually not used in its proof). \square

We shall need an alternative version of the semicontinuity result (2.16) to deal with locally Sobolev functions; we include a proof since we could not find it in the literature.

Lemma A.2. *Let $p \in (1, \infty)$ and suppose $(f_n) \subset W_{loc}^{1,2}(X_n)$ is L^p -strong converging to f_∞ . Then*

$$\|\nabla f_\infty\|_{L^2(\mathbf{m}_\infty)}^2 \leq \varliminf_{n \rightarrow \infty} \|\nabla f_n\|_{L^2(\mathbf{m}_n)}^2, \tag{A.2}$$

(meaning that, if the right hand side is finite, then $f_\infty \in W_{loc}^{1,2}(X_\infty)$ and (A.2) holds).

Proof. Since $|f_n| \rightarrow |f_\infty|$ L^p -strongly (see [8, a] in Prop. 3.3]) and $|\nabla f_n| = |\nabla|f_n||$ \mathbf{m} -a.e. for every f_n , without loss of generality we can suppose f_n, f_∞ nonnegative. If the \liminf in (A.2) is infinite, there is nothing to prove. So, let us assume that it is finite. For every $k \in \mathbb{N}$, we consider $\varphi^k \in \text{LIP}([0, \infty)$ with $\text{Lip}(\varphi^k) \leq 1$, $\varphi^k(0) = 0$, converging point-wise to the identity as $k \uparrow \infty$ and such that $\{\varphi^k(f_n)\}_n$ is L^2 -bounded. For instance we can take $\varphi^k(t) := (t - 1/k)^+ \wedge k$, indeed

$$\|\varphi^k(f_n)\|_{L^2(\mathbf{m}_n)}^2 \leq k^2 \mathbf{m}_n(\{f_n > 1/k\}) \leq k^{2+p} \|f_n\|_{L^p(\mathbf{m}_n)}^p,$$

for every $n \in \mathbb{N}$. Again by [8, a] in Prop. 3.3], we have $\varphi^k(f_n)$ is L^p -strong convergent to $\varphi^k(f_\infty)$. Moreover is also L^2 -bounded, thus it is also L^2 -weak convergent to $\varphi^k(f_\infty)$. Then, by (2.16) we have $\varphi^k(f_\infty) \in W^{1,2}(X_\infty)$ and

$$\|\nabla(\varphi^k(f_\infty))\|_{L^2(\mathbf{m}_\infty)}^2 \leq \varliminf_{n \rightarrow \infty} \|\nabla(\varphi^k(f_n))\|_{L^2(\mathbf{m}_n)}^2 \leq \varliminf_{n \rightarrow \infty} \|\nabla f_n\|_{L^2(\mathbf{m}_n)}^2 < \infty,$$

having used the fact that φ^k is 1-Lipschitz. By arbitrariness of $k > 0$ and since $\varphi^k(f_\infty) \rightarrow f_\infty$ pointwise, we see by semicontinuity (2.1) that (A.2) follows. \square

The following lemma allows extracting L_{loc}^2 -converging subsequences from $W^{1,2}$ -boundedness.

Lemma A.3. *Let $p \geq 2$ and suppose $u_n \in W_{loc}^{1,2}(X_n)$ converges L^p -weak to $u_\infty \in L^p(\mathbf{m}_\infty)$ and $\sup_n \|\nabla u_n\|_{L^2(\mathbf{m}_n)} < \infty$. Then, up to a subsequence u_n converges L_{loc}^2 -strong to $u_\infty \in W_{loc}^{1,2}(X_\infty)$ with $|\nabla u_\infty| \in L^2(\mathbf{m}_\infty)$.*

Proof. We first prove the L_{loc}^2 convergence. Consider $\varphi \in \text{Lip}_{bs}(Z)$ (recall that (Z, d) is a space realizing the convergence). Since $\sup_n \mathbf{m}_n(B_R(x_n)) < +\infty$, for every $R > 0$, by Hölder inequality we have $\sup_n \|\varphi u_n\|_{L^2(\mathbf{m}_n)} < +\infty$. Analogously using the Leibniz

rule, $\varphi u_n \in W^{1,2}(X_n)$ with $\sup_n \|\nabla(\varphi u_n)\|_{L^2(\mathfrak{m}_n)} < \infty$. Thus there exists a subsequence (n_k) (see [58, Theorem 6.3]) such that φu_{n_k} converges L^2 -strong to some v , which must be equal to φu_∞ by uniqueness of weak limits. Hence the whole sequence φu_n is L^2 -strongly convergent to φu_∞ . The fact that $u_\infty \in W^{1,2}_{loc}(X_\infty)$ follows by the Mosco convergence of the Cheeger energies (see (2.16)), indeed for every $\varphi \in \text{LIP}_{bs}(Z)$, $\text{Ch}(\varphi u_\infty) \leq \varliminf_n \text{Ch}(\varphi u_n) < \infty$. It remains to prove that $|\nabla u_\infty| \in L^2(\mathfrak{m}_\infty)$. Fix a ball $B \subset Z$ and take $\varphi \in \text{Lip}_{bs}(Z)$ equal to 1 on B . Using [8, Lemma 5.8], we have $\int_B |\nabla u_\infty|^2 \, d\mathfrak{m}_\infty = \int_B |\nabla(\varphi u_\infty)| \, d\mathfrak{m}_\infty \leq \varliminf_n \int_B |\nabla(\varphi u_n)|^2 \, d\mathfrak{m}_n \leq \sup_n \|\nabla u_n\|_{L^2(\mathfrak{m}_n)} < \infty$. Where in the first and last step we used the locality of the gradient. By the arbitrariness of B this implies $|\nabla u_\infty| \in L^2(\mathfrak{m}_\infty)$. \square

Lemma A.4. *Let $p \geq 2$ and $u_\infty \in W^{1,2}_{loc}(X_\infty) \cap L^p(\mathfrak{m}_\infty)$ with $|\nabla u_\infty| \in L^2(\mathfrak{m}_\infty)$. Then, there exists a sequence $u_n \in W^{1,2}_{loc}(X_n) \cap L^p(\mathfrak{m}_n)$ that converges L^p and L^2_{loc} -strong to u_∞ and so that $|\nabla u_n|$ converges L^2 -strong to $|\nabla u_\infty|$.*

Proof. By Lemma 3.2 there exists a sequence $u_n \in W^{1,2}(X_\infty) \cap L^p(\mathfrak{m}_\infty)$ such that $u_n \rightarrow u_\infty$ in $L^p(\mathfrak{m}_\infty)$ and $|\nabla u_n| \rightarrow |\nabla u_\infty|$ in $L^2(\mathfrak{m}_\infty)$. From [88, Lemma 6.4] (there written for compact spaces, but the same proof works in the present setting) there exists a sequence $u_n^k \in W^{1,2}(X_n)$ that converges L^p and $W^{1,2}$ -strong to u_n . By [8, Theorem 5.7] this implies that $|\nabla u_n^k|$ converges L^2 -strong to $|\nabla(\eta_k u_n)|$. The conclusion then follows via diagonal argument. Finally the L^2_{loc} -strong convergence follows from Lemma A.3. \square

We prove a convergence result for pairings (the case $p = 2$ follows from [8, Theorem 5.4]).

Lemma A.5. *Let $p \in [2, \infty)$ and $u_n, v_n \in L^p(\mathfrak{m}_n) \cap W^{1,2}_{loc}(X_n)$ be converging L^p -strong to u_∞, v_∞ respectively. Suppose that $u_\infty \in W^{1,2}_{loc}(X_\infty)$, that $\|\nabla u_n\|_{L^2(\mathfrak{m}_n)} \rightarrow \|\nabla u_\infty\|_{L^2(\mathfrak{m}_\infty)} < +\infty$ and $\overline{\lim}_n \|\nabla v_n\|_{L^2(\mathfrak{m}_n)} < +\infty$. Then $v_\infty \in W^{1,2}_{loc}(X_\infty)$, $|\nabla v_\infty| \in L^2(\mathfrak{m}_\infty)$ and*

$$\lim_{n \rightarrow \infty} \int \langle \nabla u_n, \nabla v_n \rangle \, d\mathfrak{m}_n = \int \langle \nabla u_\infty, \nabla v_\infty \rangle \, d\mathfrak{m}_\infty.$$

Proof. The fact that $v_\infty \in W^{1,2}_{loc}(X_\infty)$ with $|\nabla v_\infty| \in L^2(\mathfrak{m}_\infty)$ follows from Lemma A.2. In particular by Cauchy-Schwarz $\langle \nabla u_\infty, \nabla v_\infty \rangle \in L^1(\mathfrak{m}_\infty)$. Let $t > 0$ and notice that $u_n + tv_n$ converges L^p -strong to $u_\infty + tv_\infty$ by (2.14). Applying again Lemma A.2 we have $u_\infty + tv_\infty \in W^{1,2}_{loc}(X_\infty)$ and

$$\begin{aligned} \int 2t \langle \nabla u_\infty, \nabla v_\infty \rangle + |\nabla u_\infty|^2 + t^2 |\nabla v_\infty|^2 \, d\mathfrak{m}_\infty &= \int |\nabla(u_\infty + tv_\infty)|^2 \, d\mathfrak{m}_\infty \\ &\stackrel{(A.2)}{\leq} \varliminf_{n \rightarrow \infty} \int |\nabla(u_n + tv_n)|^2 \, d\mathfrak{m}_n \\ &\leq 2t \varliminf_{n \rightarrow \infty} \int \langle \nabla u_n, \nabla v_n \rangle \, d\mathfrak{m}_n + 2 \overline{\lim}_n \int |\nabla v_n|^2 \, d\mathfrak{m}_n + \int |\nabla u_\infty|^2 \, d\mathfrak{m}_\infty. \end{aligned}$$

Simplifying $\int |\nabla u_\infty|^2 dm_\infty$, dividing by t and sending $t \downarrow 0$ we obtain $\int \langle \nabla u_\infty, \nabla v_\infty \rangle dm_\infty \leq \underline{\lim}_{n \rightarrow \infty} \int \langle \nabla u_n, \nabla v_n \rangle dm_n$. Arguing analogously for $t < 0$, we conclude. \square

A.2. Concentration compactness principles

Here we briefly extend two concentration compactness principles from [80,81] (see also [94]) for general sequences of probabilities on metric measure spaces.

The first deal with an arbitrary sequence of probability measures on varying ambient space. Compare also with the version [12, Lemma 2.1].

Lemma A.6. *Let (Z, d) be a complete and separable metric spaces and let $\nu_n \in \mathcal{P}(Z)$, for $n \in \mathbb{N}$. Then, up to a subsequence, one of the following holds:*

- i) COMPACTNESS. *There exists $(z_n) \subset Z$ such that for all $\varepsilon > 0$, there exists $R > 0$ satisfying*

$$\nu_n(B_R(z_n)) \geq 1 - \varepsilon, \quad \forall n \in \mathbb{N}.$$

- ii) VANISHING.

$$\lim_{n \rightarrow \infty} \sup_{z \in Z} \nu_n(B_R(z)) = 0, \quad \forall R > 0.$$

- iii) DICHOTOMY. *There exists $\lambda \in (0, 1)$ with $\lambda \geq \overline{\lim}_n \sup_{z \in Z} \nu_n(B_R(z))$, for all $R > 0$, so that: there exists $R_n \uparrow \infty$, $(z_n) \subset Z$ and there are ν_n^1, ν_n^2 two non-negative Borel measures satisfying*

$$\begin{aligned} 0 &\leq \nu_n^1 + \nu_n^2 \leq \nu_n, \\ \text{supp}(\nu_n^1) &\subset B_{R_n}(z_n), \quad \text{supp}(\nu_n^2) \subset Z \setminus B_{10R_n}(z_n), \\ \overline{\lim}_{n \rightarrow \infty} |\lambda - \nu_n^1(Z)| + |(1 - \lambda) - \nu_n^2(Z)| &= 0. \end{aligned}$$

The above can be obtained arguing exactly as in [94, Lemma I in Section 4.3] and therefore its proof is omitted. We briefly comment on the difference in case iii) with respect to [94]: our formulation of case iii) using a sequence R_n follows from the one used in [94] (where R is fixed depending on a parameter $\varepsilon > 0$) with a diagonal argument (this is observed also in the proof of [94, Theorem 4.9]); the condition $\lambda \geq \overline{\lim}_n \sup_{z \in Z} \nu_n(B_R(z))$ (not present in [94]) instead can be directly checked to hold by the way λ is chosen in the proof.

The second principle is a concentration compactness result for the Sobolev embedding stating that concentration may occur only at countably-many points. With respect to [88, Lemma 6.6], here we extend the principle to deal with varying pmGH-convergent RCD spaces (hence, the difference arises when considering noncompact limit spaces).

Lemma A.7. Let (X_n, d_n, m_n, x_n) , $n \in \mathbb{N} \cup \{\infty\}$, be pointed RCD(K, N) spaces, $K \in \mathbb{R}$, $N \in (1, \infty)$ with $X_n \xrightarrow{pmGH} X_\infty$ and assume that X_n supports a Sobolev inequality **(S)** with uniformly bounded constants $A_n > 0, B_n \geq 0$.

Suppose further that $u_n \in W_{loc}^{1,2}(X_n) \cap L^{2^*}(m_n)$ with $\sup_n \|\nabla u_n\|_{L^2(m_n)} < \infty$ is L_{loc}^2 -strong converging to $u_\infty \in L^{2^*}(m_\infty)$ and suppose that $|\nabla u_n|^2 m_n \rightharpoonup \omega$, $|u_n|^{2^*} m_n \rightharpoonup \nu$ in duality with $C_{bs}(Z)$ and $C_b(Z)$, respectively (where (Z, d) is a fixed realization of the convergence).

Then, $u_\infty \in W_{loc}^{1,2}(X_\infty)$ with $|\nabla u_\infty| \in L^2(m_\infty)$ and:

- i) there exists a countable set of indices J , points $(x_j)_{j \in J} \subset X_\infty$ and weights $(\nu_j)_{j \in J} \subset \mathbb{R}^+$ so that

$$\nu = |u_\infty|^{2^*} m_\infty + \sum_{j \in J} \nu_j \delta_{x_j};$$

- ii) there exists $(\omega_j)_{j \in J} \subset \mathbb{R}^+$ satisfying $\nu_j^{2/2^*} \leq (\overline{\lim}_n A_n) \omega_j$ and such that

$$\omega \geq |\nabla u_\infty|^2 m_\infty + \sum_{j \in J} \omega_j \delta_{x_j}.$$

In particular, we have $\sum_j \nu_j^{2/2^*} < \infty$.

Proof. We subdivide the proof into two steps.

STEP 1. Suppose first that $u_\infty = 0$. Then, the conclusion follows arguing as in Step 1 of [88, Lemma 6.6] taking here φ a Lipschitz and boundedly supported (instead of only Lipschitz) cut-off and using the assumed L_{loc}^2 -strong convergence.

STEP 2. For general u_∞ , the idea is to apply the above to ‘ $u_\infty - u_n$ ’ and then use a Brezis-Lieb lemma to recover the information for u_∞ . Take \tilde{u}_n a recovery sequence given by Lemma A.4 for u_∞ . Thus, for every $\varphi \in \text{Lip}_{bs}(Z)^+$, we have φu_n is L^2 -strong to φu_∞ and L^{2^*} -bounded and $\varphi \tilde{u}_n$ is L^2 and L^{2^*} -strong convergent to φu_∞ . Therefore Lemma A.1 ensures

$$\lim_{n \rightarrow \infty} \int |\varphi|^{2^*} |u_n|^{2^*} dm_n - \int |\varphi|^{2^*} |u_n - \tilde{u}_n|^{2^*} dm_n = \int |\varphi|^{2^*} |u_\infty|^{2^*} dm_\infty. \tag{A.3}$$

Now define $v_n := u_n - \tilde{u}_n$ and notice that all the assumptions ensure that v_n is L_{loc}^2 -strong and L^{2^*} -weak convergent to zero. From the bounds $|v_n|^{2^*} \leq 2^{2^*} (|u_n|^{2^*} + |\tilde{u}_n|^{2^*})$ and $|\nabla v_n|^2 \leq 2(|\nabla u_n|^2 + |\nabla \tilde{u}_n|^2)$ by tightness we can extract a not relabeled subsequence where $|v_n|^{2^*} m_n$ converge in duality with $C_b(Z)$ to $\bar{\nu}$ and $|\nabla v_n|^2 m_n$ converge in duality with $C_{bs}(Z)$ to a finite Borel measure $\bar{\omega}$. Then from Step 1, i), ii) hold true for (v_n) , for suitable weights $(\nu_j), (\omega_j) \subset \mathbb{R}^+$ and points $(x_j) \subset X_\infty$. Then passing to the limit in (A.3)

$$\int \varphi^{2^*} d\nu - \int \varphi^{2^*} d\bar{\nu} = \int \varphi^{2^*} |u_\infty|^{2^*} dm_\infty, \quad \forall \varphi \in \text{Lip}_{bs}(Z)^+.$$

This in turn implies $\nu = |u_\infty|^{2^*} m_\infty + \bar{\nu} = |u_\infty|^{2^*} m_\infty + \sum_j \nu_j \delta_{x_j}$ that is point i). We pass to prove ii) and therefore we need to show separately that

$$\begin{aligned} \omega(\{x_j\}) &= \bar{\omega}(\{x_j\}) \geq \omega_j, \quad \forall j \in J, \\ \omega &\geq |\nabla u_\infty|^2 m_\infty. \end{aligned}$$

The first can be verified arguing exactly as in Step 2 of [88, Lemma 6.6] replacing the usage of [8, Theorem 5.7] with Lemma A.4 above. For the second, we fix $\varphi \in C_{bs}(Z)$, $\varphi \geq 0$, and $\chi \in \text{LIP}_{bs}(Z)$ be such that $\chi = 1$ in $\text{supp}(\varphi)$. It is easy to check that χu_n is $W^{1,2}$ -weak converging to χu_∞ (recall that $u_n \rightarrow u_\infty$ in L^2_{loc}). Then, [8, Lemma 5.8] ensures that

$$\begin{aligned} \int \varphi |\nabla u_\infty|^2 dm_\infty &= \int \varphi |\nabla(\chi u_\infty)|^2 dm_\infty \\ &\leq \varliminf_{n \rightarrow \infty} \int \varphi |\nabla(\chi u_n)|^2 dm_n = \varliminf_{n \rightarrow \infty} \int \varphi |\nabla u_n|^2 dm_n \end{aligned}$$

By arbitrariness of φ , we showed ii) and the proof is now concluded. \square

Appendix B. Technical results

In this appendix, we collect basic results about Sobolev inequalities and a version of the chain rule for the weak upper gradient.

Lemma B.1. *Let (X, d, m) be an $\text{RCD}(K, N)$ space, $N \in (2, \infty)$, $K \in \mathbb{R}$, satisfying for $A > 0$*

$$\|u\|_{L^{2^*}(m)} \leq A \|\nabla u\|_{L^2(m)}, \quad \forall u \in \text{LIP}_c(X), \tag{B.1}$$

where $2^* := \frac{2N}{N-2}$. Then (B.1) holds also for all $u \in W^{1,2}_{loc}(X)$ satisfying $m(\{|u| > t\}) < +\infty$ for all $t > 0$.

Proof. It is enough to prove (B.1) for non-negative functions. First note that (B.1) holds for every $u \in W^{1,2}(X)$, by density in energy of Lipschitz functions [5] and by the lower semicontinuity of the L^{2^*} -norm with respect to L^2 -convergence. For a general $u \geq 0$ as in the hypotheses, if $\int |\nabla u|^2 dm = +\infty$ there is nothing to prove, otherwise take $u_n := ((u-1/n)^+) \wedge n \in W^{1,2}(X)$ (since $u_n, |\nabla u_n| \in L^2(m)$) and then send $n \rightarrow +\infty$. \square

Lemma B.2 (Local Sobolev embedding). *Let (X, d, m) be an $\text{RCD}(K, N)$ space for some $K \in \mathbb{R}, N \in (2, \infty)$ and set $2^* := 2N/(N-2)$. Then exists $\tilde{r}_{K^-, N} > 0$ (with $\tilde{r}_{0, N} = +\infty$) such that for every $B_R(x) \subsetneq X$, $R \leq \tilde{r}_{K^-, N}$ it holds*

$$\|u\|_{L^{2^*}(\mathfrak{m})} \leq \frac{C_{N,K}R}{\mathfrak{m}(B_R(x))^{1/N}} \|\nabla u\|_{L^2(\mathfrak{m})}, \quad \forall u \in W_0^{1,2}(B_{R/2}(x)). \tag{B.2}$$

Proof. It is enough to prove the statement for $u \in \text{LIP}_c(B_{R/2}(x))$. Thanks to the uniformly locally doubling property of (X, d, \mathfrak{m}) and the validity of a local $(1, 1)$ -Poincaré inequality ([91]), from the results in [65] the following Sobolev-Poincaré inequality holds

$$\left(\int_{B_R(x)} |f - f_{B_R(x)}|^{2^*} \, d\mathfrak{m} \right)^{\frac{1}{2^*}} \leq C(N, K, R_0)R \left(\int_{B_{2R}(x)} |\nabla f|^2 \, d\mathfrak{m} \right)^{\frac{1}{2}}, \quad \forall f \in \text{LIP}(X), \tag{B.3}$$

for every $R \leq R_0$ and where $f_{B_R(x)} := \int_{B_R(x)} f \, d\mathfrak{m}$ (see also [26]). Moreover if $K \geq 0$, the constant $C(N, K, R_0)$ can be taken independent of R_0 .

Hence applying (B.3) to $u \in \text{LIP}_c(B_{R/2}(x))$ we can write

$$\begin{aligned} & \left(\int_{B_R(x)} |u|^{2^*} \, d\mathfrak{m} \right)^{\frac{1}{2^*}} \\ & \leq C_{N,K}R \frac{\mathfrak{m}(B_R(x))^{1/2^*}}{\mathfrak{m}(B_{2R}(x))^{1/2}} \left(\int_{B_{2R}(x)} |\nabla u|^2 \right)^{\frac{1}{2}} + \mathfrak{m}(B_R(x))^{1/2^*-1} \int_{B_{R/2}(x)} |u| \, d\mathfrak{m} \\ & \leq C_{N,K}R \mathfrak{m}(B_R(x))^{-1/N} \left(\int_{B_{2R}(x)} |\nabla u|^2 \right)^{\frac{1}{2}} + \frac{\mathfrak{m}(B_{R/2}(x))^{1-1/2^*}}{\mathfrak{m}(B_R(x))^{1-1/2^*}} \left(\int_{B_{R/2}(x)} |u|^{2^*} \, d\mathfrak{m} \right)^{\frac{1}{2^*}}, \end{aligned}$$

where we have used that $\text{supp}(u) \subset B_{R/2}(x)$. Thanks to the reverse doubling inequality (recall (2.5)), assuming $R \leq R_{K^-,N}$, we can absorb the rightmost term inside the left-hand side of the above to obtain (B.2) as desired. \square

A technical result needed in this note is a chain rule for the composition with an absolutely continuous function φ , which we could not find in the literature (see [55] or [59] for the classical one with φ Lipschitz).

Lemma B.3 (*Chain rule for composition with AC-functions*). *Let (X, d, \mathfrak{m}) be a proper metric measure space and $u \in \text{LIP}_{loc}(\Omega)$ with $\Omega \subset X$ open. Let $\varphi \in \text{AC}_{loc}(I)$ with I open interval such that $u(\Omega') \subset\subset I$ for every $\Omega' \subset\subset \Omega$. Suppose also that $|\varphi'(u)| |\nabla u| \in L^2_{loc}(\Omega)$.*

Then $\varphi(u) \in W^{1,2}_{loc}(\Omega)$ and $|\nabla \varphi(u)| = |\varphi'(u)| |\nabla u|$ \mathfrak{m} -a.e.

Proof. Up to subtracting a constant, we can assume that $0 \in I$ and $\varphi(0) = 0$. Then with a cut-off argument we can reduce to the case when $u \in \text{LIP}_c(X)$ and $\varphi \in \text{AC}(\mathbb{R})$ with compact support and $\varphi(0) = 0$. We argue by approximation and define functions $\varphi_n \in \text{LIP}(\mathbb{R})$ by

$$\varphi_n(t) := \int_0^t -n \vee \varphi'(s) \wedge n \, ds.$$

Clearly $\varphi_n \rightarrow \varphi$ pointwise in \mathbb{R} . By the usual chain rule for Lipschitz composition we have that $\varphi_n(u) \in W^{1,2}(X)$ with $|\nabla \varphi_n(u)| = |\varphi'_n(u)| |\nabla u| \leq |\varphi'(u)| |\nabla u|$, \mathfrak{m} -a.e., where we have used that $|\varphi'_n| \leq |\varphi'|$ a.e. In particular the sequence $|\nabla \varphi_n(u)|$ is bounded in $L^2(\mathfrak{m})$. Moreover $\varphi_n(u) \rightarrow \varphi(u)$ pointwise and from the lower semicontinuity of the minimal weak upper gradient (see, e.g., [59, Prop. 2.1.13]) we deduce that $\varphi(u) \in W^{1,2}(X)$ and

$$|\nabla \varphi(u)| \leq |\varphi'(u)| |\nabla u|, \quad \mathfrak{m}\text{-a.e.} \quad (\text{B.4})$$

The equality in (B.4) then follows with a standard argument (see e.g. [59, Theorem 2.1.28]). \square

References

- [1] V. Agostiniani, M. Fogagnolo, L. Mazzieri, Sharp geometric inequalities for closed hypersurfaces in manifolds with nonnegative Ricci curvature, *Invent. Math.* 222 (2020) 1033–1101.
- [2] L. Ambrosio, Calculus, heat flow and curvature-dimension bounds in metric measure spaces, in: *Proceedings of the International Congress of Mathematicians—Rio de Janeiro 2018. vol. I. Plenary Lectures*, World Sci. Publ., Hackensack, NJ, 2018, pp. 301–340.
- [3] L. Ambrosio, S. Di Marino, Equivalent definitions of BV space and of total variation on metric measure spaces, *J. Funct. Anal.* 266 (2014) 4150–4188.
- [4] L. Ambrosio, N. Gigli, A. Mondino, T. Rajala, Riemannian Ricci curvature lower bounds in metric measure spaces with σ -finite measure, *Trans. Am. Math. Soc.* 367 (2012) 4661–4701.
- [5] L. Ambrosio, N. Gigli, G. Savaré, Density of Lipschitz functions and equivalence of weak gradients in metric measure spaces, *Rev. Mat. Iberoam.* 29 (2013) 969–996.
- [6] L. Ambrosio, N. Gigli, G. Savaré, Metric measure spaces with Riemannian Ricci curvature bounded from below, *Duke Math. J.* 163 (2014) 1405–1490.
- [7] L. Ambrosio, N. Gigli, G. Savaré, Bakry-Émery curvature-dimension condition and Riemannian Ricci curvature bounds, *Ann. Probab.* 43 (2015) 339–404.
- [8] L. Ambrosio, S. Honda, New stability results for sequences of metric measure spaces with uniform Ricci bounds from below, in: *Measure Theory in Non-smooth Spaces*, in: *Partial Differ. Equ. Meas. Theory*, De Gruyter Open, Warsaw, 2017, pp. 1–51.
- [9] L. Ambrosio, A. Mondino, G. Savaré, On the Bakry-Émery condition, the gradient estimates and the Local-to-Global property of $RCD^*(K, N)$ metric measure spaces, *J. Geom. Anal.* 26 (2014) 1–33.
- [10] G. Antonelli, E. Bruè, M. Fogagnolo, M. Pozzetta, On the existence of isoperimetric regions in manifolds with nonnegative Ricci curvature and Euclidean volume growth, *Calc. Var. Partial Differ. Equ.* 61 (2022) 77, 40 pp.
- [11] G. Antonelli, M. Fogagnolo, M. Pozzetta, The isoperimetric problem on Riemannian manifolds via Gromov-Hausdorff asymptotic analysis, *Commun. Contemp. Math.* (2022), <https://doi.org/10.1142/S0219199722500687>.
- [12] G. Antonelli, S. Nardulli, M. Pozzetta, The isoperimetric problem *via* direct method in noncompact metric measure spaces with lower Ricci bounds, *ESAIM Control Optim. Calc. Var.* 28 (2022) 57, 32 pp.
- [13] G. Antonelli, E. Pasqualetto, M. Pozzetta, D. Semola, Asymptotic isoperimetry on non collapsed spaces with lower Ricci bounds, *Math. Ann.* (2023), <https://doi.org/10.1007/s00208-023-02674-y>.
- [14] G. Antonelli, E. Pasqualetto, M. Pozzetta, I.Y. Violo, Topological regularity of isoperimetric sets in PI spaces having a deformation property, *Proc. R. Soc. Edinb. Sect. A Math.* (2023), <https://doi.org/10.1017/prm.2023.105>, published online; arXiv:2303.01280.
- [15] T. Aubin, Équations différentielles non linéaires et problème de Yamabe concernant la courbure scalaire, *J. Math. Pures Appl.* (9) 55 (1976) 269–296.

- [16] T. Aubin, Problèmes isopérimétriques et espaces de Sobolev, *J. Differ. Geom.* 11 (1976) 573–598.
- [17] T. Aubin, *Nonlinear Analysis on Manifolds. Monge-Ampère Equations*, Grundlehren der Mathematischen Wissenschaften (Fundamental Principles of Mathematical Sciences), vol. 252, Springer-Verlag, New York, 1982.
- [18] K. Bacher, K.-T. Sturm, Localization and tensorization properties of the curvature-dimension condition for metric measure spaces, *J. Funct. Anal.* 259 (2010) 28–56.
- [19] D. Bakry, L’hypercontractivité et son utilisation en théorie des semigroupes, in: *Lectures on Probability Theory, Saint-Flour, 1992*, in: *Lecture Notes in Math.*, vol. 1581, Springer, Berlin, 1994, pp. 1–114.
- [20] D. Bakry, I. Gentil, M. Ledoux, *Analysis and Geometry of Markov Diffusion Operators*, Grundlehren der Mathematischen Wissenschaften (Fundamental Principles of Mathematical Sciences), vol. 348, Springer, Cham, 2014.
- [21] D. Bakry, M. Ledoux, Sobolev inequalities and Myers’s diameter theorem for an abstract Markov generator, *Duke Math. J.* 85 (1996) 253–270.
- [22] Z.M. Balogh, A. Kristály, Sharp isoperimetric and Sobolev inequalities in spaces with nonnegative Ricci curvature, *Math. Ann.* 385 (2023) 1747–1773.
- [23] P. Bérard, D. Meyer, Inégalités isopérimétriques et applications, *Ann. Sci. Ecole Norm. Sup. (4)* 15 (1982) 513–541.
- [24] M. Bhakta, D. Ganguly, D. Karmakar, S. Mazumdar, Sharp quantitative stability of Poincaré-Sobolev inequality in the hyperbolic space and applications to fast diffusion flows, arXiv:2207.11024, 2022.
- [25] G. Bianchi, H. Egnell, A note on the Sobolev inequality, *J. Funct. Anal.* 100 (1991) 18–24.
- [26] A. Björn, J. Björn, *Nonlinear Potential Theory on Metric Spaces*, EMS Tracts in Mathematics, vol. 17, European Mathematical Society (EMS), Zürich, 2011.
- [27] G.A. Bliss, An integral inequality, *J. Lond. Math. Soc.* 5 (1930) 40–46.
- [28] S. Brendle, Sobolev inequalities in manifolds with nonnegative curvature, *Commun. Pure Appl. Math.* (2021).
- [29] S. Brendle, F.C. Marques, Recent progress on the Yamabe problem, in: *Surveys in Geometric Analysis and Relativity*, in: *Adv. Lect. Math. (ALM)*, vol. 20, Int. Press, Somerville, MA, 2011, pp. 29–47.
- [30] H. Brézis, E. Lieb, A relation between pointwise convergence of functions and convergence of functionals, *Proc. Am. Math. Soc.* 88 (1983) 486–490.
- [31] J.E. Brothers, W.P. Ziemer, Minimal rearrangements of Sobolev functions, *J. Reine Angew. Math.* 384 (1988) 153–179.
- [32] D. Burago, Y. Burago, S. Ivanov, *A Course in Metric Geometry*, Graduate Studies in Mathematics, vol. 33, American Mathematical Society, Providence, RI, 2001.
- [33] F. Cavalletti, D. Manini, Rigidities of isoperimetric inequality under nonnegative Ricci curvature, arXiv:2207.03423, 2022.
- [34] F. Cavalletti, D. Manini, Isoperimetric inequality in noncompact MCP spaces, *Proc. Am. Math. Soc.* 150 (2022) 3537–3548.
- [35] F. Cavalletti, E. Milman, The globalization theorem for the curvature-dimension condition, *Invent. Math.* 226 (2021) 1–137.
- [36] F. Cavalletti, A. Mondino, Sharp geometric and functional inequalities in metric measure spaces with lower Ricci curvature bounds, *Geom. Topol.* 21 (2017) 603–645.
- [37] F. Cavalletti, A. Mondino, New formulas for the Laplacian of distance functions and applications, *Anal. PDE* 13 (2020) 2091–2147.
- [38] F. Cavalletti, A. Mondino, D. Semola, Quantitative Obata’s theorem, *Anal. PDE* 16 (2023) 1389–1431.
- [39] J. Cheeger, Differentiability of Lipschitz functions on metric measure spaces, *Geom. Funct. Anal.* 9 (1999) 428–517.
- [40] A. Cianchi, N. Fusco, F. Maggi, A. Pratelli, The sharp Sobolev inequality in quantitative form, *J. Eur. Math. Soc.* 11 (2009) 1105–1139.
- [41] T.H. Colding, Large manifolds with positive Ricci curvature, *Invent. Math.* 124 (1996) 193–214.
- [42] C. Debin, N. Gigli, E. Pasqualetto, Quasi-continuous vector fields on RCD spaces, *Potential Anal.* 54 (2021) 183–211.
- [43] J. Dolbeault, M.J. Esteban, Improved interpolation inequalities and stability, *Adv. Nonlinear Stud.* 20 (2020) 277–291.
- [44] L. Dupaigne, I. Gentil, S. Zugmeyer, Sobolev’s inequality under a curvature-dimension condition, *Ann. Fac. Sci. Toulouse Math. (6)* 32 (2023) 125–144.

- [45] M. Engelstein, R. Neumayer, L. Spolaor, Quantitative stability for minimizing Yamabe metrics, *Trans. Amer. Math. Soc. Ser. B* 9 (2022) 395–414.
- [46] M. Erbar, K. Kuwada, K.-T. Sturm, On the equivalence of the entropic curvature-dimension condition and Bochner’s inequality on metric measure spaces, *Invent. Math.* 201 (2014) 1–79.
- [47] A. Figalli, R. Neumayer, Gradient stability for the Sobolev inequality: the case $p \geq 2$, *J. Eur. Math. Soc.* 21 (2019) 319–354.
- [48] A. Figalli, Y.R.-Y. Zhang, Sharp gradient stability for the Sobolev inequality, *Duke Math. J.* 171 (2022) 2407–2459.
- [49] M. Fogagnolo, L. Mazzieri, Minimising hulls, p -capacity and isoperimetric inequality on complete Riemannian manifolds, *J. Funct. Anal.* 283 (2022) 109638.
- [50] E. Fontenas, Sur les constantes de Sobolev des variétés riemanniennes compactes et les fonctions extrémales des sphères, *Bull. Sci. Math.* 121 (1997) 71–96.
- [51] R.L. Frank, Degenerate stability of some Sobolev inequalities, *Ann. Inst. Henri Poincaré, Anal. Non Linéaire* 39 (2022) 1459–1484.
- [52] N. Gigli, On the heat flow on metric measure spaces: existence, uniqueness and stability, *Calc. Var. Partial Differ. Equ.* 39 (2010) 101–120.
- [53] N. Gigli, An overview of the proof of the splitting theorem in spaces with non-negative Ricci curvature, *Anal. Geom. Metric Spaces* 2 (2014) 169–213.
- [54] N. Gigli, On the differential structure of metric measure spaces and applications, *Mem. Am. Math. Soc.* 236 (2015), vi+91 pp.
- [55] N. Gigli, Nonsmooth differential geometry—an approach tailored for spaces with Ricci curvature bounded from below, *Mem. Am. Math. Soc.* 251 (2018), v+161 pp.
- [56] N. Gigli, On the regularity of harmonic maps from $\text{RCD}(K, N)$ to $\text{CAT}(0)$ spaces and related results, [arXiv:2204.04317](https://arxiv.org/abs/2204.04317), 2022.
- [57] N. Gigli, B. Han, Independence on p of weak upper gradients on RCD spaces, *J. Funct. Anal.* 271 (2014).
- [58] N. Gigli, A. Mondino, G. Savaré, Convergence of pointed non-compact metric measure spaces and stability of Ricci curvature bounds and heat flows, *Proc. Lond. Math. Soc.* 3 (111) (2015) 1071–1129.
- [59] N. Gigli, E. Pasqualetto, *Lectures on Nonsmooth Differential Geometry*, SISSA Springer Series, vol. 2, 2020.
- [60] N. Gigli, E. Pasqualetto, Behaviour of the reference measure on RCD spaces under charts, *Commun. Anal. Geom.* 29 (2021) 1391–1414.
- [61] N. Gigli, C. Rigoni, A note about the strong maximum principle on RCD spaces, *Can. Math. Bull.* 62 (2019) 259–266.
- [62] N. Gigli, I.Y. Violo, Monotonicity formulas for harmonic functions in $\text{RCD}(0, N)$ spaces, *J. Geom. Anal.* 33 (2023) 100, 89 pp.
- [63] A. Grigor’yan, J. Hu, K.-S. Lau, Heat kernels on metric spaces with doubling measure, in: *Fractal Geometry and Stochastics IV*, in: *Progr. Probab.*, vol. 61, Birkhäuser Verlag, Basel, 2009, pp. 3–44.
- [64] M. Gromov, *Metric Structures for Riemannian and Non-Riemannian Spaces*, English ed., *Modern Birkhäuser Classics*, Birkhäuser Boston Inc., Boston, MA, 2007, based on the 1981 French original, with appendices by M. Katz, P. Pansu and S. Semmes, translated from the French by Sean Michael Bates.
- [65] P. Hajłasz, P. Koskela, Sobolev met Poincaré, *Mem. Am. Math. Soc.* 145 (2000), x+101 pp.
- [66] B.-X. Han, Ricci tensor on $\text{RCD}^*(K, N)$ spaces, *J. Geom. Anal.* 28 (2018) 1295–1314.
- [67] Q. Han, F. Lin, *Elliptic Partial Differential Equations*, second ed., *Courant Lecture Notes in Mathematics*, vol. 1, Courant Institute of Mathematical Sciences/American Mathematical Society, New York/Providence, RI, 2011.
- [68] E. Hebey, *Nonlinear Analysis on Manifolds: Sobolev Spaces and Inequalities*, *Courant Lecture Notes in Mathematics*, vol. 5, New York University, Courant Institute of Mathematical Sciences/American Mathematical Society, New York/Providence, RI, 1999.
- [69] J. Heinonen, P. Koskela, From local to global in quasiconformal structures, *Proc. Natl. Acad. Sci. USA* 93 (1996) 554–556.
- [70] J. Heinonen, P. Koskela, Quasiconformal maps in metric spaces with controlled geometry, *Acta Math.* 181 (1998) 1–61.
- [71] S. Honda, Ricci curvature and L^p -convergence, *J. Reine Angew. Math.* 705 (2015) 85–154.
- [72] S. Ilias, Constantes explicites pour les inégalités de Sobolev sur les variétés riemanniennes compactes, *Ann. Inst. Fourier (Grenoble)* 33 (1983) 151–165.
- [73] R. Jiang, Cheeger-harmonic functions in metric measure spaces revisited, *J. Funct. Anal.* 266 (2014) 1373–1394.

- [74] F. Johne, Sobolev inequalities on manifolds with nonnegative Bakry-Émery Ricci curvature, arXiv preprint, arXiv:2103.08496, 2021.
- [75] M. Kell, A note on Lipschitz continuity of solutions of Poisson equations in metric measure spaces, arXiv:1307.2224, 2013.
- [76] S. Kesavan, *Symmetrization & Applications*, Series in Analysis, vol. 3, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2006.
- [77] C. Ketterer, Obata's rigidity theorem for metric measure spaces, *Anal. Geom. Metric Spaces* 3 (2015) 278–295.
- [78] J.M. Lee, T.H. Parker, The Yamabe problem, *Bull. Am. Math. Soc. (N.S.)* 17 (1987) 37–91.
- [79] E.H. Lieb, Sharp constants in the Hardy-Littlewood-Sobolev and related inequalities, *Ann. Math.* (2) 118 (1983) 349–374.
- [80] P.-L. Lions, The concentration-compactness principle in the calculus of variations. The locally compact case. I, *Ann. Inst. Henri Poincaré, Anal. Non Linéaire* 1 (1984) 109–145.
- [81] P.-L. Lions, The concentration-compactness principle in the calculus of variations. The limit case. I, *Rev. Mat. Iberoam.* 1 (1985) 145–201.
- [82] J. Lott, C. Villani, Ricci curvature for metric-measure spaces via optimal transport, *Ann. Math.* (2) 169 (2009) 903–991.
- [83] H. Lou, On singular sets of local solutions to p -Laplace equations, *Chin. Ann. Math., Ser. B* 29 (2008) 521–530.
- [84] M. Miranda, Functions of bounded variation on “good” metric spaces, *J. Math. Pures Appl.* 82 (2003) 975–1004.
- [85] A. Mondino, A. Naber, Structure theory of metric measure spaces with lower Ricci curvature bounds, *J. Eur. Math. Soc.* 21 (2019) 1809–1854.
- [86] A. Mondino, D. Semola, Polya-Szego inequality and Dirichlet p -spectral gap for non-smooth spaces with Ricci curvature bounded below, *J. Math. Pures Appl.* 9 (137) (2020) 238–274.
- [87] R. Neumayer, A note on strong-form stability for the Sobolev inequality, *Calc. Var. Partial Differ. Equ.* 59 (2020) 25, 8 pp.
- [88] F. Nobili, I.Y. Violo, Rigidity and almost rigidity of Sobolev inequalities on compact spaces with lower Ricci curvature bounds, *Calc. Var. Partial Differ. Equ.* 61 (2022) 180.
- [89] G. Pólya, G. Szegő, *Isoperimetric Inequalities in Mathematical Physics (AM-27)*, Princeton University Press, 1951.
- [90] A. Profeta, The sharp Sobolev inequality on metric measure spaces with lower Ricci curvature bounds, *Potential Anal.* 43 (2015) 513–529.
- [91] T. Rajala, Local Poincaré inequalities from stable curvature conditions on metric spaces, *Calc. Var. Partial Differ. Equ.* 44 (2012) 477–494.
- [92] R. Schoen, Conformal deformation of a Riemannian metric to constant scalar curvature, *J. Differ. Geom.* 20 (1984) 479–495.
- [93] N. Shanmugalingam, Newtonian spaces: an extension of Sobolev spaces to metric measure spaces, *Rev. Mat. Iberoam.* 16 (2000) 243–279.
- [94] M. Struwe, *Variational Methods*, fourth ed., *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics (Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics)*, vol. 34, Springer-Verlag, Berlin, 2008, Applications to nonlinear partial differential equations and Hamiltonian systems.
- [95] K.-T. Sturm, On the geometry of metric measure spaces. I, *Acta Math.* 196 (2006) 65–131.
- [96] K.-T. Sturm, On the geometry of metric measure spaces. II, *Acta Math.* 196 (2006) 133–177.
- [97] G. Talenti, Best constant in Sobolev inequality, *Ann. Mat. Pura Appl.* (4) 110 (1976) 353–372.
- [98] N.S. Trudinger, Remarks concerning the conformal deformation of Riemannian structures on compact manifolds, *Ann. Sc. Norm. Super. Pisa, Cl. Sci.* (3) 22 (1968) 265–274.
- [99] H. Yamabe, On a deformation of Riemannian structures on compact manifolds, *Osaka Math. J.* 12 (1960) 21–37.