



SCUOLA INTERNAZIONALE SUPERIORE DI STUDI AVANZATI

SISSA Digital Library

Continuity of some non-local functionals with respect to a convergence of the underlying measures

Original

Continuity of some non-local functionals with respect to a convergence of the underlying measures / Braides, A; Dal Maso, G. - In: JOURNAL DE MATHÉMATIQUES PURES ET APPLIQUÉES. - ISSN 0021-7824. - 170:(2023), pp. 136-149. [10.1016/j.matpur.2022.12.005]

Availability:

This version is available at: 20.500.11767/135816 since: 2023-12-10T14:34:43Z

Publisher:

Published

DOI:10.1016/j.matpur.2022.12.005

Terms of use:

Testo definito dall'ateneo relativo alle clausole di concessione d'uso

Publisher copyright

Elsevier

This version is available for education and non-commercial purposes.

note finali coverpage

(Article begins on next page)

Continuity of some non-local functionals with respect to a convergence of the underlying measures

Andrea Braides and Gianni Dal Maso
SISSA, via Bonomea 265, Trieste, Italy

Abstract

We study some non-local functionals on the Sobolev space $W_0^{1,p}(\Omega)$ involving a double integral on $\Omega \times \Omega$ with respect to a measure μ . We introduce a suitable notion of convergence of measures on product spaces which implies a stability property in the sense of Γ -convergence of the corresponding functionals.

Keywords: non-local functionals, Γ -convergence, Mosco convergence, graphons, cut norm

AMS Class: 49J45, 46E35, 28A33, 28A35

1 Introduction

In this paper we give a contribution to the study of Γ -limits of non-local integral functional, for which only few results are available in the literature (see for instance [13]). We consider sequences of integrals of the type

$$\int_{\Omega \times \Omega} f(u(x), u(y)) d\mu_k(x, y) + \int_{\Omega} g_k(x, \nabla u(x)) dx, \quad (1.1)$$

where Ω is a bounded open subset of \mathbb{R}^d , with $d \geq 1$. These functionals have a non-local term

$$F_k(u) := \int_{\Omega \times \Omega} f(u(x), u(y)) d\mu_k(x, y)$$

depending on a fixed function $f: \mathbb{R} \times \mathbb{R} \rightarrow [0, +\infty)$ and varying positive bounded measures μ_k on $\Omega \times \Omega$, while the local term

$$G_k(u) := \int_{\Omega} g_k(x, \nabla u(x)) dx$$

depends on a function $g_k: \Omega \times \mathbb{R}^d \rightarrow [0, +\infty)$. These functionals are defined for u in the Sobolev space $W_0^{1,p}(\Omega)$.

We assume that the functions g_k satisfy usual growth conditions and that the integral functionals G_k Γ -converge in the weak topology in $W_0^{1,p}(\Omega)$ to a functional G of the same form, with integrand g .

We address the question of the stability for functionals in (1.1); more precisely, we focus on a notion of convergence on μ_k such that the functionals $F_k + G_k$ Γ -converge with respect to the weak topology in $W_0^{1,p}(\Omega)$ to a functional of the form

$$\int_{\Omega \times \Omega} f(u(x), u(y)) d\mu(x, y) + \int_{\Omega} g(x, \nabla u(x)) dx \quad (1.2)$$

for a limit measure μ . Under some additional assumptions, we also obtain the convergence of $F_k + G_k$ in the sense of Mosco convergence in $W_0^{1,p}(\Omega)$.

If $p = 2$, $g_k(x, \cdot)$ are quadratic forms, and $f(s, t) = |t - s|^2$, then the study of such functionals can be framed within the theory of Dirichlet Forms [8], where the Beurling-Deny formula ensures, under suitable assumptions, that the Γ -limit of $F_k + G_k$ can be represented analogously (see [13]). The extension of that theory is not immediate in a non-quadratic setting or when f is an arbitrary continuous function.

Note that, under suitable growth conditions on f , stability is easily proved under the strong assumption of convergence of μ_k to μ in the space $W^{-1,q}(\Omega \times \Omega)$ dual to $W_0^{1,p}(\Omega \times \Omega)$. However, this result is not satisfactory, since such a space may fail to contain relevant measures μ , depending on the value of p , such as Dirac deltas if $p < 2d$.

We introduce a wider space of measures on $\Omega \times \Omega$, together with a new notion of norm, inspired by a convergence that is used in the theory of *graphons* [5, 11]. The latter can be seen as limits of Dirichlet forms on dense graphs (for an interpretation in terms of Γ -convergence we refer to [2]).

We prove that, if μ_k are non-negative measures that converge to μ with respect to that ‘graphon’ norm and $\mu_k(\Omega \times \Omega) \rightarrow \mu(\Omega \times \Omega)$, then the functionals defined by (1.1) Γ -converge under the only assumption that f be continuous and a very mild technical assumption (see (5.2)).

We now describe more in detail the content of the paper. In Section 2 we recall some preliminaries on Sobolev functions and introduce the quasicontinuous representative \tilde{u} of a function $u \in W_0^{1,p}(\Omega)$, which is needed in the precise definition of the functionals (1.1) and (1.2), when μ_k or μ are not absolutely continuous with respect to the Lebesgue measure.

In Section 3 we introduce the space $\mathcal{M}^{1,p}(\Omega \times \Omega)$ of Radon measures on $\Omega \times \Omega$ with finite ‘Sobolev cut norm’, defined as

$$\|\mu\|_{\square} := \sup \left\{ \left| \int_{\Omega \times \Omega} \varphi(x) \psi(y) d\mu(x, y) \right| : \varphi, \psi \in C_c^\infty(\Omega), \|\varphi\|_{1,p}, \|\psi\|_{1,p} \leq 1 \right\},$$

where $\|\cdot\|_{1,p}$ is the norm in $W_0^{1,p}(\Omega)$. We prove that, if $\mu \in \mathcal{M}^{1,p}(\Omega \times \Omega)$ and $u, v \in$

$W_0^{1,p}(\Omega) \cap L^\infty(\Omega)$ with compact support in Ω , we have

$$\left| \int_{\Omega \times \Omega} \tilde{u}(x) \tilde{v}(y) d\mu(x, y) \right| \leq \|\mu\|_{\square} \|u\|_{1,p} \|v\|_{1,p},$$

and the integral does not depend on the choice of the quasicontinuous representatives (see Theorem 3.4).

In Section 4 we prove some continuity results for double integrals. We consider $\mu_k, \mu \in \mathcal{M}_+^{1,p}(\Omega \times \Omega)$ with $\mu_k(\Omega \times \Omega) < +\infty$, $\mu(\Omega \times \Omega) < +\infty$, and

$$\|\mu_k - \mu\|_{\square} \rightarrow 0 \quad \text{and} \quad \mu_k(\Omega \times \Omega) \rightarrow \mu(\Omega \times \Omega).$$

If u_k, v_k are sequences in $W_0^{1,p}(\Omega) \cap L^\infty(\Omega)$ converging to u, v weakly in $W_0^{1,p}(\Omega)$ and with $\sup_k (\|u_k\|_\infty + \|v_k\|_\infty) < +\infty$, then we have

$$\lim_{k \rightarrow +\infty} \int_{\Omega \times \Omega} f(\tilde{u}_k(x), \tilde{v}_k(y)) d\mu_k(x, y) = \int_{\Omega \times \Omega} f(\tilde{u}(x), \tilde{v}(y)) d\mu(x, y)$$

for all continuous functions $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ (see Corollary 4.5). This result is obtained by considering first the special case $f(s, t) = st$ (see Corollary 4.2), from which the result can be obtained for a general polynomial (see Proposition 4.4), and eventually for an arbitrary continuous function by approximation.

Finally, in Section 5 we consider non-negative continuous functions f , and prove the above-mentioned Γ -convergence results (see Theorem 5.1). Note that no convexity assumption on f is needed.

2 Preliminaries on fine properties of Sobolev functions

Throughout the paper we fix $1 < p < +\infty$, $d \geq 1$, and a bounded open subset Ω of \mathbb{R}^d . We consider the Sobolev space $W_0^{1,p}(\Omega)$ endowed with the norm

$$\|u\|_{1,p} = \left(\int_{\Omega} |\nabla u|^p dx \right)^{1/p}.$$

Its dual is denoted by $W^{-1,q}(\Omega)$, where $\frac{1}{p} + \frac{1}{q} = 1$, and this norm is denoted by $\|\cdot\|_{-1,q}$.

For all subset $A \subset \Omega$ the *capacity* of in Ω is defined as

$$C_{1,p}(A) := \inf \left\{ \int_{\Omega} |\nabla u|^p dx : u \in W_0^{1,p}(\Omega), u \geq 1 \text{ a.e. in a neighbourhood of } A \right\}.$$

A function $f: \Omega \rightarrow \mathbb{R}$ is *quasicontinuous* if for every $\varepsilon > 0$ there exists an open set $U \subset \Omega$ with $C_{1,p}(U) < \varepsilon$ such that the restriction of to $\Omega \setminus U$ is continuous. It is known (see e.g. [10, 7]) that each function $u \in W_0^{1,p}(\Omega)$ has a *Borel quasicontinuous representative*,

which we denote by \tilde{u} , in the sense that \tilde{u} is Borel measurable and quasicontinuous and $\tilde{u} = u$ almost everywhere in Ω . Such a representative is unique up to sets of zero capacity.

Let $\mathcal{M}(\Omega)$ denote the space of signed Radon measures on Ω , which can be identified with the dual of $C_c^0(\Omega)$. We say that $\mu \in \mathcal{M}(\Omega)$ belongs to $W^{-1,q}(\Omega)$ if there exists $C \geq 0$ such that

$$\left| \int_{\Omega} \varphi d\mu \right| \leq C \|\varphi\|_{1,p} \quad \text{for all } \varphi \in C_c^\infty(\Omega).$$

In this case there exists a unique $T_\mu \in W^{-1,q}(\Omega)$ such that

$$\langle T_\mu, \varphi \rangle = \int_{\Omega} \varphi d\mu \quad \text{for all } \varphi \in C_c^\infty(\Omega), \quad (2.1)$$

where $\langle \cdot, \cdot \rangle$ denotes the duality product between $W^{-1,q}(\Omega)$ and $W_0^{1,p}(\Omega)$. In other words μ and T_μ coincide as distributions on Ω . For notational convenience, we shall sometimes directly write μ in the place of T_μ when the distinction between the two is not relevant.

In the following theorem we recall a property of sets of zero $C_{1,p}$ -capacity (see [9]) and an integral representation of T_μ that can be deduced from a result of Brezis and Browder [3].

In order to simplify the notation, we introduce the space

$$W_c^{1,p}(\Omega) := \{u \in W_0^{1,p}(\Omega) : u \text{ has compact support in } \Omega\}.$$

Theorem 2.1. *Let $\mu \in \mathcal{M}(\Omega) \cap W^{-1,q}(\Omega)$. Then for every $A \subset \Omega$ with $C_{1,p}(A) = 0$ -capacity A is $|\mu|$ -measurable and*

$$|\mu|(A) = 0 \quad (2.2)$$

If $u \in W_c^{1,p}(\Omega) \cap L^\infty(\Omega)$, then

$$\langle T_\mu, u \rangle = \int_{\Omega} \tilde{u} d\mu. \quad (2.3)$$

If, in addition, $\mu \geq 0$, then, for all $u \in W_0^{1,p}(\Omega)$, we have $\tilde{u} \in L^1(\Omega; \mu)$ and (2.3) holds.

3 Sobolev graphons

Graphons are functions ρ defined on $(0,1) \times (0,1)$ introduced to study functionals of the form

$$\int_{(0,1) \times (0,1)} f(u(x), u(y)) \rho(x, y) dx dy,$$

defined for $u \in L^\infty(0,1)$, especially when $f(u, v) = |u - v|^2$. The functionals above are introduced as a generalization, and in some sense a limit, of energies on dense graphs [5, 11]. To that end, the space of graphons is equipped with the so-called *cut norm*

$$\|\rho\|_{\square} := \sup_{\varphi, \psi: (0,1) \rightarrow [0,1]} \left| \int_{(0,1) \times (0,1)} \varphi(x) \psi(y) \rho(x, y) dx dy \right|. \quad (3.1)$$

We extend the definition of cut norm to arbitrary measures defined on $\Omega \times \Omega$, with Ω in the place of $(0, 1)$. For our purpose it is convenient to take test functions in the Sobolev space $W_0^{1,p}(\Omega)$.

Definition 3.1. *The space $\mathcal{M}^{1,p}(\Omega \times \Omega)$ is defined as*

$$\mathcal{M}^{1,p}(\Omega \times \Omega) := \{\mu \in \mathcal{M}(\Omega \times \Omega) : \|\mu\|_{\square} < +\infty\}, \quad (3.2)$$

where the Sobolev cut norm of μ is defined as

$$\|\mu\|_{\square} := \sup \left\{ \left| \int_{\Omega \times \Omega} \varphi(x)\psi(y)d\mu(x,y) \right| : \varphi, \psi \in C_c^{\infty}(\Omega), \|\varphi\|_{1,p}, \|\psi\|_{1,p} \leq 1 \right\}. \quad (3.3)$$

We let $\mathcal{M}_+^{1,p}(\Omega \times \Omega)$ denote the cone of the positive measures in $\mathcal{M}^{1,p}(\Omega \times \Omega)$.

Remark 3.2. We make some observations on the convergence in the space $\mathcal{M}^{1,p}(\Omega \times \Omega)$, and in particular we compare it with weak* convergence in $\mathcal{M}(\Omega \times \Omega)$ and with the convergence $W^{-1,q}(\Omega \times \Omega)$.

(i) $\|\cdot\|_{\square}$ defines a norm on $\mathcal{M}^{1,p}(\Omega \times \Omega)$;

(ii) if μ belongs to $W^{-1,q}(\Omega \times \Omega)$, then $\|\mu\|_{\square} \leq \|\mu\|_{-1,q}$. Indeed in such a case the function $\Phi(x, y) = \varphi(x)\psi(y)$ belongs to $W_0^{1,p}(\Omega \times \Omega)$ and

$$\begin{aligned} \int_{\Omega \times \Omega} |\nabla \Phi|^p dx dy &= \int_{\Omega \times \Omega} |\psi(y)\nabla \varphi(x) + \varphi(x)\nabla \psi(y)|^p dx dy \\ &\leq C \int_{\Omega \times \Omega} |\psi(y)|^p |\nabla \varphi(x)|^p + |\varphi(x)|^p |\nabla \psi(y)|^p dx dy \leq C \end{aligned}$$

so that

$$\left| \int_{\Omega \times \Omega} \varphi(x)\psi(y)d\mu(x,y) \right| \leq C \|\mu\|_{-1,q};$$

(iii) if $\mu_k, \mu \in \mathcal{M}^{1,p}(\Omega \times \Omega)$ and $\|\mu_k - \mu\|_{\square} \rightarrow 0$, then

$$\int_{\Omega \times \Omega} \varphi(x)\psi(y)d\mu_k(x,y) \rightarrow \int_{\Omega \times \Omega} \varphi(x)\psi(y)d\mu(x,y) \quad (3.4)$$

for all $\varphi, \psi \in C_c^{\infty}(\Omega)$. If in addition $\sup_k |\mu_k|(\Omega \times \Omega) < +\infty$, then (3.4) and a density argument imply that μ_k converges to μ weakly* in $\mathcal{M}(\Omega \times \Omega)$;

(iv) if μ_k^1 and μ_k^2 are such that μ_k^j converge to μ^j in $W^{-1,q}(\Omega)$, then the measures $\mu_k = \mu_k^1 \otimes \mu_k^2$ converge to $\mu = \mu^1 \otimes \mu^2$;

(v) if $\sup_k \|\mu_k\|_{\square} < +\infty$ and μ_k converges to some μ weakly* in $\mathcal{M}(\Omega \times \Omega)$, then $\mu \in \mathcal{M}^{1,p}(\Omega \times \Omega)$ and $\|\mu\|_{\square} \leq \liminf_k \|\mu_k\|_{\square}$;

Note that $\mathcal{M}^{1,p}(\Omega \times \Omega)$ is strictly larger than $\mathcal{M}(\Omega \times \Omega) \cap W^{-1,q}(\Omega \times \Omega)$, and in particular its convergence is weaker than convergence in $W^{-1,q}(\Omega \times \Omega)$. Some examples of this inclusion are given below.

Example 3.3. (i) The first example is simply a Dirac delta $\mu = \delta(x_0, y_0)$ for $x_0, y_0 \in \Omega$. Indeed if $d < p < 2d$, then μ does not belong to $W^{-1,q}(\Omega \times \Omega)$, while

$$\left| \int_{\Omega \times \Omega} \varphi(x)\psi(y)d\mu(x, y) \right| = |\varphi(x_0)\psi(y_0)| \leq C\|\varphi\|_{1,p}\|\psi\|_{1,p}$$

where the inequality follows from the embedding of $W_0^{1,p}(\Omega)$ into $L^\infty(\Omega)$.

(ii) We can also exhibit an example where μ is absolutely continuous with respect to \mathcal{L}^d with density of the form $m(x)m(y)$. We choose $d = 1$, $p = 2$, $\Omega = (-1, 1)$, and

$$m(x) = \frac{1}{|x| \log |x| \log |\log x| \log^2 |\log |\log x||}.$$

Note that $m \in L^1(0, 1)$ so that $\mu \in \mathcal{M}^{1,2}(\Omega \times \Omega)$. If we take

$$w(x, y) = \begin{cases} \left| \log |\log \sqrt{x^2 + y^2}| \right| & \text{if } \sqrt{x^2 + y^2} < 1/e, \\ 0 & \text{otherwise,} \end{cases}$$

then $w \in W_0^{1,2}((-1, 1)^2)$ but $\int_{(-1,1)^2} w d\mu = +\infty$. By Theorem 2.1, this shows that $\mu \notin W^{-1,2}((-1, 1)^2)$.

Theorem 3.4. *Let $\mu \in \mathcal{M}^{1,p}(\Omega \times \Omega)$. Then for every $A \subset \Omega$ with $C_{1,p}(A) = 0$ the sets $A \times \Omega$ and $\Omega \times A$ are $|\mu|$ -measurable and*

$$|\mu|(A \times \Omega) = |\mu|(\Omega \times A) = 0. \quad (3.5)$$

Moreover, for all $u, v \in W_c^{1,p}(\Omega) \cap L^\infty(\Omega)$ we have

$$\left| \int_{\Omega \times \Omega} \tilde{u}(x)\tilde{v}(y)d\mu(x, y) \right| \leq \|\mu\|_{\square} \|u\|_{1,p} \|v\|_{1,p}. \quad (3.6)$$

Proof. Fix $\psi \in C_c^\infty(\Omega)$, and let $\mu_\psi \in \mathcal{M}(\Omega)$ be defined by

$$\mu_\psi(B) = \int_{B \times \Omega} \psi(y)d\mu(x, y) \text{ for all Borel sets } B \subset \subset \Omega. \quad (3.7)$$

Since

$$\int_{\Omega} \varphi(x)d\mu_\psi(x) = \int_{\Omega \times \Omega} \varphi(x)\psi(y)d\mu(x, y) \text{ for all } \varphi \in C_c^\infty(\Omega),$$

by the definition of $\|\mu\|_{\square}$ we have

$$\left| \int_{\Omega} \varphi(x)d\mu_\psi(x) \right| \leq \|\mu\|_{\square} \|\psi\|_{1,p} \|\varphi\|_{1,p} \text{ for all } \varphi \in C_c^\infty(\Omega). \quad (3.8)$$

Hence, $\mu_\psi \in \mathcal{M}(\Omega) \cap W^{-1,q}(\Omega)$.

Let $B \subset \Omega$ be a Borel set with $C_{1,p}(B) = 0$. Let Ω' be an open set with $\Omega' \subset\subset \Omega$, and let ψ_k be a non-decreasing sequence of non-negative functions in $C_c^\infty(\Omega')$ converging to the constant 1 in Ω . By the definition of μ_{ψ_k} and Theorem 2.1, we have

$$\int_{B \times \Omega} \psi_k(y) d\mu(x, y) = \mu_{\psi_k}(B) = 0.$$

Letting $k \rightarrow +\infty$ we deduce $\mu(B \times \Omega') = 0$. Since this holds for all Borel subsets of B we deduce that $|\mu|(B \times \Omega') = 0$. By the arbitrariness of $\Omega' \subset\subset \Omega$ we finally obtain $|\mu|(B \times \Omega) = 0$. For a general $A \subset \Omega$ with $C_{1,p}(A) = 0$ it is sufficient to observe that there exists a Borel set B such that $A \subset B \subset \Omega$ and $C_{1,p}(B) = 0$. This implies that $A \times \Omega$ and is $|\mu|$ -measurable and $|\mu|(A \times \Omega) = 0$. A similar argument proves the same result for $\Omega \times A$.

Let $T_{\mu_\psi} \in W^{-1,q}(\Omega)$ be defined as in (2.1) with μ replaced by μ_ψ . Fix $u \in W_c^{1,p}(\Omega) \cap L^\infty(\Omega)$. Then, by Theorem 2.1 and the definition of μ_ψ , we have

$$\langle T_{\mu_\psi}, u \rangle = \int_{\Omega} \tilde{u}(x) d\mu_\psi(x) = \int_{\Omega \times \Omega} \tilde{u}(x) \psi(y) d\mu(x, y).$$

By (3.8) we have $\|T_{\mu_\psi}\|_{-1,q} \leq \|\mu\|_{\square} \|\psi\|_{1,p}$, so that the previous equality gives

$$\left| \int_{\Omega \times \Omega} \tilde{u}(x) \psi(y) d\mu(x, y) \right| \leq \|\mu\|_{\square} \|\psi\|_{1,p} \|u\|_{1,p} \quad (3.9)$$

for all $u \in W_c^{1,p}(\Omega) \cap L^\infty(\Omega)$ and every $\psi \in C_c^\infty(\Omega)$.

Fix $u \in W_c^{1,p}(\Omega) \cap L^\infty(\Omega)$ and let $\mu^u \in \mathcal{M}(\Omega)$ be defined by

$$\mu^u(B) := \int_{\Omega \times B} \tilde{u}(x) d\mu(x, y) \text{ for all Borel sets } B \subset\subset \Omega. \quad (3.10)$$

Since

$$\int_{\Omega} \psi(y) d\mu^u(y) = \int_{\Omega \times \Omega} \tilde{u}(x) \psi(y) d\mu(x, y) \text{ for all } \psi \in C_c^\infty(\Omega),$$

thanks to (3.9) we then have

$$\left| \int_{\Omega} \psi(y) d\mu^u(y) \right| \leq \|\mu\|_{\square} \|\psi\|_{1,p} \|u\|_{1,p} \text{ for all } \psi \in C_c^\infty(\Omega).$$

Hence, $\mu^u \in \mathcal{M}(\Omega) \cap W^{-1,q}(\Omega)$, and, using the notation above, we obtain

$$\|T_{\mu^u}\|_{-1,q} \leq \|\mu\|_{\square} \|u\|_{1,p}.$$

By Theorem 2.1 and the definition of μ^u , we then have

$$\langle T_{\mu^u}, v \rangle = \int_{\Omega} \tilde{v}(y) d\mu^u(y) = \int_{\Omega \times \Omega} \tilde{u}(x) \tilde{v}(y) d\mu(x, y).$$

Together with the previous inequality this gives

$$\left| \int_{\Omega \times \Omega} \tilde{u}(x)\tilde{v}(y)d\mu(x, y) \right| \leq \|\mu\| \square \|u\|_{1,p} \|v\|_{1,p},$$

which concludes the proof of (3.6). \square

4 Continuity properties of some double integrals

In this section we find conditions on f, u_k, v_k, u, v, μ_k , and μ which imply the convergence

$$\lim_{k \rightarrow +\infty} \int_{\Omega \times \Omega} f(\tilde{u}_k(x), \tilde{v}_k(y))d\mu_k(x, y) = \int_{\Omega \times \Omega} f(\tilde{u}(x), \tilde{v}(y))d\mu(x, y).$$

We begin with the case $f(s, t) = st$. Subsequently, we consider the case when f is a polynomial, and finally an arbitrary continuous function by approximation.

Lemma 4.1. *Let $\mu \in \mathcal{M}_+^{1,p}(\Omega \times \Omega)$ and let u_k, v_k be sequences in $W_c^{1,p}(\Omega) \cap L^\infty(\Omega)$ converging to u, v weakly in $W^{1,p}(\Omega)$. Assume that there exist a compact set $K \subset \Omega$ and a constant M such that*

$$\text{supp}(u_k) \cup \text{supp}(v_k) \subset K \text{ and } \|u_k\|_\infty + \|v_k\|_\infty \leq M, \quad (4.1)$$

where $\|\cdot\|_\infty$ denotes the norm in $L^\infty(\Omega)$. Then

$$\lim_{k \rightarrow +\infty} \int_{\Omega \times \Omega} \tilde{u}_k(x)\tilde{v}_k(y)d\mu(x, y) = \int_{\Omega \times \Omega} \tilde{u}(x)\tilde{v}(y)d\mu(x, y). \quad (4.2)$$

Proof. We write

$$\begin{aligned} & \int_{\Omega \times \Omega} \tilde{u}_k(x)\tilde{v}_k(y)d\mu(x, y) - \int_{\Omega \times \Omega} \tilde{u}(x)\tilde{v}(y)d\mu(x, y) \\ &= \int_{\Omega \times \Omega} (\tilde{u}_k(x) - \tilde{u}(x))\tilde{v}_k(y)d\mu(x, y) + \int_{\Omega \times \Omega} \tilde{u}(x)(\tilde{v}_k(y) - \tilde{v}(y))d\mu(x, y). \end{aligned} \quad (4.3)$$

The first integral in (4.3) is estimated by

$$\begin{aligned} \left| \int_{\Omega \times \Omega} (\tilde{u}_k(x) - \tilde{u}(x))\tilde{v}_k(y)d\mu(x, y) \right| &\leq M \int_{\Omega \times \Omega} |\tilde{u}_k(x) - \tilde{u}(x)|\psi(y)d\mu(x, y) \\ &= M \langle T_{\mu_\psi}, |u_k - u| \rangle = o(1) \end{aligned} \quad (4.4)$$

as $k \rightarrow +\infty$, where ψ is any function in $C_c^\infty(\Omega)$ with $0 \leq \psi \leq 1$ in Ω and $\psi = 1$ on K , and μ_ψ is defined in (3.7), and the convergence to 0 follows from the fact that $|u_k - u| \rightharpoonup 0$ weakly in $W_0^{1,p}(\Omega)$. Moreover, if μ^u is defined in (3.10), we have

$$\int_{\Omega \times \Omega} \tilde{u}(x)(\tilde{v}_k(y) - \tilde{v}(y))d\mu(x, y) = \langle T_{\mu^u}, v_k - v \rangle = o(1) \quad (4.5)$$

as $k \rightarrow +\infty$. The convergence to 0 follows from the fact that $v_k - v \rightharpoonup 0$ weakly in $W_0^{1,p}(\Omega)$. \square

Corollary 4.2. *Under the same hypotheses of Lemma 4.1, let $\mu_k \in \mathcal{M}^{1,p}(\Omega \times \Omega)$ with $\|\mu_k - \mu\|_{\square} \rightarrow 0$. Then*

$$\lim_{k \rightarrow +\infty} \int_{\Omega \times \Omega} \tilde{u}_k(x) \tilde{v}_k(y) d\mu_k(x, y) = \int_{\Omega \times \Omega} \tilde{u}(x) \tilde{v}(y) d\mu(x, y). \quad (4.6)$$

Proof. It suffices to remark that

$$\left| \int_{\Omega \times \Omega} \tilde{u}_k(x) \tilde{v}_k(y) d\mu_k(x, y) - \int_{\Omega \times \Omega} \tilde{u}_k(x) \tilde{v}_k(y) d\mu(x, y) \right| \leq \|\mu_k - \mu\|_{\square} \|u_k\|_{1,p} \|v_k\|_{1,p}$$

and that the right-hand side tends to 0. The conclusion then follows from Lemma 4.1. \square

Proposition 4.3. *Let $P: \mathbb{R}^2 \rightarrow \mathbb{R}$ be a polynomial function and let $\varphi, \psi \in C_c^\infty(\Omega)$. Let $\mu_k, \mu \in \mathcal{M}_+^{1,p}(\Omega \times \Omega)$ with $\|\mu_k - \mu\|_{\square} \rightarrow 0$, let u_k, v_k be sequences in $W_0^{1,p}(\Omega) \cap L^\infty(\Omega)$ converging to u, v weakly in $W_0^{1,p}(\Omega)$ and equibounded in $L^\infty(\Omega)$. Then*

$$\lim_{k \rightarrow +\infty} \int_{\Omega \times \Omega} \varphi(x) \psi(y) P(\tilde{u}_k(x), \tilde{v}_k(y)) d\mu_k(x, y) = \int_{\Omega \times \Omega} \varphi(x) \psi(y) P(\tilde{u}(x), \tilde{v}(y)) d\mu(x, y). \quad (4.7)$$

Proof. It is sufficient to consider $P(s, t) = s^m t^n$ for some non-negative integers m, n . We define $w_k(x) = \varphi(x) u_k(x)^m$ and $z_k(y) = \psi(y) v_k(y)^n$. Observe that w_k, z_k satisfy the hypotheses of Lemma 4.1, weakly converging in $W_0^{1,p}(\Omega)$ to w, z given by $w(x) = \varphi(x) u(x)^m$ and $z(y) = \psi(y) v(y)^n$. Since

$$\varphi(x) \psi(y) P(\tilde{u}_k(x), \tilde{v}_k(y)) = \tilde{w}_k(x) \tilde{z}_k(y), \quad \varphi(x) \psi(y) P(\tilde{u}(x), \tilde{v}(y)) = \tilde{w}(x) \tilde{z}(y),$$

the claim then follows by Corollary 4.2. \square

In the next proposition we consider a stronger condition on the convergence of μ_k to μ which allows to avoid the multiplication by the cut-off functions in the previous proposition. Note that the second condition in (4.8) is necessary for the validity of (4.9) when P is a constant.

Proposition 4.4. *Let $P: \mathbb{R}^2 \rightarrow \mathbb{R}$ be a polynomial function. Let $\mu_k, \mu \in \mathcal{M}_+^{1,p}(\Omega \times \Omega)$ with $\mu_k(\Omega \times \Omega) < +\infty$ and $\mu(\Omega \times \Omega) < +\infty$. Suppose that*

$$\|\mu_k - \mu\|_{\square} \rightarrow 0 \quad \text{and} \quad \mu_k(\Omega \times \Omega) \rightarrow \mu(\Omega \times \Omega). \quad (4.8)$$

Let u_k, v_k be sequences in $W_0^{1,p}(\Omega) \cap L^\infty(\Omega)$ converging to u, v weakly in $W_0^{1,p}(\Omega)$ and with $\sup_k (\|u_k\|_\infty + \|v_k\|_\infty) = M < +\infty$. Then

$$\lim_{k \rightarrow +\infty} \int_{\Omega \times \Omega} P(\tilde{u}_k(x), \tilde{v}_k(y)) d\mu_k(x, y) = \int_{\Omega \times \Omega} P(\tilde{u}(x), \tilde{v}(y)) d\mu(x, y). \quad (4.9)$$

Proof. By Remark 3.2(iii) the first condition in (4.8) implies that $\mu_k \rightharpoonup \mu$ weakly* in $\mathcal{M}(\Omega \times \Omega)$. This, together with the second condition in (4.8), gives $\mu_k(B) \rightarrow \mu(B)$ for all Borel sets in $\Omega \times \Omega$ such that $\mu(\partial B \cap (\Omega \times \Omega)) = 0$. Then, for every $\varepsilon > 0$ there exists a compact set K_ε of Ω such that

$$\mu((\Omega \times \Omega) \setminus (K_\varepsilon \times K_\varepsilon)) < \varepsilon \quad \text{and} \quad \mu_k((\Omega \times \Omega) \setminus (K_\varepsilon \times K_\varepsilon)) < \varepsilon \quad \text{for every } k. \quad (4.10)$$

Let $\varphi_\varepsilon \in C_c^\infty(\Omega)$ with $0 \leq \varphi_\varepsilon \leq 1$ in Ω and $\varphi_\varepsilon = 1$ on K_ε , and let

$$C_M = \max\{P(s, t) : s, t \in [-M, M]\}.$$

With this choice

$$\left| \int_{\Omega \times \Omega} P(\tilde{u}_k(x), \tilde{v}_k(y)) d\mu_k(x, y) - \int_{\Omega \times \Omega} \varphi_\varepsilon(x) \varphi_\varepsilon(y) P(\tilde{u}_k(x), \tilde{v}_k(y)) d\mu_k(x, y) \right| \leq C_M \varepsilon$$

and

$$\left| \int_{\Omega \times \Omega} P(\tilde{u}(x), \tilde{v}(y)) d\mu(x, y) - \int_{\Omega \times \Omega} \varphi_\varepsilon(x) \varphi_\varepsilon(y) P(\tilde{u}(x), \tilde{v}(y)) d\mu(x, y) \right| \leq C_M \varepsilon.$$

By (4.7) with $\varphi = \psi = \varphi_\varepsilon$ we then deduce

$$\limsup_{k \rightarrow +\infty} \left| \int_{\Omega \times \Omega} P(\tilde{u}_k(x), \tilde{v}_k(y)) d\mu_k(x, y) - \int_{\Omega \times \Omega} P(\tilde{u}(x), \tilde{v}(y)) d\mu(x, y) \right| \leq 2C_M \varepsilon,$$

from which we obtain (4.9) by the arbitrariness of ε . \square

We are now ready to prove the result for an arbitrary continuous function f .

Corollary 4.5. *Let $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ be a continuous function. Under the assumptions of Proposition 4.4 we have*

$$\lim_{k \rightarrow +\infty} \int_{\Omega \times \Omega} f(\tilde{u}_k(x), \tilde{v}_k(y)) d\mu_k(x, y) = \int_{\Omega \times \Omega} f(\tilde{u}(x), \tilde{v}(y)) d\mu(x, y). \quad (4.11)$$

Proof. It suffices to approximate uniformly f by polynomials on $[-M, M]^2$ and apply the previous proposition. \square

Finally, if f is bounded we can remove the hypothesis that u_k and v_k are bounded in L^∞ .

Theorem 4.6. *Let $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ be a bounded continuous function. Let $\mu_k, \mu \in \mathcal{M}_+^{1,p}(\Omega \times \Omega)$ with $\mu_k(\Omega \times \Omega) < +\infty$ and $\mu(\Omega \times \Omega) < +\infty$. Suppose that*

$$\|\mu_k - \mu\|_{\square} \rightarrow 0 \quad \text{and} \quad \mu_k(\Omega \times \Omega) \rightarrow \mu(\Omega \times \Omega). \quad (4.12)$$

Let u_k, v_k be sequences in $W_0^{1,p}(\Omega)$ converging to u, v weakly in $W_0^{1,p}(\Omega)$, then

$$\lim_{k \rightarrow +\infty} \int_{\Omega \times \Omega} f(\tilde{u}_k(x), \tilde{v}_k(y)) d\mu_k(x, y) = \int_{\Omega \times \Omega} f(\tilde{u}(x), \tilde{v}(y)) d\mu(x, y). \quad (4.13)$$

Proof. For all $\lambda > 0$ we define the truncation operator

$$\tau^\lambda(s) := (s \vee (-\lambda)) \wedge \lambda. \quad (4.14)$$

With fixed $\lambda > 1$ we set $u_k^\lambda(x) := \tau^\lambda(u_k(x))$ and $v_k^\lambda(y) := \tau^\lambda(v_k(y))$, and correspondingly, $u^\lambda(x) := \tau^\lambda(u(x))$ and $v^\lambda(y) := \tau^\lambda(v(y))$. By the uniqueness of the quasicontinuous representatives, we deduce from (3.5) that $\tilde{u}_k^\lambda(x) = \tau^\lambda(\tilde{u}_k(x))$ and $\tilde{v}_k^\lambda(x) = \tau^\lambda(\tilde{v}_k(x))$ for μ_k -almost every $(x, y) \in \Omega \times \Omega$. Similarly, we have $\tilde{u}^\lambda(x) = \tau^\lambda(\tilde{u}(x))$ and $\tilde{v}^\lambda(x) = \tau^\lambda(\tilde{v}(x))$ for μ -almost every $(x, y) \in \Omega \times \Omega$.

Since $u_k^\lambda \rightharpoonup u^\lambda$ and $v_k^\lambda \rightharpoonup v^\lambda$ weakly in $W_0^{1,p}(\Omega)$, we have

$$\lim_{k \rightarrow +\infty} \int_{\Omega \times \Omega} f(\tau^\lambda(\tilde{u}_k(x)), \tau^\lambda(\tilde{v}_k(y))) d\mu_k(x, y) = \int_{\Omega \times \Omega} f(\tau^\lambda(\tilde{u}(x)), \tau^\lambda(\tilde{v}(y))) d\mu(x, y) \quad (4.15)$$

by Corollary 4.5. To conclude the proof of the result it suffices to estimate

$$A_k^\lambda := \left| \int_{\Omega \times \Omega} f(\tau^\lambda(\tilde{u}_k(x)), \tau^\lambda(\tilde{v}_k(y))) d\mu_k(x, y) - \int_{\Omega \times \Omega} f(\tilde{u}_k(x), \tilde{v}_k(y)) d\mu_k(x, y) \right|, \quad (4.16)$$

$$A^\lambda := \left| \int_{\Omega \times \Omega} f(\tau^\lambda(\tilde{u}(x)), \tau^\lambda(\tilde{v}(y))) d\mu(x, y) - \int_{\Omega \times \Omega} f(\tilde{u}(x), \tilde{v}(y)) d\mu(x, y) \right|. \quad (4.17)$$

Let $M_0 = \sup |f|$. Since

$$A_k^\lambda \leq 2M_0 \mu_k(\{(x, y) \in \Omega \times \Omega : (\tilde{u}_k(x), \tilde{v}_k(y)) \notin [-\lambda, \lambda]^2\}),$$

it is enough to separately estimate

$$\mu_k(\{(x, y) \in \Omega \times \Omega : \tilde{u}_k(x) > \lambda\}), \quad \mu_k(\{(x, y) \in \Omega \times \Omega : \tilde{u}_k(x) < -\lambda\}), \quad (4.18)$$

$$\mu_k(\{(x, y) \in \Omega \times \Omega : \tilde{v}_k(y) > \lambda\}), \quad \mu_k(\{(x, y) \in \Omega \times \Omega : \tilde{v}_k(y) < -\lambda\}). \quad (4.19)$$

For fixed $\varepsilon > 0$ let K_ε be the compact sets introduced at the beginning of the proof of Proposition 4.4. By (4.10) we have

$$\mu_k(\{(x, y) \in \Omega \times \Omega : \tilde{u}_k(x) > \lambda\}) \leq \mu_k(\{(x, y) \in K_\varepsilon \times K_\varepsilon : \tilde{u}_k(x) > \lambda\}) + \varepsilon. \quad (4.20)$$

Let $\psi_\varepsilon \in C_c^\infty(\Omega)$ with $0 \leq \psi_\varepsilon \leq 1$ and $\psi_\varepsilon = 1$ on K_ε . Let T_ε be the element in $W^{-1,q}(\Omega)$ defined by

$$\langle T_\varepsilon, v \rangle := \int_{\Omega \times \Omega} \tilde{v}(x) \psi_\varepsilon(y) d\mu(x, y) \text{ for every } v \in W_0^{1,p}(\Omega).$$

Since u_k are equibounded in $W_0^{1,p}(\Omega)$, by (3.6) there exists $C_\varepsilon > 0$ and $k_\varepsilon \in \mathbb{N}$ such that

$$\begin{aligned} & \mu_k(\{(x, y) \in K_\varepsilon \times K_\varepsilon : \tilde{u}_k(x) > \lambda\}) \leq \int_{\Omega \times \Omega} [\tilde{u}_k(x) - \lambda + 1]^+ \psi_\varepsilon(x) \psi_\varepsilon(y) d\mu_k(x, y) \\ & \leq \left| \int_{\Omega \times \Omega} [\tilde{u}_k(x) - \lambda + 1]^+ \psi_\varepsilon(x) \psi_\varepsilon(y) d(\mu_k - \mu)(x, y) \right| + \int_{\Omega \times \Omega} [\tilde{u}_k(x) - \lambda + 1]^+ \psi_\varepsilon(y) d\mu(x, y) \\ & \leq C_\varepsilon \|\mu_k - \mu\|_\square + \langle T_{\psi_\varepsilon}, [u_k - \lambda + 1]^+ \rangle = \varepsilon + \langle T_{\psi_\varepsilon}, [u - \lambda + 1]^+ \rangle \end{aligned}$$

for all $k \geq k_\varepsilon$. Now, since $\lim_{\lambda \rightarrow +\infty} \langle T\psi_\varepsilon, [u - \lambda + 1]^+ \rangle = 0$, we can choose $\lambda_\varepsilon > 0$ such that

$$\mu_k(\{(x, y) \in K_\varepsilon \times K_\varepsilon : \tilde{u}_k(x) > \lambda\}) \leq 2\varepsilon$$

for all $k \geq k_\varepsilon$ and $\lambda \geq \lambda_\varepsilon$. By (4.20) this in turn gives

$$\mu_k(\{(x, y) \in \Omega \times \Omega : \tilde{u}_k(x) > \lambda\}) \leq 3\varepsilon$$

for all $k \geq k_\varepsilon$ and $\lambda \geq \lambda_\varepsilon$.

In the same way, we can prove analogue estimates for the other measures in (4.18) and (4.19), which we may assume to hold for the same K_ε and λ_ε , and conclude that $A_k^\lambda \leq 24M_0\varepsilon$ for all $k \geq k_\varepsilon$ and $\lambda \geq \lambda_\varepsilon$. Similarly we can prove that $A^\lambda \leq 24M_0\varepsilon$ for $\lambda \geq \lambda_\varepsilon$. From these estimates, by (4.15)–(4.17) the claim follows by the arbitrariness of ε . \square

We finally prove a lower bound for limits of double integrals.

Corollary 4.7. *Let $f: \mathbb{R}^2 \rightarrow [0, +\infty)$ be a continuous function. Let $\mu_k, \mu \in \mathcal{M}_+^{1,p}(\Omega \times \Omega)$ with $\mu_k(\Omega \times \Omega) < +\infty$, $\mu(\Omega \times \Omega) < +\infty$, satisfying (4.12). Let u_k, v_k be sequences in $W_0^{1,p}(\Omega)$ converging to u, v weakly in $W_0^{1,p}(\Omega)$. Then*

$$\liminf_{k \rightarrow +\infty} \int_{\Omega \times \Omega} f(\tilde{u}_k(x), \tilde{v}_k(y)) d\mu_k(x, y) \geq \int_{\Omega \times \Omega} f(\tilde{u}(x), \tilde{v}(y)) d\mu(x, y). \quad (4.21)$$

Proof. For every $\lambda > 0$ let $f^\lambda := \tau^\lambda(f)$, where τ^λ is the truncation operator as in (4.14). By Theorem 4.6 we then have

$$\begin{aligned} \liminf_{k \rightarrow +\infty} \int_{\Omega \times \Omega} f(\tilde{u}_k(x), \tilde{v}_k(y)) d\mu_k(x, y) &\geq \lim_{k \rightarrow +\infty} \int_{\Omega \times \Omega} f^\lambda(\tilde{u}_k(x), \tilde{v}_k(y)) d\mu_k(x, y) \\ &= \int_{\Omega \times \Omega} f^\lambda(\tilde{u}(x), \tilde{v}(y)) d\mu(x, y). \end{aligned}$$

We then conclude by letting $\lambda \rightarrow +\infty$. \square

5 Γ -convergence

In this final section we shall prove the Γ -convergence and the Mosco convergence of sequences of functionals as in (1.1).

Let $f: \mathbb{R}^2 \rightarrow [0, +\infty)$ be a continuous function, and let $\mu_k, \mu \in \mathcal{M}_+^{1,p}(\Omega \times \Omega)$. We define $F_k, F: W_0^{1,p}(\Omega) \rightarrow [0, +\infty]$ by

$$F_k(u) := \int_{\Omega \times \Omega} f(\tilde{u}(x), \tilde{u}(y)) d\mu_k(x, y) \quad \text{and} \quad F(u) := \int_{\Omega \times \Omega} f(\tilde{u}(x), \tilde{u}(y)) d\mu(x, y). \quad (5.1)$$

We assume that f satisfies the following condition: there exists an unbounded set $\Lambda \subset [0, +\infty)$ and two constants $a, b \geq 0$ such that

$$f(\tau^\lambda(s), \tau^\lambda(t)) \leq a f(s, t) + b \quad \text{for all } s, t \in \mathbb{R} \text{ and for all } \lambda \in \Lambda, \quad (5.2)$$

where τ^λ is the truncation operator defined in (4.14). Note that this condition is valid if f is bounded (with $a = 0$) or when f is decreasing by truncations (with $a = 1$ and $b = 0$).

Let $g_k, g: \Omega \times \mathbb{R}^d \rightarrow \mathbb{R}$ be Carathéodory functions satisfying the growth conditions

$$c_0 |\xi|^p \leq g_k(x, \xi) \leq c_1 |\xi|^p + a(x), \quad c_0 |\xi|^p \leq g(x, \xi) \leq c_1 |\xi|^p + a(x) \quad \text{for all } (x, \xi) \in \Omega \times \mathbb{R}^d, \quad (5.3)$$

for some constants $c_0, c_1 > 0$ and some function $a \in L^1(\Omega)$. Let $G_k, G: W_0^{1,p}(\Omega) \rightarrow [0, +\infty)$ be defined by

$$G_k(u) := \int_{\Omega} g_k(x, \nabla u) dx \quad \text{and} \quad G(u) := \int_{\Omega} g(x, \nabla u) dx. \quad (5.4)$$

Theorem 5.1. *Let $\mu_k, \mu \in \mathcal{M}_+^{1,p}(\Omega \times \Omega)$ with $\mu_k(\Omega \times \Omega) < +\infty$ and $\mu(\Omega \times \Omega) < +\infty$. Let F_k, F be defined as in (5.1) with $f: \mathbb{R}^2 \rightarrow [0, +\infty)$ a continuous function satisfying (5.2). Let G_k, G be defined as in (5.4) with $g_k, g: \Omega \times \mathbb{R}^d \rightarrow \mathbb{R}$ Carathéodory functions satisfying (5.3). Suppose that*

$$\|\mu_k - \mu\|_{\square} \rightarrow 0 \quad \text{and} \quad \mu_k(\Omega \times \Omega) \rightarrow \mu(\Omega \times \Omega), \quad (5.5)$$

$$G = \Gamma\text{-}\lim_{k \rightarrow +\infty} G_k \quad \text{with respect to the weak convergence in } W_0^{1,p}(\Omega). \quad (5.6)$$

Then $F + G = \Gamma\text{-}\lim_{k \rightarrow +\infty} (F_k + G_k)$ with respect to the weak convergence in $W_0^{1,p}(\Omega)$.

Proof. By [6, Proposition 8.10] we have to prove that for all $u \in W_0^{1,p}(\Omega)$ the following properties hold:

(i) for all u_k converging to u weakly in $W_0^{1,p}(\Omega)$ we have

$$F(u) + G(u) \leq \liminf_{k \rightarrow +\infty} (F_k(u_k) + G_k(u_k));$$

(ii) there exist u_k converging to u weakly in $W_0^{1,p}(\Omega)$ such that

$$F(u) + G(u) = \lim_{k \rightarrow +\infty} (F_k(u_k) + G_k(u_k)).$$

Claim (i) follows from Corollary 4.7 with $v_k = u_k$ and from the liminf inequality for G_k , which follows from (5.6).

If $u \in W_0^{1,p}(\Omega) \cap L^\infty(\Omega)$, we deduce from (5.6) that there exists a sequence u_k converging to u weakly in $W_0^{1,p}(\Omega)$ such that $G(u_k) \rightarrow G(u)$ and $\|u_k\|_\infty \leq M$ for some constant M and

for all k (see for instance [4, Proposition 2.5]). Setting $\lambda = \max\{f(s, t) : |s| \leq M, |t| \leq M\}$, we may apply Theorem 4.6 with $v_k = u_k$ and $\tau^\lambda(f)$ in the place of f , obtaining (ii).

Let now $u \in W_0^{1,p}(\Omega)$. If $F(u) = +\infty$, then claim (ii) follows from claim (i). Suppose then that $F(u) < +\infty$. By the validity of claim (ii) for $\tau^\lambda(u)$ we obtain

$$F(\tau^\lambda(u)) + G(\tau^\lambda(u)) = \left(\Gamma\text{-lim sup}_{k \rightarrow +\infty} (F_k + G_k) \right) (\tau^\lambda(u)). \quad (5.7)$$

By (5.3) we have $G(\tau^\lambda(u)) \rightarrow G(u)$ as $\lambda \rightarrow +\infty$. Since $F(u) < +\infty$, by (5.2) and the Dominated Convergence Theorem we have $F(\tau^\lambda(u)) \rightarrow F(u)$ as $\lambda \rightarrow +\infty$ with $\lambda \in \Lambda$. By (5.7), using the lower semicontinuity of the Γ -limsup we get

$$F(u) + G(u) \geq \left(\Gamma\text{-lim sup}_{k \rightarrow +\infty} (F_k + G_k) \right) (u).$$

By [6, Proposition 8.10] we then obtain an inequality in claim (ii). The proof is completed by using claim (i). \square

Remark 5.2 (Mosco convergence). If the functionals G_k converge to G in the sense of the Mosco convergence in $W_0^{1,p}(\Omega)$; that is, for all $u \in W_0^{1,p}(\Omega)$ we have

(i) for all u_k converging to u weakly in $W_0^{1,p}(\Omega)$ we have

$$G(u) \leq \liminf_{k \rightarrow +\infty} G_k(u_k);$$

(ii) there exist u_k converging to u strongly in $W_0^{1,p}(\Omega)$ such that

$$G(u) = \lim_{k \rightarrow +\infty} G_k(u_k),$$

then also $F_k + G_k$ converges in the sense of the Mosco convergence in $W_0^{1,p}(\Omega)$.

Acknowledgements. This paper is based on work supported by the National Research Project (PRIN 2017) ‘‘Variational Methods for Stationary and Evolution Problems with Singularities and Interfaces’’, funded by the Italian Ministry of University, and Research. The authors are members of the Gruppo Nazionale per l’Analisi Matematica, la Probabilit a e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM).

References

- [1] A. Braides: *Γ -convergence for Beginners*. Oxford University Press, Oxford, 2002.
- [2] A. Braides, P. Cermelli, S. Dovetta: Γ -limit of the cut functional on dense graph sequences. *ESAIM Control Optim. Calc. Var.* **26** (2020), 26.

- [3] H. Brezis, F. Browder: A property of Sobolev spaces. *Comm. Partial Differential Equations* **4** (1979), 1077–1083.
- [4] G. Buttazzo, G. Dal Maso: Γ -limits of integral functionals. *J. Analyse Math.* **37** (1980), 145–185.
- [5] F.R. Chung, R.L. Graham, R.M. Wilson: Quasi-random graphs. *Combinatorica* **9** (1989), 345–362.
- [6] G. Dal Maso: *An Introduction to Γ -convergence*. Birkhäuser, Boston, 1993.
- [7] L.C. Evans, R.F. Gariepy: *Measure Theory and Fine Properties of Functions. Revised edition*. Textbooks in Mathematics. CRC Press, Boca Raton, FL, 2015.
- [8] M. Fukushima, Y. Oshima, M. Takeda: *Dirichlet Forms and Symmetric Markov Processes. Second revised and extended edition*. De Gruyter Studies in Mathematics, 19. Walter de Gruyter & Co., Berlin, 2011.
- [9] M. Grun-Rehomme: Caractérisation du sous-différentiel d'intégrales convexes dans les espaces de Sobolev. *J. Math. Pures Appl.* **56** (1977), 149–156.
- [10] J. Heinonen, T. Kilpeläinen, O. Martio: *Nonlinear Potential Theory of Degenerate Elliptic Equations. Unabridged republication of the 1993 original*. Dover Publications Inc., Mineola, NY, 2006.
- [11] L. Lovász: *Large Networks and Graph Limits*. American Mathematical Society Publications, Providence, RI, 2012.
- [12] U. Mosco: Convergence of convex sets and of solutions of variational inequalities. *Advances in Math.* **3** (1969), 510–585.
- [13] U. Mosco: Composite media and asymptotic Dirichlet forms, *J. Funct. Anal.* **123** (1994), 368–421,