



SCUOLA INTERNAZIONALE SUPERIORE DI STUDI AVANZATI

SISSA Digital Library

ADHM in 8d, coloured solid partitions and Donaldson-Thomas invariants on orbifolds

*Original*

ADHM in 8d, coloured solid partitions and Donaldson-Thomas invariants on orbifolds / Bonelli, G.; Fasola, N.; Tanzini, A.; Zenkevich, Y.. - In: JOURNAL OF GEOMETRY AND PHYSICS. - ISSN 0393-0440. - 191:(2023). [10.1016/j.geomphys.2023.104910]

*Availability:*

This version is available at: 20.500.11767/133610 since: 2023-08-15T17:07:31Z

*Publisher:*

*Published*

DOI:10.1016/j.geomphys.2023.104910

*Terms of use:*

Testo definito dall'ateneo relativo alle clausole di concessione d'uso

*Publisher copyright*

note finali coverpage

(Article begins on next page)

# ADHM in $8d$ , coloured solid partitions and Donaldson-Thomas invariants on orbifolds

---

**G. Bonelli,<sup>a,b,c</sup> N. Fasola,<sup>a,b,c</sup> A. Tanzini,<sup>a,b,c</sup> and Y. Zenkevich<sup>a,b,c,d,e,f</sup>**

<sup>a</sup>*S.I.S.S.A. – Scuola Internazionale di Studi Scientifici Avanzati  
via Bonomea, 265 - 34136 Trieste, Italy*

<sup>b</sup>*I.N.F.N. – Sezione di Trieste*

<sup>c</sup>*I.G.A.P. – Institute for Geometry and Physics  
via Beirut, 4 - 34100 Trieste, Italy*

<sup>d</sup>*ITEP – Institute for Theoretical and Experimental Physics  
Bolshaya Cheremushkinskaya street 25, 117218 Moscow, Russia*

<sup>e</sup>*ITMP MSU – Institute for Theoretical and Mathematical Physics  
Leninskie gory 1, 119991 Moscow, Russia*

<sup>f</sup>*MIPT – Moscow Institute of Physics and Technology  
Institutski per. 9, 141701 Dolgoprudny, Russia*

*E-mail: [bonelli@sissa.it](mailto:bonelli@sissa.it), [nfasola@sissa.it](mailto:nfasola@sissa.it), [tanzini@sissa.it](mailto:tanzini@sissa.it),  
[yegor.zenkevich@gmail.com](mailto:yegor.zenkevich@gmail.com)*

**ABSTRACT:** We study the moduli space of  $SU(4)$  invariant BPS conditions in supersymmetric gauge theory on non-commutative  $\mathbb{C}^4$  by means of an ADHM-like quiver construction and we classify the invariant solutions under the natural toric action in terms of solid partitions. In the orbifold case  $\mathbb{C}^4/G$ ,  $G$  being a finite subgroup of  $SU(4)$ , the classification is given in terms of coloured solid partitions. The statistical weight for their counting is defined through the associated equivariant cohomological gauge theory. We explicitly compute its partition function on  $\mathbb{C}^4$  and  $\mathbb{C}^2 \times (\mathbb{C}^2/\mathbb{Z}_2)$  which conjecturally provides the corresponding orbifold Donaldson-Thomas invariants.

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Topological gauge theory in eight dimensions</b>	<b>4</b>
<b>3</b>	<b>ADHM construction in eight dimensions</b>	<b>6</b>
3.1	Eight-dimensional equations	7
3.2	Derrick's theorem and noncommutativity	8
3.3	$8d$ ADHM construction	9
3.4	Formulation in complex coordinates	10
3.5	Matrix formulation	12
3.6	U(1) one-instanton	12
3.7	U(1) multi-instanton	13
<b>4</b>	<b>Counting solid partitions on orbifolds</b>	<b>14</b>
4.1	Orbifolding the quiver	14
4.2	Orbifold partition function	18
4.3	An example: $\mathbb{C}^2/\mathbb{Z}_2 \times \mathbb{C}^2$	21
<b>A</b>	<b>Review of the <math>4d</math> ADHM construction</b>	<b>24</b>
A.1	Self-dual connections in $4d$	24
A.2	ADHM data and equations	25
A.3	Spinor formalism	26
A.4	Hyperkähler reduction and complex structure(s)	27
A.5	Non-commutative deformation.	29
A.6	Matrix form of the non-commutative self-duality equations.	29
A.7	U(1) instantons and ideals	31
A.8	$4d$ fixed point instantons	32
<b>B</b>	<b><math>8d</math> instantons and sheaf cohomology</b>	<b>33</b>
B.1	Moduli spaces of $8d$ instantons via Beilinson's Theorem	33
B.2	Orbifold instantons and ADHM data decomposition	36

---

# 1 Introduction

The aim of this paper is the study of the moduli space of solutions of an eight dimensional analog of the celebrated self-duality equation  $F = \star F$  for the gauge theory curvature in four dimensions [1]. The equation in eight dimensions reads

$$(1.1) \quad F \wedge T = \star F$$

where  $F = dA + A \wedge A$  is the curvature of the gauge bundle,  $T$  is an invariant closed four-form and  $\star$  is the Hodge star operator with respect to a given Riemannian structure on the eight dimensional manifold. Equation (1.1) was introduced in [2] in 1982. The very existence of the invariant four-form  $T$  restricts the holonomy group of the eight dimensional manifold  $X$  to be contained in  $\text{Spin}(7)$  [3].

We provide the eight dimensional analog of the ADHM [4] construction for (1.1) with  $U(N)$  gauge group for  $X = \mathbb{C}^4$  and its discrete Calabi-Yau quotients. As we will discuss, differently from the real four dimensional case, genuine solutions to the above equation exist only on eight dimensional spaces whose local model is a non-commutative deformation of  $\mathbb{C}^4$ . The latter is obtained by deforming one of the moment maps à la Nekrasov-Schwarz [5]. This further breaks the holonomy group  $\text{Spin}(7) \rightarrow \text{Spin}(6) \simeq \text{SU}(4)$ , implying that  $X$  is a Calabi-Yau fourfold. The corresponding gauge theory can be engineered as the low energy limit of a  $D(-1)/D7$  system in a stabilising non trivial  $B$ -field background [6], aligned along the invariant 2-form associated to the deformed moment map. The more general configuration that we will study includes also a set of  $\overline{D7}$ s which act as a source of matter field/observables, as in [7, 8]. By resorting to our higher dimensional ADHM construction, we provide explicit solutions of equation (1.1) in the abelian case. Let us notice that  $\mathbb{C}^4$  admits a  $(\mathbb{C}^*)^3$  toric action compatible with its (trivial) Calabi-Yau structure, which naturally lifts to the moduli space of solutions to (1.1). We describe the invariant solutions supported at the fixed points of this toric action, by making crucial use of the non-commutative deformation, and find that these are classified by solid partitions. These are a four dimensional analog of Young diagrams built with hypercubes accumulating on the corner of  $\mathbb{R}_+^4$ . This provides a lift to four complex dimensions of the statistical crystal melting model based on plane partitions, see [9] for the  $U(1)$  case and [10, 11] for  $U(N)$ . All this construction has a natural extension on discrete quotients of  $\mathbb{C}^4$ , the fixed points being described in this case by coloured solid partitions.<sup>1</sup>

As it is well known, the ADHM construction for four-dimensional instantons is at the root of an isomorphism with the moduli space of framed torsion free coherent sheaves on  $\mathbb{P}^2$  [12]. We provide here an analog isomorphism between the moduli space of solution of (1.1) on non-commutative  $\mathbb{C}^4$  and the moduli space of framed

---

<sup>1</sup>Let us remark that on (partial) resolutions of  $\mathbb{C}^4$  orbifolds one can also construct abelian instantons whose gauge flux is along non-trivial two cycles.

torsion free coherent sheaves on  $\mathbb{P}^4$ , extending it also to the orbifold case by adapting the Kronheimer-Nakajima construction [13]. In this context the fixed points are described by ideal sheaves on  $\mathbb{C}^4$  and its quotients.

Building on equation (1.1), one can construct [14] a (semi)Topological Field Theory<sup>2</sup>, which is indeed a topological twist of the eight dimensional gauge theory describing the  $D(-1)/D7$  system at low energy [15, 16]. This provides the setting for BPS-bound states counting whose mathematical counterpart is given by Donaldson-Thomas theory on four-folds [17]. Let us remark that major progresses have been recently obtained on the compactification of the moduli space of solutions of (1.1) and a rigorous definition of the associated enumerative invariants [18–22]. A natural extension is to consider the theory on  $S^1 \times X$  computing the Witten index of  $D0/D8/\overline{D8}$  bound states whose mathematical counterpart is the lift to K-theory. On toric manifolds one can study the equivariant extension of the sTQFT so providing a geometrically motivated statistical weight for counting solid partitions which describe the fixed points of the gauge theory moduli space [8]. Chiral ring observables can be introduced via descent equations as in the four dimensional case [14]. Their explicit evaluation on  $\mathbb{C}^4$  via equivariant localisation recently appeared in [8, 21].

Let us remark that M-theory on local four-folds provides a geometric engineering description of supersymmetric gauge theories in three dimensions [23], analogously to the much better known case of local three-folds [24] which instead describes five dimensional supersymmetric gauge theories. Moreover, interesting classes of  $(0, 2)$  supersymmetric models in two-dimensions arise from  $D1$ -branes probing toric Calabi-Yau four-fold singularities [25]. It is thus interesting to study the eight dimensional BPS counting problem on some examples of local CY four-fold geometries. To this end, in this paper we also provide the generalisation of the eight-dimensional ADHM-like quiver to orbifolds  $\mathbb{C}^4/G$ , where  $G$  is a discrete subgroup of  $SU(4)$ . The fixed points in this case are classified by coloured solid partitions whose statistical weight depends on the representation of  $G$ . This boils down to count  $G$ -coloured hypercubes configurations whose colouring rules are dictated by the specific action of  $G$  on  $\mathbb{C}^4$ . This provides an eight dimensional analog of instanton counting on four-dimensional ALE spaces [26–33]. As an example, we explicitly address the associated K-theoretic counting problem on  $\mathbb{C}^2 \times \mathbb{C}^2/\mathbb{Z}_2$ .

There are a number of interesting problems to be addressed. Supersymmetric gauged linear sigma models in two dimensions modelled on the ADHM-like quivers presented in this paper can be studied via localisation technique. In the sphere case this could possibly shed light on the associated quantum cohomology and its relation with quantum integrable hydrodynamics analogously to the four dimensional ADHM quiver studied in [34–36]. The torus case would allow to compute the elliptic genus

---

<sup>2</sup>The "semi" refers to the dependence on the four-form  $T$ , which calibrates the volume of the four-cycles in the eight dimensional manifold.

of the eight dimensional ADHM moduli space and as such to provide an elliptic lift of Donaldson-Thomas invariants on fourfolds analogous to the one studied in [37] on three-folds. Finally defect operators can be investigated by a generalisation of the ADHM-like quivers similar to the one studied in [38–40] for the four-dimensional case and also by the eight dimensional generalisation of the *nested instantons* studied in [41, 42].

The structure of the paper is the following. In section 2 we discuss the equivariant extension of the eight-dimensional sTQFT, set up the gauge theory framework to count its BPS states and describe the relevant equivariant observables. In section 3 we provide the ADHM-like construction solving equation (1.1) on  $\mathbb{C}^4$  with  $B$ -field and describe the explicit solutions at the fixed points of the toric action in terms of solid partitions. In section 4 we extend the  $8d$  ADHM quiver construction to orbifolds  $\mathbb{C}^4/G$  and provide the computation of the equivariant partition function on the explicit example of  $\mathbb{C}^2 \times \mathbb{C}^2/\mathbb{Z}_2$ . In Appendix A we collect basic facts on the classical ADHM construction focusing on the points relevant for its  $8d$  generalisation. Appendix B is devoted to the discussion of the relation between  $8d$  instantons and sheaf cohomology: in B.1 we explain the relation between the two relevant moduli spaces via Beilinson’s theorem, while in B.2 we extend it to the orbifold case.

## 2 Topological gauge theory in eight dimensions

Let  $X$  be a real Riemannian eight dimensional differentiable manifold with a torsion free  $\text{Spin}(7)$  structure [43]. This is determined by a real covariantly constant spinor  $\psi_+$

$$T = \psi_+^T \Gamma^{[\wedge 4]} \psi_+ \in \Lambda^4(X)$$

where  $\Gamma \equiv \Gamma_\mu dx^\mu$  and  $\{\Gamma_\mu\}_{\mu=1,\dots,8}$  the  $\text{SO}(8)$   $\Gamma$ -matrices. We assume  $\psi_+$  to be of positive chirality and normalised as  $\psi_+^T \psi_+ = 1$ .

On  $X$  the vector spaces of  $p$ -forms  $\Lambda^p(X)$  split in irreducible representations of the holonomy group. In particular one has  $\Lambda^2(X) = \Lambda_7^2(X) \oplus \Lambda_{21}^2(X)$ . This split corresponds to the projections on  $\omega \in \Lambda^2(X)$  given by  $T \wedge \omega = -3\star\omega$  and  $T \wedge \omega = \star\omega$  respectively.

The Spin-bundles on  $X$  are isomorphic to  $S^+(X) \sim \Lambda^1(X)$  and  $S^-(X) \sim \Lambda^0(X) \oplus \Lambda_7^2(X)$ . A supersymmetric gauge theory can be formulated on a  $\text{Spin}(7)$  manifold via a topological twist which uses these isomorphisms [14–16]. The corresponding twisted supersymmetry transformations of the gauge theory can be made equivariant with respect to an isometry  $V$  and read

$$(2.1) \quad \begin{aligned} QA &= \Psi, & Q\Psi &= \iota_V F - iD\Phi, & Q\Phi &= \iota_V \Psi \\ Q\chi_7 &= B_7, & QB_7 &= \mathcal{L}_V \chi_7 + i[\Phi, \chi_7], & Q\eta &= \bar{\Phi}, & Q\bar{\Phi} &= \iota_V D\eta + i[\Phi, \eta], \end{aligned}$$

where  $V$  is any isometry of the  $\text{Spin}(7)$  structure, that is  $\star L_V = L_V \star$  and  $L_V T = 0$ , where  $L_V \equiv \iota_V d + d\iota_V$  is the Lie derivative. In (2.1),  $\mathcal{L}_V \equiv \iota_V D + D\iota_V$  is the

covariant Lie derivative. Notice that in (2.1),  $\Psi \in S^+(X) \sim \Lambda^1(X)$  and in the second line  $(\eta, \chi_7)$  and  $(\bar{\Phi}, B_7) \in S^-(X) \sim \Lambda^0(X) \oplus \Lambda_7^2(X)$ .

The supersymmetric action after the topological twist can be written as a topological term plus a Q-exact one as

$$(2.2) \quad S = \int_X T \wedge \text{Tr}(F \wedge F) + Q\nu,$$

where

$$(2.3) \quad \nu = \int_X \text{Tr} [i \star \chi_7 \wedge F + \Psi \wedge \star(Q\Psi)^\dagger + g^2 \chi_7 \wedge \star(Q\chi_7)^\dagger + \eta \wedge \star(Q\eta)^\dagger],$$

where  $g$  is the Yang-Mills coupling constant which in the topological theory is a gauge fixing parameter. In the path-integral, in the  $\delta$ -gauge  $g = 0$ , the  $B_7$  field appears as the Lagrange multiplier for the Spin(7)-instanton equation

$$(2.4) \quad F_7 = 0,$$

which is nothing but a rewriting of eq (1.1). In the following Section we will provide an ADHM-like description of the solutions to the above equation and their moduli space. This will turn out to have positive (virtual) dimension, inducing a  $U(1)_R$ -anomaly due to the presence of chiral fermionic zero-modes. In order to have a non-vanishing result one has thus to insert non-trivial observables in analogy with the well known case of Donaldson theory in four real dimensions [44]. The observables are given by non-trivial cohomology classes of the twisted supersymmetry (2.1), and can be obtained from an equivariant version of the usual descent equations, see [45] for the four-dimensional case. Indeed, the supersymmetry transformations (2.1) can be rewritten as the equivariant Bianchi identity for the curvature  $\mathcal{F} = F + \Psi + \Phi$  of the universal bundle as [46]

$$(2.5) \quad \mathcal{D}\mathcal{F} \equiv (-Q + D + i\nu_V)(F + \Psi + \Phi) = 0,$$

and expanding in the de Rham form degree. Picking an ad-invariant polynomial  $\mathcal{P}$  on the Lie algebra of the gauge group, we have

$$(2.6) \quad Q\mathcal{P}(\mathcal{F}) = (d + i\nu_V)\mathcal{P}(\mathcal{F}),$$

so that one can build the equivariant observables as intersection of the above with elements of the equivariant cohomology of the manifold,  $\Omega \in H_V^\bullet(X)$  as

$$(2.7) \quad \mathcal{O}(\Omega, \mathcal{P}) \equiv \int_X \Omega \wedge \mathcal{P}(\mathcal{F}).$$

In the path integral formulation of the gauge theory, we will actually consider the generating function of the equivariant observables through the determinant bundle

$$(2.8) \quad \mathcal{O}_{\text{det}}(\Omega) \equiv \int_X \Omega \wedge \det(m\mathbf{1} + \mathcal{F}),$$

where  $m$  is a generating parameter of the observables. In the calculations of the following Sections, we will consider the K-theoretic uplift of the above, or in other terms the index of the equivariant theory on  $X \times S^1$ .

### 3 ADHM construction in eight dimensions

In this section we describe an eight dimensional generalisation of the classical ADHM construction in four dimensions and show that it describes the moduli space of solutions to (1.1). For the sake of completeness we recall in Appendix A the four-dimensional ADHM construction and highlight few aspects of the latter which the reader could find useful to follow the eight dimensional generalisation.

Let us start by fixing the spinorial notation to write equation(1.1) and the ADHM representation of its solutions. The Cliff(8) gamma matrices can be chosen as  $16 \times 16$  real matrices of the form

$$(3.1) \quad \Gamma^\mu = \begin{pmatrix} 0 & \Sigma^\mu \\ \bar{\Sigma}^\mu & 0 \end{pmatrix},$$

where  $\Sigma^0 = \bar{\Sigma}^0 = 1_{8 \times 8}$  and  $\Sigma^i = -\bar{\Sigma}^i$  for  $i = 1, \dots, 7$ . The latter are real anti-symmetric matrices (they are in fact  $\sqrt{-1}$  times purely imaginary Cliff(7) gamma-matrices). Let  $S_\pm$  denote eight-dimensional real irreducible Majorana-Weyl spinor representations of Spin(8) of positive and negative chirality respectively. Since the representations  $S_\pm$  are real, the matrices of Spin(8) generators

$$(3.2) \quad \Gamma^{\mu\nu} = \frac{1}{2}[\Gamma^\mu, \Gamma^\nu] = \frac{1}{2} \begin{pmatrix} \Sigma^{[\mu} \bar{\Sigma}^{\nu]} & 0 \\ 0 & \bar{\Sigma}^{[\mu} \Sigma^{\nu]} \end{pmatrix}$$

are real and antisymmetric and so are the  $8 \times 8$  blocks  $\Sigma^{\mu\nu} = \frac{1}{2}\Sigma^{[\mu} \bar{\Sigma}^{\nu]}$  and  $\bar{\Sigma}^{\mu\nu} = \frac{1}{2}\bar{\Sigma}^{[\mu} \Sigma^{\nu]}$ . Formulated differently, we have (cf. (A.16)) an isomorphism of three Spin(8) representations spaces, each represented by real and antisymmetric  $8 \times 8$  matrices:

$$(3.3) \quad \Lambda^2 S_+ = \Lambda^2 S_- = \Lambda^2 \mathbb{R}^8 = \text{adj}_{SO(8)}.$$

The triality of Spin(8) permutes  $S_+$ ,  $S_-$  and  $\mathbb{R}^8$ , which is the defining representation of  $SO(8)$ . Notice that each of the two sets of 28 matrices  $(\Sigma^{\mu\nu})_{\alpha\beta}$  or  $(\bar{\Sigma}^{\mu\nu})_{\alpha\beta}$  form a basis in the space of real antisymmetric matrices<sup>3</sup>. Due to this fact, the following Fierz identities hold:

$$(3.4) \quad (\Sigma^{\mu\nu})_{\alpha\beta} (\Sigma^{\mu\nu})_{\gamma\delta} = (\bar{\Sigma}^{\mu\nu})_{\alpha\beta} (\bar{\Sigma}^{\mu\nu})_{\gamma\delta} = -8(\delta_{\alpha\delta} \delta_{\beta\gamma} - \delta_{\beta\delta} \delta_{\alpha\gamma}).$$

The coefficient in the l.h.s. of Eq. (3.4) can be obtained by contracting  $\beta$  and  $\gamma$  indices and using the relations of the Clifford algebra to get  $\Gamma^{\mu\nu} \Gamma^{\mu\nu} = -56 \cdot 1_{16 \times 16}$ .

---

<sup>3</sup>A third basis of matrices corresponding to the representation  $\Lambda^2 \mathbb{R}^8$  is given by  $(\delta_\alpha^\mu \delta_\beta^\nu - \delta_\beta^\mu \delta_\alpha^\nu)$ .



### 3.1 Eight-dimensional equations

There is no way to formulate a first order equation for  $F_{\mu\nu}$  in  $8d$  in an  $SO(8)$ -invariant manner. Another way to formulate this is to say that there is no  $SO(8)$ -invariant four-index tensor  $T^{\mu\nu\lambda\rho}$ , which can be used to write

$$(3.5) \quad \lambda F_{\mu\nu} = \frac{1}{2} T^{\mu\nu\lambda\rho} F_{\lambda\rho}.$$

with  $\lambda$  being some eigenvalue. However, if we make a choice of a constant spinor on  $\mathbb{R}^8$  we can build from it a tensor  $T^{\mu\nu\lambda\rho}$  invariant under  $\text{Spin}(7) \subset \text{Spin}(8)$ . This is the largest possible symmetry subgroup which can be preserved by equations of the form (3.5) in eight space-time dimensions. For this construction let us fix  $\psi_+ \in S_+$  such that  $\psi_+^T \psi_+ = 1$  and write

$$(3.6) \quad T^{\mu\nu\lambda\rho} = \psi_+^T \Gamma^{\mu\nu\lambda\rho} \psi_+,$$

where

$$(3.7) \quad \Gamma^{\mu\nu\lambda\rho} = \frac{1}{4!} \Gamma^{[\mu} \Gamma^\nu \Gamma^\lambda \Gamma^{\rho]}$$

The tensor  $T^{\mu\nu\lambda\rho}$  then satisfies the  $8d$  self-duality equation<sup>4</sup>

$$(3.8) \quad T^{\mu\nu\lambda\rho} = \epsilon^{\mu\nu\lambda\rho\alpha\beta\gamma\delta} T^{\alpha\beta\gamma\delta},$$

since

$$(3.9) \quad \epsilon^{\mu\nu\lambda\rho\alpha\beta\gamma\delta} \Gamma^{\alpha\beta\gamma\delta} = \Gamma^{\mu\nu\lambda\rho} \Gamma^9.$$

For definiteness we take<sup>5</sup>  $\psi_+^\alpha = \delta_0^\alpha$ . The choice of the spinor  $\psi_+$  allows us to split  $S_+$  into the one-dimensional subspace proportional to  $\psi_+$  and the seven-dimensional orthogonal complement which we call  $\tilde{S}_+$ . In a group-theoretical language this corresponds to the splitting of the representation  $S_+$  into  $\mathbf{1} \oplus \mathbf{7}$  under the subgroup  $\text{Spin}(7) \subset \text{Spin}(8)$ .

The irreducible two-form representation  $\Lambda^2 \mathbb{R}^8 = \mathbf{28}$  of  $\text{Spin}(8)$  splits into a sum of two irreps  $\mathbf{7} \oplus \mathbf{21}$  of  $\text{Spin}(7)$ . These two irreps correspond to two different eigenvalues  $\lambda = 1$  and  $\lambda = -3$  respectively in the first order field equations (3.5). In this way the splitting allows us to write two different  $\text{Spin}(7)$ -invariant conditions<sup>6</sup> on the field strength  $F_{\mu\nu}$ . The conditions correspond to the vanishing of the component

---

<sup>4</sup>We could have started with a negative chirality spinor  $\psi_-$  corresponding to anti-self-dual  $T^{\mu\nu\lambda\rho}$ . The resulting construction is isomorphic due to the triality of  $SO(8)$ .

<sup>5</sup>In our conventions the spinor indices run from 0 to 7 similarly to the indices of the  $\mathbb{R}^8$  vectors.

<sup>6</sup>These two conditions may be viewed as analogues of the self-duality and anti-self-duality conditions in  $4d$ . However, the latter are more similar to choosing the opposite chirality spinor  $\psi_-$  instead of  $\psi_+$ .

of  $F_{\mu\nu}$  lying in one of the two irreps of  $\text{Spin}(7)$ , or equivalently to the eigenspaces corresponding to two different eigenvalues in Eq. (3.5). The choice  $\lambda = 1$  gives Eq. (1.1).

As discussed in [2] (1.1) reads then in spinorial form as

$$(3.10) \quad (\Sigma^{\mu\nu})_{\alpha\beta}\psi_+^\beta F_{\mu\nu} = (\Sigma^{\mu\nu})_{\alpha 0}F_{\mu\nu} = 0,$$

which imposes  $7N^2$  equations, so that – together with  $N^2$  gauge fixing conditions for the gauge group  $U(N)$  – eliminate all functional degrees of freedom from  $A_\mu(x)$  and therefore a finite dimensional moduli space  $\mathcal{M}_{N,k}$  of solutions remains. Similarly to the  $4d$  case, one can view  $\mathcal{M}_{N,k}$  as an *octonionic* quotient of the space of connections  $\mathcal{A}$  by the gauge group  $\mathcal{G}$ . Indeed we can introduce seven natural symplectic forms  $\omega_{\mu\nu}^{(A)} = (\Sigma^{\mu\nu})_{A0}$  on  $\mathbb{R}^8$  and use them to write seven symplectic structures on  $\mathcal{A}$ :

$$(3.11) \quad \Omega_A[\delta_1 A_\mu(x), \delta_2 A_\mu(y)] = \int_{\mathbb{R}^8} T \wedge \omega^{(A)} \wedge \delta_1 A(x) \wedge \delta_2 A(x),$$

where  $T$  denotes the four-form with components  $T^{\mu\nu\lambda\rho}$ . Then the  $7N^2$  conditions (3.10) correspond to the vanishing of the seven moment maps

$$(3.12) \quad \mu^A[\phi(x)] = \int_{\mathbb{R}^8} T \wedge \omega^{(A)} \wedge \text{tr} \phi(x) F,$$

and we have the "octonionic" quotient

$$(3.13) \quad \mathcal{M}_{N,k} = \mathcal{A}////////\mathcal{G}.$$

### 3.2 Derrick's theorem and noncommutativity

Any solution of the first order equations (3.5) is automatically a solution of the  $8d$  Yang-Mills equations. Indeed,

$$(3.14) \quad D_\mu F_{\mu\nu} = \frac{1}{2} T_{\mu\nu\lambda\rho} D_\mu F_{\lambda\rho} = \frac{1}{2} T_{\mu\nu\lambda\rho} D_{[\mu} F_{\lambda\rho]} = 0$$

where we have used the Bianchi identity for  $F_{\mu\nu}$  and the fact that  $T_{\mu\nu\lambda\rho}$  is totally antisymmetric. We are looking for localized solutions, *i.e.* for those sufficiently rapidly decaying at infinity in order to have finite action. Then, the solutions of (3.5), if any exist, should be true extrema of the Yang-Mills action. The well-known Derrick's theorem states that in dimensions greater than four no such localized solution are possible. The idea of the proof is to provide for any non-singular field configuration a continuous family of configurations with lower action, so that no true minimum can exist. The family of configurations is obtained by scaling the initial configuration into smaller and smaller volume. A simple power counting then shows that the action on the scaled down configuration is lower.

Thus, classically, the moduli space  $\mathcal{M}_{N,k}$  of solutions is empty. However, there is a natural way to deform the problem to get a non-empty space of solutions by introducing noncommutativity. The commutation relations for the coordinates are similar to those of the  $4d$  case (cf. (A.32)):

$$(3.15) \quad [x^\mu, x^\nu] = i\zeta(\omega^{-1})^{\mu\nu},$$

where  $\zeta$  is a real parameter and  $\omega_{\mu\nu}$  is a non-degenerate constant 2-form on  $\mathbb{R}^8$ . In this case Derrick's theorem doesn't apply, since the coordinates cannot be rescaled without affecting their commutation relations (3.15). To put it another way, the non-commutativity introduces an additional fundamental scale  $\sqrt{\zeta}$  into the problem, which puts a limit on how much one can scale down localized field configurations. So, even when no classical non-singular solutions to field equations exist, additional solutions to the non-commutative version of the problem having typical size  $\sqrt{\zeta}$  may appear. This is exactly the situation we have with Eq. (3.5), where there is a finite-dimensional moduli space of solutions after the non-commutative deformation. This will be our definition of  $\mathcal{M}_{N,k}$ . In the next section we will describe this moduli space using an analogue of the ADHM construction.

### 3.3 $8d$ ADHM construction

The ADHM equations for the  $8d$  instantons were written in [7, 8]. They correspond to bound states of  $k$  D0 and  $N$  D8 branes in a suitable  $B$ -field background. As in sec. A.5, the  $B$ -field introduces non-commutativity into the  $8d$  gauge theory. Also, as we have explained in sec. 3.2 it allows for the very existence of solutions to the first order equations (3.5), corresponding to stable low-energy bound states of D-branes.

To introduce the non-commutativity we pick one of the seven complex structures on  $\mathbb{R}^8$ , or correspondingly one of the symplectic forms  $\omega_{\mu\nu}^{(A)}$  to represent the Kähler form. For definiteness we choose  $\omega_{\mu\nu}^{(1)}$  and denote the projection of the two-form  $B$  on  $\omega_{\mu\nu}^{(1)}$  by  $\zeta$ .

The choice of the complex structure breaks down the Spin(7) symmetry of the seven first order equations (3.10) further to Spin(6) = SU(4). Equivalently we can say that by choosing a complex structure we introduce one more fixed chiral spinor  $\chi_+^\alpha = \delta_1^\alpha$  into our theory (which corresponds to the index 1 in  $\omega_{\mu\nu}^{(1)}$ ). The seven equations (3.10), transforming as a Spin(7) spinor, split into an SU(4) singlet (corresponding to the component in the direction of  $\chi_+$ ) and further six equations lying in the representation

$$(3.16) \quad \mathbf{6} = \mathbb{R}_{SO(6)}^6 = (\Omega^{(2,0)} \oplus \Omega^{(0,2)}\mathbb{C}^4)_+,$$

of complex two-forms obeying  $\alpha_{z_i z_j} = \epsilon_{z_i z_j z_k z_l} \alpha_{\bar{z}_k \bar{z}_l}$ .

Unlike in the  $4d$  case the  $B$ -field not only adds a constant to the value of one of the seven moment maps  $\mu^A$ , but introduces new degrees of freedom corresponding to the rectangular matrix  $I$ , which only appears in one of the moment map equations.

Somewhat similarly to the  $4d$  case we introduce a  $(8k + N) \times 8k$  matrix  $\Delta(x)$ , which can be written as<sup>7</sup>

$$(3.17) \quad \Delta(x) = \begin{pmatrix} (B_\mu - x_\mu 1_{k \times k}) \otimes \bar{\Sigma}^\mu \\ I^\dagger \otimes (\psi_+^\dagger + i\chi_+^\dagger) \end{pmatrix},$$

where  $B_\mu$  are eight Hermitian  $k \times k$  matrices,  $\Sigma^\mu$  are defined in Eq. (3.1),  $I^\dagger$  is an  $N \times k$  matrix.

We will be looking for solutions of

$$(3.18) \quad \Delta^\dagger(x)U(x) = 0,$$

with  $\Delta$  satisfying certain moment map conditions. However, differently from the  $4d$  case, these conditions do not imply that  $\Delta^\dagger\Delta = 1_{8 \times 8} \otimes f_{k \times k}^{-1}$ <sup>8</sup>. Indeed, to solve Eq.(3.10), it is enough to impose

$$(3.20) \quad \Delta^\dagger\Delta\psi_+ = \psi_+ \otimes f_{k \times k}^{-1}.$$

In our convention for  $\psi_+$  and  $\chi_+$  Eq. (3.20) has explicit components

$$(3.21) \quad (\Delta^\dagger\Delta)_{A0} = 0, \quad A = 1, \dots, 7,$$

or more explicitly

$$(3.22) \quad i(\omega^{(A)})_{\mu\nu}^{-1}[B_\mu, B_\nu] + \delta^{A,1}II^\dagger = \zeta\delta^{A,1}, \quad A = 1, \dots, 7.$$

### 3.4 Formulation in complex coordinates

By explicitly using the complex structure, we can rewrite Eq.(3.22) in the form given in [7]. The  $8d$  ADHM data contains four complex  $k \times k$  matrices  $B_a$  and a complex  $k \times N$  matrix  $I$  and the equations read

$$(3.23) \quad \sum_{a=1}^4 [B_a, B_a^\dagger] + II^\dagger = \zeta 1_{k \times k},$$

$$(3.24) \quad [B_a, B_b] - \frac{1}{2}\epsilon_{abcd}[B_c^\dagger, B_d^\dagger] = 0.$$

---

<sup>7</sup>Notice that the combination  $\psi_+ + i\chi_+$  transforms in the complex one-dimensional Weyl spinor representation of the  $\text{Spin}(2) = U(1)$  part of  $\text{Spin}(6) \times \text{Spin}(2) \subset \text{Spin}(8)$ .

<sup>8</sup>Indeed, since  $(\Sigma^{\mu\nu})_{\alpha\beta}$  is a complete basis of real antisymmetric  $8 \times 8$  matrices,

$$\begin{aligned} \Delta^\dagger(x)\Delta(x) &= II^\dagger \otimes (\psi_+ + i\chi_+)(\psi_+^T - i\chi_+^T) + \\ &+ \frac{1}{2}([B_\mu, B_\nu] + i\zeta\omega_{\mu\nu}^{(1)} 1_{k \times k}) \otimes \Sigma^{\mu\nu} + (B_\mu - x_\mu 1_{k \times k})(B_\mu - x_\mu 1_{k \times k}) \otimes 1_{8 \times 8} = 1_{8 \times 8} \otimes f_{k \times k}^{-1} \end{aligned}$$

implies (recall that  $\omega_{\mu\nu}^{(1)} = (\Sigma^{\mu\nu})_{01}$ )

$$(3.19) \quad [B_\mu, B_\nu] + i\zeta\omega_{\mu\nu}^{(1)} \otimes 1_{k \times k} + i\omega_{\mu\nu}^{(1)}II^\dagger = 0,$$

which gives 28 matrix equations, instead of just seven.

The matrix  $\Delta^\dagger$  defined in Eq. (3.17) can be written quite explicitly in complex coordinates. Indeed, the main ingredient of  $\Delta^\dagger$  is the matrix  $x_\mu \Sigma^\mu : S_- \rightarrow S_+$ , which acts from one Majorana-Weyl representation of Spin(8) to another. Under the SU(4) subgroup the spinor representations  $S_\pm$  split into sums of even and odd parts of the exterior algebra:

$$(3.25) \quad S_+ = \mathbf{1} \oplus \mathbf{6} \oplus \mathbf{1} = \Lambda^0 \mathbb{C}^4 \oplus \Lambda^2 \mathbb{C}^4 \oplus \Lambda^4 \mathbb{C}^4,$$

$$(3.26) \quad S_- = \mathbf{4} \oplus \bar{\mathbf{4}} = \Lambda^1 \mathbb{C}^4 \oplus \Lambda^3 \mathbb{C}^4.$$

The matrix  $x_\mu \Sigma^\mu$  then acts on the exterior powers as an operator

$$(3.27) \quad \iota_{z_a} \partial_{z_a} + \bar{z}_a dz_a,$$

where  $\iota_{z_a} \partial_{z_a}$  is the substitution of the vector field  $z_a \partial_{z_a}$ . In this way we get:

$$(3.28) \quad \Delta^\dagger = \left( \begin{array}{cccc|cccc|c} b_1 & b_2 & b_3 & b_4 & 0 & 0 & 0 & 0 & I \\ \hline b_2^\dagger & -b_1^\dagger & 0 & 0 & 0 & 0 & b_4 & -b_3 & 0 \\ b_3^\dagger & 0 & -b_1^\dagger & 0 & 0 & -b_4 & 0 & b_2 & 0 \\ b_4^\dagger & 0 & 0 & -b_1^\dagger & 0 & b_3 & -b_2 & 0 & 0 \\ \hline 0 & b_3^\dagger & -b_2^\dagger & 0 & b_4 & 0 & 0 & -b_1 & 0 \\ 0 & -b_4^\dagger & 0 & -b_2^\dagger & b_3 & 0 & -b_1 & 0 & 0 \\ 0 & 0 & b_4^\dagger & -b_3^\dagger & b_2 & -b_1 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & b_1^\dagger & b_2^\dagger & b_3^\dagger & b_4^\dagger & 0 \end{array} \right),$$

where we have abbreviated  $b_a = B_a - z_a$ . Notice that unlike in the four-dimensional case both representations  $S_\pm$  are real, i.e. they admit outer automorphisms which square to one. Explicitly these automorphisms are given by the matrices

$$(3.29) \quad \tau_+ = \left( \begin{array}{cccccccc|c} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ \hline 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \hline 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right), \quad \tau_- = \left( \begin{array}{c|c} 0 & 1_{4 \times 4} \\ \hline 1_{4 \times 4} & 0 \end{array} \right).$$

The square part of the matrix  $\Delta$  commutes with the automorphisms in the sense that

$$(3.30) \quad \tau_+ \Delta_{\text{square}}^\dagger \tau_- = \Delta_{\text{square}}^*.$$

### 3.5 Matrix formulation

After choosing the non-commutative deformation (3.15), the original instanton equations (3.10) become:

$$(3.31) \quad F_{ab} = \frac{1}{2} \epsilon_{abcd} F_{\bar{c}\bar{d}},$$

$$(3.32) \quad \sum_{a=1}^4 F_{a\bar{a}} = 0.$$

Let us rewrite these equations in terms of the matrix variables analogously to the noncommutative  $4d$  case recalled in sec. A.6. Plugging the  $Z_a$  variables in the instanton equations (3.31), (3.32) we get:

$$(3.33) \quad [Z_a, Z_b] = \frac{1}{2} \epsilon_{abcd} [Z_c^\dagger, Z_d^\dagger],$$

$$(3.34) \quad \sum_{a=1}^4 [Z_a^\dagger, Z_a] = 2\zeta.$$

Again the nontrivial r.h.s. in Eq. (3.34) arises because of the noncommutativity of  $z_a$  and  $z_a^\dagger$ . The vacuum solution of Eqs. (3.33), (3.34) is given by

$$(3.35) \quad Z^a = z^a,$$

and corresponds to vanishing gauge potential  $A_a$ . Let us now discuss the simplest non trivial solutions.

### 3.6 U(1) one-instanton

Nontrivial solutions to Eqs. (3.33), (3.34) correspond to nontrivial ideals in the ring of polynomials in four variables. Let us consider the simplest solution corresponding to a single abelian instanton sitting at the origin of  $\mathbb{C}^4$ . In this case (cf. also [47, Eq. (3.18)])

$$(3.36) \quad Z_a = S_{[[[1]]]} z_a f_{[[[1]]]}(N) S_{[[[1]]]}^\dagger,$$

where  $N = \sum_{a=1}^4 a_a^\dagger a_a$  and

$$(3.37) \quad f_{[[[1]]]}(N) = \left( 1 - \frac{24}{N(N+1)(N+2)(N+3)} \right)^{\frac{1}{2}},$$

and  $S_{[[[1]]]}$  is the partial isometry of the Hilbert space (which is  $\mathcal{H} = \mathbb{C}[z_1, z_2, z_3, z_4]$ ), satisfying

$$(3.38) \quad S_{[[[1]]]} S_{[[[1]]]}^\dagger = 1, \quad S_{[[[1]]]}^\dagger S_{[[[1]]]} = 1 - |0, 0, 0, 0\rangle \langle 0, 0, 0, 0| = P_{\mathcal{H} \setminus \{0,0,0,0\}}.$$

Notice that  $N$  in the denominator of  $f_{[[[1]]]}(N)$  is never zero, because the state  $|0, 0, 0, 0\rangle$  is projected out by the partial isometry .

### 3.7 U(1) multi-instanton

By taking the square of equations (3.33) we deduce that the operators  $Z_a$  commute with each other:

$$(3.39) \quad [Z_a, Z_b] = 0.$$

As we have noticed above the multi-instanton solutions correspond to ideals in the ring  $\mathcal{H} = \mathbb{C}[z_1, z_2, z_3, z_4]$ . Having such an ideal  $\mathcal{I}$ , we define a partial isometry  $S_{\mathcal{I}}$ , which satisfies

$$(3.40) \quad S_{\mathcal{I}} S_{\mathcal{I}}^\dagger = 1, \quad S_{\mathcal{I}}^\dagger S_{\mathcal{I}} = P_{\mathcal{I}},$$

where  $P_{\mathcal{I}}$  is the projection operator on the ideal  $\mathcal{I}$ . The matrix  $\Delta^\dagger$  contains the information about the resolution of the ideal corresponding to the solution of the ADHM equations. Consider for example the 1-instanton solution (3.36). It corresponds to the ideal  $\mathcal{I}_{[[[1]]]}$  of polynomials without constant terms. The resolution of this ideal is written as the following exact sequence:

$$(3.41) \quad 0 \rightarrow \mathcal{O} \xrightarrow{\mu_4} \mathcal{O}^{\oplus 4} \xrightarrow{\mu_3} \mathcal{O}^{\oplus 6} \xrightarrow{\mu_2} \mathcal{O}^{\oplus 4} \xrightarrow{\mu_1} \mathcal{O} \xrightarrow{p} \mathcal{I}_{[[[1]]]} \rightarrow 0,$$

where  $\mathcal{O} = \mathbb{C}[z_1, z_2, z_3, z_4]$ ,  $p$  is the projection and the linear operators  $\mu_i$  are

$$(3.42) \quad \mu_1 = (z_1 \ z_2 \ z_3 \ z_4),$$

$$(3.43) \quad \mu_2 = \begin{pmatrix} z_2 & z_3 & z_4 & 0 & 0 & 0 \\ -z_1 & 0 & 0 & z_3 & -z_4 & 0 \\ 0 & -z_1 & 0 & -z_2 & 0 & z_4 \\ 0 & 0 & -z_1 & 0 & z_2 & -z_3 \end{pmatrix},$$

$$(3.44) \quad \mu_3 = \begin{pmatrix} 0 & 0 & z_4 & z_3 \\ 0 & -z_4 & 0 & z_2 \\ 0 & z_3 & -z_2 & 0 \\ z_4 & 0 & 0 & -z_1 \\ z_3 & 0 & -z_1 & 0 \\ z_2 & -z_1 & 0 & 0 \end{pmatrix}, \quad \mu_4 = \begin{pmatrix} z_1 \\ z_2 \\ z_3 \\ z_4 \end{pmatrix}.$$

Notice that these operators are very similar to those featuring in  $\Delta^\dagger$  in Eq. (3.28). This similarity seems to be a generic property of any ADHM-like construction.

We are interested in the solutions which are fixed points of the  $U(1)^3 \subset SU(4)$  action on  $\mathbb{C}^4$ . Those correspond to monomial ideals in the ring  $\mathbb{C}[z_1, z_2, z_3, z_4]$  and are enumerated by solid partitions. We denote the ideal corresponding to a solid partition  $\sigma$  by  $\mathcal{I}_\sigma$ . The fixed point multi-instanton solutions can be obtained with an ansatz similar to Eq. (3.36), but now the function  $f(r)$  does not need to be symmetric in  $N_a = a_a^\dagger a_a$  (no summation over  $a$ ), so that

$$(3.45) \quad Z_a = U_\sigma z_a f_a^{(\sigma)}(N_1, N_2, N_3, N_4) U_\sigma^\dagger$$

We thus have to determine four functions  $f_a^{(\sigma)}(N_1, N_2, N_3, N_4)$  of four variables. Eqs. (3.39), (3.34) imply the following recurrence relations<sup>9</sup> for  $f_a$

$$(3.46) \quad f_a(N_b + 1)f_b(N) = f_b(N_a + 1)f_a(N),$$

$$(3.47) \quad \sum_{a=1}^4 \left\{ (f_a(N))^2(N_a + 1) - (f_a(N_a - 1))^2 N_a \right\} = 4.$$

Eqs. (3.46) are “flatness” conditions for  $f_a$  and can be solved explicitly. Indeed, one can see that

$$(3.48) \quad f_a(N) = \frac{h(N)}{h(N_a + 1)}$$

solves Eqs. (3.46) for any function  $h(N)$ . There remains a single recurrence relation (3.47) for  $h(r)$ , which reads

$$(3.49) \quad \sum_{a=1}^4 \left\{ \frac{h(N)^2}{h(N_a + 1)^2} (N_a + 1) - \frac{h(N_a - 1)^2}{h(N)^2} N_a \right\} = 4.$$

An example of instanton solution with second Chern class equal to  $\frac{k(k+1)(k+2)(k+3)}{24}$  is given by

$$(3.50) \quad Z_a = S_k z_a f^{(k)}(N) S_k^\dagger,$$

where  $S_k$  now avoids all the states corresponding to monomials of degree at most  $k$  and

$$(3.51) \quad f^{(k)}(N) = \left( 1 - \frac{k(k+1)(k+2)(k+3)}{N(N+1)(N+2)(N+3)} \right)^{\frac{1}{2}}.$$

Notice that all four functions  $f_a(N)$  are in this case equal to each other. The solutions (3.51) correspond to “pentachoron” solid partitions (decreasing sequences of tetrahedral plane partitions), e.g. [[[1]]] for  $k = 1$  or [[[2, 1], [1]], [[1]]] for  $k = 2$ .


## 4 Counting solid partitions on orbifolds

### 4.1 Orbifolding the quiver

As we showed in section 3, eight-dimensional instanton dynamics is encoded in the representation theory of the quiver depicted in Fig.1, with relations

$$(4.1) \quad [B_a, B_b] = 0, \quad 1 \leq a < b \leq 4$$





**Figure 1:** Local model for the Quot scheme of points.

and a stability condition.<sup>10</sup> The moduli space of its stable representations  $\mathcal{M}_{k,N}$  is isomorphic to the quot scheme of points  $\text{Quot}_{\mathbb{A}^4}(\mathcal{O}^{\oplus N}, k)$  or, equivalently, to the moduli space of framed torsion free sheaves on  $\mathbb{P}^4$ , as we briefly show in appendix B.1. This isomorphism follows from an application of Beilinson’s theorem or, equivalently, from an infinitesimal argument due to [48].

The next natural step is then to study eight-dimensional instanton dynamics on orbifolds of  $\mathbb{C}^4$ . Then we let  $G$  be a finite subgroup  $G \subset \text{SU}(4)$ , and study instantons on  $\mathbb{C}^4/G$ . From the open string theory perspective this amounts to consider twisted representations of the Chan-Paton factors under the discrete group  $G$ , which manifest in the low energy quiver dynamics of fractional and regular branes [49]. The mathematical counterpart of the quiver ADHM-like description for orbifold instantons on  $\mathbb{C}^4$  can be obtained as an application of Beilinson’s theorem which is outlined in appendix B.2.

The useful thing to point out here is that the monad description for the moduli space  $\mathcal{M}_{k,N}^G$  of orbifold instantons, which can be obtained by means of homological algebra, is then given in terms of a sequence of maps between vector spaces. These maps can be easily understood as an equivariant decomposition (in terms of the  $G$ -action) of the maps and vector spaces arising in the quiver description of  $\mathcal{M}_{k,N}$ . If we introduce the action of  $G$  on the coordinates  $z_\alpha$  in  $\mathbb{C}^4$  by  $r_\alpha z_\alpha$ , we have a decomposition of the fundamental representation  $Q = \rho_{r_1} \oplus \cdots \oplus \rho_{r_4}$ , where  $\rho_{r_\alpha}$  denotes the irreducible representation of  $G$  with weight  $r_\alpha$ . This decomposition also defines a coloring  $\mathbb{N}^{\oplus 4} \rightarrow G$  by

$$(n_1, n_2, n_3, n_4) \mapsto \rho_{r_1}^{\otimes n_1} \otimes \rho_{r_2}^{\otimes n_2} \otimes \rho_{r_3}^{\otimes n_3} \otimes \rho_{r_4}^{\otimes n_4}.$$

Correspondingly we also have decompositions of the vector spaces

$$W = \bigoplus_r W_r \otimes \rho_r^\vee, \quad V = \bigoplus_r V_r \otimes \rho_r^\vee,$$

<sup>9</sup>We denote by  $f_a(N_b + 1)$  the function  $f_a(N)$  with  $N_b \rightarrow N_b + 1$ .

<sup>10</sup>Strictly speaking we showed that the moduli space of  $\text{SU}(4)$  instantons can be identified with a space of matrices cut by equations (4.1) plus an additional real constraint, modulo gauge symmetry. Though it has not rigorously been proved as of this writing, it is believed that the last real condition can be traded for a stability condition, so that the moduli space of instantons may be identified with stable representations of a quiver with relations.

where all of the  $W_r, V_r$  are finite dimensional vector spaces carrying a trivial  $G$ -action. The corresponding decomposition of the dimensions  $k = \dim_{\mathbb{C}} V$ ,  $N = \dim_{\mathbb{C}} W$  is then induced as

$$k = \sum_r k_r = \sum_r \dim_{\mathbb{C}} V_r, \quad N = \sum_r N_r = \sum_r \dim_{\mathbb{C}} W_r.$$

Here  $k_r = \dim_{\mathbb{C}} V_r$  represents the fractional instanton charge in the  $\rho_r^{\vee}$  representation of  $G$ , which, from an equivariant localisation point of view, will specify the number of boxes of  $r$ -th type in a  $G$ -colored solid partition. On the other hand, the gauge sheaf at infinity transforms in a given representation  $\rho$  of  $G$ , and the  $N_r$  dimensions determine the multiplicities of the decomposition of  $\rho$  in irreducible representations. If the theory is abelian, *i.e.*  $N = 1$ , only one of the  $N_r$  is not zero, and equal to one, while in the case of a non-abelian theory one is given with a plethora of different possibilities. We will restrict our attention to the abelian case for the moment. The decomposition of  $V$  then induces a decomposition of the linear maps  $B_{\alpha} \in \text{Hom}_G(V, Q \otimes V)$  as

$$B = \bigoplus_r (B_1^r, B_2^r, B_3^r, B_4^r),$$

so that  $B_{\alpha}^r : V_r \rightarrow V_{r+r_{\alpha}}$ . This decomposition immediately gives us the orbifold generalisation of the 8d ADHM equations as

$$(4.2) \quad B_{\beta}^{r+r_{\alpha}} B_{\alpha}^r = B_{\alpha}^{r+r_{\beta}} B_{\beta}^r.$$

In general, even for abelian theories, one should consider different cases (corresponding in the abelian case to which one of the  $W_r$  vector spaces has non-vanishing dimension). These different choices correspond to instanton configurations with different asymptotics at infinity. One can however argue along the lines of [50] that moduli spaces corresponding to different asymptotics are isomorphic, so that when computing partition functions we will just consider the distinguished boundary condition  $\overline{N} = (1, 0, \dots, 0)$ .

As in the lower dimensional cases (see [13, 50, 51]) all the information about the moduli space of orbifold instantons can be encoded in the datum of a quiver generalising the McKay quiver, which is determined by the representation theory data of the  $G$ -action. This quiver will moreover encode the decomposition of the usual ADHM data according to the  $G$ -action and it will have the orbifold ADHM equations as relations. One then starts considering all the irreducible representations  $\widehat{G}$  of  $G \subset \text{SL}(4, \mathbb{C})$ . To each representation in  $\widehat{G}$ , including the trivial one, we associate a node in the quiver, while a node  $r$  is connected to a node  $s$  by a number of arrows  $a_{rs}$  which is determined by the decomposition

$$Q \otimes \rho_r = \bigoplus_s a_{rs} \rho_s, \quad a_{rs} = \dim_{\mathbb{C}} \text{Hom}_G(\rho_s, Q \otimes \rho_r).$$


To the resulting quiver  $\mathbf{Q}$  we also associate the framed quiver  $\mathbf{Q}^f$ , its path algebra  $\mathbb{C}\mathbf{Q}^f$ , and the bounded quiver  $(\mathbf{Q}^f, \mathbf{R})$  determined by an ideal  $\langle \mathbf{R} \rangle$  of relations in  $\mathbb{C}\mathbf{Q}^f$ . Its representations form the category  $\text{Rep}(\mathbf{Q}^f, \mathbf{R})$ , which is equivalent to the  $\mathbf{A}$ -mod category of left  $\mathbf{A}$ -modules for the factor path algebra  $\mathbf{A} = \mathbb{C}\mathbf{Q}^f / \langle \mathbf{R} \rangle$ . Moreover, to each vertex  $v$  of  $\mathbf{Q}$  one can associate a simple module  $\mathbf{D}_v$ , defined to be the representation  $V_v \cong \mathbb{C}$  and  $V_w = 0$ , for  $v \neq w$ . Projective resolutions of the simple modules  $\mathbf{D}_v$  can be constructed by means of the submodule  $\mathbf{P}_v$  of  $\mathbf{A}$  generated by paths beginning on vertex  $v$

$$\cdots \longrightarrow \bigoplus_w d_{w,v}^p \mathbf{P}_w \longrightarrow \cdots \longrightarrow \bigoplus_w d_{w,v}^1 \mathbf{P}_w \longrightarrow \mathbf{P}_v \longrightarrow \mathbf{D}_v \longrightarrow 0,$$

where

$$d_{v,w}^p = \dim_{\mathbb{C}} \text{Ext}_{\mathbf{A}}^p(\mathbf{D}_v, \mathbf{D}_w).$$

In the lower dimensional case a special role is played by representations  $\text{Rep}(\mathbf{Q}, \mathbf{R})$  of the bounded quiver with dimensions  $k_r = 1$  and  $\dim_{\mathbb{C}} W = 1$ , as it turns out they correspond to smooth crepant resolutions of toric singularities. In some cases also the path algebra  $\mathbf{A}$  is a different desingularisation by itself, known as the noncommutative crepant resolution of the toric singularity, which contains the coordinate ring of the singularity as its center. In four complex dimensions, however, a crepant resolution of the orbifold singularity is not even granted to exist, though in some simple classes of examples this is known to be the case [52, 53]. Take as an example the case of  $\mathbb{C}^4 / \mathbb{Z}_4$  with the diagonal action  $(z_1, z_2, z_3, z_4) \mapsto (\zeta z_1, \zeta z_2, \zeta z_3, \zeta z_4)$ , where  $\zeta = e^{2\pi i/4} = i$ . As  $Q = \rho_1 \oplus \rho_1 \oplus \rho_1 \oplus \rho_1$ ,  $r_\alpha = 1$  for each  $\alpha = 1, \dots, 4$  and the relevant associated quiver  $\mathbf{Q}$  is then



and the maps  $B_\alpha^r : \mathbf{D}_r \rightarrow \mathbf{D}_{r+r_\alpha \bmod 4}$  are of the form

$$\begin{aligned} B_\alpha^0 &: \mathbf{D}_0 \rightarrow \mathbf{D}_1, \\ B_\alpha^1 &: \mathbf{D}_1 \rightarrow \mathbf{D}_2, \\ B_\alpha^2 &: \mathbf{D}_2 \rightarrow \mathbf{D}_3, \\ B_\alpha^3 &: \mathbf{D}_3 \rightarrow \mathbf{D}_0. \end{aligned}$$

The relevant relations for the unframed quiver are obtained by decomposing accordingly the ADHM equations, thus obtaining

$$\begin{aligned}
B_2^1 B_1^0 &= B_1^1 B_2^0, & B_3^1 B_1^0 &= B_1^1 B_3^0, & B_4^1 B_1^0 &= B_1^1 B_4^0, & B_3^1 B_2^0 &= B_2^1 B_3^0, \\
B_4^1 B_2^0 &= B_2^1 B_4^0, & B_4^1 B_3^0 &= B_3^1 B_4^0, & B_2^2 B_1^1 &= B_1^2 B_2^1, & B_3^2 B_1^1 &= B_1^2 B_3^1, \\
B_4^2 B_1^1 &= B_1^2 B_4^1, & B_3^2 B_2^1 &= B_2^2 B_3^1, & B_4^2 B_2^1 &= B_2^2 B_4^1, & B_4^2 B_3^1 &= B_3^2 B_4^1, \\
B_2^3 B_1^2 &= B_1^3 B_2^2, & B_3^3 B_1^2 &= B_1^3 B_3^2, & B_4^3 B_1^2 &= B_1^3 B_4^2, & B_3^3 B_2^2 &= B_2^3 B_3^2, \\
B_4^3 B_2^2 &= B_2^3 B_4^2, & B_4^3 B_3^2 &= B_3^3 B_4^2, & B_2^0 B_1^3 &= B_1^0 B_2^3, & B_3^0 B_1^3 &= B_1^0 B_3^3, \\
B_4^0 B_1^3 &= B_1^0 B_4^3, & B_3^0 B_2^3 &= B_2^0 B_3^3, & B_4^0 B_2^3 &= B_2^0 B_4^3, & B_4^0 B_3^3 &= B_3^0 B_4^3.
\end{aligned}$$

The center  $Z(\mathbf{A})$  of the path algebra  $\mathbf{A}$  associated to the bounded quiver is generated as a ring by elements

$$x_{\alpha\beta\gamma\delta} = B_\alpha^3 B_\beta^2 B_\gamma^1 B_\delta^0, \quad \alpha \leq \beta \leq \gamma \leq \delta.$$

As the  $G$ -action is chosen to be diagonal one can identify the generators  $x_{\alpha\beta\gamma\delta}$  with the invariant elements in  $\mathbb{C}[z_1, z_2, z_3, z_4]$  by  $x_{\alpha\beta\gamma\delta} \rightsquigarrow z_\alpha z_\beta z_\gamma z_\delta$ , so that

$$\mathrm{Spec} Z(\mathbf{A}) \cong \mathbb{C}^4 / \mathbb{Z}_4$$

and the factor path algebra  $\mathbf{A}$  is a resolution of the orbifold singularity  $\mathbb{C}^4 / \mathbb{Z}_4$ .

## 4.2 Orbifold partition function

The K-theoretic instanton partition function in eight dimensions has been studied in [7] by means of supersymmetric localisation in terms of the quantum mechanics of a D0-D8 system. In the abelian case the moduli space of BPS vacua is identified with the Hilbert scheme of points of  $\mathbb{C}^4$ . In the general case of a proper Calabi-Yau fourfold  $X$ , the Hilbert scheme of points  $X^{[n]}$  is known to carry an obstruction theory, though not perfect. The mathematical definition of the DT-like invariants corresponding to the instanton partition function is made difficult precisely by the latter fact. It is known that they depend on the choice of an orientation of the virtual tangent space and that they need insertions in order to be defined properly. Indeed, if  $\mathcal{Z} \subseteq X^{[n]} \times X$  denotes the universal object, then the virtual tangent space to  $X^{[n]}$  can be written as

$$T_{X^{[n]}}^{\mathrm{vir}} = \mathbf{R}\mathcal{H}\mathrm{om}_{\pi_{X^{[n]}}} (I_{\mathcal{Z}}, I_{\mathcal{Z}})_0[1] = \mathbf{R}\pi_{X^{[n],*}} \circ \mathbf{R}\mathcal{H}\mathrm{om}(I_{\mathcal{Z}}, I_{\mathcal{Z}})_0[1],$$

and this obstruction theory is not perfect. However, the machinery put forward by the work of Borisov-Joyce, [54], and more recently by Thomas [22], one can still construct a virtual fundamental class  $[X^{[n]}]_{o(\mathcal{L})}^{\mathrm{vir}}$  depending on the choice of a square root of the isomorphism  $Q : \mathcal{L} \times \mathcal{L} \rightarrow \mathcal{O}$ , where  $\mathcal{L} = \det \mathbf{R}\mathcal{H}\mathrm{om}_{\pi_{X^{[n]}}} (I_{\mathcal{Z}}, I_{\mathcal{Z}})$ . As, however, we are interested in the case of a quasi-projective variety, namely  $(\mathbb{C}^4)^{[n]}$ , the

previous observations don't provide direct access to definitions of relevant invariants. Equivariant localisation (with respect to the action of  $\mathbb{T} = \{(t_1, t_2, t_3, t_4) \in (\mathbb{C}^*)^4 : t_1 t_2 t_3 t_4 = 1\} \subset (\mathbb{C}^*)^4$ ), however, does provide an easy way out. For a thorough description of this procedure, see [21]. For us, let it suffice to say that, given any  $\mathbb{T}$ -equivariant line bundle  $L$  on  $X$ , one can define the following K-theoretic invariant, with a slight abuse of notation

$$\begin{aligned} Z_X^K(L, y) &= \chi \left( X^{[n]}, \widehat{\mathcal{O}}^{\text{vir}} \otimes \frac{\Lambda^\bullet(L^{[n]} \otimes y^{-1})}{\det^{1/2}(L^{[n]} \otimes y^{-1})} \right) \\ &= \sum_{S \in (X^{[n]})^\tau} (-1)^{o(\mathcal{L})|_S} e \left( \sqrt{\text{Ob}_{X^{[n]}}|_S}^{\text{fix}} \right) \frac{\text{ch} \left( \sqrt{K_{X^{[n]}}^{\text{vir}}|_S}^{1/2} \right)}{\text{ch} \left( \Lambda^\bullet \sqrt{N^{\text{vir}}|_S}^\vee \right)} \\ &\quad \cdot \frac{\text{ch} \left( \Lambda^\bullet(L^{[n]} \otimes y^{-1}) \right)}{\text{ch} \left( \det^{1/2}(L^{[n]} \otimes y^{-1}) \right)} \text{td} \left( \sqrt{T_{X^{[n]}}^{\text{vir}}|_S}^{\text{fix}} \right), \end{aligned}$$

where  $L^{[n]} = \mathbf{R}\pi_{X^{[n]},*}(\mathbf{R}\pi_X^* L \otimes \mathcal{O}_{\mathcal{Z}})$  and the choice of the square root of the virtual tangent space at a fixed point  $S$  induces those of  $\text{Ob}_{X^{[n]}}|_S = h^1(T_{X^{[n]}}^{\text{vir}})$ ,  $K_{X^{[n]}}^{\text{vir}}|_S = \det(T_{X^{[n]}}^{\text{vir}}|_S^\vee)$  and  $N^{\text{vir}}|_S = (T_{X^{[n]}}^{\text{vir}}|_S)^{\text{mov}}$ . Before moving on, let us also notice that the choice that square roots is not unique, so that the invariants at hand are defined only up to a sign. Precisely this definition of  $Z_X^K(L, y)$  is what the partition function of the D0-D8 system computes, with a given prescription for the orientation choice.

All these considerations translate into the physical treatment of the problem, where the supersymmetric measure corresponding to the bulk contribution to the Witten index manifest ghost number anomaly, reminiscent of the positive virtual dimension of the underlying moduli space. Then, in the absence of the  $\Omega$  deformation, the Witten index is vanishing unless observables matching the ghost number anomaly are inserted. This can be neatly done by adding auxiliary hypermultiplets representing the matter deformation necessary in order to cure the anomaly. A similar story goes for the non-abelian case, which generalises the moduli space to the Quot scheme of points of  $\mathbb{C}^4$ , and was studied in [8]. In general the partition function takes the form

$$Z_N^{D8} = \sum_k Z_{N,k}^{D8} d^k,$$

where  $Z_{N,k}^{D8}$  is computed by the JK integration

$$Z_{N,k}^{D8} = \int_{\text{JK}} Z_{N,k}^{1\text{-loop}} d^k u = \frac{1}{k!} \sum_{u_* \in \mathfrak{M}_{\text{sing}}} \text{JK-Res}_{u_*, \zeta} \chi_k d^k u$$

and the instanton measure  $\chi_k$  is defined by

$$\chi_k \propto \prod_{i>j} \frac{\sin^2(u_i - u_j) \prod_{a<b}^3 \sin(u_i - u_j - \epsilon_a - \epsilon_b)}{\prod_{a=1}^3 \sin(u_i - u_j - \epsilon_a) \sin(u_j - u_i - \epsilon_a)} \prod_{i=1}^k \prod_{\alpha=1}^N \frac{\sin(u_i - m_\alpha)}{\sin(u_i - a_\alpha)}.$$

It is then known that poles contributing to the JK integration are only those corresponding to  $N$ -tuples of solid partitions  $\bar{\pi} = (\pi_1, \dots, \pi_N)$ , such that  $|\bar{\pi}| = |\pi_1| + \dots + |\pi_N| = k$ . It turns out it is more convenient to work with exponential variables  $t_a = e^{2i\epsilon_a}$ ,  $x_i = e^{2iu_i}$ ,  $\nu_\alpha = e^{2ia_\alpha}$  and  $\mu_\alpha = e^{2im_\alpha}$ , in which case we have  $Z_{N,k}^{D8} = \sum_{|\bar{\pi}|=k} M_{\mathbf{z}}(\bar{\pi})$ , with

$$M_{\mathbf{z}}(\bar{\pi}) = \text{Res}_{x=x_{\bar{\pi}}} \chi_k \prod_i \frac{dx_i}{x_i}$$

and

$$\chi_k = \prod_{i \neq j} \frac{(x_j - x_i) \prod_{a < b}^3 (x_j - x_i t_a t_b)}{\prod_{a=1}^4 (x_j - x_i t_a)} \prod_{i=1}^k \prod_{\alpha=1}^N \frac{\mu_\alpha - x_i}{\nu_\alpha - x_i}$$

up to a normalisation constant. It also turns out that a more geometric interpretation for the index computation is available. Indeed, in the rank one case, if  $\mathbf{Q}$  denotes the character of a solid partition and  $\sqrt{\mathbf{V}} = \sum_\mu t^\mu - \sum_\nu t^\nu \in K_0^{\mathbb{T}}(\text{pt})$  is the character of the square root of the virtual tangent space to the BPS moduli space at a fixed point, one has

$$\sqrt{\mathbf{V}} = \mathbf{Q} - \bar{P}_{123} \bar{\mathbf{Q}} \mathbf{Q},$$

where  $P_{123} = (1 - t_1)(1 - t_2)(1 - t_3)$  and the involution acts on the generators of  $K_0^{\mathbb{T}}(\text{pt})$  as  $\bar{t}_i = t_i^{-1}$ . Then

$$M_{\mathbf{z}}(\pi) = (-1)^{h(\pi)} \left[ -\sqrt{\widetilde{\mathbf{V}}} \right],$$

with

$$h(\pi) = |\pi| + \#\{(a, d) : (a, a, a, d) \in \pi \text{ and } a \leq d\},$$

$\sqrt{\widetilde{\mathbf{V}}} = \sqrt{\mathbf{V}} - y \bar{\mathbf{Q}}$ , while the action of the brackets operator  $[\bullet]$  on  $\bullet \in K_0^{\mathbb{T}}(\text{pt})$  is defined as

$$[\mathbf{V}] = \frac{\prod_\mu [t^\mu]}{\prod_\nu [t^\nu]} = \frac{\prod_\mu (t^{\mu/2} - t^{-\mu/2})}{\prod_\nu (t^{\nu/2} - t^{-\nu/2})}.$$

The same procedure might be followed in order to compute partition functions for orbifold instantons. In this case, however, the bosonic field content is the one associated to the morphisms of the quiver specifying the natural crepant resolution of the orbifold singularity, provided it exists. Let us notice here that as  $G \subset \text{SL}(4, \mathbb{C})$  is contained in the localising torus  $\mathbb{T}$  the locus on which the computation localises can be identified with the  $G$ -invariant part of the  $\mathbb{T}$ -fixed one. Moreover, as the geometrical interpretation is that of an equivariant count of  $G$ -equivariant zero-dimensional schemes, in order to perform computation one can proceed by simply extracting the  $G$ -invariant part  $\sqrt{\widetilde{\mathbf{V}}}^G$  of  $\sqrt{\widetilde{\mathbf{V}}}$ .

In the case of an orbifold theory the JK residue form gets easily generalised. Let us consider for the sake of simplicity the case of  $\mathbb{C}^2/\mathbb{Z}_n \times \mathbb{C}^2$ . The bosonic field content

is now encoded in the relevant quiver describing the resolution of the orbifold. Let then  $\mathbf{Q}_0$  and  $\mathbf{Q}_1$  be the node set and the edge set of the quiver. We have

$$Z_{\bar{N}}^{\text{orb}} = \sum_{\bar{k}} \underline{q}^{\bar{k}} Z_{\bar{N},\bar{k}}^{\text{orb}},$$

with  $\underline{q}^{\bar{k}} = \prod_{a \in \mathbf{Q}_0} q_a^{k_a}$  and

$$Z_{\bar{N},\bar{k}}^{\text{orb}} = \int_{\text{JK}} Z_{\bar{N},\bar{k}}^{1\text{-loop}} d^{\bar{k}}\bar{x} = \sum_{u_* \in \mathfrak{M}_{\text{sing}}} \text{JK-Res}_{u_*, \zeta} \chi_{\bar{k}}^{\text{orb}} d^{\bar{k}}\bar{x},$$

where we denote by  $\bar{x}$  the collection of coordinates associated to the gauge nodes in  $\mathbf{Q}^f$ , *i.e.*  $\bar{x} = (x_1^{(0)}, \dots, x_{k_0}^{(0)}, \dots)$  and

$$d^{\bar{k}}\bar{x} = \prod_{a \in \mathbf{Q}_0} \prod_{i=1}^{k_a} \frac{dx_i^{(a)}}{x_i^{(a)}}.$$

The orbifold instanton measure  $\chi_{\bar{k}}^{\text{orb}}$  can be easily read off the edge set  $\mathbf{Q}_1$  of the quiver, and we have:

$$\chi_{\bar{k}}^{\text{orb}} \propto \prod_{a \in \mathbf{Q}_0} \frac{1}{k_a!} Z_{\text{f/af}}^{(a)} Z_{\text{adj}}^{(a)} Z_{\text{bif}}^{(a)},$$

where  $Z_{\text{f/af}}^{(a)}$ ,  $Z_{\text{adj}}^{(a)}$ ,  $Z_{\text{bif}}^{(a)}$  encode the bosonic field content of the theory in the fundamental/antifundamental, adjoint and bifundamental representation of the  $a$ -th node respectively. In particular we have

$$\begin{aligned} Z_{\text{f/af}}^{(a)} &= \prod_{i=1}^{k_a} \prod_{\alpha=1}^{N_a} \frac{\mu_{\alpha}^{(a)} - x_i^{(a)}}{\nu_{\alpha}^{(a)} - x_i^{(a)}}, \\ Z_{\text{adj}}^{(a)} &= \prod_{i \neq j}^{k_a} \frac{(x_j^{(a)} - x_i^{(a)})(x_j^{(a)} - x_i^{(a)} t_1 t_2)}{(x_j^{(a)} - x_i^{(a)} t_3)(x_j^{(a)} - x_i^{(a)} t_4)}, \\ Z_{\text{bif}}^{(a)} &= \prod_{i=1}^{k_a} \prod_{j=1}^{k_{a+1}} \frac{(x_i^{(a)} - x_j^{(a+1)} t_1 t_3)(x_i^{(a)} - x_j^{(a-1)} t_2 t_3)}{(x_i^{(a)} - x_j^{(a+1)} t_1)(x_i^{(a)} - x_j^{(a-1)} t_2)}, \end{aligned}$$


with the node indices being understood to be  $a \pmod{n}$ .

*Remark 4.1.* The JK integral formula for the 4-fold orbifold partition function immediately reduces to the integral formula for orbifold counting on 3-folds after the specialisation  $\mu \rightsquigarrow t_4$ , as is to be expected.  $\blacktriangleleft$

### 4.3 An example: $\mathbb{C}^2/\mathbb{Z}_2 \times \mathbb{C}^2$

As an example let us consider a local  $\mathbb{P}^1$  realised as  $\text{Tot}_{\mathbb{P}^1}(\mathcal{O}(-2) \oplus \mathcal{O}^{\oplus 2})$ . This can be understood as the canonical crepant resolution of the orbifold singularity  $\mathbb{C}^4/\mathbb{Z}_2 \cong$

$\mathbb{C}^2/\mathbb{Z}_2 \times \mathbb{C}^2$ , with the  $\mathbb{Z}_2$ -action defined as  $(z_1, z_2, z_3, z_4) \mapsto (\zeta z_1, \zeta^{-1} z_2, z_3, z_4)$ . The ADHM data associated to the orbifolded  $\mathbb{C}^2$  directions can be decomposed according to the  $\mathbb{Z}_2$ -action in irreducible  $\mathbb{Z}_2$  representations as we described in section 4.1. The relevant quiver then takes the following form



**Figure 2:** Orbifolded  $\mathbb{C}^4/\mathbb{Z}_2^{(1,-1,0,0)}$  quiver

For the sake of simplicity we will restrict to the case  $N = 1$ , which enforces  $\overline{N} = (1, 0)$  by the construction in [13] and the observations in 4.1. Supersymmetric localisation can be exploited in order to compute partition functions. As  $\mathbb{Z}_2$  is a subgroup of the localising torus  $\mathbb{T} \cong (\mathbb{C}^*)^3$ , the relevant fixed points will simply be identified to be the  $\mathbb{Z}_2$ -invariant locus of the  $\mathbb{T}$ -fixed one. In particular, as the  $\mathbb{T}$  and the  $\mathbb{Z}_2$ -actions commute and as the  $\mathbb{T}$ -fixed locus of the theory on  $\mathbb{C}^4$  is into bijective correspondence with solid partitions, if we take the framing to be in the trivial representation of  $\mathbb{Z}_2$  their  $\mathbb{Z}_2$ -analogue will be identified with solid partitions  $\pi_{\mathbb{Z}_2}$  decorated by a  $\mathbb{Z}_2$ -coloring, which must be compatible with the action of  $\mathbb{Z}_2$  on  $\mathbb{C}^4$ . By identifying solid partitions themselves with  $\mathbb{Z}_2$  representations, this coloring is in fact induced by the coloring  $(n_1, n_2, n_3, n_4) \mapsto \rho_{r_1}^{\otimes n_1} \otimes \rho_{r_2}^{\otimes n_2} \otimes \rho_{r_3}^{\otimes n_3} \otimes \rho_{r_4}^{\otimes n_4}$  we described in section 4.1. One way we can construct these  $\mathbb{Z}_2$ -colored solid partitions goes as follows: let  $|\pi| = k$  and fix an ordinary partition  $\mu = (\mu_1, \mu_2, \dots, \mu_\ell)$ ,  $|\mu| \leq k$ , colored according to  $\mathbb{Z}_2$ -action, then associate to each one of the boxes of the Young diagram of  $\mu$  another partition  $\lambda_s$ , so that  $\sum_{s \in \mu} |\lambda_s| = n$ , and the partitions  $\lambda_s$  must satisfy a nesting relation in either direction of  $\mu$ . Graphically we have

$$p \in \text{fixed locus} \longleftrightarrow \begin{array}{cccc} \boxed{\lambda_{1,1}} & \boxed{\lambda_{1,2}} & \cdots & \boxed{\lambda_{1,\mu_1}} \\ \boxed{\lambda_{1,1}} & \boxed{\lambda_{1,2}} & \cdots & \\ \vdots & \vdots & \vdots & \\ \boxed{\lambda_{\ell,1}} & \cdots & \boxed{\lambda_{\ell,\mu_\ell}} & \end{array},$$

with  $\lambda_{i,j} \supset \lambda_{i,j+1}$  and  $\lambda_{i,j} \supset \lambda_{i+1,j}$ , for  $(i, j) \in \mu$ . The coloring of the resulting solid partition is then induced by a coloring of the Young diagram  $\mu$ , where each  $\lambda_s$  acquires the same color as the underlying box  $s \in \mu$ . The main difference from the standard instanton counting consists then in the fact that only  $\mathbb{Z}_2$ -invariant boxes in  $\pi_{\mathbb{Z}_2}$  are now going to contribute to the computation of the partition functions.



**Example 4.1.** Consider the case  $N = 1$ ,  $k = 2$  of the cohomological limit of the  $K$ -theoretic partition function we discussed in the previous sections. This cohomological limit can be interpreted geometrically as follows: the Chern character provides a natural transformation from the  $\mathbb{T}$ -equivariant  $K$ -theory to the  $\mathbb{T}$ -equivariant Chow group with rational coefficient by  $t_i \mapsto e^{s_i}$ , with  $s_i = c_1^{\mathbb{T}}(t_i)$ . This natural map can be extended to complex coefficients as  $t_i^b \mapsto e^{bs_i}$ ,  $b \in \mathbb{C}$ , and it gives a simple linearisation of the  $K$ -theoretic brackets  $[\bullet] = \bullet^{1/2} - \bullet^{-1/2}$  operator  $\text{ch}[t^{b\mu}] = be_{\mathbb{T}}(t^\mu) + O(b^2)$ . This linearisation property can be employed to define a map  $Z_{\overline{N}, \overline{k}}^{D8} \mapsto Z_{\overline{N}, \overline{k}}^{D8, \text{coh}}$ , which can in turn be identified with the rational limit of the trigonometric partition function, from a physical standpoint. The rational partition function is also given an integral representation in terms of  $JK$  residues, which also depends on all possible decompositions of  $(k, N)$  in  $(\overline{k}, \overline{N})$ .

Coming back to the particular case of  $(k, N) = (2, 1)$ , one can see that the only possible solid partitions with two boxes are

$$(4.3) \quad \pi_1 = \begin{array}{|c|c|} \hline [1^1] & [1^1] \\ \hline \end{array}, \quad \pi_2 = \begin{array}{|c|} \hline [1^1] \\ \hline [1^1] \\ \hline \end{array}, \quad \pi_3 = \begin{array}{|c|} \hline [2^1] \\ \hline \end{array}, \quad \pi_4 = \begin{array}{|c|} \hline [1^2] \\ \hline \end{array}.$$

The same solid partitions may also be visualised in the following way:

$$(4.4) \quad \pi_1 = \begin{array}{c} \begin{array}{|c|c|} \hline 1 & 1 \\ \hline \end{array} \\ \begin{array}{|c|c|} \hline 1 & 1 \\ \hline \end{array} \end{array}, \quad \pi_2 = \begin{array}{c} \begin{array}{|c|} \hline 1 \\ \hline \end{array} \\ \begin{array}{|c|} \hline 1 \\ \hline \end{array} \end{array}, \quad \pi_3 = \begin{array}{c} \begin{array}{|c|} \hline 1 \\ \hline \end{array} \\ \begin{array}{|c|} \hline 1 \\ \hline \end{array} \end{array}, \quad \pi_4 = \begin{array}{c} \begin{array}{|c|} \hline 2 \\ \hline \end{array} \end{array},$$

where the number on the  $(i, j, k)$  box of each plane partition denotes the height of a pile of boxes stacked on  $(i, j, k)$  along the  $\epsilon_4$  direction. The corresponding fixed points labelled with the coloring corresponding to the orbifold action will then be

$$(4.5) \quad \pi_1 = \begin{array}{|c|c|} \hline [1^1] & [1^1] \\ \hline \end{array}, \quad \pi_2 = \begin{array}{|c|} \hline [1^1] \\ \hline [1^1] \\ \hline \end{array}, \quad \pi_3 = \begin{array}{|c|} \hline [2^1] \\ \hline \end{array}, \quad \pi_4 = \begin{array}{|c|} \hline [1^2] \\ \hline \end{array}.$$

The partition function (equivariant under  $\mathbb{T}_0 = (\mathbb{C}^*)^3|_{s_1+s_2+s_3+s_4=0}$ ) for the non orbifolded theory then reads

$$(4.6) \quad Z_{1,2}^{D8}(\epsilon; q) = \frac{q^2}{2} \left( \frac{m^2(\epsilon_1 + \epsilon_2)^2(\epsilon_1 + \epsilon_3)^2(\epsilon_2 + \epsilon_3)^2}{\epsilon_1^2 \epsilon_2^2 \epsilon_3^2 \epsilon_4^2} + 5 \frac{m(\epsilon_1 + \epsilon_2)(\epsilon_1 + \epsilon_3)(\epsilon_2 + \epsilon_3)}{\epsilon_1 \epsilon_2 \epsilon_3 \epsilon_4} \right),$$

while the contribution from the regular instanton sector to the orbifold partition function will be

$$\begin{aligned} Z_{\text{reg}}^{D8, \text{orb}}(\epsilon; q_0, q_1) &= Z_{(1,0),(2,0)}^{D8, \text{orb}}(\epsilon; q_0, q_1) + Z_{(1,0),(1,1)}^{D8, \text{orb}}(\epsilon; q_0, q_1) \\ &= \frac{q_0^2 m \epsilon_{12}}{2 \epsilon_3 \epsilon_{123} (\epsilon_{12} + 2 \epsilon_3)} \left( \frac{(\epsilon_3 - m)(\epsilon_{12} - \epsilon_3)}{\epsilon_3} + \frac{(\epsilon_{123} + m)(2 \epsilon_{12} + \epsilon_3)}{\epsilon_{123}} \right) \\ &\quad + \frac{q_0 q_1 m \epsilon_{12}}{2 \epsilon_3 \epsilon_{123} (\epsilon_2 - \epsilon_1)} \left( \frac{(2 \epsilon_2 + \epsilon_3)(\epsilon_{13} - \epsilon_2)}{\epsilon_2} - \frac{(2 \epsilon_1 + \epsilon_3)(\epsilon_{23} - \epsilon_1)}{\epsilon_1} \right), \end{aligned}$$

with  $\epsilon_i = c_1^\mathbb{T}(t_i)$ ,  $\epsilon_{ij} = \epsilon_i + \epsilon_j$  and similarly  $\epsilon_{123} = \epsilon_1 + \epsilon_2 + \epsilon_3$ . The previous formula can be either obtained through an integral formula and iterated residues, or by generalising the geometric correspondence of [8] to the orbifold case

$$Z_{\text{reg}}^{D8, \text{orb}}(\epsilon; q_0, q_1) = \sum_{|\pi|=2} (-1)^{h(\pi)} e_{\mathbb{T}} \left[ - \left( \sqrt{T_{\pi}^{\text{vir}}} \right)^{\mathbb{Z}_2} \right] q_0^{|\pi|_0} q_1^{|\pi|_1},$$

where  $\sqrt{T_{\pi}^{\text{vir}}}$  is a square root of the virtual tangent space  $T_{(\mathbb{C}^4)^{[k]}|\pi}^{\text{vir}}$  and  $|\pi|_i$  denotes the number of boxes in  $\pi$ , seen as  $\mathbb{Z}_2$ -modules, transforming in the  $\rho_i$ -th representation of  $\mathbb{Z}_2$ .

## Acknowledgments

We would like to thank M. Cirafici, A. Gorsky, S. Mironov, S. Monavari and Al. Morozov for interesting discussions. The research of G.B. is partly supported by the INFN Iniziativa Specifica ST&FI and by the PRIN project ‘‘Non-perturbative Aspects Of Gauge Theories And Strings’’. The research of N.F. and A.T. is partly supported by the INFN Iniziativa Specifica GAST. The work of A.T. is partly supported by the PRIN project ‘‘Geometria delle varietà algebriche’’. Y.Z. is partly supported by RSF grant 18-71-10073.

## A Review of the 4d ADHM construction

### A.1 Self-dual connections in 4d

In this appendix we recall some basic facts about the standard ADHM construction [4] of the solutions to the self-duality equation on  $\mathbb{R}^4$ :

$$(A.1) \quad F_{\mu\nu} = \frac{1}{2} \epsilon_{\mu\nu\lambda\rho} F_{\lambda\rho},$$

where  $F_{\mu\nu} = \partial_{\mu} A_{\nu} - \partial_{\nu} A_{\mu} + [A_{\mu}, A_{\nu}]$  is the curvature of a  $U(N)$  connection. The instanton number is defined as

$$(A.2) \quad k = \frac{1}{16\pi^2} \int d^4x \epsilon_{\mu\nu\lambda\rho} \text{tr}(F_{\mu\nu} F_{\lambda\rho})$$

We denote the moduli space of self-dual  $U(N)$  connections of instanton number  $k$  by  $\mathcal{M}_{k,N}$ . The (virtual) dimension of the moduli space of instantons is by definition the number of independent deformations

$$(A.3) \quad (\delta_{\mu}^{\alpha} \delta_{\nu}^{\beta} - \frac{1}{2} \epsilon^{\alpha\beta\mu\nu}) D_{\mu} \delta A_{\nu} = 0,$$

minus the number of gauge degrees of freedom:

$$(A.4) \quad \delta A_{\mu} \sim D_{\mu} \phi,$$

where  $D_\mu$  is the covariant derivative in the background  $A_\mu$ . This difference can be understood as the index of the elliptic complex (this requires certain vanishing theorem for the cohomologies in degrees 0 and 2):

$$(A.5) \quad \mathcal{C} : 0 \rightarrow \Omega^0(\mathbb{R}^4, \mathfrak{su}(N)) \xrightarrow{D_A} \Omega^1(\mathbb{R}^4, \mathfrak{su}(N)) \xrightarrow{(1-* )D_A} \Omega_+^2(\mathbb{R}^4, \mathfrak{su}(N)) \rightarrow 0,$$

which by the Atiyah-Singer index theorem is given by

$$(A.6) \quad \dim \mathcal{M}_{k,N} = \text{ind } \mathcal{C} = 4Nk.$$

## A.2 ADHM data and equations

In the ADHM construction we consider a complex  $(2k + N) \times N$  matrix  $U$  satisfying the equation

$$(A.7) \quad \Delta^\dagger(x)U(x) = 0,$$

where  $\Delta(x)$  is a  $(2k + N) \times 2k$  complex matrix

$$(A.8) \quad \Delta(x) = \begin{pmatrix} B_2^\dagger - z_2^* & -B_1 + z_1 \\ B_1^\dagger - z_1^* & B_2 - z_2 \\ I^\dagger & J \end{pmatrix},$$

and  $z_1 = x_2 + ix_1$ ,  $z_2 = x_4 + ix_3$ . The auxiliary matrix  $\Delta(x)$  is required to satisfy the moment map equation

$$(A.9) \quad \Delta^\dagger(x)\Delta(x) = 1_{2 \times 2} \otimes f_{k \times k}^{-1}(x),$$

where  $f(x)$  is a  $k \times k$  invertible matrix. Eq. (A.9) leads to the well-known conditions on  $B_{1,2}$ ,  $I$  and  $J$ :

$$(A.10) \quad \mu_{\mathbb{C}} = [B_1, B_2] + IJ = 0, \quad \mu_{\mathbb{R}} = [B_1, B_1^\dagger] + [B_2, B_2^\dagger] - II^\dagger + J^\dagger J = 0.$$

One has to normalize  $U$  so that

$$(A.11) \quad U^\dagger U = 1_{N \times N}.$$

In other words, the matrix  $U$  defines a basis of  $N$  orthonormal vectors inside  $\mathbb{C}^{2k+N}$ . The columns of the matrix  $\Delta(x)$  form a set of  $k$  linearly independent vectors inside  $\mathbb{C}^{2k+N}$  (the linear independence is guaranteed by Eq. (A.9)). Eq. (A.7) means that the basis defined by  $U$  is orthogonal to the set of  $2k$  vectors inside  $\mathbb{C}^{2k+N}$  defined by  $\Delta(x)$ . Together the columns of  $U$  and  $\Delta$  form a complete basis in  $\mathbb{C}^{2k+N}$ , and therefore:

$$(A.12) \quad 1_{(2k+N) \times (2k+N)} - U(x)U^\dagger(x) = \Delta(x)(1_{2 \times 2} \otimes f_{k \times k}(x))\Delta^\dagger(x).$$

The self-dual connection  $A_\mu(x)$ , corresponding to the matrix  $U(x)$  is given by

$$(A.13) \quad A_\mu(x) = U^\dagger \partial_\mu U(x).$$

It is easy to show that the connection (A.13) is indeed self-dual:

$$(A.14) \quad \begin{aligned} F_{\mu\nu} &= \partial_\mu U^\dagger \partial_\nu U(x) - \partial_\nu U^\dagger \partial_\mu U(x) + [U^\dagger \partial_\mu U(x), U^\dagger \partial_\nu U(x)] = \\ &= (\partial_{[\mu} U^\dagger)(1 - UU^\dagger)(\partial_{\nu]} U^\dagger) = U^\dagger (\partial_{[\mu} \Delta(x))(1_{2 \times 2} \otimes f_{k \times k}(x))(\partial_{\nu]} \Delta^\dagger(x))U = \\ &= U^\dagger (\sigma_{\mu\nu} \otimes f_{k \times k}(x))U, \end{aligned}$$

where

$$(A.15) \quad \sigma_{\mu\nu} = \sigma_\mu \bar{\sigma}_\nu - \sigma_\nu \bar{\sigma}_\mu = i \eta_{\mu\nu}^a \sigma_a,$$

and  $\sigma_\mu = (1, i\vec{\sigma})$ ,  $\bar{\sigma}_\mu = \sigma_\mu^\dagger = (1, -i\vec{\sigma})$ ,  $\eta_{\mu\nu}^a$  is the 't Hooft symbol. The matrices  $(\sigma_{\mu\nu})_\alpha^\beta$  when seen as matrices with indices  $\mu\nu$  are known to be self-dual. In fact they give an intertwining operator between the three-dimensional representation  $\Lambda_\pm^2 \mathbb{R}^4$  of  $SO(4)$  and the adjoint representation of  $SU(2)_+$ , which is part of  $\text{Spin}(4) = SU(2)_+ \times SU(2)_-$ , so that

$$(A.16) \quad \Lambda_\pm^2 \mathbb{R}^4 = \text{adj}_{SU(2)_\pm}.$$

The moduli space  $\mathcal{M}_{k,N}$  of solutions to the ADHM equations is the space of matrices  $(B_1, B_2, I, J)$  obeying  $\mu_{\mathbb{C}} = 0$  and  $\mu_{\mathbb{R}} = 0$  quotiented by the action of  $U(k)$  group (indeed, constant  $U(k)$  transformations don't change the connection (A.13)). The resulting dimension of the moduli space is

$$(A.17) \quad \dim_{\mathbb{R}} \mathcal{M}_{k,N} = \underbrace{2k^2}_{B_1} + \underbrace{2k^2}_{B_2} + \underbrace{2Nk}_I + \underbrace{2Nk}_J - \underbrace{2k^2}_{\mu_{\mathbb{C}}} - \underbrace{\mu_{\mathbb{R}}}_{k^2} - \underbrace{k^2}_{/U(k)} = 4Nk,$$

is equal to the dimension of the tangent (A.6), as it should.

### A.3 Spinor formalism

Let us see how the ADHM equations can be formulated using spinors. Let  $\gamma^\mu$ ,  $\mu = 0, \dots, 4$  be four-dimensional gamma-matrices generating the algebra  $\text{Cliff}(4)$ . Then we have

$$(A.18) \quad \gamma^\mu \gamma^\nu \gamma^\lambda \gamma^\rho = \epsilon^{\mu\nu\lambda\rho} \gamma_5,$$

where  $\gamma_5$  is the chirality operator. Let  $\psi_\pm$  be a positive (resp. negative) chirality spinor of  $\text{Spin}(4)$

$$(A.19) \quad \gamma_5 \psi_\pm = \pm \psi_\pm,$$

normalized so that  $\psi_{\pm}^{\dagger}\psi_{\pm} = 1$ . We can then trivially write that

$$(A.20) \quad \psi_{\pm}^{\dagger}\gamma^{\mu}\gamma^{\nu}\gamma^{\lambda}\gamma^{\rho}\psi_{\pm} = \pm\epsilon^{\mu\nu\lambda\rho}.$$

Notice that this identity is independent of the concrete value of  $\psi_{\pm}$ . We can thus write the self-duality condition for  $F_{\mu\nu}$  as

$$(A.21) \quad F_{\mu\nu} = \frac{1}{2}\psi_{+}^{\dagger}\gamma^{\mu}\gamma^{\nu}\gamma^{\lambda}\gamma^{\rho}\psi_{+}F_{\lambda\rho}.$$

Alternatively, one can notice that

$$(A.22) \quad \gamma^{\mu\nu} = \frac{1}{2}[\gamma^{\mu}\gamma^{\nu}] = \begin{pmatrix} \sigma_{\mu\nu} & 0 \\ 0 & \bar{\sigma}_{\mu\nu} \end{pmatrix} = \begin{pmatrix} \eta_{\mu\nu}^a i\sigma^a & 0 \\ 0 & \bar{\eta}_{\mu\nu}^a i\sigma^a \end{pmatrix},$$

where  $\bar{\sigma}_{\mu\nu} = \bar{\sigma}_{[\mu}\sigma_{\nu]}$ , so that the projection on the self-dual part of  $F_{\mu\nu}$  can be written as

$$(A.23) \quad \gamma^{\lambda}\gamma^{\rho}\psi_{-}F_{\lambda\rho} = 0, \quad \text{or} \quad \bar{\sigma}_{\lambda\rho}F_{\lambda\rho} = 0.$$

Eq. (A.23) contains three independent equations on the components of  $A_{\mu}(x)$ . The fourth equation is provided by the gauge fixing condition, e.g.  $\partial_{\mu}A_{\mu} = 0$ , so that the total number of components is equal to the total number of (first order) equations. After imposing the equations no *functional* degrees of freedom in  $A_{\mu}(x)$  remain, and the moduli space of solutions  $\mathcal{M}_{k,N}$  is finite-dimensional.

#### A.4 Hyperkähler reduction and complex structure(s)

We have considered self-duality equations in  $\mathbb{R}^4$  without any reference to a specific choice of the complex structure. However, we can incorporate complex structure in our construction.  $\mathbb{R}^4$  is a hyperkähler manifold with three basis complex structures  $I, J$  and  $K$ <sup>11</sup> satisfying the usual quaternionic relations. Correspondingly, there are three Kähler forms  $\omega^I, \omega^J$  and  $\omega^K$ .

These forms can be used to give the structure of the hyperkähler manifold to the space  $\mathcal{A}$  of  $U(N)$  connections on  $\mathbb{R}^4$ . Indeed, the metric on the space of connections is induced from the flat metric on  $\mathbb{R}^4$

$$(A.24) \quad ds^2[\delta_1 A_{\mu}(x), \delta_2 A_{\mu}(y)] = \int_{\mathbb{R}^4} \text{tr} \delta_1 A_{\mu}(x) \delta_2 A_{\mu}(x),$$

and we can write the symplectic forms on  $\mathcal{A}$  as follows

$$(A.25) \quad \Omega^a[\delta_1 A_{\mu}(x), \delta_2 A_{\mu}(y)] = \int_{\mathbb{R}^4} \omega^a \wedge \text{tr} \delta_1 A(x) \wedge \delta_2 A(x),$$

---

<sup>11</sup>There is in fact a  $S^2$  worth of complex structures. Indeed,  $aI + bJ + cK$  is a complex structure as long as  $a^2 + b^2 + c^2 = 1$ .

where  $a = I, J, K$ . All three  $\omega^a$  are actually *self-dual* two-forms. One can then verify that the action of  $\mathcal{G}$  on  $\mathcal{A}$  is Hamiltonian for all three symplectic forms  $\Omega^a$ , and the corresponding three moment maps are

$$(A.26) \quad \mu^a[\phi(x)] = \int_{\mathbb{R}^4} \omega^a \wedge \text{tr}(\phi(x)F(x)),$$

where  $F = dA + [A, A]$  is the field strength. Requiring moment maps to vanish we get precisely the self-duality equations:

$$(A.27) \quad \omega^a \wedge F = 0.$$

Indeed, Eq. (A.27) mean that the *anti-self-dual* part  $F$  vanishes. The space of self-dual connections can be thought of as the hyperkähler reduction of  $\mathcal{A}$  by the group  $\mathcal{G}$  of all gauge transformations. If we introduce  $\mathcal{A}_k$  as the space of connections with instanton number  $k$  then we can write

$$(A.28) \quad \mathcal{M}_{k,N} = \mathcal{A}_k // \mathcal{G}.$$

Let us choose a complex structure  $I$ , such that  $z_1$  and  $z_2$  defined as in Eq. (A.8) are the holomorphic coordinates. Only a subgroup  $U(2) \subset SO(4)$  of rotations preserves this choice. The choice of the complex structure  $I$  singles out one of the Kähler forms  $\omega^I$ , which can be written as

$$(A.29) \quad \omega^I = \omega_{\mathbb{R}} = dz^1 \wedge d\bar{z}^1 + dz^2 \wedge d\bar{z}^2.$$

The other two forms  $\omega^J$  and  $\omega^K$  can be recast into the holomorphic symplectic form (and its conjugate)

$$(A.30) \quad \omega^J + i\omega^K = \omega_{\mathbb{C}} = dz^1 \wedge dz^2.$$

The subgroup of  $SO(4)$ , which preserves both the complex structure  $I$  and  $\omega_{\mathbb{C}}$  is  $SU(2) \subset U(2)$  (the remaining  $U(1) \subset U(2)$  rotates the phase of  $\omega_{\mathbb{C}}$ ). The field strength  $F_{\mu\nu}$  breaks into the following irreducible representations of  $SU(2)$  (we also list the number of components):

$$(A.31) \quad F_{\mu\nu} = \begin{cases} F_{z_1 z_2}^{(2,0)} = F_{\mu\nu}(\omega_{\mathbb{C}}^{-1})^{\mu\nu}, & \dim = 1, \\ F_{\bar{z}_1 \bar{z}_2}^{(0,2)} = F_{\mu\nu}(\bar{\omega}_{\mathbb{C}}^{-1})^{\mu\nu}, & \dim = 1, \\ F_{\omega}^{(1,1)} = F_{\mu\nu}(\omega_{\mathbb{R}}^{-1})^{\mu\nu}, & \dim = 1, \\ (F_0^{(1,1)})_{z_i \bar{z}_j} = F_{z_i \bar{z}_j}^{(1,1)} - \frac{1}{2} F_{z_k \bar{z}_l}^{(1,1)}(\omega_{\mathbb{R}}^{-1})^{z_k \bar{z}_l}(\omega_{\mathbb{R}})_{z_i \bar{z}_j}, & \dim = 3. \end{cases}$$

The first three one-dimensional pieces turn out to be self-dual (they are projections on  $\omega^a$ ), while the three-dimensional representation is anti-self-dual. Having these identifications it is easy to see that the moment map equation (A.9) is just the requirement that the  $\Delta^\dagger \Delta$  lies in the self-dual part  $\mathbf{1}$  of the tensor product of  $SU(2)$  representations  $\bar{\mathbf{2}} \otimes \mathbf{2} = \mathbf{3} \oplus \mathbf{1}$ .

## A.5 Non-commutative deformation.

The instanton moduli space  $\mathcal{M}_{k,N}$  contains singularities, which correspond to instantons of zero size. A clever way to regularize the moduli space is to consider instantons on the non-commutative spacetime  $\mathbb{R}^4$  with coordinates satisfying:

$$(A.32) \quad [x^\mu, x^\nu] = i\theta^{\mu\nu},$$

where  $\theta^{\mu\nu}$  is a constant 2-form. We further assume that  $\theta^{\mu\nu}$  is anti-self-dual, then with an  $\text{SO}(4)$  rotation it can be aligned with one of the three basis Kähler structures, e.g.

$$(A.33) \quad \theta^{\mu\nu} = \frac{\zeta}{4} \bar{\eta}_{\mu\nu}^3,$$

where  $\zeta$  is a real non-commutativity parameter which we take to be positive.

As shown in [5], the ADHM construction for the non-commutative case requires only a minor update. The expression (A.8) for  $\Delta(x)$  is still valid and the conditions (A.9) still holds. However, since the coordinates no longer commute, one of the moment maps is modified:

$$(A.34) \quad \mu_{\mathbb{R}} = [B_1^\dagger, B_1] + [B_2^\dagger, B_2] + II^\dagger - J^\dagger J = \zeta 1_{k \times k}.$$

Notice that after the non-commutative deformation, the instanton solutions also appear in the  $\text{U}(1)$  gauge theory.

In string theory language, the self-dual connections we are studying correspond to bound states of D0 and D4 branes (or, more generally  $Dp$  and  $D(p+4)$  branes). Non-commutativity arises if we turn on the nonzero  $B$ -field background along the D4 brane.

## A.6 Matrix form of the non-commutative self-duality equations.

In the noncommutative setting it will be convenient for us to recast the self-duality equations in matrix form. To this end we introduce the operator analogues of covariant derivatives

$$(A.35) \quad X^\mu = x^\mu + i\theta^{\mu\nu} A_\nu,$$

The commutator of covariant derivatives gives the field strength:

$$(A.36) \quad [X^\mu, X^\nu] = i\theta^{\mu\nu} + \theta^{\mu\lambda}\theta^{\nu\rho} F_{\lambda\rho},$$

where the second term in the r.h.s. is due to Eq. (A.32).

The self-duality equations can be rewritten as an equation for the operators  $X^\mu$

$$(A.37) \quad [X^\mu, X^\nu] - i\theta^{\mu\nu} = \frac{1}{2} \epsilon_{\mu\nu\lambda\rho} ([X^\lambda, X^\rho] - i\theta^{\lambda\rho}).$$

Let us choose a Kähler structure on  $\mathbb{R}^4$  proportional to the  $B$ -field. In the complex structure compatible with  $B$  the noncommutativity reads

$$(A.38) \quad [z_a, \bar{z}_b] = -\frac{\zeta}{2}\delta_{ab}, \quad [z_a, z_b] = [\bar{z}_a, \bar{z}_b] = 0.$$

We introduce complex covariant derivatives  $Z_a$  corresponding to complex coordinates  $z_a$ . Eqs. (A.37) are then written as one real and one complex equation for  $Z_a$  operators:

$$(A.39) \quad [Z_1, Z_2] = 0, \quad [Z_1^\dagger, Z_2^\dagger] = 0,$$

$$(A.40) \quad [Z_1^\dagger, Z_1] + [Z_2^\dagger, Z_2] = \zeta.$$

The commutation relations (A.38) are of course nothing but a pair of standard Heisenberg algebras of creation and annihilation operators, so that

$$(A.41) \quad z_a = \sqrt{\frac{2}{\zeta}}a_a^\dagger, \quad \bar{z}_a = \sqrt{\frac{2}{\zeta}}a_a,$$

and

$$(A.42) \quad [a_a, a_b^\dagger] = \delta_{ab}, \quad [a_a, a_b] = [a_a^\dagger, a_b^\dagger] = 0.$$

The operators  $a_a, a_a^\dagger$  act on the Hilbert space  $\mathcal{H}$ , which is spanned by the eigenstates  $|n, m\rangle$  of the number operators:

$$(A.43) \quad N_a = a_a^\dagger a_a, \quad N_1|n, m\rangle = n|n, m\rangle, \quad N_2|n, m\rangle = m|n, m\rangle.$$

$\mathcal{H}$  can be identified with the ring of polynomials in  $a_a^\dagger$  (or in  $z_a$ ) acting on the vacuum  $|0, 0\rangle$ .

The simplest solution to Eqs. (A.39) is the vacuum solution

$$(A.44) \quad Z_a = z_a,$$

which corresponds to zero  $A_\mu$  and vanishing instanton charge.

Another example of a solution is the non-commutative  $U(1)$  instanton sitting at the origin of  $\mathbb{C}^2$ :

$$(A.45) \quad Z_a = \sqrt{\frac{2}{\zeta}}S_{[1]}^\dagger a_a^\dagger \sqrt{\frac{N(N+3)}{(N+1)(N+2)}}S^\dagger = \sqrt{\frac{2}{\zeta}}S \sqrt{\frac{N+2}{N}}a_a^\dagger \sqrt{\frac{N}{N+2}}S_{[1]}^\dagger,$$

where  $N = a_1^\dagger a_1 + a_2^\dagger a_2$  and  $S_{[1]}^\dagger$  acts on  $\mathcal{H}$  by relabelling the (infinite number of) basis vectors, so that the state  $|0, 0\rangle$  does not belong to its image. An example of such a transformation is

$$(A.46) \quad S_{[1]}^\dagger|n, m\rangle = \begin{cases} |n, m\rangle & m \neq 0, \\ |n+1, m\rangle & m = 0. \end{cases}$$



The solution (A.45) is non-singular (all of its matrix elements are well-defined) and invariant under  $U(2)$  rotations of the space-time  $\mathbb{C}^2$  (the operator  $S_{[1]}$  is invariant only up to a unitary transformation, which can be viewed as a gauge transformation). We give some more examples of instanton solutions invariant under  $U(1)^2 \subset U(2)$  in A.8.

### A.7 $U(1)$ instantons and ideals

Let us recall the correspondence between  $U(1)$  noncommutative instantons and ideals in the ring of polynomials  $\mathbb{C}[z_1, z_2]$ . The matrix  $U$  for gauge group  $U(1)$  is a  $(2k+1)$ -dimensional column vector, which we write as

$$(A.47) \quad U(z) = \begin{pmatrix} \psi_1(z) \\ \psi_2(z) \\ \xi(z) \end{pmatrix},$$

where  $\psi_{1,2}(z)$  are two  $k$ -dimensional column vectors of polynomials and  $\xi(z)$  is a polynomial. Eq. (A.7) for  $U$  is then equivalent to two equations

$$(A.48) \quad I\xi(z) = (z_2 - B_2)\psi_1(z) + (z_1 - B_1)\psi_2(z),$$

$$(A.49) \quad 0 = (B_1^\dagger - \bar{z}_1)\psi_1(z) + (\bar{z}_2 - B_2^\dagger)\psi_2(z).$$

Eq. (A.48) implies (see e.g.[55] for details) that  $\xi(z)$  belongs to an ideal  $\mathcal{I} \subset \mathbb{C}[z_1, z_2]$  of polynomials which vanish when one substitutes in them matrices  $B_{1,2}$  instead of the variables  $z_{1,2}$ . Indeed, acting with Eq. (A.48) on the states  $|n, m\rangle \in \mathcal{H}$  we get

$$(A.50) \quad \xi(z)|n, m\rangle I = (z_2 - B_2)\psi_1(z)|n, m\rangle + (z_1 - B_1)\psi_2(z)|n, m\rangle.$$

After the substitution  $z_{1,2} \rightarrow B_{1,2}$  we get

$$(A.51) \quad \xi(B_1, B_2)|n, m\rangle I = 0.$$

Acting with  $B_1^k B_2^l$  on Eq. (A.51) and using the commutativity of  $B_1$  and  $B_2$  we see that  $\xi(B_1, B_2)|n, m\rangle$  is zero on the whole  $\mathbb{C}^k$ , so it is a zero matrix and thus  $\xi(B_1, B_2) = 0$  as a matrix equation. This determines the ideal completely. Vice versa, each element  $\xi(z)$  of  $\mathcal{I}$  by definition can be written as a linear combination from Eq. (A.48), but not uniquely: one can shift

$$(A.52) \quad \psi_1(z) \rightarrow \psi_1(z) + (z_1 - B_1)\chi(z),$$

$$(A.53) \quad \psi_2(z) \rightarrow \psi_2(z) + (z_2 - B_2)\chi(z),$$

for any column-vector  $\chi(z)$ . To fix a unique decomposition we have to gauge fix the symmetry (A.52), (A.53), which is done by requiring Eq. (A.49) (note that  $\bar{z}_a$  act as the derivatives in  $z_a$ ).

For the single  $U(1)$  instanton solution described in the previous section we have  $B_1 = B_2 = 0$ ,  $I = \sqrt{\zeta}$  and

$$(A.54) \quad U(z) = \begin{pmatrix} \bar{z}_2 \\ \bar{z}_1 \\ \frac{1}{\sqrt{\zeta}}(z_1\bar{z}_1 + z_2\bar{z}_2) \end{pmatrix} \lambda(z).$$

For any  $\lambda(z)$  the vector (A.54) solves (A.7). However to satisfy the normalization condition (A.11), we have to put the following normalization factor instead of  $\lambda(z)$ :

$$(A.55) \quad \mathcal{N} = \frac{1}{\sqrt{(z_1\bar{z}_1 + z_2\bar{z}_2)(z_1\bar{z}_1 + z_2\bar{z}_2 + \zeta)}} S^\dagger.$$

The solution (A.45) is obtained from the normalized  $U(z)$  as

$$(A.56) \quad Z_a = U^\dagger z_a U, \quad Z_a^\dagger = U^\dagger \bar{z}_a U.$$

### A.8 4d fixed point instantons

We give here some examples of noncommutative instanton solutions on  $\mathbb{R}^4$  which are equivariant under the  $U(1)^2$  subgroup of  $SO(4)$ . The ideals in  $\mathbb{C}[z_1, z_2]$  corresponding to these solutions are monomial ideals enumerated by Young diagrams.

All the solutions have the form

$$(A.57) \quad Z_a = \sqrt{\frac{\zeta}{2}} S_Y h_Y(N_1, N_2)^{-1} a_a^\dagger h_Y(N_1, N_2) S_Y^\dagger,$$

where  $S_Y$  is the partial isometry associated with the monomial ideal labelled by the Young diagram  $Y$ ,  $N_a = a_a^\dagger a_a$  and

$$(A.58) \quad h_Y(N_1, N_2) = \sqrt{\frac{g_Y(N_1, N_2)}{g_Y(N_1 + 1, N_2 + 1)}}.$$

Let us display the functions  $g_Y(N_1, N_2)$  for some elementary Young diagrams

$$(A.59) \quad g_{[1]} = N,$$

$$(A.60) \quad g_{[2]} = N^2 - N_1 + N_2,$$

$$(A.61) \quad g_{[1,1]} = N^2 + N_1 - N_2,$$

$$(A.62) \quad g_{[3]} = N(N^2 - 3(N_1 - N_2) + 2),$$

$$(A.63) \quad g_{[1,1,1]} = N(N^2 + 3(N_1 - N_2) + 2),$$

$$(A.64) \quad g_{[2,1]} = (N - 1)N(N + 1),$$

$$(A.65) \quad g_{[3,2,1]} = (N - 2)(N - 1)N^2(N + 1)(N + 2).$$

where  $N = N_1 + N_2$ . The first solution (A.59) is the one-instanton solution discussed in sec. A.6. One can check that for any function  $g_Y$  given above the matrix equations (A.39), (A.40) are indeed satisfied. In fact Eq. (A.39) is satisfied automatically

by the ansatz (A.57), and it is only necessary to check a single recurrence relation for the function  $h(N_1, N_2)$ :

$$(A.66) \quad \sum_{a=1}^2 \left\{ \frac{h(N)^2}{h(N_a + 1)^2} (N_a + 1) - \frac{h(N_a - 1)^2}{h(N)^2} N_a \right\} = 2,$$

which turns out to be true. Of the solutions given above only (A.59) and (A.65) are invariant under the full  $SU(2)$  rotation symmetry. There exists an infinite family of such fully symmetric solutions corresponding to triangular Young diagrams with

$$(A.67) \quad h_{[k, k-1, \dots, 1]}(N_1, N_2) = \left( 1 - \frac{k(k+1)}{N(N+1)} \right)^{\frac{1}{2}} = \left( \frac{(N-k)(N+k+1)}{N(N+1)} \right)^{\frac{1}{2}}.$$

Notice that  $g_Y(i, j)$  vanishes if the box  $(i, j)$  lies on the border of the Young diagram  $Y$ . This guarantees that the action of  $Z_a$  and  $Z_a^\dagger$  from Eq. (A.57) is well-defined.

## B 8d instantons and sheaf cohomology

### B.1 Moduli spaces of 8d instantons via Beilinson's Theorem

Here we will briefly study moduli spaces of framed torsion-free sheaves on  $\mathbb{P}^4$  and their relation to spaces of  $SU(4)$  instantons (and their generalisations) on  $\mathbb{C}^4$ . Let us first notice that, in general, if  $\mathcal{E}$  is a torsion-free sheaf of rank  $N$  on  $\mathbb{P}^4$  with  $\text{ch}(\mathcal{E}) = (N, 0, 0, 0, -k)$ , framed along a divisor  $D$ , there exists a natural sequence of sheaves (for the proof of this result, see [48])

$$0 \rightarrow \mathcal{E} \hookrightarrow \mathcal{O}_{\mathbb{P}^d}^{\oplus N} \twoheadrightarrow \mathcal{Q} \rightarrow 0,$$

and  $\mathcal{Q}$  has finite support in  $\mathbb{A}^4 \cong \mathbb{P}^4/D$ . Consider then the moduli space

$$(B.1) \quad \left\{ \begin{array}{l} \mathcal{E} \in \text{Coh}(\mathbb{P}^4), \mathcal{E} \text{ torsion-free, } \text{rk}(\mathcal{E}) = N, \\ \mathcal{E} \hookrightarrow \mathcal{O}_{\mathbb{P}^4}^{\oplus N} \twoheadrightarrow \mathcal{Q} \text{ s.t.} \\ \mathcal{Q} \text{ of finite support in } \mathbb{A}^4 \cong \mathbb{P}^4/D \end{array} \right\} / \text{iso},$$

parametrising torsion-free sheaves on  $\mathbb{P}^4$  fitting in a given short exact sequence. Of course the previous moduli space is identified with the quot scheme of points in  $\mathbb{A}^4$ , as the Grothendieck moduli functor

$$\text{Quot}_{\mathbb{P}^4}(\mathcal{O}^{\oplus N}, k) : \text{Sch}_{\mathbb{C}}^{\text{op}} \rightarrow \text{Sets}$$

contains an open subfunctor  $\text{Quot}_{\mathbb{A}^4}(\mathcal{O}^{\oplus N}, k) \hookrightarrow \text{Quot}_{\mathbb{P}^4}(\mathcal{O}^{\oplus N}, k)$  parametrising precisely the quotients above. What one might want to do is then to study a model for

deformations of the moduli space of sheaves

$$\mathcal{M}_{r,n}(\mathbb{P}^4) = \left\{ \begin{array}{l} \mathcal{E} \in \text{Coh}(\mathbb{P}^4), \mathcal{E} \text{ torsion-free, } \text{rk}(\mathcal{E}) = N, \\ c_1(\mathcal{E}) = c_2(\mathcal{E}) = c_3(\mathcal{E}) = 0, c_4(\mathcal{E}) = k, \mathcal{E}|_{\wp_\infty^i} \cong \mathcal{O}_{\mathbb{P}^4}^{\oplus N} \\ H^3(\mathbb{P}^4, \mathcal{E}(-3)) = 0, H^2(\mathbb{P}^4, \mathcal{E}(-\ell)) = 0, \forall \ell \end{array} \right\} / \text{iso},$$

where  $\wp_\infty^i$ ,  $i = 1, \dots, 4$ , are hyperplanes at infinity in  $\mathbb{P}^4$  defined by  $z_i = 0$  in homogeneous coordinates. The aim is to generalize the monad construction for the usual  $\text{SU}(N)$  instantons to the case of  $\mathbb{C}^4$ . Let then  $\mathcal{E} \in \text{Coh}(\mathbb{P}^4)$  a sheaf on  $\mathbb{P}^4$  as in the definition and consider  $\mathbb{P}^4 \times \mathbb{P}^4$  with the projections on the two factors

$$\begin{array}{ccc} & \mathbb{P}^4 \times \mathbb{P}^4 & \\ & \swarrow p_1 \quad \searrow p_2 & \\ \mathbb{P}^4 & & \mathbb{P}^4 \end{array}$$

One then has the Koszul resolution of  $\mathcal{O}_\Delta$ ,  $\Delta$  being the diagonal  $\Delta \cong \mathbb{P}^4 \hookrightarrow \mathbb{P}^4 \times \mathbb{P}^4$ :

$$\begin{array}{ccccccc} 0 & \longrightarrow & \Lambda^4 \widehat{\mathcal{O}} & \longrightarrow & \Lambda^3 \widehat{\mathcal{O}} & \longrightarrow & \Lambda^2 \widehat{\mathcal{O}} \\ & & & & & & \downarrow \\ & & & & & & \widehat{\mathcal{O}} \\ & & & & & & \downarrow \\ & & & & & & \widehat{\mathcal{O}} \\ & & & & & & \downarrow \\ & & & & & & \mathcal{O}_{\mathbb{P}^4 \times \mathbb{P}^4} \\ & & & & & & \downarrow \\ & & & & & & \mathcal{O}_\Delta \\ & & & & & & \downarrow \\ & & & & & & 0, \end{array}$$

where  $\widehat{\mathcal{O}} = \mathcal{O}_{\mathbb{P}^4}(1) \boxtimes \mathcal{Q}^\vee$ ,<sup>12</sup> and  $\mathcal{Q} \cong T_{\mathbb{P}^4}(-1)$ . The previous sequence tells us that  $[\mathcal{O}_\Delta] \cong [\Lambda^\bullet(\mathcal{O}_{\mathbb{P}^4}(-1) \boxtimes \mathcal{Q}^\vee)]$  in the derived category. We then define

$$C^p = \bigwedge^{-i} (\mathcal{O}_{\mathbb{P}^4}(-1) \boxtimes \mathcal{Q}^\vee),$$

by means of which we will define the Beilinson spectral sequence in this case. As  $\mathcal{E} \in \text{Coh}(\mathbb{P}^4)$ , we have the trivial identity  $p_{1*}(p_2^* \mathcal{E} \otimes \mathcal{O}_\Delta) = \mathcal{E}$ , and if we replace  $\mathcal{O}_\Delta$  by its Koszul resolution we get the double complex for the hyperdirect image, which can be expressed in terms of the Fourier-Mukai transform

$$\mathbf{R}^* p_{1,*} (p_2^* \mathcal{E} \otimes C^\bullet).$$

There are then two different spectral sequences that can be taken for the Fourier-Mukai transform. One of them gives back the trivial identity we started with, while the other one has  $E_1$ -term given by

$$E_1^{p,q} = R^q p_{1,*} (p_2^* \mathcal{E} \otimes C^p),$$

and as  $C^p = \mathcal{F}_1^p \boxtimes \mathcal{F}_2^p$ , the  $E_1$ -term can be written as

$$E_1^{p,q} = \mathcal{F}_1^p \otimes H^q(\mathbb{P}^4, \mathcal{E} \otimes \mathcal{F}_2^p).$$

<sup>12</sup>For any two coherent sheaves  $\mathcal{F}, \mathcal{G}$  on  $\mathbb{P}^4$ , we define a sheaf on  $\mathbb{P}^4 \times \mathbb{P}^4$  as  $\mathcal{F} \boxtimes \mathcal{G} = p_1^* \mathcal{F} \otimes p_2^* \mathcal{G}$ .



virtue of a version in families of Beilinson's theorem. Indeed, let  $S$  be a scheme over  $\mathbb{C}$ . Given any coherent sheaf  $\mathcal{E}$  on  $\mathbb{P}^m \times S$  there exists a spectral sequence  $E_i^{p,q}$  whose  $E_1^{p,q}$  term is

$$E_1^{p,q} = \mathcal{O}_{\mathbb{P}^m}(p) \boxtimes R^q p_{2,*} \left( \mathcal{E} \otimes \Omega_{\mathbb{P}^4 \times S/S}^{-p}(-p) \right),$$

where  $p_2$  is the projection of  $\mathbb{P}^m \times S$  on the second factor. The spectral sequence  $E_i^{p,q}$  converges to

$$E_\infty^{q,p} = \begin{cases} \mathcal{E}, & \text{if } q + p = 0 \\ 0, & \text{otherwise.} \end{cases}$$

This approach was studied in much greater generality in [56], and more details about the proof can be found also in [57, 58]. Moreover, the scheme-theoretic isomorphism between moduli spaces of framed torsion-free sheaves on  $\mathbb{P}^m$  and quot schemes of points on  $\mathbb{A}^m$  was recently proved by means of an infinitesimal argument in [48].

*Remark B.1.* The choice of conditions constraining the sheaves  $\mathcal{E} \in \text{Coh}_{\mathbb{P}^4}$  are different from the instanton sheaf conditions found in [59] as the latter would exclude certain sheaf configurations, such as ideal sheaves of points, which are instead known to be interesting to the problem at hand.  $\blacktriangleleft$

## B.2 Orbifold instantons and ADHM data decomposition

It is interesting to notice here that in general it is not known whether a Gorenstein orbifold in dimension 4 admits a crepant resolution, and many singularities are terminal. Nonetheless some classification results, though a bit scattered, are available. It is known, for example, that orbifolds of the form  $\mathbb{C}^r/G$  admit crepant resolutions in the case of finite *abelian* subgroups  $G \subset \text{SL}(r, \mathbb{C})$  for which  $\mathbb{C}^r/G$  is a complete intersection, [60, 61], while some arithmetic condition have been derived in the case of some series of cyclic quotient singularities [62, 63]. Moreover, contrary to the lower dimensional cases, here the Hilbert-Chow morphism

$$\text{Hilb}^G(\mathbb{C}^r) \xrightarrow{\pi} \mathbb{C}^r/G$$

does not necessarily provide a crepant resolution, as there are even cases in which  $\text{Hilb}^G(\mathbb{C}^r)$  is singular despite  $\mathbb{C}^r/G$  being known to have projective crepant resolutions. Here  $\text{Hilb}^G(\mathbb{C}^r)$  denotes the  $G$ -Hilbert scheme of zero-dimensional  $G$ -invariant subschemes  $Z \subset \mathbb{C}^r$  of length  $|G|$  such that  $H^0(\mathcal{O}_Z)$  is the regular representation of  $G$ . It turns out, however, that in some cases which will be of interest to us a canonical crepant (projective) resolution of the orbifold singularity is indeed provided by the  $G$ -Hilbert scheme, and toric geometry techniques are well suited to check whether this is the case or not, [52, 53]. Let's assume for the moment that  $X = \text{Hilb}^G(\mathbb{C}^4)$  is a crepant resolution of  $\mathbb{C}^4/G$ : this will enable us to justify the ADHM data decomposition which we will use in the following in order to study the orbifold instantons

in eight dimensions. Take then the universal object  $\mathcal{Z} \subset X \times \mathbb{C}^4$  with the projections  $\pi_i$  on the  $i$ -th factor.

$$\begin{array}{ccc}
 & \mathcal{Z} & \\
 & | & \\
 & X \times \mathbb{C}^4 & \\
 \swarrow \pi_1 & & \searrow \pi_2 \\
 X & & \mathbb{C}^4
 \end{array}$$

We can then introduce the tautological bundle  $\mathcal{T}$  on  $X$  by pushing forward

$$\mathcal{T} = \pi_{1*} \mathcal{O}_{\mathcal{Z}}.$$

Under the  $G$ -action,  $\mathcal{T}$  transforms under the regular representation, and it is easy to show that its fibers are of the form  $\mathbb{C}[z_1, z_2, z_3, z_4]/I \cong H^0(\mathcal{O}_Y)$ , where  $I$  is a  $G$ -invariant ideal corresponding to the zero-dimensional subscheme  $Y \subseteq \mathbb{C}^4$  of length  $|G|$ . Multiplication by the coordinates along the fibers of  $\mathcal{T}$  induces a  $G$ -invariant homomorphism  $B \in Q \otimes \text{End}(\mathcal{T})$  (where  $Q$  denotes the regular representation of  $G$ ), which is representable by a quadruple of endomorphisms  $(B_1, \dots, B_4)$ , such that  $B \wedge B = \sum_{a < b} [B_a, B_b] = 0 \in \text{Hom}_G(\mathcal{T}, \wedge^2 Q \otimes \mathcal{T})$ . As we noticed in section 4.1 the regular representation  $Q$  may be decomposed in irreducible representations of  $G$ . This induces a decomposition of the tautological bundle as

$$\mathcal{T} = \bigoplus_r \mathcal{T}_r \otimes \rho_r.$$

The monad construction for the ADHM representation of orbifold instantons then follows from a generalisation of [13, 50, 51]. One starts from the resolution of the diagonal in  $X \times X$ , which is

$$\begin{array}{c}
 (\mathcal{T} \boxtimes \mathcal{T}^\vee \otimes \wedge^4 Q^\vee)^G \rightarrow (\mathcal{T} \boxtimes \mathcal{T}^\vee \otimes \wedge^3 Q^\vee)^G \rightarrow (\mathcal{T} \boxtimes \mathcal{T}^\vee \otimes \wedge^2 Q^\vee)^G \\
 \left. \begin{array}{c} \phantom{(\mathcal{T} \boxtimes \mathcal{T}^\vee \otimes \wedge^4 Q^\vee)^G} \\ \phantom{(\mathcal{T} \boxtimes \mathcal{T}^\vee \otimes \wedge^3 Q^\vee)^G} \end{array} \right\} \\
 \phantom{(\mathcal{T} \boxtimes \mathcal{T}^\vee \otimes \wedge^4 Q^\vee)^G} \longrightarrow (\mathcal{T} \boxtimes \mathcal{T}^\vee \otimes Q^\vee)^G \longrightarrow (\mathcal{T} \boxtimes \mathcal{T}^\vee)^G \longrightarrow \mathcal{O}_\Delta.
 \end{array}$$

We then compactify  $X$  to  $\overline{X}$  by compactifying  $\mathbb{C}^4/G$  to  $\mathbb{P}^4/G$  and resolving the singularity at the origin. We can do this by defining  $\overline{X} = X \sqcup \mathcal{P}_\infty$ , where  $\mathcal{P}_\infty \cong \mathbb{P}^3/G$ . This is useful as we can then glue objects on  $X$  and  $G$ -invariant objects on  $\mathcal{P}_\infty$  so as to get globally defined objects on  $\overline{X}$ . A globally defined resolution of

the diagonal sheaf in  $\overline{X} \times \overline{X}$  can then be obtained in this way, and we get

$$\begin{array}{c}
0 \rightarrow (\mathcal{I}(-4\mathcal{P}_\infty) \boxtimes \mathcal{I}^\vee \otimes \wedge^4 \mathcal{Q}^\vee)^G \rightarrow (\mathcal{I}(-3\mathcal{P}_\infty) \boxtimes \mathcal{I}^\vee \otimes \wedge^3 \mathcal{Q}^\vee)^G \\
\left\{ \begin{array}{l} \longrightarrow (\mathcal{I}(-2\mathcal{P}_\infty) \boxtimes \mathcal{I}^\vee \otimes \wedge^2 \mathcal{Q}^\vee)^G \longrightarrow (\mathcal{I}(-\mathcal{P}_\infty) \boxtimes \mathcal{I}^\vee \otimes \mathcal{Q}^\vee)^G \\ \longrightarrow (\mathcal{I} \boxtimes \mathcal{I}^\vee)^G \longrightarrow \mathcal{O}_\Delta \longrightarrow 0. \end{array} \right.
\end{array}$$

The construction then follows the same strategy adopted in the previous section. In particular, if  $\mathcal{E}$  is a coherent sheaf on  $\overline{X}$ , one can take its Fourier-Mukai transform, where now  $C^\bullet$  is the resolution of the diagonal  $\mathcal{O}_\Delta$  in  $\overline{X}$ . Then, associated to the sheaf  $\mathcal{E}(-l) = \mathcal{E} \otimes \mathcal{O}_{\overline{X}}(-l\mathcal{P}_\infty)$  there is a spectral sequence  $E_i^{p,q}$ , whose  $E_1^{p,q}$ -term is given by

$$E_1^{p,q} = \left( \mathcal{I}(p) \otimes H^q(\overline{X}, \mathcal{E}(-l) \times \mathcal{I}^\vee \otimes \wedge^{-p} \mathcal{Q}^\vee) \right)^G.$$

The tensor product decomposition of the tautological bundle  $\mathcal{I}$  as  $\mathcal{I} = \bigoplus \mathcal{I}_r \otimes \rho_r$  makes it possible to reduce the homological algebra coming from the orbifold spectral sequence to the easier  $\mathbb{P}^4$  case. Indeed, cohomology groups relevant to the computations all take the form  $H^q(\overline{X}, \mathcal{E}(-l) \otimes \mathcal{I}_r)$ , which can in turn be interpreted as the equivariant decomposition of the ADHM datum.

## References

- [1] A. Belavin, A. M. Polyakov, A. Schwartz and Y. Tyupkin, *Pseudoparticle Solutions of the Yang-Mills Equations*, *Phys. Lett. B* **59** (1975) 85.
- [2] E. Corrigan, C. Devchand, D. B. Fairlie and J. Nuyts, *First-order equations for gauge fields in spaces of dimension greater than four*, *Nuclear Physics. B. Theoretical, Phenomenological, and Experimental High Energy Physics. Quantum*
- [3] D. D. Joyce, *Compact 8-manifolds with holonomy Spin(7)*, *Invent. Math.* **123** (1996) 507.
- [4] M. F. Atiyah, N. J. Hitchin, V. G. Drinfel'd and Y. I. Manin, *Construction of instantons*, *Phys. Lett. A* **65** (1978) 185.
- [5] N. Nekrasov and A. S. Schwarz, *Instantons on noncommutative  $R^{**}_4$  and  $(2,0)$  superconformal six-dimensional theory*, *Commun. Math. Phys.* **198** (1998) 689 [[hep-th/9802068](https://arxiv.org/abs/hep-th/9802068)].
- [6] E. Witten, *BPS Bound states of D0 - D6 and D0 - D8 systems in a B field*, *JHEP* **04** (2002) 012 [[hep-th/0012054](https://arxiv.org/abs/hep-th/0012054)].
- [7] N. Nekrasov, "Magnificent Four." <https://arxiv.org/abs/1712.08128>, 2017.



- [8] N. Nekrasov and N. Piazzalunga, *Magnificent four with colors*, *Comm. Math. Phys.* **372** (2019) 573.
- [9] A. Iqbal, N. Nekrasov, A. Okounkov and C. Vafa, *Quantum foam and topological strings*, *JHEP* **04** (2008) 011 [[hep-th/0312022](#)].
- [10] M. Cirafici, A. Sinkovics and R. J. Szabo, *Instantons, Quivers and Noncommutative Donaldson-Thomas Theory*, *Nucl. Phys. B* **853** (2011) 508 [[1012.2725](#)].
- [11] N. Fasola, S. Monavari and A. T. Ricolfi, *Higher rank K-theoretic Donaldson-Thomas theory of points*, [2003.13565](#).
- [12] S. K. Donaldson, *Instantons and geometric invariant theory*, *Comm. Math. Phys.* **93** (1984) 453.
- [13] P. B. Kronheimer and H. Nakajima, *Yang-Mills instantons on ALE gravitational instantons*, *Mathematische Annalen* **288** (1990) 263.
- [14] L. Baulieu, H. Kanno and I. Singer, *Special quantum field theories in eight-dimensions and other dimensions*, *Commun. Math. Phys.* **194** (1998) 149 [[hep-th/9704167](#)].
- [15] B. S. Acharya, M. O’Loughlin and B. J. Spence, *Higher dimensional analogs of Donaldson-Witten theory*, *Nucl. Phys. B* **503** (1997) 657 [[hep-th/9705138](#)].
- [16] M. Blau and G. Thompson, *Euclidean SYM theories by time reduction and special holonomy manifolds*, *Phys. Lett. B* **415** (1997) 242 [[hep-th/9706225](#)].
- [17] S. K. Donaldson and R. P. Thomas, *Gauge theory in higher dimensions*, in *The geometric universe (Oxford, 1996)*, pp. 31–47. Oxford Univ. Press, Oxford, 1998.
- [18] Y. Cao and N. Conan Leung, *Donaldson-Thomas theory for Calabi-Yau 4-folds*, *arXiv e-prints* (2014) arXiv:1407.7659 [[1407.7659](#)].
- [19] Y. Cao and N. C. Leung, *Orientability for gauge theories on Calabi-Yau manifolds*, *Adv. Math.* **314** (2017) 48.
- [20] Y. Cao and M. Kool, *Zero-dimensional Donaldson-Thomas invariants of Calabi-Yau 4-folds*, *Adv. Math.* **338** (2018) 601 [[1712.07347](#)].
- [21] Y. Cao, M. Kool and S. Monavari, “K-theoretic DT/PT correspondence for toric Calabi–Yau 4-folds.” <https://arxiv.org/abs/1906.07856>, 2019.
- [22] J. Oh and R. P. Thomas, *Counting sheaves on Calabi-Yau 4-folds, I*, [2009.05542](#).
- [23] K. Intriligator, H. Jockers, P. Mayr, D. R. Morrison and M. Plesser, *Conifold Transitions in M-theory on Calabi-Yau Fourfolds with Background Fluxes*, *Adv. Theor. Math. Phys.* **17** (2013) 601 [[1203.6662](#)].
- [24] T. J. Hollowood, A. Iqbal and C. Vafa, *Matrix models, geometric engineering and elliptic genera*, *JHEP* **03** (2008) 069 [[hep-th/0310272](#)].
- [25] S. Franco, S. Lee and R.-K. Seong, *Brane Brick Models, Toric Calabi-Yau 4-Folds and 2d (0,2) Quivers*, *JHEP* **02** (2016) 047 [[1510.01744](#)].

- [26] F. Fucito, J. F. Morales and R. Poghossian, *Multi instanton calculus on ALE spaces*, *Nucl. Phys. B* **703** (2004) 518 [[hep-th/0406243](#)].
- [27] G. Bonelli, K. Maruyoshi and A. Tanzini, *Gauge Theories on ALE Space and Super Liouville Correlation Functions*, *Lett. Math. Phys.* **101** (2012) 103 [[1107.4609](#)].
- [28] G. Bonelli, K. Maruyoshi and A. Tanzini, *Instantons on ALE spaces and Super Liouville Conformal Field Theories*, *JHEP* **08** (2011) 056 [[1106.2505](#)].
- [29] A. Belavin, M. Bershtein, B. Feigin, A. Litvinov and G. Tarnopolsky, *Instanton moduli spaces and bases in coset conformal field theory*, *Commun. Math. Phys.* **319** (2013) 269 [[1111.2803](#)].
- [30] G. Bonelli, K. Maruyoshi, A. Tanzini and F. Yagi,  *$N=2$  gauge theories on toric singularities, blow-up formulae and  $W$ -algebras*, *JHEP* **01** (2013) 014 [[1208.0790](#)].
- [31] A. Belavin, M. Bershtein and G. Tarnopolsky, *Bases in coset conformal field theory from AGT correspondence and Macdonald polynomials at the roots of unity*, *JHEP* **03** (2013) 019 [[1211.2788](#)].
- [32] U. Bruzzo, M. Pedrini, F. Sala and R. J. Szabo, *Framed sheaves on root stacks and supersymmetric gauge theories on ALE spaces*, *Adv. Math.* **288** (2016) 1175 [[1312.5554](#)].
- [33] U. Bruzzo, F. Sala and R. J. Szabo,  *$\mathcal{N} = 2$  Quiver Gauge Theories on A-type ALE Spaces*, *Lett. Math. Phys.* **105** (2015) 401 [[1410.2742](#)].
- [34] G. Bonelli, A. Sciarappa, A. Tanzini and P. Vasko, *The Stringy Instanton Partition Function*, *JHEP* **01** (2014) 038 [[1306.0432](#)].
- [35] G. Bonelli, A. Sciarappa, A. Tanzini and P. Vasko, *Vortex partition functions, wall crossing and equivariant Gromov-Witten invariants*, *Commun. Math. Phys.* **333** (2015) 717 [[1307.5997](#)].
- [36] G. Bonelli, A. Sciarappa, A. Tanzini and P. Vasko, *Six-dimensional supersymmetric gauge theories, quantum cohomology of instanton moduli spaces and  $gl(N)$  Quantum Intermediate Long Wave Hydrodynamics*, *JHEP* **07** (2014) 141 [[1403.6454](#)].
- [37] F. Benini, G. Bonelli, M. Poggi and A. Tanzini, *Elliptic non-Abelian Donaldson-Thomas invariants of  $\mathbb{C}^3$* , *JHEP* **07** (2019) 068 [[1807.08482](#)].
- [38] H. Kanno and Y. Tachikawa, *Instanton counting with a surface operator and the chain-saw quiver*, *JHEP* **06** (2011) 119 [[1105.0357](#)].
- [39] H. Nakajima, *Handsaw quiver varieties and finite  $W$ -algebras*, *Moscow Math. J.* **12** (2012) 633 [[1107.5073](#)].
- [40] S. Jeong and N. Nekrasov, *Riemann-Hilbert correspondence and blown up surface defects*, [2007.03660](#).
- [41] G. Bonelli, N. Fasola and A. Tanzini, *Defects, nested instantons and comet shaped quivers*, [1907.02771](#).

- [42] G. Bonelli, N. Fasola and A. Tanzini, *Flags of sheaves, quivers and symmetric polynomials*, [1911.12787](#).
- [43] D. Joyce, *Compact 8-manifolds with holonomy spin (7)*., *Inventiones mathematicae* **123** (1996) 507.
- [44] E. Witten, *Topological quantum field theory*, *Comm. Math. Phys.* **117** (1988) 353.
- [45] M. Bershtein, G. Bonelli, M. Ronzani and A. Tanzini, *Exact results for  $\mathcal{N} = 2$  supersymmetric gauge theories on compact toric manifolds and equivariant Donaldson invariants*, *JHEP* **07** (2016) 023 [[1509.00267](#)].
- [46] L. Baulieu, G. Bossard and A. Tanzini, *Topological vector symmetry of BRSTQFT and construction of maximal supersymmetry*, *JHEP* **08** (2005) 037 [[hep-th/0504224](#)].
- [47] N. A. Nekrasov, *Lectures on open strings, and noncommutative gauge fields*, in *Les Houches Summer School: Session 76: Euro Summer School on Unity of Fundamental Physics: Gravity, Gauge Theory and Strings*, pp. 477–495, 3, 2002, [hep-th/0203109](#).
- [48] A. Cazzaniga and A. T. Ricolfi, *Framed sheaves on projective space and quot schemes*, [2004.13633v2](#).
- [49] M. R. Douglas and G. W. Moore, *D-branes, quivers, and ALE instantons*, [hep-th/9603167](#).
- [50] M. Cirafici, A. Sinkovics and R. J. Szabo, *Instantons, quivers and noncommutative Donaldson–Thomas theory*, *Nuclear Physics. B. Theoretical, Phenomenological, and Experimental High Energy Physics. Quantum*
- [51] H. Nakajima, *Sheaves on ALE spaces and quiver varieties*, *Moscow Mathematical Journal* **7** (2007) 699.
- [52] K. Sato, *Existence of crepant resolution for abelian quotient singularities by order  $p$  elements in dimension 4*, *Saitama Math. J.* **27** (2010) 9.
- [53] Y. Sato, *Crepant resolutions and  $\text{Hilb}^G(\mathbb{C}^4)$  for certain abelian subgroups for  $SL(4, \mathbb{C})$* , *arXiv e-prints* (2019) arXiv:1905.06244 [[1905.06244](#)].
- [54] D. Borisov and D. Joyce, *Virtual fundamental classes for moduli spaces of sheaves on Calabi-Yau four-folds*, *Geometry & Topology* **21** (2017) 3231.
- [55] K. Furuuchi, *Instantons on noncommutative  $R^{**}4$  and projection operators*, *Prog. Theor. Phys.* **103** (2000) 1043 [[hep-th/9912047](#)].
- [56] A. A. Henni and D. M. Guimarães, *A note on the adhm description of quot schemes of points on affine spaces*, *arXiv e-prints* (2017) arXiv:1712.04369.
- [57] A. A. Henni and M. Jardim, *Commuting matrices and the Hilbert scheme of points on affine spaces*, *Adv. Geom.* **18** (2018) 467.
- [58] A. A. Henni, M. Jardim and R. V. Martins, *ADHM construction of perverse instanton sheaves*, *Glasgow Mathematical Journal* **57** (2015) 285.

- [59] M. Jardim, *Instanton sheaves on complex projective spaces*, *Universitat de Barcelona. Collectanea Mathematica* **57** (2006) 69.
- [60] D. I. Dais, M. Henk and G. M. Ziegler, *All abelian quotient C.I.-singularities admit projective crepant resolutions in all dimensions*, *Advances in Mathematics* **139** (1998) 194.
- [61] D. I. Dais, C. Haase and G. M. Ziegler, *All toric local complete intersection singularities admit projective crepant resolutions*, *Tohoku Math. J. (2)* **53** (2001) 95.
- [62] D. I. Dais, U.-U. Haus and M. Henk, *On crepant resolutions of 2-parameter series of Gorenstein cyclic quotient singularities*, *Results Math.* **33** (1998) 208.
- [63] D. I. Dais, M. Henk and G. M. Ziegler, *On the existence of crepant resolutions of Gorenstein abelian quotient singularities in dimensions  $\geq 4$* , in *Algebraic and geometric combinatorics*, vol. 423 of *Contemp. Math.*, pp. 125–193. Amer. Math. Soc., Providence, RI, 2006. [DOI](#).