

Research Article

Andrea Braides* and Margherita Solci

Compactness by Coarse-Graining in Long-Range Lattice Systems

<https://doi.org/10.1515/ans-2020-2100>

Received December 17, 2019; accepted July 4, 2020

Abstract: We consider energies on a periodic set \mathcal{L} of the form $\sum_{i,j \in \mathcal{L}} a_{ij}^\varepsilon |u_i - u_j|$, defined on spin functions $u_i \in \{0, 1\}$, and we suppose that the typical range of the interactions is R_ε with $R_\varepsilon \rightarrow +\infty$, i.e., if $|i - j| \leq R_\varepsilon$, then $a_{ij}^\varepsilon \geq c > 0$. In a discrete-to-continuum analysis, we prove that the overall behavior as $\varepsilon \rightarrow 0$ of such functionals is that of an interfacial energy. The proof is performed using a coarse-graining procedure which associates to scaled functions defined on $\varepsilon\mathcal{L}$ with equibounded energy a family of sets with equibounded perimeter. This agrees with the case of equibounded R_ε and can be seen as an extension of coerciveness result for short-range interactions, but is different from that of other long-range interaction energies, whose limit exits the class of surface energies. A computation of the limit energy is performed in the case $\mathcal{L} = \mathbb{Z}^d$.

Keywords: Homogenization, Lattice Systems, Long-Range Interactions, Interfacial Energies, Coarse Graining

MSC 2010: 49J45, 49Q20, 35B27, 82B20

Communicated by: Aldo Pratelli

1 Introduction

In this paper, we give a contribution to the general problem of the asymptotic analysis of systems of lattice interactions of the form

$$\sum_{i,j \in \mathcal{L}} a_{ij}^\varepsilon |u_i - u_j|,$$

where \mathcal{L} is a periodic lattice in \mathbb{R}^d , $\varepsilon > 0$ is a parameter tending to 0 and a_{ij}^ε are non-negative coefficients. These functionals depend on (scalar) “spin functions” with $u_i \in \{0, 1\}$, somehow related to ferromagnetic energies in the terminology of statistical mechanics (where usually $u_i \in \{-1, 1\}$).

We investigate coerciveness properties related to such energies in a discrete-to-continuum process, where the values u_i^ε are identified as the values $u^\varepsilon(\varepsilon i)$ of a function defined on $\varepsilon\mathcal{L}$. In this way, a continuum limit of u^ε can be defined as a limit of their piecewise-constant interpolations, e.g., defined as $u^\varepsilon(x) = u_i^\varepsilon$ if the point of minimum distance of $\varepsilon\mathcal{L}$ from x is εi . Coerciveness is established by exhibiting scales s_ε such that if u_i^ε are such that

$$\sum_{i,j \in \mathcal{L}} a_{ij}^\varepsilon |u_i^\varepsilon - u_j^\varepsilon| \leq s_\varepsilon,$$

then the interpolations u^ε are precompact in some topology and their limit points are in general non-trivial. This can be expressed by proving that the domain of the Γ -limit of the scaled energies

$$\frac{1}{s_\varepsilon} \sum_{i,j \in \mathcal{L}} a_{ij}^\varepsilon |u_i^\varepsilon - u_j^\varepsilon|$$

in that topology is not trivial.

*Corresponding author: **Andrea Braides**, Dipartimento di Matematica, Università di Roma Tor Vergata, via della ricerca scientifica 1, 00133 Roma, Italy, e-mail: braides@mat.uniroma2.it

Margherita Solci, DADU, Università di Sassari, piazza Duomo 6, 07041 Alghero (SS), Italy, e-mail: margherita@uniss.it

The simplest case that has been previously treated [1, 16] is nearest-neighbor interactions; i.e., when a_{ij}^ε are strictly positive only when i, j are nearest neighbors (n.n. for short) in the Delaunay triangulation of \mathcal{L} (e.g., $|i - j| = 1$ if $\mathcal{L} = \mathbb{Z}^d$). In this case, choosing $s_\varepsilon = \varepsilon^{1-d}$ gives that the scaled energies

$$\sum_{i,j \in \mathcal{L} \text{ i.n.n.}} \varepsilon^{d-1} a_{ij}^\varepsilon |u_i^\varepsilon - u_j^\varepsilon|$$

can be directly seen as a (possibly anisotropic) perimeter of the sets $\{x : u^\varepsilon(x) = 1\}$ defined through the piecewise-constant interpolation of u^ε from the scaled lattice $\varepsilon\mathcal{L}$. Then the compactness properties of sets of equibounded perimeter ensure the coerciveness in $L^1_{\text{loc}}(\mathbb{R}^d)$ and the limits are characteristic functions. Moreover, the Γ -limit of the energies can be described by an energy defined on sets of finite perimeter A , which, in the simplest homogeneous case, takes the form

$$\int_{\partial A} \varphi(v) d\mathcal{H}^{d-1}. \tag{1.1}$$

The same scaling works for finite-range interactions, i.e., when a_{ij}^ε is 0 if $|i - j| > R$ for some R , even though the energies in that case must be interpreted as a non-local perimeter [14]. The finiteness of the range of the interactions can be weakened to a decay condition that can be quantified as

$$\sup \left\{ \sum_{j \in \mathcal{L} \setminus \{i\}} a_{ij}^\varepsilon |j - i| : i \in \mathcal{L}, \varepsilon > 0 \right\} < +\infty, \tag{1.2}$$

even though the limit energies may have a non-local part if the “tails” of these series are not uniformly negligible [2]. We note that such analysis is valid beyond pair potentials and generalizes to classes of many-point interactions (see [12]).

If the decay assumptions (1.2) do not hold, then the “natural” scaling for the energies may be different from the “surface scaling” ε^{d-1} , and we might exit the class of interfacial energies. An extreme case is that of “dense graphs”, which is better stated in a bounded domain, i.e., when considering energies

$$\sum_{i,j \in \mathcal{L} \cap \frac{1}{\varepsilon}Q} a_{ij}^\varepsilon |u_i - u_j|,$$

with Q a cube in \mathbb{R}^d , and suppose that $a_{ij}^\varepsilon \geq c > 0$ for (a positive percentage of) all interactions. In that case, the scaling is $s_\varepsilon = \varepsilon^{-2d}$, and the limit behavior is described by a more abstract limit functional called a “graphon” energy [3, 5, 20, 21], which can be viewed as a relaxation of a double integral on $(0, 1)$ of the form

$$\int_{(0,1) \times (0,1)} W(x, y) |v(x) - v(y)| dx dy$$

defined on $BV((0, 1); \{0, 1\})$ after a complex and rather abstract relabeling procedure and identification of functions defined on Q with functions defined on $(0, 1)$ (see [9]). For sparse graphs (i.e., graphs which are not dense according to the definition above) and interactions not satisfying decay conditions (1.2), the correct scaling, the relative convergence and the form of the Γ -limit is a complex open problem. In [8], an example is given of one-dimensional energies with range $R_\varepsilon = 1/\sqrt{\varepsilon}$ such that a non-trivial Γ -limit exists for $s_\varepsilon = 1/\sqrt{\varepsilon}$ with respect to the L^∞ -weak* convergence, but it is defined on *all* functions of bounded variation with values in $[0, 1]$ (and not only those with values in $\{0, 1\}$). In that example, a crucial issue is the topology of the graph of the connections where $a_{ij}^\varepsilon \neq 0$.

In this paper, we consider an intermediate case, i.e., when the decay condition described above does not hold, and $a_{ij}^\varepsilon \geq c > 0$ when $|i - j| \leq R_\varepsilon$ with $R_\varepsilon \gg 1$ but the topology of the interactions within that range is that of a “dense” graph. We further make the assumption $\varepsilon R_\varepsilon \ll 1$ so that the discrete-to-continuum process makes sense. We note that this latter condition is not restrictive upon a redefinition of ε in terms of R_ε , e.g. taking $R_\varepsilon^{-1/2}$ in the place of ε . We keep the dependence of our system on R_ε and ε separate since these parameters may be defined independently in applications. Under these conditions, we have

$$s_\varepsilon = \frac{R_\varepsilon^{d+1}}{\varepsilon^{d-1}},$$

and with this scaling, functions of equibounded energy interpolated on the lattice $\varepsilon\mathcal{L}$ converge to a characteristic function of a set of finite perimeter. The main argument for obtaining this result is by coarse-graining. Namely, we average the values of u^ε for interaction on cubes with side length of order R_ε so that we can think of those averages as labeled on $\varepsilon R_\varepsilon \mathbb{Z}^d$. We prove first that those labels for which averages are not essentially close to 0 and 1 are negligible; hence, we may regard such functions as spin functions defined on a cubic lattice. Then we show that the arguments used for nearest-neighbor interactions of [1] can be adapted for the interpolated functions of the averages. Once a limit set of finite perimeter is obtained, we can prove the convergence of the interpolations of the original functions to the same set.

As an application of this scaling argument, we show that, for

$$a_{ij}^\varepsilon = a\left(\frac{i-j}{R_\varepsilon}\right),$$

where a is a positive function with $\int a(\xi)|\xi| d\xi$ finite, the Γ -limit of the energies

$$F_\varepsilon(u) = \frac{\varepsilon^{d-1}}{R_\varepsilon^{d+1}} \sum_{i,j \in \mathbb{Z}^d} a_{ij}^\varepsilon |u_i - u_j|,$$

defined on the cubic lattice of \mathbb{R}^d , is given by an energy as in (1.1) with

$$\varphi(v) = \int_{\mathbb{R}^d} a(\xi)|\langle \xi, v \rangle| d\xi. \tag{1.3}$$

In particular, if a is radially symmetric, then (1.1) is simply a multiple of the perimeter of A . It is interesting to note that, in a sense, the case $R_\varepsilon \rightarrow +\infty$ can be seen as a limit of the case of R_ε finite, for which the Γ -limit is of form (1.1) with the integrand $\varphi(v)$ given by a discretization of the integral in (1.3) (as seen in [4, 11, 17] in a slightly different context). This convergence can be re-obtained using the results in [19], where transportation maps are used to transform discrete energies in convolution functionals.

2 A Compactness Result

We denote by $Q_R = [-\frac{R}{2}, \frac{R}{2})^d$ the (semi-open) coordinate cube centered in 0 and with side length R in \mathbb{R}^d , by B_R the open ball centered in 0 and with side length R in \mathbb{R}^d , and by e_1, \dots, e_d the vectors of the canonical basis of \mathbb{R}^d . Moreover, \mathcal{H}^{d-1} denotes the $d - 1$ -dimensional Hausdorff measure and $|\cdot|$ the Lebesgue d -dimensional measure.

Let $\mathcal{L} \subset \mathbb{R}^d$ be a discrete periodic set. We can suppose without loss of generality that it is periodic in the coordinate directions with period 1; i.e., $\mathcal{L} + e_i = \mathcal{L}$ for all $i \in \{1, \dots, d\}$. The Voronoi cells of \mathcal{L} are defined as

$$V_i = \{x \in \mathbb{R}^d : |x - i| < |x - j| \text{ for all } j \in \mathcal{L}, j \neq i\}.$$

By the periodicity of \mathcal{L} , there exists a constant $C_{\mathcal{L}} > 0$ such that

$$\frac{1}{C_{\mathcal{L}}} \leq |V_i| \leq C_{\mathcal{L}}, \quad \frac{1}{C_{\mathcal{L}}} \leq \mathcal{H}^{d-1}(\partial V_i) \leq C_{\mathcal{L}}. \tag{2.1}$$

Each spin function $u : \varepsilon\mathcal{L} \rightarrow \{0, 1\}$ is identified with its piecewise-constant interpolation, which is the L^∞ function defined by

$$u(x) = u(\varepsilon i) \quad \text{if } x \in \varepsilon V_i; \quad \text{i.e., } |x - \varepsilon i| < |x - \varepsilon j| \quad \text{for all } j \in \mathcal{L}, j \neq i.$$

Note that, by (2.1), the L^1 norm of such u is equivalent to $\varepsilon^d \#\{i : u_i \neq 0\}$.

In this section, we prove coerciveness properties for energies E_ε defined on spin functions $u : \varepsilon\mathcal{L} \rightarrow \{0, 1\}$ by

$$E_\varepsilon(u) = \frac{\varepsilon^{2d}}{\eta^{d+1}} \sum_{i,j \in \mathcal{L}, i-j \in Q_{\eta/\varepsilon}} |u_i - u_j|, \tag{2.2}$$

where we denote $u_i = u(\varepsilon i)$, and $\eta = \eta_\varepsilon$ are such that

$$\lim_{\varepsilon \rightarrow 0} \eta_\varepsilon = \lim_{\varepsilon \rightarrow 0} \frac{\varepsilon}{\eta_\varepsilon} = 0. \tag{2.3}$$

Lemma 1 (Compactness). *Let u^ε be spin functions such that $E_\varepsilon(u^\varepsilon)$ is equibounded. Then, up to subsequences, the corresponding piecewise-constant interpolations, still denoted by u^ε , converge in $L^1_{\text{loc}}(\mathbb{R}^d)$ to $u = \chi_A$, where A is a set of finite perimeter.*

Proof. The idea of the proof is to subdivide the set of indices \mathcal{L} into disjoint cubes of side length $\frac{\eta}{4\varepsilon}$. The factor 4 is chosen so that if we consider i, j indices belonging to two neighboring cubes with this side length, respectively, then $i - j \in Q_{\eta/\varepsilon}$ so that they interact in energy E_ε . In such a way, we can associate to each u^ε and each such smaller cube the value 0 or 1 of the “majority phase” if such majority phase is sufficiently close to 0 and 1, respectively, while we prove that the remaining cubes can be neglected. In this way, we will construct coarse-grained functions for which the energy E_ε can be viewed as a standard nearest-neighbor ferromagnetic energy, and the compactness then follows by interpreting spin functions as sets of finite perimeter.

For any $k \in \mathbb{Z}^d$, we set

$$Q_k^\varepsilon = \frac{\eta}{4\varepsilon} k + Q_{\eta/(4\varepsilon)}.$$

For $u: \varepsilon\mathcal{L} \rightarrow \{0, 1\}$, we define

$$D(\varepsilon, k)(u) = \frac{|\#\{i \in Q_k^\varepsilon \cap \mathcal{L} : u_i = 1\} - \#\{i \in Q_k^\varepsilon \cap \mathcal{L} : u_i = 0\}|}{\#(Q_k^\varepsilon \cap \mathcal{L})}.$$

Note that $D(\varepsilon, k)(u)$ measures how close the function u is to its majority phase; more precisely, $D(\varepsilon, k)(u) = 1$ if u is constant on $Q_k^\varepsilon \cap \mathcal{L}$, while $D(\varepsilon, k)(u) = 0$ if the values of u are equally distributed between 0 and 1 in $Q_k^\varepsilon \cap \mathcal{L}$.

With fixed $\delta \in (0, 1)$, we define $\mathcal{B}^\varepsilon(u) = \{k \in \mathbb{Z}^d : D(\varepsilon, k)(u) < 1 - \delta\}$. The Q_k^ε corresponding to $k \in \mathcal{B}^\varepsilon(u)$ will be considered as the cubes where u is not close to a phase 1 or 0. We will first show that such cubes are negligible. Indeed, note that, thanks to the first inequality in (2.1), the number of pairs of indices i, j within Q_k^ε are of order $(\frac{\eta}{\varepsilon})^{2d}$, and hence, there exists $C_\delta > 0$ such that if $k \in \mathcal{B}^\varepsilon(u)$, then the number of “interactions within the cube” Q_k^ε is at least $C_\delta(\frac{\eta}{\varepsilon})^{2d}$, namely, $\#\{(i, j) : i, j \in Q_k^\varepsilon \cap \mathcal{L}, u_i \neq u_j\} \geq C_\delta(\frac{\eta}{\varepsilon})^{2d}$.

Hence, if u^ε are as in the hypotheses of the lemma, that is, $F_\varepsilon(u^\varepsilon) \leq c$, we have $\#\mathcal{B}^\varepsilon(u^\varepsilon) \leq \frac{c}{C_\delta} \eta^{1-d}$. We can estimate the measures

$$\begin{aligned} \left| \bigcup_{k \in \mathcal{B}^\varepsilon(u^\varepsilon)} \varepsilon Q_k^\varepsilon \right| &= \#\mathcal{B}^\varepsilon(u^\varepsilon) \frac{\eta^d}{4^d} \leq \frac{c}{4^d C_\delta} \eta, \\ \mathcal{H}^{d-1}\left(\partial \bigcup_{k \in \mathcal{B}^\varepsilon(u^\varepsilon)} \varepsilon Q_k^\varepsilon\right) &= \#\mathcal{B}^\varepsilon(u^\varepsilon) \frac{2d\eta^{d-1}}{4^{d-1}} \leq \frac{2cd}{4^{d-1} C_\delta}. \end{aligned} \tag{2.4}$$

As for the indices such that $D(\varepsilon, k)(u) \geq 1 - \delta$, we subdivide them into the sets

$$\begin{aligned} \mathcal{A}_1^\varepsilon(u) &= \{k \in \mathbb{Z}^d : D(\varepsilon, k)(u) \geq 1 - \delta, \#\{i \in Q_k^\varepsilon : u_i = 1\} > \#\{i \in Q_k^\varepsilon : u_i = 0\}\}, \\ \mathcal{A}_0^\varepsilon(u) &= \{k \in \mathbb{Z}^d : D(\varepsilon, k)(u) \geq 1 - \delta, \#\{i \in Q_k^\varepsilon : u_i = 1\} < \#\{i \in Q_k^\varepsilon : u_i = 0\}\} \end{aligned}$$

and define

$$K_j^\varepsilon(u) = \bigcup_{k \in \mathcal{A}_j^\varepsilon(u)} \varepsilon Q_k^\varepsilon \quad \text{for } j = 0, 1.$$

In order to estimate the measure of the boundary of $K_1^\varepsilon(u)$, we estimate the number of cubes Q_k^ε with $k \in \mathcal{A}_1^\varepsilon(u)$ which have a side in common with a cube $Q_{k'}^\varepsilon$ with $k' \in \mathcal{A}_0^\varepsilon(u)$, parameterized on the set

$$\mathcal{A}^\varepsilon(u) := \{k \in \mathcal{A}_1^\varepsilon(u) : k + e_j \in \mathcal{A}_0^\varepsilon(u) \text{ for some } j = 1, \dots, d\}.$$

To that end, note that if $D(\varepsilon, k)(u) \geq 1 - \delta$ and $k \in \mathcal{A}_1^\varepsilon(u)$, then

$$\#\{i \in Q_k^\varepsilon \cap \mathcal{L} : u_i = 1\} \geq \left(1 - \frac{\delta}{2}\right) \#\{i \in Q_k^\varepsilon \cap \mathcal{L}\}$$

so that, again recalling the first inequality in (2.1), each site $i \in Q_k^\varepsilon$ such that $u_i = 1$ interacts with $C'_\delta(\frac{\eta}{\varepsilon})^d$ and conversely for each site $i \in Q_{k'}^\varepsilon$ such that $u_i = 0$. Hence, the interacting pairs $(i, j) \in Q_k^\varepsilon \times Q_{k'}^\varepsilon$ are at least $C'_\delta(\frac{\eta}{\varepsilon})^{2d}$.

Hence,

$$E_\varepsilon(u) \geq \frac{\varepsilon^{2d}}{\eta^{d+1}} \#A^\varepsilon(u) C''_\delta \left(\frac{\eta}{\varepsilon}\right)^{2d} = C''_\delta \#A^\varepsilon(u) \eta^{d-1}$$

so that

$$\#A^\varepsilon(u) \leq \frac{1}{C''_\delta} E_\varepsilon(u) \eta^{1-d}.$$

For the functions u^ε , we then obtain

$$\#A^\varepsilon(u^\varepsilon) \leq \frac{c}{C''_\delta} \eta^{1-d}$$

so that

$$\begin{aligned} \mathcal{H}^{d-1}(\partial K_1^\varepsilon(u^\varepsilon)) &\leq 2d \left(\mathcal{H}^{d-1} \left(\partial \bigcup_{k \in \mathcal{A}^\varepsilon(u^\varepsilon)} \varepsilon Q_k^\varepsilon \right) + \mathcal{H}^{d-1} \left(\partial \bigcup_{k \in \mathcal{B}^\varepsilon(u^\varepsilon)} \varepsilon Q_k^\varepsilon \right) \right) \\ &\leq 2d \left(\#A^\varepsilon(u^\varepsilon) \frac{2d\eta^{d-1}}{4^{d-1}} + \frac{2cd}{4^{d-1}C_\delta} \right) \leq C'''_\delta \end{aligned}$$

where C'''_δ is a positive constant depending only on d , c and δ . By the compactness of sets of equibounded perimeter, this shows that the characteristic functions of the sets $K_1^\varepsilon(u^\varepsilon)$ are compact in $L^1_{\text{loc}}(\mathbb{R}^d)$. The symmetric argument shows also that $K_0^\varepsilon(u^\varepsilon)$ are compact in $L^1_{\text{loc}}(\mathbb{R}^d)$. Moreover, if we denote a limit of the sets $K_j^\varepsilon(u^\varepsilon)$ by K_j , then we have

$$|\mathbb{R}^d \setminus (K_0 \cup K_1)| = 0 \quad (2.5)$$

by (2.4). We highlight the possible dependence of the sets obtained by this procedure on δ by renaming them K_1^δ and K_0^δ .

Note that if $\delta < \delta'$, then $K_1^{\delta'} \subset K_1^\delta$ and $K_0^{\delta'} \subset K_0^\delta$. Since in both cases (2.5) holds, we must have $K_1^{\delta'} = K_1^\delta$ and $K_0^{\delta'} = K_0^\delta$ so that these sets are independent of δ and we may go back to denoting them by K_1 and K_0 .

We can now prove the convergence of u^ε . Fixed $\delta < 1$ as above, we write

$$u^\varepsilon = u^\varepsilon \chi_{K_1^\varepsilon(u^\varepsilon)} + u^\varepsilon \chi_{K_0^\varepsilon(u^\varepsilon)} + u^\varepsilon \chi_{\mathbb{R}^d \setminus (K_1^\varepsilon(u^\varepsilon) \cup K_0^\varepsilon(u^\varepsilon))}.$$

By (2.4), the last term converges to 0 in $L^1(\mathbb{R}^d)$. As for the other two terms, we localize the convergence by restricting to a cube Q_R . Note that, for $k \in \mathcal{A}_1^\varepsilon(u^\varepsilon)$, we have

$$\|u^\varepsilon - 1\|_{L^1(\varepsilon Q_k^\varepsilon)} \leq C(1 - \delta)\eta^d$$

so that

$$\|u^\varepsilon \chi_{K_1^\varepsilon(u^\varepsilon) \cap Q_R} - \chi_{K_1^\varepsilon(u^\varepsilon) \cap Q_R}\|_{L^1(\mathbb{R}^d)} \leq C(1 - \delta)R^d,$$

where C denotes a positive constant not depending on δ . Analogously, for $k \in \mathcal{A}_0^\varepsilon(u^\varepsilon)$, we have

$$\|u^\varepsilon\|_{L^1(\varepsilon Q_k^\varepsilon)} \leq C(1 - \delta)\eta^d,$$

and hence,

$$\|u^\varepsilon \chi_{K_0^\varepsilon(u^\varepsilon) \cap Q_R}\|_{L^1(\mathbb{R}^d)} \leq C(1 - \delta)R^d.$$

We then have, by the local convergence of $K_j^\varepsilon(u^\varepsilon)$,

$$\begin{aligned} &\limsup_{\varepsilon \rightarrow 0} \|u^\varepsilon \chi_{K_1^\varepsilon(u^\varepsilon) \cap Q_R} - \chi_{K_1 \cap Q_R}\|_{L^1(\mathbb{R}^d)} \\ &\leq \limsup_{\varepsilon \rightarrow 0} (\|u^\varepsilon \chi_{K_1^\varepsilon(u^\varepsilon) \cap Q_R} - \chi_{K_1^\varepsilon(u^\varepsilon) \cap Q_R}\|_{L^1(\mathbb{R}^d)} + \|\chi_{K_1^\varepsilon(u^\varepsilon) \cap Q_R} - \chi_{K_1 \cap Q_R}\|_{L^1(\mathbb{R}^d)}) \leq C(1 - \delta)R^d \end{aligned}$$

and

$$\limsup_{\varepsilon \rightarrow 0} \|u^\varepsilon \chi_{K_0^\varepsilon(u^\varepsilon) \cap Q_R}\|_{L^1(\mathbb{R}^d)} \leq C(1 - \delta)R^d$$

so that, by the arbitrariness of δ , u^ε converge locally to χ_{K_1} . \square

Remark 2. The proof of Lemma 1 works exactly in the same way if we suppose that “almost all” pairs of indices of \mathcal{L} within $Q_{\eta/\varepsilon}$ interact; namely, if in place of energy (2.2), we consider

$$E_\varepsilon(u) = \frac{\varepsilon^{2d}}{\eta^{d+1}} \sum_{i,j \in \mathcal{L}: i-j \in Q_{\eta/\varepsilon}} a_{ij}^\varepsilon |u_i - u_j|,$$

with the requirement that there exists $c > 0$ such that

$$\lim_{\varepsilon \rightarrow 0} \frac{\#\{(i, j) : i, j \in x + Q_{\eta/\varepsilon} : a_{ij}^\varepsilon \geq c\}}{\#\{(i, j) : i, j \in x + Q_{\eta/\varepsilon}\}} = 1 \tag{2.6}$$

uniformly in $x \in \mathbb{R}^d$. Condition (2.6) is trivially satisfied by energies (2.2) for $c = 1$.

Note that condition (2.6) cannot be relaxed to “having a proportion” of pairs of indices of \mathcal{L} within $Q_{\eta/\varepsilon}$ interacting, however large this proportion may be below 1; i.e., it is not sufficient that

$$\lim_{\varepsilon \rightarrow 0} \frac{\#\{(i, j) : i, j \in x + Q_{\eta/\varepsilon} : a_{ij}^\varepsilon \geq c\}}{\#\{(i, j) : i, j \in x + Q_{\eta/\varepsilon}\}} \geq \lambda \tag{2.7}$$

for any $\lambda < 1$. To check this, we may consider the following example: choose $\mathcal{L} = \mathbb{R}^d$, fix $N \in \mathbb{N}$, and define

$$a_{ij}^\varepsilon = \begin{cases} 1 & \text{if } i - j \in Q_{\eta/\varepsilon} \text{ and both } i, j \notin N\mathbb{Z}^d, \\ 0 & \text{otherwise.} \end{cases}$$

Then (2.7) holds for $\lambda = (1 - \frac{1}{N^d})^2$, but if we define

$$u_i^\varepsilon = \begin{cases} 1 & \text{if } i \notin N\mathbb{Z}^d, \\ 0 & \text{if } i \in N\mathbb{Z}^d, \end{cases}$$

then u^ε converge weakly in $L^1_{\text{loc}}(\mathbb{R}^d)$ to the constant $1 - \frac{1}{N^d}$. Since $E_\varepsilon(u_\varepsilon) = 0$, this shows that Lemma 1 does not hold.

In this example, the subset $N\mathbb{Z}^d$ of \mathbb{Z}^d can be considered as a “perforation” of the domain and can be treated as such, considering convergence only of the restriction of the functions to $\mathbb{Z}^d \setminus N\mathbb{Z}^d$ (see [10, Section 3]). However, the situation can be more complicated if we take

$$a_{ij}^\varepsilon = \begin{cases} 1 & \text{if } i - j \in Q_{\eta/\varepsilon} \text{ and both } i, j \notin N\mathbb{Z}, \\ c_\varepsilon & \text{otherwise,} \end{cases}$$

that can be regarded as representing a “high-contrast medium”, for which the effect of the “perforation” cannot be neglected and for some values of c_ε may give a “double porosity” effect [10].

3 Homogenization of Long-Range Lattice Systems

Let $a: \mathbb{R}^d \rightarrow [0, +\infty)$ be such that $a(\xi)|\xi|$ is Riemann integrable on bounded sets and such that

$$\int_{\mathbb{R}^d} a(\xi)|\xi| d\xi < +\infty,$$

and

$$a(\xi) \geq c_0 \quad \text{if } |\xi| \leq r_0 \tag{3.1}$$

for some $c_0, r_0 > 0$.

Given $\varepsilon, \eta = \eta_\varepsilon$ satisfying (2.3), we define the coefficients

$$a_{ij}^\varepsilon = a_{i-j}^\varepsilon = a\left(\frac{\varepsilon(i-j)}{\eta}\right)$$

for $i, j \in \mathbb{Z}^d$, and the energies

$$F_\varepsilon(u) = \frac{\varepsilon^{2d}}{\eta^{d+1}} \sum_{i,j \in \mathbb{Z}^d} a_{i-j}^\varepsilon |u_i - u_j|. \tag{3.2}$$

Definition 3. A family $\{u^\varepsilon\}$ of functions $u^\varepsilon : \varepsilon\mathbb{Z}^d \rightarrow \{0, 1\}$ converges to a set $A \subset \mathbb{R}^d$ if the piecewise-constant interpolations of u^ε converge to the characteristic function χ_A in $L^1_{\text{loc}}(\mathbb{R}^d)$ as $\varepsilon \rightarrow 0$.

By hypothesis (3.1), we may apply the compactness lemma (Lemma 1), obtaining that the family $\{F_\varepsilon\}$ is coercive with respect to this convergence.

Proposition 4. Let $\{u^\varepsilon\}$ be such that $\sup_\varepsilon F_\varepsilon(u^\varepsilon) < +\infty$. Then, up to subsequences, there exists a set of finite perimeter A such that u^ε converge to A in the sense of Definition 3.

This coerciveness property justifies the computation of the Γ -limit of F_ε with respect to the convergence in Definition 3. We use standard notation in the theory of sets of finite perimeter (see, e.g., [6, 22]).

Theorem 5 (Homogenization). The functionals defined in (3.2) Γ -converge with respect to the convergence in Definition 3 to the functional F defined on sets of finite perimeter by

$$F(A) = \int_{\partial^* A} \varphi_a(\nu) d\mathcal{H}^{d-1},$$

where $\partial^* A$ denotes the reduced boundary of A , ν the outer normal to A and φ_a is given by

$$\varphi_a(\nu) = \int_{\mathbb{R}^d} a(\xi) |\langle \xi, \nu \rangle| d\xi.$$

Proof. In order to better illustrate the proof in the general d -dimensional case, we first deal with the one-dimensional case, in which we may highlight the coarse-graining procedure without the technical complexities of the higher-order geometry. In this case, we may rewrite the energies as

$$F_\varepsilon(u) = \frac{\varepsilon^2}{\eta^2} \sum_{\xi \in \mathbb{Z}} \sum_{i \in \mathbb{Z}} a_\xi^\varepsilon |u_{i+\xi} - u_i|.$$

The relevant computation is that of the lower bound for the target $A = [0, +\infty)$. Let u^ε converge to A . For each $\xi \in \mathbb{Z} \setminus \{0\}$ and $i \in \{1, \dots, |\xi|\}$, we consider the function u^ε restricted to $\varepsilon i + \varepsilon \xi \mathbb{Z}$. By the L^1_{loc} convergence, we may suppose that each such restriction changes value; i.e., there exists some $k_{i,\xi} \in \mathbb{Z}$ such that

$$u^\varepsilon(\varepsilon i + \varepsilon k_{i,\xi} \xi) = 0 \quad \text{and} \quad u^\varepsilon(\varepsilon i + \varepsilon(k_{i,\xi} + 1)\xi) = 1.$$

The set of ξ and i for which this does not hold is negligible for $\varepsilon \rightarrow 0$; the precise proof is directly given for the d -dimensional functionals below. For each ξ , we then have

$$\sum_{i=1}^{|\xi|} \sum_{k \in \mathbb{Z}} a_\xi^\varepsilon |u_{i+(k+1)\xi}^\varepsilon - u_{i+k\xi}^\varepsilon| \geq |\xi| a\left(\frac{\varepsilon}{\eta} \xi\right)$$

so that

$$\liminf_{\varepsilon \rightarrow 0} F_\varepsilon(u^\varepsilon) \geq \liminf_{\varepsilon \rightarrow 0} \frac{\varepsilon^2}{\eta^2} \sum_{\xi \in \mathbb{Z}} |\xi| a\left(\frac{\varepsilon}{\eta} \xi\right) = \liminf_{\varepsilon \rightarrow 0} \sum_{\xi \in \mathbb{Z}} \frac{\varepsilon}{\eta} a\left(\frac{\varepsilon}{\eta} \xi\right) \left|\frac{\varepsilon}{\eta} \xi\right|,$$

the latter being a Riemann sum giving the integral $\int_{\mathbb{R}} a(\xi) |\xi| d\xi$, which is $F(A)$.

We now deal with the d -dimensional case. The proof of the lower bound follows the argument above, but is more complex since we must take into account the direction of the interaction vectors ξ . We prove the inequality by applying the blow-up technique (see [13, 18], and for instance [14, 15, 23] for the discrete setting).

We assume that the sequence $\{F_\varepsilon(u^\varepsilon)\}$ is equibounded and that u^ε converge in $L^1_{\text{loc}}(\mathbb{R}^d)$ to $u = \chi_A$, where A is a set of finite perimeter. Up to subsequences, we can assume that $\liminf_{\varepsilon \rightarrow 0} F_\varepsilon(u^\varepsilon) = \lim_{\varepsilon \rightarrow 0} F_\varepsilon(u^\varepsilon)$. We define the localized energy on an open set U by

$$F_\varepsilon(u^\varepsilon; U) = \frac{\varepsilon^{2d}}{\eta^{d+1}} \sum_{i \in U} \sum_{j \in \mathbb{Z}^d} a_{i-j}^\varepsilon |u_i^\varepsilon - u_j^\varepsilon|,$$

and define the measures $\mu_\varepsilon(U) = F_\varepsilon(u^\varepsilon; U)$; since the family $\{\mu_\varepsilon\}$ is equibounded, we can assume that $\mu_\varepsilon \xrightarrow{*} \mu$ up to subsequences. Now, let $\lambda = \mathcal{H}^{d-1} \llcorner \partial^* A$; the lower bound inequality follows if we show that, for \mathcal{H}^{d-1} -a.a. $x \in \partial^* A$, we have

$$\frac{d\mu}{d\lambda}(x) \geq \varphi_\alpha(v),$$

where $\frac{d\mu}{d\lambda}$ denotes the Radon–Nikodym derivative of μ with respect to the Hausdorff $d - 1$ -dimensional measure λ . By the Besicovitch derivation theorem, for \mathcal{H}^{d-1} -a.a. $x \in \partial^* A$, we have

$$\frac{d\mu}{d\lambda}(x) = \lim_{\varrho \rightarrow 0} \frac{\mu(Q_\varrho^v(x))}{\lambda(Q_\varrho^v(x))},$$

where λ is the measure $\mathcal{H}^{d-1} \llcorner \partial^* A$, v is the normal vector to $\partial^* A$ at x and $Q_\varrho^v(x)$ is a cube centered in x with side length ϱ and a face orthogonal to v . We can fix $x = 0$ and denote $Q_\varrho^v(0)$ by Q_ϱ^v . Hence, the lower bound follows if we show that

$$\lim_{\varrho \rightarrow 0} \liminf_{\varepsilon \rightarrow 0} \frac{1}{\varrho^{d-1}} F_\varepsilon(u^\varepsilon; Q_\varrho^v) \geq \varphi_\alpha(v). \tag{3.3}$$

We may therefore assume that $\varrho = \varrho_\varepsilon$ be such that $\frac{\varepsilon}{\varrho} \rightarrow 0$ and the scaled functions $u^\varepsilon(\frac{\varepsilon}{\varrho}i)$ interpolated on the lattice $\frac{\varepsilon}{\varrho}\mathbb{Z}^d$ converge to the characteristic function of the half space $H^v = \{x : \langle x, v \rangle < 0\}$ on Q_v^1 . We define

$$A_\varepsilon := \{x \in Q_v^1 : u^\varepsilon(x) \neq \chi_{H^v}(x)\}$$

so that $|A_\varepsilon| \rightarrow 0$.

If we define

$$I_{\varepsilon/\varrho}^\xi = \left\{ i \in \mathbb{Z}^d : \frac{\varepsilon}{\varrho}i, \frac{\varepsilon}{\varrho}(i + \xi) \in Q_v^1 \right\},$$

then

$$F_\varepsilon(u^\varepsilon; Q_v^\varrho) \geq \frac{\varepsilon^{2d}}{\eta^{d+1}} \sum_{\xi \in \mathbb{Z}^d} a\left(\frac{\varepsilon}{\eta}\xi\right) \sum_{i \in I_{\varepsilon/\varrho}^\xi} \left| u^\varepsilon\left(\frac{\varepsilon}{\varrho}(i + \xi)\right) - u^\varepsilon\left(\frac{\varepsilon}{\varrho}i\right) \right|.$$

We begin by estimating

$$\sum_{i \in I_{\varepsilon/\varrho}^\xi} \left| u^\varepsilon\left(\frac{\varepsilon}{\varrho}(i + \xi)\right) - u^\varepsilon\left(\frac{\varepsilon}{\varrho}i\right) \right| = \#\left\{ i \in I_{\varepsilon/\varrho}^\xi : u^\varepsilon\left(\frac{\varepsilon}{\varrho}(i + \xi)\right) \neq u^\varepsilon\left(\frac{\varepsilon}{\varrho}i\right) \right\}.$$

With fixed $\alpha \in (0, 1)$ for each $\xi \in \mathbb{Z}^d$ satisfying

$$\left| \left\langle \frac{\xi}{|\xi|}, v \right\rangle \right| \geq \frac{\alpha}{\sqrt{1 + \alpha^2}}, \tag{3.4}$$

we define

$$P^{\alpha, \xi} = \left\{ y \in \Pi_v \cap Q_v^1 : y \pm \frac{\alpha}{2|\langle \xi, v \rangle|} \xi \in Q_v^1 \right\},$$

which is not empty by (3.4), and

$$R^{\alpha, \xi} = \left\{ x \in Q_v^1 : x = y + t\xi, y \in P^{\alpha, \xi}, -\frac{\alpha}{2|\langle \xi, v \rangle|} \leq t \leq \frac{\alpha}{2|\langle \xi, v \rangle|} \right\}$$

(see Figure 1). Furthermore, we fix β with

$$\beta > \frac{\alpha}{\sqrt{1 + \alpha^2}}. \tag{3.5}$$

Since we will restrict our arguments to sets $P^{\alpha, \xi}$ and $R^{\alpha, \xi}$ above with ξ satisfying

$$\left| \left\langle \frac{\xi}{|\xi|}, v \right\rangle \right| \geq \beta, \tag{3.6}$$

we omit the dependence of the sets $P^{\alpha, \xi}$ and $R^{\alpha, \xi}$ on v since the estimates we will obtain will be independent on v .




Figure 1: The set $R^{\alpha, \xi}$

As in the one-dimensional case, we consider the functions restricted to the discrete lines $\frac{\varepsilon}{\rho}i + \frac{\varepsilon}{\rho}\xi\mathbb{Z}$. The parameter α is introduced so as to estimate the number of sites of such discrete lines inside Q_v^1 . We then set

$$B_{\varepsilon/\rho}^{\alpha, \xi} = \left\{ i \in \mathbb{Z}^d : \frac{\varepsilon}{\rho}i \in R^{\alpha, \xi} \text{ and } u^\varepsilon \text{ is not constant in } \left(\frac{\varepsilon}{\rho}i + \frac{\varepsilon}{\rho}\xi\mathbb{Z} \right) \cap R^{\alpha, \xi} \right\}.$$

Note that if $i \in B_{\varepsilon/\rho}^{\alpha, \xi}$, then $i + k\xi \in B_{\varepsilon/\rho}^{\alpha, \xi}$ for all k with $\frac{\varepsilon}{\rho}(i + k\xi) \in R^{\alpha, \xi}$ so that if we define the equivalence relation $i \sim i'$ if $i - i' \in \xi\mathbb{Z}$, we may set

$$\tilde{B}_{\varepsilon/\rho}^{\alpha, \xi} = B_{\varepsilon/\rho}^{\alpha, \xi} / \sim$$

getting

$$\#\left\{ i \in I_{\varepsilon/\rho}^{\xi} : u^\varepsilon\left(\frac{\varepsilon}{\rho}(i + \xi)\right) \neq u^\varepsilon\left(\frac{\varepsilon}{\rho}i\right) \right\} \geq \#\tilde{B}_{\varepsilon/\rho}^{\alpha, \xi}.$$

We can estimate the number of “discrete lines” intersecting $R^{\alpha, \xi}$ as

$$\begin{aligned} \#\left(\left\{ i \in \mathbb{Z}^d : \frac{\varepsilon}{\rho}i \in R^{\alpha, \xi} \right\} / \sim\right) &\geq \frac{|R^{\alpha, \xi}|}{\left(\frac{\varepsilon}{\rho}\right)^d} \frac{1}{\frac{\alpha}{\varepsilon|\langle \xi, v \rangle|}} - C_\alpha |\xi| \left(\frac{\varepsilon}{\rho}\right)^{2-d} \\ &\geq \mathcal{H}^{d-1}(P^{\alpha, \xi}) |\langle \xi, v \rangle| \left(\frac{\varepsilon}{\rho}\right)^{1-d} - C_\alpha |\xi| \left(\frac{\varepsilon}{\rho}\right)^{2-d}, \end{aligned}$$

where the last term is an error term accounting for the cubes intersecting the boundary of $R^{\alpha, \xi}$.

Note that to every element of the complement of $\tilde{B}_{\varepsilon/\rho}^{\alpha, \xi}$ there correspond at least $\lfloor \frac{\alpha}{\varepsilon|\langle \xi, v \rangle|} \rfloor$ points in $\frac{\varepsilon}{\rho}\mathbb{Z}^d \cap A_\varepsilon$ so that, for ε sufficiently small, we get

$$\begin{aligned} \#\left(\left\{ i \in \mathbb{Z}^d : \frac{\varepsilon}{\rho}i \in R^{\alpha, \xi} \text{ and } u^\varepsilon \text{ constant in } \left(\frac{\varepsilon}{\rho}i + \frac{\varepsilon}{\rho}\xi\mathbb{Z} \right) \cap R^{\alpha, \xi} \right\} / \sim\right) \\ \leq \frac{|A_\varepsilon|}{\left(\frac{\varepsilon}{\rho}\right)^d} \frac{\varepsilon|\langle \xi, v \rangle|}{\alpha} + C'_\alpha |\xi| \left(\frac{\varepsilon}{\rho}\right)^{2-d} = \frac{1}{\alpha} |A_\varepsilon| \left(\frac{\varepsilon}{\rho}\right)^{1-d} |\langle \xi, v \rangle| + C'_\alpha |\xi| \left(\frac{\varepsilon}{\rho}\right)^{2-d}, \end{aligned}$$

with C'_α again a positive constant accounting for boundary cubes, and hence,

$$\#\tilde{B}_{\varepsilon/\rho}^{\alpha, \xi} \geq \mathcal{H}^{d-1}(P^{\alpha, \xi}) |\langle \xi, v \rangle| \left(\frac{\varepsilon}{\rho}\right)^{1-d} - \frac{1}{\alpha} |A_\varepsilon| \left(\frac{\varepsilon}{\rho}\right)^{1-d} |\langle \xi, v \rangle| - (C_\alpha + C'_\alpha) |\xi| \left(\frac{\varepsilon}{\rho}\right)^{2-d}. \tag{3.7}$$

By (3.6), we can estimate

$$\mathcal{H}^{d-1}(P^{\alpha, \xi}) \geq \left(1 - \frac{\alpha}{2|\langle \xi, v \rangle|} |\xi - \langle \xi, v \rangle v|\right)^{d-1} \geq \left(1 - \frac{\alpha}{2} \sqrt{\frac{1}{\beta^2} - 1}\right)^{d-1} \tag{3.8}$$

(see also Figure 1), and hence, upon fixing $R > 0$ and introducing the set

$$\Xi_\varepsilon^v(R, \beta) = \left\{ \xi \in \mathbb{Z}^d : |\xi| \leq \frac{\eta}{\varepsilon} R, \left| \left\langle \frac{\xi}{|\xi|}, v \right\rangle \right| \geq \beta \right\},$$




Figure 2: Upper-bound construction

by (3.7) and (3.8), we have

$$\begin{aligned} \frac{1}{\varrho^{d-1}} F_\varepsilon(u^\varepsilon; Q_v^\varrho) &\geq \frac{1}{\varrho^{d-1}} \frac{\varepsilon^{2d}}{\eta^{d+1}} \sum_{\xi \in \Xi_\varepsilon^v(R, \beta)} a\left(\frac{\varepsilon}{\eta} \xi\right) \mathcal{H}^{d-1}(P^{a, \xi}) |\langle \xi, v \rangle| \left(\frac{\varepsilon}{\varrho}\right)^{1-d} \\ &\quad - \frac{1}{\varrho^{d-1}} \frac{\varepsilon^{2d}}{\eta^{d+1}} \sum_{\xi \in \Xi_\varepsilon^v(R, \beta)} a\left(\frac{\varepsilon}{\eta} \xi\right) \frac{1}{\alpha} |A_\varepsilon| \left(\frac{\varepsilon}{\varrho}\right)^{1-d} |\langle \xi, v \rangle| \\ &\quad - (C_\alpha + C'_\alpha) \left(\frac{\varepsilon}{\varrho}\right)^{2-d} \frac{1}{\varrho^{d-1}} \frac{\varepsilon^{2d}}{\eta^{d+1}} \sum_{\xi \in \Xi_\varepsilon^v(R, \beta)} a\left(\frac{\varepsilon}{\eta} \xi\right) |\xi| \\ &\geq \left(1 - \frac{\alpha}{2} \sqrt{\frac{1}{\beta^2} - 1}\right)^{d-1} \sum_{\xi \in \Xi_\varepsilon^v(R, \beta)} \left(\frac{\varepsilon}{\eta}\right)^d a\left(\frac{\varepsilon}{\eta} \xi\right) \left|\left\langle \frac{\varepsilon}{\eta} \xi, v \right\rangle\right| \\ &\quad - \frac{1}{\alpha} |A_\varepsilon| \sum_{\xi \in \Xi_\varepsilon^v(R, \beta)} \left(\frac{\varepsilon}{\eta}\right)^d a\left(\frac{\varepsilon}{\eta} \xi\right) \left|\left\langle \frac{\varepsilon}{\eta} \xi, v \right\rangle\right| \\ &\quad - (C_\alpha + C'_\alpha) \frac{\varepsilon}{\varrho} \sum_{\xi \in \Xi_\varepsilon^v(R, \beta)} \left(\frac{\varepsilon}{\eta}\right)^d a\left(\frac{\varepsilon}{\eta} \xi\right) \left|\frac{\varepsilon}{\eta} \xi\right|. \end{aligned}$$

Since $|A_\varepsilon| \rightarrow 0$ and

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \sum_{\xi \in \Xi_\varepsilon^v(R, \beta)} \left(\frac{\varepsilon}{\eta}\right)^d a\left(\frac{\varepsilon}{\eta} \xi\right) \left|\left\langle \frac{\varepsilon}{\eta} \xi, v \right\rangle\right| &= \int_{\{|\xi| \leq R, |\langle \xi / |\xi|, v \rangle| \geq \beta\}} a(\xi) |\langle \xi, v \rangle| d\xi, \\ \lim_{\varepsilon \rightarrow 0} \sum_{\xi \in \Xi_\varepsilon^v(R, \beta)} \left(\frac{\varepsilon}{\eta}\right)^d a\left(\frac{\varepsilon}{\eta} \xi\right) \left|\frac{\varepsilon}{\eta} \xi\right| &= \int_{\{|\xi| \leq R, |\langle \xi / |\xi|, v \rangle| \geq \beta\}} a(\xi) |\xi| d\xi, \end{aligned}$$

we get

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{\varrho^{d-1}} F_\varepsilon(u^\varepsilon; Q_v^\varrho) \geq \left(1 - \frac{\alpha}{2} \sqrt{\frac{1}{\beta^2} - 1}\right)^{d-1} \int_{\{|\xi| \leq R, |\langle \xi / |\xi|, v \rangle| \geq \beta\}} a(\xi) |\langle \xi, v \rangle| d\xi.$$

Note that, by (3.5), we may let first $\alpha \rightarrow 0$ and then $\beta \rightarrow 0$. We eventually obtain

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{\varrho^{d-1}} F_\varepsilon(u^\varepsilon; Q_v^\varrho) \geq \int_{\{|\xi| \leq R\}} a(\xi) |\langle \xi, v \rangle| d\xi,$$

which, by the arbitrariness of R , gives (3.3).

The upper bound is obtained by a density argument (see [7, Section 1.7]). Hence, it suffices to treat the case of A polyhedral. In this case, it suffices to take (the interpolations) $u_i^\varepsilon = \chi_A(\varepsilon i)$ for $i \in \mathbb{Z}^d$. Indeed, we write ∂A as a union of N $d - 1$ -dimensional polytopes Σ_k , and we denote by ν_k the outer normal to Σ_k and by K the $d - 2$ -dimensional skeleton of A .

We note that there exists a constant C depending only on A such that, for any $\eta, R > 0$, after removing the closed neighborhood $K + \overline{B}_{C\eta R}$ from ∂A , we obtain a disjoint collection $\tilde{\Sigma}_1, \dots, \tilde{\Sigma}_N$ with $\tilde{\Sigma}_k \subset \Sigma_k$ such that

$$(\tilde{\Sigma}_k + B_{\eta R}) \cap (\tilde{\Sigma}_{k'} + B_{\eta R}) = \emptyset \quad \text{for any } k \neq k'$$

(see Figure 2). Hence, for any $\xi \in \mathbb{Z}^d$ with $|\varepsilon\xi| \leq \eta R$, $k \in \{1, \dots, N\}$, and $j \in \mathbb{Z}^d$ such that the line $\varepsilon j + \varepsilon\xi\mathbb{R}$ intersects $\tilde{\Sigma}_k$, the values u_i^ε change only once on the points of the discrete lines $\varepsilon j + \varepsilon\xi\mathbb{Z}$ which lie in an ηR neighborhood of $\tilde{\Sigma}_k$. We note that, for lines intersecting Σ_k at a point of distance not larger than $C\eta R$ from K , such changes of value are at most N ; then, repeating the counting argument used in the lower bound, we obtain

$$\begin{aligned} \limsup_{\varepsilon \rightarrow 0} \frac{\varepsilon^{2d}}{\eta^{d+1}} \sum_{\substack{\xi \in \mathbb{Z}^d \\ |\varepsilon\xi| \leq \eta R}} a\left(\frac{\varepsilon}{\eta}\xi\right) \sum_{i \in \mathbb{Z}^d} |u_{i+\xi}^\varepsilon - u_i^\varepsilon| &\leq \limsup_{\varepsilon \rightarrow 0} \left(\sum_{k=1}^N \mathcal{H}^{d-1}(\Sigma_k) \frac{\varepsilon^d}{\eta^d} \sum_{\substack{\xi \in \mathbb{Z}^d \\ |\varepsilon\xi| \leq \eta R}} a\left(\frac{\varepsilon}{\eta}\xi\right) \left| \left\langle \frac{\varepsilon}{\eta}\xi, \nu_k \right\rangle \right| + O(\eta R) \right) \\ &\leq \limsup_{\varepsilon \rightarrow 0} \sum_{k=1}^N \mathcal{H}^{d-1}(\Sigma_k) \int_{\{|\xi| \leq R\}} a(\xi) |\langle \xi, \nu_k \rangle| d\xi \leq \int_{\partial A} \varphi_a(\nu) d\mathcal{H}^{d-1}. \end{aligned}$$

Since also for $|\varepsilon\xi| \geq \eta R$ the changes of value of u_i^ε are at most N , we then get

$$\limsup_{\varepsilon \rightarrow 0} \frac{\varepsilon^{2d}}{\eta^{d+1}} \sum_{\substack{\xi \in \mathbb{Z}^d \\ |\varepsilon\xi| > \eta R}} a\left(\frac{\varepsilon}{\eta}\xi\right) \sum_{i \in \mathbb{Z}^d} |u_{i+\xi}^\varepsilon - u_i^\varepsilon| \leq N \mathcal{H}^{d-1}(\partial A) \int_{\{|\xi| > R\}} a(\xi) |\xi| d\xi.$$

Since this term vanishes as $R \rightarrow +\infty$, the upper bound follows. □

Example 6. If a is radially symmetric, then we have

$$F(A) = \sigma \mathcal{H}^{d-1}(\partial^* A),$$

where σ is given by

$$\sigma = \int_{\mathbb{R}^d} a(\xi) |\xi_1| d\xi.$$

In particular, we may take $a = \chi_{B_1}$ the characteristic function of the unit ball in \mathbb{R}^d . In this case, the limit of

$$F_\varepsilon(u) = \frac{\varepsilon^{2d}}{\eta^{d+1}} \sum_{i,j \in \mathbb{Z}^d, |i-j| < \eta/\varepsilon} |u_i - u_j|$$

is given by

$$\sigma = \int_{B_1} |\xi_1| d\xi.$$

Remark 7 (Local Version). If $\Omega \subset \mathbb{R}^d$ is an open set with Lipschitz boundary, we may define

$$F_\varepsilon(u) = \frac{\varepsilon^{2d}}{\eta^{d+1}} \sum_{i,j \in \mathbb{Z}^d \cap \frac{1}{\varepsilon}\Omega} a_{i-j}^\varepsilon |u_i - u_j|.$$

Then the Γ -limit is

$$F(A) = \int_{\Omega \cap \partial^* A} \varphi_a(\nu) d\mathcal{H}^{d-1},$$

with minor modifications in the proof.

Acknowledgment: We thank the anonymous referee of [8], who drew our attention to the problem in this paper.

Funding: Andrea Braides acknowledges the MIUR Excellence Department Project awarded to the Department of Mathematics, University of Rome Tor Vergata, CUP E83C18000100006. Margherita Solci acknowledges the project “Fondo di Ateneo per la ricerca 2019”, funded by the University of Sassari.

References

- [1] R. Alicandro, A. Braides and M. Cicalese, Phase and anti-phase boundaries in binary discrete systems: A variational viewpoint, *Netw. Heterog. Media* **1** (2006), no. 1, 85–107.
- [2] R. Alicandro and M. S. Gelli, Local and nonlocal continuum limits of Ising-type energies for spin systems, *SIAM J. Math. Anal.* **48** (2016), no. 2, 895–931.
- [3] I. Benjamini and O. Schramm, Recurrence of distributional limits of finite planar graphs, *Electron. J. Probab.* **6** (2001), Paper No. 23.
- [4] X. Blanc and C. Le Bris, Définition d'énergies d'interfaces à partir de modèles atomiques, *C. R. Math. Acad. Sci. Paris* **340** (2005), no. 7, 535–540.
- [5] C. Borgs, J. T. Chayes, L. Lovász, V. T. Sós and K. Vesztegombi, Convergent sequences of dense graphs II. Multiway cuts and statistical physics, *Ann. of Math. (2)* **176** (2012), no. 1, 151–219.
- [6] A. Braides, *Approximation of Free-discontinuity Problems*, Lecture Notes in Math. 1694, Springer, Berlin, 1998.
- [7] A. Braides, *Γ -convergence for Beginners*, Oxford Lecture Ser. Math. Appl. 22, Oxford University, Oxford, 2002.
- [8] A. Braides, A. Causin and M. Solci, Asymptotic analysis of a ferromagnetic Ising system with “diffuse” interfacial energy, *Ann. Mat. Pura Appl. (4)* **197** (2018), no. 2, 583–604.
- [9] A. Braides, P. Cermelli and S. Dovetta, Γ -limit of the cut functional on dense graph sequences, *ESAIM Control Optim. Calc. Var.* **26** (2020), Paper No. 26.
- [10] A. Braides, V. Chiadò Piat and M. Solci, Discrete double-porosity models for spin systems, *Math. Mech. Complex Syst.* **4** (2016), no. 1, 79–102.
- [11] A. Braides and M. S. Gelli, Limits of discrete systems with long-range interactions, *J. Convex Anal.* **9** (2002), no. 2, 363–399.
- [12] A. Braides and L. Kreutz, An integral-representation result for continuum limits of discrete energies with multibody interactions, *SIAM J. Math. Anal.* **50** (2018), no. 2, 1485–1520.
- [13] A. Braides, M. Maslennikov and L. Sigalotti, Homogenization by blow-up, *Appl. Anal.* **87** (2008), no. 12, 1341–1356.
- [14] A. Braides and A. Piatnitski, Homogenization of surface and length energies for spin systems, *J. Funct. Anal.* **264** (2013), no. 6, 1296–1328.
- [15] A. Braides and M. Solci, Interfacial energies on Penrose lattices, *Math. Models Methods Appl. Sci.* **21** (2011), no. 5, 1193–1210.
- [16] L. A. Caffarelli and R. de la Llave, Planelike minimizers in periodic media, *Comm. Pure Appl. Math.* **54** (2001), no. 12, 1403–1441.
- [17] A. Chambolle, Finite-differences discretizations of the Mumford-Shah functional, *M2AN Math. Model. Numer. Anal.* **33** (1999), no. 2, 261–288.
- [18] I. Fonseca and S. Müller, Quasi-convex integrands and lower semicontinuity in L^1 , *SIAM J. Math. Anal.* **23** (1992), no. 5, 1081–1098.
- [19] N. García Trillos and D. Slepčev, Continuum limit of total variation on point clouds, *Arch. Ration. Mech. Anal.* **220** (2016), no. 1, 193–241.
- [20] L. Lovász, *Large Networks and Graph Limits*, Amer. Math. Soc. Colloq. Publ. 60, American Mathematical Society, Providence, 2012.
- [21] L. Lovász and B. Szegedy, Limits of dense graph sequences, *J. Combin. Theory Ser. B* **96** (2006), no. 6, 933–957.
- [22] F. Maggi, *Sets of Finite Perimeter and Geometric Variational Problems. An Introduction to Geometric Measure Theory*, Cambridge Stud. Adv. Math. 135, Cambridge University Press, Cambridge, 2012.
- [23] S. Neukamm, M. Schäffner and A. Schlömerkemper, Stochastic homogenization of nonconvex discrete energies with degenerate growth, *SIAM J. Math. Anal.* **49** (2017), no. 3, 1761–1809.