



SCUOLA INTERNAZIONALE SUPERIORE DI STUDI AVANZATI

SISSA Digital Library

Unstable blowups

Original

Unstable blowups / Stoppa, Jacopo. - In: JOURNAL OF ALGEBRAIC GEOMETRY. - ISSN 1056-3911. - 19:1(2010), pp. 1-17. [10.1090/S1056-3911-08-00503-1]

Availability:

This version is available at: 20.500.11767/16650 since:

Publisher:

Published

DOI:10.1090/S1056-3911-08-00503-1

Terms of use:

Testo definito dall'ateneo relativo alle clausole di concessione d'uso

Publisher copyright

AMS - American Mathematical Society

This version is available for education and non-commercial purposes.

note finali coverpage

(Article begins on next page)

Unstable blowups

Jacopo Stoppa

Abstract

Let (X, L) be a polarised manifold. We show that K-stability and asymptotic Chow stability of the blowup of X along a 0-dimensional cycle are closely related to Chow stability of the cycle itself, for polarisations making the exceptional divisors small. This can be used to give (almost) a converse to the results of Arezzo and Pacard in [1], [2] and to give new examples of Kähler classes with no constant scalar curvature representatives.

1 Introduction

The theme of this paper is to construct and study *test configurations* (i.e. particular degenerations) for blowups of a polarised manifold. More precisely we compute the *Donaldson-Futaki invariant* of these configurations (Theorem 1.3). These concepts are recalled in sections 2 and 4 respectively, but in essence the Donaldson-Futaki invariant is a rational number attached to our degeneration which morally plays the role of the Hilbert-Mumford weight in Geometric Invariant Theory. The main application is to nonexistence of constant scalar curvature Kähler (cscK for brevity) metrics contained in particular Kähler classes (1.9). The search for such manifolds has played an important role in the development of the theory (see [15] for more details) and the method presented here yields infinitely many (which we can take to be rational, see 1.10).

A more algebraic application is to asymptotically Chow unstable polarisations, see 1.4.

Test configurations for a polarised manifold (M, L) together with generalised Futaki invariants were introduced by Donaldson in [8] following work of Tian [19]. Let $S(\omega)$ denote the scalar curvature of a Kähler metric $\omega \in c_1(L)$ with average $\widehat{S} = 2\pi n \frac{c_1(M) \cup c_1(L)^{n-1}}{c_1(L)^n}$. In [9] Donaldson proves that the *Calabi functional*

$$Ca(\omega) = \int_M (S(\omega) - \widehat{S})^2 \frac{\omega^n}{n!}$$

is bounded below in all of $c_1(L)$ by the negative of the generalised Futaki invariant $F(\mathcal{M})$ (divided by a positive term) for *any* test configuration $(\mathcal{M}, \mathcal{L})$ relative to (M, L) . Thus if $c_1(L)$ contains a cscK metric, M must be *K-semistable* with respect to L , meaning precisely $F(\mathcal{M}) \geq 0$ for all $(\mathcal{M}, \mathcal{L})$. Moreover it is expected that $F(\mathcal{M}) = 0$ if and only if \mathcal{M} is isomorphic to the product $M \times \mathbb{C}$, i.e. a cscK manifold should be *K-polystable*. That the converse holds as well is the content of the conjectural Hitchin-Kobayashi correspondence for manifolds (or Yau-Tian-Donaldson conjecture).

Now to our case. Let X be a compact connected complex manifold, $\dim(X) = n$. We assume that X is polarised, that is we fix an ample line bundle $L \rightarrow X$. Then for any subscheme $Z \subset X$, the blowup $\text{Bl}_Z X$ with exceptional divisor E is endowed with the line bundle $L^\gamma - E$ which is ample for all large enough positive integers γ . Suppose now that $\text{Aut}(X)$ contains a nontrivial compact connected subgroup, so that there is a 1-parameter subgroup (1-PS) of automorphisms of X (i.e. a group homomorphism $\alpha : \mathbb{C}^* \hookrightarrow \text{Aut}(X)$). Then taking the limit of the action of α on X as $t \rightarrow 0$ induces in a natural way a test configuration \mathcal{X} for $(\text{Bl}_Z X, L^\gamma - E)$ intuitively by making the components of E move around and possibly collide. In general one would like to make this rigorous and to understand this configuration as much as is needed to compute the first terms of the asymptotic expansion of its Futaki invariant $F(\mathcal{X})$ as $\gamma \rightarrow \infty$.

Remark 1.1 It is important to note that we cannot expect the central fibre \mathcal{X}_0 to be the blowup of X along the limit Z_0 of Z as $t \rightarrow 0$ in the relevant Hilbert scheme. This is explained at the end of section 2. In general we can only say that $\text{Bl}_{Z_0} X$ is an irreducible component of \mathcal{X}_0 .

Our main result is that we can carry out this program completely when Z is a 0-dimensional cycle, say

$$Z = \sum_i a_i p_i \quad (p_i \in X, a_i > 0).$$

Remark 1.2 We emphasise that by blowing up $a_i p_i$ we mean blowing up the ideal $\mathcal{I}_{p_i}^{a_i}$, so that the *algebraic* multiplicity of p_i is the length of $\mathcal{O}_{a_i p_i}$:

$$\text{len}(\mathcal{O}_{a_i p_i}) = \binom{n + a_i - 1}{a_i - 1}$$

while a_i is the multiplicity of p_i in the cycle. Note that by [10], II Exercise 7.11, there is an isomorphism of polarised schemes

$$(\text{Bl}_Z X, L^\gamma - E) \cong (\text{Bl}_{\{p_i\}} X, L^\gamma - \sum_i a_i E_i)$$

where E_i is the component of the exceptional divisor over p_i .

In particular, we will relate $F(\mathcal{X})$ to two fundamental weights associated to the action of α on X : on the one hand, the classical Futaki invariant $F(X)$ for the holomorphic vector field which generates α , on the other, the natural GIT weight for the action of α on the Chow variety of 0-cycles of total multiplicity $m = \sum_i a_i^{n-1}$ on (X, L^γ) (i.e. the symmetric product $X^{(m)}$ with polarisation induced by $(L^\gamma)^{\boxtimes m}$). We denote this weight by $\mathcal{CH}(\sum_i a_i^{n-1} p_i, \alpha)$. The few GIT notions we need (including Chow stability for 0-cycles) are recalled in section 3. The following results are proved in section 4.

Theorem 1.3

$$F(\mathcal{X}) = F(X) \gamma^n - \mathcal{CH}(\sum_i a_i^{n-1} p_i, \alpha) \frac{\gamma}{2(n-2)!} + O(1).$$

As a consequence if (X, L) is K -polystable, the blowup of X along a Chow-unstable 0-cycle is K -unstable when the exceptional divisors are small enough (i.e. $\gamma \gg 0$).

Corollary 1.4 *If (X, L) is asymptotically Chow polystable its blowup along a Chow unstable 0-cycle is asymptotically Chow unstable, when the exceptional divisors are small enough.*

Remark 1.5 Thus when we blow up the cycle $\sum_i a_i p_i$ it is the GIT stability of the cycle $\sum_i a_i^{n-1} p_i$ that naturally shows up in $F(\mathcal{X})$. To interpret this difference note that the volume of the weighted exceptional divisor $a_i E_i$ over p_i with respect to $L^\gamma - E$ is $\gamma^{1-n} a_i^{n-1}$ (up to a dimensional constant). Then an example in [17] 27–28 illustrating Donaldson’s theory of balanced metrics from [7] suggests that when blowing up we are perturbing the centre of mass of X in $\mathfrak{su}(H_X^0(L^{\gamma r})^*)$ (for $r, \gamma \gg 0$) by attaching a small weight proportional to $\gamma^{1-n} a_i^{n-1}$ over the point p_i .

We must mention at this point that our interest in this topic came from trying to find an algebro-geometric counterpart to the results of Arezzo and Pacard on blowing up and desingularizing cscK metrics contained in [1], [2]. As blowing up is such a fundamental tool it is not surprising that the Arezzo-Pacard theorem plays a key role in the proofs of many recent results in the field. Among them we recall a theorem of Shu [20] stating that any compact complex surface with $b_1 = 0$ (except the blowup of \mathbb{P}^2 in 1 or 2 points) is deformation equivalent to one bearing a cscK metric and the construction of Einstein metrics which are conformally Kähler on \mathbb{P}^2 blown up in 2 points due to Chen-LeBrun-Weber [5]. By analogy it seems that understanding the

behaviour of the Donaldson-Futaki invariant under blowing up may be useful for the algebraic side of the story.

Working in the Kähler setting, Arezzo and Pacard prove that if ω is cscK on X , and if the cycle $\sum_i a_i p_i$ ($a_i \in \mathbb{R}^+$) satisfies the three conditions recalled below, then for all small positive ϵ the Kähler class on the blowup of X at $\{p_i\}$ given by

$$\pi^*[\omega] - \epsilon \left(\sum a_i [E_i] \right)$$

contains a cscK form ω_ϵ . Moreover ω_ϵ converges to ω in the C^∞ sense over $X - \sum_i p_i$. These conditions are expressed in terms of a moment map

$$\mu : X \rightarrow \mathfrak{ham}^*(X, J, \omega)$$

for the action of the Hamiltonian isometries of the Kähler manifold X , namely

1. $\{\mu(p_i)\}$ span $\mathfrak{ham}^*(X, J, \omega)$;
2. no nonzero element of $\mathfrak{ham}(X, J, \omega)$ vanishes at all the $\{p_i\}$;
3. $\sum_i a_i^{n-1} \mu(p_i) = 0$.

Remark 1.6 Condition 2 is only needed to get rid of residual automorphisms on the blowup, and without it the theorem continues to hold in a slightly modified form. It is also expected that 1 is not necessary, but rather a side effect of the analytic argument used in the proof (see [3] for more details). Thus 3 (which we may call a balanced- or stability- condition) seems to be heart of the matter. This fits in well with the results of this paper.

We are now going to recast the Arezzo-Pacard theorem in terms of Chow stability. This bears on the projective case when $\omega = c_1(L)$ and the a_i are positive integers. The Kempf-Ness theorem shows that the cycle $\sum_i a_i^{n-1} p_i$ can be modified by elements of $\text{Aut}(X)$ to satisfy condition 3 if and only if it is Chow polystable with respect to the action of $\text{Aut}(X)$ (for a moment we refer to this new cycle as the balanced image).

Remark 1.7 One must be careful in applying the Kempf-Ness theorem here since the relevant symmetric product $X^{(m)}$ is a singular variety when $n > 1$, $m > 1$. However (see section 3) $X^{(m)}$ is defined as the geometric quotient X^m / Σ_m . Also the polarisation on $X^{(m)}$ is induced by $(L^\gamma)^{\boxtimes m}$ on X^m . Since all the points of X^m are stable under the action of Σ_m , we can lift a cycle in $X^{(m)}$ to the product X^m , apply Kempf-Ness there, and project back.

In the polystable case we might leave $\sum_i a_i^{n-1} p_i$ fixed and pullback ω by some automorphism instead so that 3 holds. Note that we can restrict to the action of the connected component of the identity $\text{Aut}^0(X)$ in the Kempf-Ness theorem, and this preserves the Kähler class. This gives the following version of the theorem.

Theorem 1.8 (*Arezzo-Pacard, projective case*) *Suppose ω is cscK, the cycle $\sum_i a_i^{n-1} p_i$ is Chow polystable and conditions 1, 2 hold on its balanced image. Then for all rational $\epsilon > 0$ small enough, the class*

$$\pi^*[\omega] - \epsilon \left(\sum_i a_i E_i \right)$$

contains a cscK metric ω_ϵ . Moreover there exists $\phi \in \text{Aut}^0(X)$ such that ω_ϵ converges to ϕ^ω in the C^∞ sense over $X - \sum_i p_i$.*

Now observe that when ω is cscK F vanishes identically on $\mathfrak{Lie} \text{Aut}(X)$. By the Hilbert-Mumford criterion we also know that $\sum_i a_i^{n-1} p_i$ is semistable if and only if $\mathcal{CH}(\sum_i a_i^{n-1} p_i, \alpha) \leq 0$ for all 1-PS $\alpha \hookrightarrow \text{Aut}(X)$, so Theorem 1.3 almost implies that condition 3 is necessary. A small discrepancy comes from the fact that *unstable* means *not semistable* which is stronger than *not polystable*. More precisely, our asymptotic expansion 1.3 together with Donaldson's lower bound on the Calabi functional immediately give

Theorem 1.9 *If ω is cscK and the cycle $\sum_i a_i^{n-1} p_i$ is Chow unstable then for all rational $\epsilon > 0$ small enough, the class*

$$\pi^*[\omega] - \epsilon \left(\sum_i a_i E_i \right)$$

does not contain a cscK metric.

Example 1.10 *Projective space.* There is a very nice geometric criterion for Chow stability of points in \mathbb{P}^n (see for example [13] 231–235). A cycle $Z = \sum_i m_i p_i$ is Chow *unstable* if and only if for some proper subspace $V \subset \mathbb{P}^n$ one has

$$\frac{|V \cap Z|}{\dim(V) + 1} > \frac{\sum_i m_i}{n + 1}.$$

So already in the case of \mathbb{P}^2 1.3 gives infinitely many new examples of K-unstable classes: it suffices that more than 2/3 of the points (counted with multiplicities) are aligned to get a K-unstable blowup. This gives a full generalisation of Example 5.30 in [16] where it is shown that when $m_1 \gg$

m_j ($j > 2$) the blowup is K-unstable (actually *slope unstable* with respect to E_1).

The following examples are straightforward applications of the theory in section 3.

Example 1.11 *Products.* Consider the product $X \times Y$ of (X, L_X) and (Y, L_Y) polarised by $L_X \boxtimes L_Y$. Then for $\alpha : \mathbb{C}^* \hookrightarrow \text{Aut}(X)$,

$$\mathcal{CH}(\sum_i a_i p_i, \alpha \times 1) = \mathcal{CH}(\sum_i a_i \pi_X(p_i), \alpha).$$

Thus we may apply the above geometric criterion to the fibres of a product $\mathbb{P}^n \times Y$. For example if Y is K-polystable the blowup of $\mathbb{P}^n \times Y$ along an unstable cycle supported at a single \mathbb{P}^n -fibre will be K-unstable. A special case is the product $\mathbb{P}^n \times \mathbb{P}^m$ polarised by $\mathcal{O}(1) \boxtimes \mathcal{O}(1)$. A 0-cycle will be unstable whenever its projection to one of the two factors is, e.g. in the case of 3 distinct point, when 2 of them lie on a vertical or horizontal fibre. This gives more examples of unstable blowups.

Example 1.12 \mathbb{P}^1 *bundles.* Similarly we can consider the projective completion X of some line bundle L over a polarised manifold. In this case the so-called momentum construction yields many examples of cscK metrics, see [11]. Any polarisation \mathcal{L} on X restricts to $\mathcal{O}_{\mathbb{P}^1}(k)$ on all the fibres for some k . There is a natural \mathbb{C}^* -action on L given by complex multiplication on the fibres, and this extends to a \mathbb{C}^* -action α on X so that points lying on the zero (resp. infinity) section X_0 (X_∞) are fixed, but with weight k (resp. $-k$) on the line above them. By acting with α^{-1} instead if necessary, we conclude that whenever more than half the points lie on X_0 or X_∞ , the corresponding 0-cycle is Chow unstable (i.e. its Chow weight is $> ck$ for some positive constant c). We see that the conclusion is really independent of k and so the blowup along such a cycle will be K- and asymptotically Chow- unstable for *any* polarisation on X making the base cscK.

Notation. We will often suppress pullback maps and use the same letter to denote a divisor and the associated line bundle. Consequently we mix additive and multiplicative notation as necessary.

Acknowledgements. This work has been made possible by the generosity and patience of my supervisor R. Thomas. The idea of this project is due to him and S. K. Donaldson who I also thank for his kind interest. I would like to thank J. Ross, G. Székelyhidi and the Referee for many important suggestions. Finally I am grateful to the geometry groups at Imperial and

Pavia and in particular to G. P. Pirola and A. Ghigi for their support and for teaching me so much.

2 Test configurations coming from automorphisms of the base

Let $Z = \sum_i a_i p_i$ be some 0-dimensional cycle on X , that is the closed subscheme supported at the points $\{p_i\}$ with nonreduced structure $\prod_i \mathcal{I}_{p_i}^{a_i}$. In this section we construct a *test configuration* for $(\mathrm{Bl}_Z X, L^\gamma - E)$ naturally associated to a 1-PS $\alpha \hookrightarrow \mathrm{Aut}(X)$. A test configuration for a polarised manifold (M, L) is given by a polarised flat family $(\mathcal{M}, \mathcal{L}) \rightarrow \mathbb{C}$ endowed with an \mathcal{L} -linearised \mathbb{C}^* -action covering the usual action of \mathbb{C}^* on \mathbb{C} , and such that for any $t \neq 0$, $(\mathcal{M}_t, \mathcal{L}|_t) \cong (M, L^s)$ for some exponent s (called the exponent of the test configuration). In our case this is the natural flat family induced by the \mathbb{C}^* -action on X and so on cycles and exceptional divisors. It will be useful to introduce the flat family of closed subschemes of X given by $\{Z_t = \alpha(t)Z, t \in \mathbb{C}^*\}$.

Lemma 2.1 *There is a flat family $p : \mathcal{X} \rightarrow \mathbb{C}$ such that $\mathcal{X}_t \cong \mathrm{Bl}_{Z_t} X$ for all $t \in \mathbb{C} - \{0\}$. This is endowed with an induced action of α covering the usual \mathbb{C}^* action on \mathbb{C} .*

Proof. We see Z as a point of the Hilbert scheme of closed subschemes of X with *constant* Hilbert polynomial equal to the length of \mathcal{O}_Z as a module over itself. From the general theory we know that the flat family $(Z_t, t) \subset X \times (\mathbb{C} - \{0\})$ has a unique flat closure i.e. there exists a unique closed subscheme $Y \subset X \times \mathbb{C}$, flat over \mathbb{C} , with fibres $Y_t = Z_t$ for all $t \in \mathbb{C} - \{0\}$. As total scheme of our test configuration we take $\mathcal{X} = \mathrm{Bl}_Y(X \times \mathbb{C})$, with projection $p : \mathcal{X} \rightarrow \mathbb{C}$ given by the composition of the blowup map $\pi : \mathcal{X} \rightarrow X \times \mathbb{C}$ with the projection onto the second factor. By [10], II Proposition 7.16 \mathcal{X} is reduced, irreducible, and p is a dominant (in fact surjective) morphism. The base \mathbb{C} is regular, 1-dimensional and dominated by every irreducible component of \mathcal{X} so by [10], III Proposition 9.7 we see that p is a flat morphism. Since Y is preserved by α there is an induced action of α on \mathcal{X} covering the usual action of \mathbb{C}^* on \mathbb{C} . □

Since $\mathcal{X} = \mathrm{Proj}(\bigoplus_r \mathcal{I}_Y^r)$, it is naturally endowed with an invertible sheaf $\mathcal{O}(1)$. Let $p_X : \mathcal{X} \rightarrow X$ be the composition $\mathcal{X} \rightarrow X \times \mathbb{C} \rightarrow X$. Define a line bundle on \mathcal{X} by $\mathcal{L} = p_X^* L^\gamma \otimes \mathcal{O}(1)$. A slight modification of the above argument then proves

Lemma 2.2 *For all large γ , \mathcal{L} is a α -linearised ample line bundle \mathcal{L} on \mathcal{X} such that $\mathcal{L}|_t \cong L^\gamma - E$ for $t \neq 0$.*

Next we need to study the central fibre \mathcal{X}_0 of our test configuration. It will be useful to write Z_0 for the limit of Z_t in the Hilbert scheme as $t \rightarrow 0$ as in the proof of 2.1, and to define

$$\widehat{X} = \mathrm{Bl}_{Z_0} X$$

and

$$E_0 = \text{the exceptional divisor of } \widehat{X} \rightarrow X.$$

We claim that there is a closed immersion $\widehat{X} \hookrightarrow \mathcal{X}_0$. To see this, consider the closed immersion $i : X \cong X \times \{0\} \hookrightarrow X \times \mathbb{C}$. By flatness we see that

Lemma 2.3 *The inverse image ideal sheaf $i^{-1}\mathcal{I}_Y \cdot \mathcal{O}_X$ is \mathcal{I}_{Z_0} .*

Lemma 2.4 *The inclusion $i : X \times 0 \hookrightarrow X \times \mathbb{C}$ induces a closed immersion*

$$\widehat{i} : \widehat{X} \hookrightarrow \mathcal{X}_0.$$

Proof. By [10], II Corollary 7.15, there is an induced closed immersion

$$\widehat{i} : \mathrm{Bl}_{i^{-1}\mathcal{I}_Y \cdot \mathcal{O}_X} X \hookrightarrow \mathrm{Bl}_Y(X \times \mathbb{C}).$$

Since the image lies in \mathcal{X}_0 we conclude by 2.3. □

Next we define a closed subscheme P of \mathcal{X}_0 by

$$P = (\mathcal{X}_0 - \widehat{X})^-.$$

In general \widehat{i} is not an isomorphism as the exceptional set P may well be a component of \mathcal{X}_0 . To motivate this recall that Z_0 is the central fibre of the flat family $\{Z_t, t \in \mathbb{C}\}$. But a family of thickenings $\{rZ_t, t \in \mathbb{C}\}$ will *not* be flat in general. The generic fibre of \mathcal{X} is $\mathcal{X}_t = \mathrm{Proj}(\bigoplus_r \mathcal{I}_{rZ_t})$ for $t \neq 0$. As \mathcal{X} itself is flat we see that \mathcal{X}_0 cannot in general be $\mathrm{Proj}(\bigoplus_r \mathcal{I}_{rZ_0})$. This means there is some extra closed subscheme P inside \mathcal{X}_0 , which we will have to take into account in section 4 when computing the Futaki invariant.

Example 2.5 As an affine example consider the ideal

$$\mathcal{I}_Y = (x(x-t), xy, y(y-t)) \subset \mathbb{C}[x, y, t]$$

which describes 3 points colliding along 2 orthogonal directions in \mathbb{A}^2 . One can show that \mathcal{X} is the closed subscheme of $\text{Spec } \mathbb{C}[x, y, t] \times \text{Proj } \mathbb{C}[\xi_0, \xi_1, \xi_2]$ defined by

$$\mathcal{I}_{\mathcal{X}} = ((x - t)\xi_1 - y\xi_0, (y - t)\xi_1 - x\xi_2)$$

with central fibre (sitting inside $\text{Spec } \mathbb{C}[x, y] \times \text{Proj } \mathbb{C}[\xi_0, \xi_1, \xi_2]$)

$$\mathcal{I}_{\mathcal{X}_0} = (x\xi_1 - y\xi_0, y\xi_1 - x\xi_2).$$

In this case \widehat{X} is the closed subscheme of \mathcal{X}_0 given by

$$\mathcal{I}_{\widehat{X}} = (x\xi_1 - y\xi_0, y\xi_1 - x\xi_2, \xi_1^2 - \xi_1\xi_2)$$

that is a whole component of \mathcal{X}_0 , while the exceptional component P is $\text{Proj } \mathbb{C}[\xi_0, \xi_1, \xi_2] \cong \mathbb{P}^2$.

In any case the restriction of \mathcal{L}_0 to \widehat{X} is the expected one.

Lemma 2.6

$$\mathcal{L}_{0|\widehat{X}} = L^\gamma - E_0.$$

The easy proof is left to the reader.

3 Chow stability for 0-dimensional cycles

In this section we recall the few GIT notions we need in the case of the Chow variety of points on X . For much more on this see [14] Chap. 3. Let the symmetric group Σ_d on d letters acts on the d -fold product X^d . The symmetric product $X^{(d)} = X^d/\Sigma_d$ is a projective variety. The points of $X^{(d)}$ are the orbits of the d -tuples of points of X under permutation and so can be identified with effective 0-cycles $\sum n_i[x_i]$ with $x_i \in X$, $n_i > 0$ and $\sum n_i = d$. This shows that $X^{(d)}$ is actually the Chow variety of length d 0-cycles on X . The construction of an $\text{Aut}(X)$ -linearised ample line on $X^{(d)}$ can be made very explicit as follows. Let $V = H^0(X, L^\gamma)^*$ for some large γ and embed $X \hookrightarrow \mathbb{P}(V)$. Denote by $\mathbb{P}(V^*)$ the projective space of hyperplanes in $\mathbb{P}(V)$, and by $\text{Div}^d(\mathbb{P}(V^*))$ the projective space of effective divisors of degree d in $\mathbb{P}(V^*)$. For any $p \in \mathbb{P}(V)$ consider the hyperplane in $\mathbb{P}(V^*)$ given by

$$H_p := \{l \in \mathbb{P}(V^*) : p \in l\}.$$

Define a morphism $ch : (\mathbb{P}(V))^d \rightarrow \text{Div}^d(\mathbb{P}(V^*))$ by

$$ch(x_1, \dots, x_d) := \sum_i H_{x_i}.$$

This is the *Chow form* of $\{x_1, \dots, x_d\}$: the divisor of hyperplanes whose intersection with $\{x_1, \dots, x_d\}$ is nonempty. As for X^d we have the composition

$$ch : X^d \hookrightarrow \mathbb{P}(V)^d \hookrightarrow \text{Div}^d(\mathbb{P}(V^*))$$

induced by the product line $(L^\gamma)^{\boxtimes d}$. Now ch is Σ_d equivariant and by the universal property of the geometric quotient factors through $\mathbb{P}(V)^{(d)}$ defining a morphism

$$ch : X^{(d)} \rightarrow \text{Div}^d(\mathbb{P}(V^*)).$$

One can check that ch defines an isomorphism on its image. We can thus identify $X^{(d)}$ with its image $ch(X^{(d)})$ in $\text{Div}^d(\mathbb{P}(V^*))$. By the usual identifications

$$\text{Div}^d(\mathbb{P}(V^*)) \cong \mathbb{P}(H^0(\mathbb{P}(V^*), \mathcal{O}(d))) \cong \mathbb{P}(S^d V)$$

we see ch as a map with values in $\mathbb{P}(S^d V)$:

$$ch : X^{(d)} \hookrightarrow \mathbb{P}(S^d V).$$

Under these identifications then ch is the map

$$X^{(d)} \ni \{[x_1], \dots, [x_d]\} \mapsto [x_1 \cdot \dots \cdot x_d] \quad (3.1)$$

defined via the embedding $X \hookrightarrow \mathbb{P}(V)$.

Remark 3.1 It is important to emphasise that ch is given by the descent of $(L^\gamma)^{\boxtimes d}$ under the action of Σ_d . This means that the Chow line $\mathcal{O}_{\text{Div}^d(\mathbb{P}(V^*))}(1)|_{X^{(d)}}$ pulls back to $(L^\gamma)^{\boxtimes d}$ under the quotient map. This holds because by 3.1 $\mathcal{O}_{\text{Div}^d(\mathbb{P}(V^*))}(1)$ pulls back to the line $\mathcal{O}_{\mathbb{P}(V)}(1)^{\boxtimes d}$ on $\mathbb{P}(V)^d$ under ch and this in turn pulls back to $(L^\gamma)^{\boxtimes d}$.

Now we assume that $\alpha \hookrightarrow \text{Aut}(X)$ acts through a 1-PS $\alpha \hookrightarrow \text{Sl}(V)$ and we come to the definition of the GIT weight for the action of α on X . Recall that we can find a basis of eigenvectors so that $\alpha(t)$ acts as $\text{diag}(t^{\lambda_0}, \dots, t^{\lambda_N})$, where $\dim(V) = N + 1$. In these projective coordinates on $\mathbb{P}(V)$, writing $X \ni x = [v_0 : \dots : v_n]$, we define the Mumford weight as

$$\lambda(x, \alpha) := \min\{\lambda_i : v_i \neq 0\}.$$

By definition of V this is the weight of the induced action on the line $(L^\gamma)^*$ over the limit x_0 of $\alpha(t)x$ as $t \rightarrow 0$ in the \mathbb{C}^* -action.

In the same way we can define the Mumford weight for the induced action of α on $X^{(d)}$. In fact $\alpha \hookrightarrow \text{Sl}(V)$ naturally induces a 1-PS $\alpha \hookrightarrow \text{Sl}(S^d V)$. Thus the embedding ch described above gives a natural linearisation for this

action. We write \mathcal{CH} for this Mumford weight. Then by (3.1) we immediately obtain the relation

$$\mathcal{CH}(\sum_i m_i x_i, \alpha) = \sum_i m_i \lambda(x_i, \alpha)$$

for any $\sum_i m_i x_i \in X^{(d)}$.

Remark 3.2 While the numerical value of the Mumford weight depends on the power L^γ we take, the fact that a cycle $\sum_i m_i p_i$ is semistable is independent of γ . This is immediate from the definition of stability in terms of invariant sections.

4 The Donaldson-Futaki invariant

In this section we compute the Donaldson-Futaki invariant of the induced \mathbb{C}^* action on the central fibre \mathcal{X}_0 . But first we recall Donaldson's definition of the Futaki invariant of a \mathbb{C}^* -action on a variety (or scheme) M of dimension n endowed with a linearised ample line bundle L . We write A_k for the infinitesimal generator of the induced \mathbb{C}^* -action on $H^0(M, L^k)$.

Remark 4.1 The lifting of the \mathbb{C}^* -action to L is *not* unique, so A_1 is not well defined. However for any other lifting, there is $\lambda \in \mathbb{Z}$ such that

$$A'_1 = A_1 + \lambda I_1.$$

where I_1 denotes the identity matrix on $H^0(M, L)$. As a consequence,

$$A'_k = A_k + k\lambda I_k$$

where I_k denotes the identity matrix on $H^0(M, L^k)$. Using this transformation rule one can check that the Futaki invariant defined below is independent of the choice of lifting.

By Riemann-Roch and its equivariant version there are expansions

$$h^0(M, L^k) = c_0 k^n + c_1 k^{n-1} + O(k^{n-2}),$$

$$\mathrm{tr}(A_k) = b_0 k^{n+1} + b_1 k^n + O(k^{n-1})$$

valid for all large k . The Futaki invariant is defined as

$$F = \frac{c_1 b_0}{c_0} - b_1. \tag{4.1}$$

According to this general definition, in our case we need to compute $h^0(\mathcal{X}_0, \mathcal{L}_0^r)$ and the trace of the induced action on $H^0(\mathcal{X}_0, \mathcal{L}_0^r)$ for all large r . It is important to keep in mind that \mathcal{L}_0 depends in turn on the parameter γ ; i.e. \mathcal{L}_0 comes from picking the line $L^\gamma - E$ on the generic fibre.

Remark 4.2 We emphasise that with this choice of notation γ^{-1} measures the volume of the exceptional divisors on the generic fibre, while r is the scale parameter needed to compute the Futaki invariant. Also in the proof of the following lemma and in many other places below, we need the vanishing of higher cohomology groups. This will always hold for $r \geq r_0 = r_0(\gamma)$, however increasing γ only makes \mathcal{L} (and other related line bundles) more positive, so at least for $\gamma \geq \gamma_0$ we can assume that r_0 is a fixed constant.

Lemma 4.3

$$h^0(\mathcal{X}_0, \mathcal{L}_0^r) = h^0(X, L^{\gamma r}) - \left(\sum_i a_i^n \right) \frac{r^n}{n!} - \left(\sum_i a_i^{n-1} \right) \frac{r^{n-1}}{2(n-2)!} + O(r^{n-2}).$$

Proof. By flatness $h^0(\mathcal{X}_0, \mathcal{L}_0^r) = h^0(\text{Bl}_Z X, L^{\gamma r} - rE)$. To compute the coefficients use the asymptotic Riemann-Roch formula, keeping Remark 1.2 in mind. □

Now for the trace. To try to keep the notation light in what follows we will write $\text{tr}(U)$ for the trace of the induced action on some vector space U . We start with the restriction \mathbb{C}^* -equivariant exact sequence

$$0 \rightarrow H_P^0(\mathcal{I}_{E_0}^r \mathcal{L}_{0|P}^r) \rightarrow H_{\mathcal{X}_0}^0(\mathcal{L}_0^r) \rightarrow H_{\widehat{X}}^0(L^{\gamma r} - rE_0) \rightarrow 0 \quad (4.2)$$

which holds for large r . So we see

$$\text{tr}(H_{\mathcal{X}_0}^0(\mathcal{L}_0^r)) = \text{tr}(H_{\widehat{X}}^0(L^{\gamma r} - rE_0)) + \text{tr}(H_P^0(\mathcal{I}_{E_0}^r \mathcal{L}_{0|P}^r)). \quad (4.3)$$

To compute the first term we turn to the natural isomorphism (for $r \gg 0$)

$$H_{\widehat{X}}^0(L^{\gamma r} - rE_0) \cong H_X^0(\mathcal{I}_{Z_0}^r L^{\gamma r}).$$

The exact sheaf sequence on X

$$0 \rightarrow \mathcal{I}_{Z_0}^r L^{\gamma r} \rightarrow L^{\gamma r} \rightarrow \mathcal{O}_{rZ_0} \otimes_{\mathbb{C}} L^{\gamma r}|_{Z_0} \rightarrow 0 \quad (4.4)$$

is \mathbb{C}^* -equivariant and gives an exact sequence of sections (for large r):

$$0 \rightarrow H_X^0(\mathcal{I}_{Z_0}^r L^{\gamma r}) \rightarrow H_X^0(L^{\gamma r}) \rightarrow \mathcal{O}_{rZ_0} \otimes_{\mathbb{C}} L^{\gamma r}|_{Z_0} \rightarrow 0. \quad (4.5)$$

Here we used that \mathcal{O}_{rZ_0} is a skyscraper sheaf supported at Z_0 . So we see

$$\mathrm{tr}(H_{\widehat{X}}^0(L^{\gamma r} - rE_0)) = \mathrm{tr}(H_X^0(L^{\gamma r})) - \sum_q \mathrm{tr}(\mathcal{O}_{rZ_0,q} \otimes L^{\gamma r}|_q) \quad (4.6)$$

where we are summing over the components of $(Z_0)_{\mathrm{red}}$. Substituting in (4.3) we get

$$\mathrm{tr}(H_{\mathcal{X}_0}^0(\mathcal{L}_0^r)) = \mathrm{tr}(H_X^0(L^{\gamma r})) - \sum_q \mathrm{tr}(\mathcal{O}_{rZ_0,q} \otimes L^{\gamma r}|_q) + \sum_q \mathrm{tr}(H_{P_q}^0(\mathcal{I}_{E_0,q}^r \mathcal{L}_{0|P_q}^r)) \quad (4.7)$$

where we write P_q for the component of P which projects to q via p_X and similarly for $E_{0,q}$.

Definition 4.4 For any $q \in (Z_0)_{\mathrm{red}}$, we denote by $\lambda(q)$ the weight of the induced \mathbb{C}^* -action on the line $L|_q$. Note that the weight of the induced action on $L^m|_q$ is then $m\lambda(q)$ for any $m > 0$. As we already observed $\lambda(q)$ depends on the choice of a lifting of α to L , but remember that this choice will not affect the Futaki invariant. For any other lifting the new weights are

$$\lambda'(q) = \lambda(q) + \lambda$$

for some $\lambda \in \mathbb{Z}$. These weights should not be confused with the relevant Chow weights, which require α to act through $\mathrm{Sl}(H^0(X, L^m))$ (for the relevant power m) in their definition. This difference will turn out to be important for our purposes.

The crucial step to get an asymptotic expansion for $F(\mathcal{X})$ is the following rough estimate, ignoring any term which is independent of γ .

Lemma 4.5

$$\mathrm{tr}(\mathcal{O}_{rZ_0,q} \otimes L^{r\gamma}|_q) = (r\gamma)\lambda(q)\dim(\mathcal{O}_{rZ_0,q}) + O(\gamma^0 r^{n+1}).$$

Proof. This is just the statement that the induced action on $\mathcal{O}_{rZ_0,q}$ as a \mathbb{C} -vector space does not depend on the parameter γ .

□

There is a similar estimate for the action on the components P_q of P .

Lemma 4.6

$$\mathrm{tr}(H_{P_q}^0(\mathcal{I}_{E_0,q}^r \mathcal{L}_{0|P_q}^r)) = (r\gamma)\lambda(q)h_{P_q}^0(\mathcal{I}_{E_0,q}^r \mathcal{L}_{0|P_q}^r) + O(\gamma^0 r^{n+1}).$$

Proof. Note that

$$\mathcal{I}_{E_0,q}^r \mathcal{L}_{0|P_q}^r \cong L^{\gamma^r}|_q \otimes_{\mathbb{C}} \mathcal{I}_{E_0,q}^r \mathcal{O}(r)|_{P_q}$$

and that $L|_q$ is the trivial line on P_q (since it is pulled back from Z_0) acted on by \mathbb{C}^* with weight $\lambda(q)$.

□

For the following lemma we introduce the sets

$$A_q = \{p_i \in Z_{\text{red}} : \lim_{t \rightarrow 0} \alpha(t)p_i = q\}.$$

Lemma 4.7

$$\dim(\mathcal{O}_{rZ_0,q}) = h_{P_q}^0(\mathcal{I}_{E_0,q}^r \mathcal{L}_{0|P_q}^r) + \left(\sum_{p_i \in A_q} a_i^n \right) \frac{r^n}{n!} + \left(\sum_{p_i \in A_q} a_i^{n-1} \right) \frac{r^{n-1}}{2(n-2)!} + O(r^{n-2}).$$

Proof. This follows from local versions of 4.3, (4.2) and (4.5) around q (in the analytic topology).

□

Putting these results together we can finally compute the trace on the central fibre.

Lemma 4.8

$$\begin{aligned} \text{tr}(H_{\mathcal{X}_0}^0(\mathcal{L}_0^r)) &= \text{tr}(H_X^0(L^{\gamma^r})) \\ &- \left(\gamma \sum_q \lambda(q) \left(\sum_{p_i \in A_q} a_i^n \right) \frac{r^{n+1}}{n!} + \gamma \sum_q \lambda(q) \left(\sum_{p_i \in A_q} a_i^{n-1} \right) \frac{r^n}{2(n-2)!} \right) \\ &+ O(\gamma^0 r^{n+1}). \end{aligned}$$

Proof. Substitute the results of lemma 4.5 and lemma 4.6 into (4.7) using 4.7 to compute the missing dimension.

□

Remark 4.9 The reader may notice that this result depends crucially on the cancellation of the terms arising from $h_{P_q}^0(\mathcal{I}_{E_0,q}^r \mathcal{L}_{0|P_q}^r)$ over the various points q . These are essentially (to higher order in γ) the terms encoding the singularities which form in \mathcal{X} as $t \rightarrow 0$, over which we have little control. A first reason why $F(\mathcal{X})$ should not “see” these singularities is the argument with balanced metrics [17] 27–28 already mentioned in 1.5 which we do not reproduce here. An alternative differential-geometric argument is sketched in 4.12 below.

The results obtained so far can be put in a form which makes applying definition (4.1) easier. Define co-efficients b_i, c_i by $h_X^0(L^{\gamma r}) = c_0 \gamma^n r^n + c_1 \gamma^{n-1} r^{n-1} + O(r^{n-2})$, $\text{tr}(H_X^0(L^{\gamma r})) = b_0 \gamma^{n+1} r^{n+1} + b_1 \gamma^n r^n + O(r^{n-1})$. Similarly we define b'_i, c'_i by $h_{\mathcal{X}_0}^0(\mathcal{L}_0^r) = c'_0(\gamma) r^n + c'_1(\gamma) r^{n-1} + O(r^{n-2})$, $\text{tr}(H^0(\mathcal{X}_0, \mathcal{L}_0^r)) = b'_0(\gamma) r^{n+1} + b'_1(\gamma) r^n + O(r^{n-1})$.

Corollary 4.10

$$\begin{aligned} b'_0 &= b_0 \gamma^{n+1} - \sum_q \lambda(q) \left(\sum_{p_i \in A_q} a_i^n \right) \frac{\gamma}{n!} + O(1), \\ c'_1 &= c_1 \gamma^{n-1} - \frac{1}{2(n-2)!} \sum_i a_i^{n-1}, \\ c'_0 &= c_0 \gamma^n - \frac{1}{n!} \sum_i a_i^n, \\ b'_1 &= b_1 \gamma^n - \sum_q \lambda(q) \left(\sum_{p_i \in A_q} a_i^{n-1} \right) \frac{\gamma}{2(n-2)!} + O(1). \end{aligned}$$

Proof. This is a restatement of 4.3 and 4.8. □

With these preliminary computations in place we can now prove our main result.

Proof of 1.3. By (4.1)

$$\begin{aligned} F(\mathcal{X}) &= \frac{b'_0 c'_1}{c'_0} - b'_1 = \\ &= \left(\frac{b_0 c_1}{c_0} - b_1 \right) \gamma^n \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{2(n-2)!} \left(\sum_q \lambda(q) \left(\sum_{p_i \in A_q} a_i^{n-1} \right) - \frac{b_0}{c_0} \sum_i a_i^{n-1} \right) \gamma + O(1) = \\
& = F(X) \gamma^n + \frac{1}{2(n-2)!} \left(\sum_q \sum_{p_i \in A_q} a_i^{n-1} \left(\lambda(q) - \frac{b_0}{c_0} \right) \right) \gamma + O(1).
\end{aligned}$$

It remains to make the connection with Chow stability. Recall that to define the Chow weights with respect to the line L^γ on X we need α to act through $\mathrm{Sl}(H_X^0(L^\gamma)^*)$. Choosing any infinitesimal generator A_γ for the action on $H_X^0(L^\gamma)^*$ we need to solve for a correction parameter λ_γ

$$\mathrm{tr}(A_\gamma) + \gamma \lambda_\gamma h_X^0(L^\gamma) = 0$$

so we get

$$\lambda_\gamma = -\frac{\mathrm{tr}(A_\gamma)}{\gamma h_X^0(L^\gamma)} = -\frac{b_0}{c_0} + O(\gamma^{-1}).$$

Note that after pulling back the family \mathcal{X} by a finite covering of \mathbb{C} (i.e. $t \mapsto t^k$ for some k) we may assume $\lambda_\gamma \in \mathbb{Z}$. So by substituting $\lambda_\gamma + O(\gamma^{-1})$ for $-\frac{b_0}{c_0}$ the expansion above may be read as

$$F(\mathcal{X}) = F(X) \gamma^n + \frac{1}{2(n-2)!} \left(\sum_q \sum_{p_i \in A_q} a_i^{n-1} \lambda'(q) \right) \gamma + O(1)$$

where $\lambda'(q)$ are the new special linear weights

$$\lambda'(q) = \lambda(q) + \lambda_\gamma.$$

By the discussion in section 3 we see

$$\sum_q \sum_{p_i \in A_q} a_i^{n-1} \lambda'(q) = -\mathcal{CH}(\sum_i a_i^{n-1} p_i, \alpha)$$

where \mathcal{CH} stands for the Chow weight relative to the polarisation on $X^{(d)}$ induced by $(L^\gamma)^{\boxtimes d}$.

□

Remark 4.11 In view of the proof we should write \mathcal{CH}_γ in theorem 1.3, but we drop the dependence on γ motivated by Remark 3.2. We should also write \mathcal{X}_γ but this dependence is not really serious since we are only pulling back \mathcal{X} by a finite covering.

Proof of 1.4. By [15] Theorem 3.9 and Theorem 4.33 we know that if (X, L) is asymptotically Chow stable then it is K-semistable. In particular this implies $F(X) = 0$. Now we blowup along a Chow unstable cycle and apply 1.3 to get $F(\mathcal{X}) < 0$ for $\gamma \gg 0$, i.e. $(\text{Bl}_Z X, L^\gamma - E)$ is K-unstable, and so in turn asymptotically Chow unstable.

□

Remark 4.12 We now give the argument promised in 4.9. For this we need to recall the definition of the K -energy functional (due to Mabuchi [12]). Consider the space of Kähler potentials with respect to a fixed Kähler form ω , $\mathcal{H} = \{\phi \in C^\infty(M, \mathbb{R}) : \omega_\phi = \omega + i\partial\bar{\partial}\phi > 0\}$. K -energy is the unique functional \mathcal{M} on \mathcal{H} (up to an additive constant) such that the derivative $\frac{d}{dt}\mathcal{M}(t)$ along any path $\omega_t = \omega + i\partial\bar{\partial}\phi_t$ is given by $\int_M \left(\frac{d}{dt}\phi_t\right) (S(\omega_t) - \widehat{S}) \omega_t^n$. Now by the general theory (markedly the moment map picture in [6]) one expects that the Donaldson-Futaki invariant for a test configuration \mathcal{X} can be computed via any path of metrics ω_t , $t \in (0, 1]$ which is “adapted” to \mathcal{X} (we will not try to make this precise here). The prediction is then $\lim_{t \rightarrow 0} t \frac{d}{dt}\mathcal{M}(t) = F(\mathcal{X})$. Let us spell out what this means in our case, starting with a Kähler form ω on the base X and a holomorphic vector field α with flow $\alpha(t)$. Choose small enough *disjoint* coordinate balls $B_i(2\epsilon)$ around each p_i . At a fixed time t , we consider the metric $\alpha(t)^*\omega$ on X ; remove a ball $B_i(\epsilon)$ and glue in a metric on a large open neighborhood of the zero section of $\mathcal{O}(-1) \rightarrow \mathbb{P}^{n-1}$, such that the volume of the zero section is $\epsilon^{n-1} a_i^{n-1}$ (up to a constant). This will require deforming $\alpha(t)^*\omega$, but we can leave it *unchanged* outside $B_i(2\epsilon)$. This construction can be made to yield a sequence of smooth metrics on $\text{Bl}_{\{p_i\}} X$ with the required cohomology class. Then taking the limit as $t \rightarrow 0$ morally yields $F(\mathcal{X})$. On the other hand, it should be clear by construction and the definition of $\frac{d}{dt}\mathcal{M}(t)$ that this limit only depends on the action of $\alpha(t)$ in a neighborhood of each p_i for small t ; that is, the result does *not* depend on mutual interaction of the points $\{p_i\}$ we blowup. While this conjectural picture could be overly difficult to make precise, it gives some geometric meaning to the cancellation of the $h_{P_q}^0(\mathcal{I}_{E_0,q}^r \mathcal{L}_{0|P_q}^r)$ terms.

Remark 4.13 A striking feature of 1.3 is that it naturally suggests that higher order contributions to $F(\mathcal{X})$ should arise from blowing up higher dimensional subschemes.

Remark 4.14 Note that 1.3 still applies when the points $\{p_i\}$ are fixed under the \mathbb{C}^* -action; for example one can apply 1.10 to fixed points with appropriate multiplicities. Moreover, we expect that 1.3 still holds for arbitrarily singular fixed points on a polarised variety (X, L) ; of course $F(X)$ will

need to be replaced with the Donaldson-Futaki invariant for the \mathbb{C}^* -action on the base.

Remark 4.15 A very interesting generalisation of 1.3 has been found by Della Vedova [4]. Roughly speaking this applies to *extremal* metrics, i.e. metrics whose scalar curvature has holomorphic $(1, 0)$ -gradient. See also [3] for more details.

References

- [1] Arezzo, C. and Pacard, F. *Blowing up and desingularizing constant scalar curvature Kähler manifolds*. arXiv:math/0411522v3 [math.DG] (2004).
- [2] Arezzo, C. and Pacard, F. *Blowing up Kähler manifolds with constant scalar curvature II*. arXiv:math/0504115v1 (2005). To appear in Annals of Math. 2007.
- [3] Arezzo, C. and Pacard, F. *On the Kähler classes of constant scalar curvature metrics on blow ups* arXiv:0706.1838v1 [math.DG] (2007).
- [4] Della Vedova, A. *Relative stability of points and relative K-stability*, in preparation.
- [5] Chen, X., LeBrun, C. and Weber, B. *On conformally Kähler, Einstein metrics*, arXiv:0705.0710v2 [math.DG] (2007).
- [6] S. K. Donaldson, *Remarks on gauge theory, complex geometry and 4-manifold topology*, Fields Medallists' Lectures (M. F. Atiyah and D. Iagolnitzer, eds.), World Sci. Publ., Singapore, 1997, pp.384-403.
- [7] Donaldson, S. K. *Scalar curvature and projective embeddings I*. J. Differential Geom. 59 (2001), no. 3, 479–522.
- [8] Donaldson, S.K. *Scalar curvature and stability of toric varieties*. Jour. Differential Geom. 62 (2002) 289-349
- [9] Donaldson, S. K. *Lower bounds on the Calabi functional*. J. Differential Geom. 70 (2005), no. 3, 453–472.
- [10] Hartshorne R. *Algebraic Geometry*. Springer-Verlag, Heidelberg 1977.
- [11] Hwang A. and Singer, M. *A momentum construction for circle-invariant Kähler metrics*, Trans. Americ. Math. Soc. 354 (2002), no. 6, 2285-2325.

- [12] Mabuchi, T. *K-energy maps integrating Futaki invariants*, Tohoku Math. J. (2) 38 (1986), no. 4, 575–593.
- [13] Mukai, S. *An introduction to invariants and moduli*. Cambridge University Press 2003.
- [14] Mumford, D., Fogarty, J. and Kirwan, F. *Geometric Invariant Theory*. Springer-Verlag, Heidelberg 1994.
- [15] Ross, J. and Thomas, R. P. *A study of the Hilbert-Mumford criterion for the stability of projective varieties*. J. Algebraic Geom. 16 (2007) 201–255. arXiv:math/0412519v2 (2004)
- [16] Ross, J. and Thomas, R. P. *An obstruction to the existence of constant scalar curvature Kähler metrics*. J. Differential Geom. 72 (2006), 429–466
- [17] Thomas, R. P. *Notes on GIT and symplectic reduction for bundles and varieties*. Surveys in Differential Geometry X (S.T. Yau editor) International Press, Somerville 2006. arXiv:math/0512411v3 [math.AG]
- [18] Székelyhidi, G. *Extremal metrics and K-stability*., PhD Thesis, Imperial College London. arXiv:math/0611002v1 [math.DG] (2007)
- [19] Tian, G. *Kähler-Einstein metrics with positive scalar curvature*. Invent. Math. 137 (1997), 1–37.
- [20] Shu, Y. *Compact complex surfaces and Constant Scalar Curvature Kähler metrics*. arXiv:math/0612013v1 [math.DG] (2006)

Università di Pavia, Via Ferrata 1 27100 Pavia, ITALY
 and
 Imperial College, London SW7 2AZ, UK.
E-mail: jacopo.stoppa@unipv.it