



SCUOLA INTERNAZIONALE SUPERIORE DI STUDI AVANZATI

SISSA Digital Library

On the two-state problem for general differential operators

Original

On the two-state problem for general differential operators / De Philippis, Guido; Palmieri, Luca; Rindler, Filip. - In: NONLINEAR ANALYSIS. - ISSN 0362-546X. - 177, Part B:(2018), pp. 387-396.
[10.1016/j.na.2018.03.015]

Availability:

This version is available at: 20.500.11767/85686 since: 2018-12-21T19:15:23Z

Publisher:

Published

DOI:10.1016/j.na.2018.03.015

Terms of use:

Testo definito dall'ateneo relativo alle clausole di concessione d'uso

Publisher copyright

Elsevier

This version is available for education and non-commercial purposes.

note finali coverpage

(Article begins on next page)

ON THE TWO-STATE PROBLEM FOR GENERAL DIFFERENTIAL OPERATORS

GUIDO DE PHILIPPIS, LUCA PALMIERI, AND FILIP RINDLER

To Carlo Sbordone, for his 70th birthday.

ABSTRACT. In this note we generalize the Ball-James rigidity theorem for gradient differential inclusions to the setting of a general linear differential constraint. In particular, we prove the rigidity for approximate solutions to the two-state inclusion with incompatible states for merely L^1 -bounded sequences. In this way, our theorem can be seen as a result of compensated compactness in the linear-growth setting.

1. INTRODUCTION

In [2] Ball and James proved the following rigidity property for gradient differential inclusions:

Theorem 1.1 (Ball–James 1987 [2]). *Let $\Omega \subset \mathbb{R}^d$ be an open, bounded, and connected set and let $A, B \in \mathbb{R}^m \otimes \mathbb{R}^d$ be $(m \times d)$ -matrices with $\text{rk}(A - B) \geq 2$. Then:*

(A) *If $u \in W^{1,\infty}(\Omega; \mathbb{R}^m)$ satisfies the differential inclusion*

$$Du(x) \in \{A, B\} \quad \text{for a.e. } x \in \Omega, \tag{1.1}$$

then either $Du \equiv A$ or $Du \equiv B$.

(B) *Let $(u_j) \subset W^{1,\infty}(\Omega; \mathbb{R}^m)$ be a uniformly norm-bounded sequence of maps such that*

$$\text{dist}(Du_j, \{A, B\}) \rightarrow 0 \quad \text{in measure.}$$

Then, up to extracting a subsequence, either

$$\int_{\Omega} |Du_j(x) - A| \, dx \rightarrow 0 \quad \text{or} \quad \int_{\Omega} |Du_j(x) - B| \, dx \rightarrow 0 \quad \text{as } j \rightarrow \infty.$$

The first part is known as *rigidity for exact solutions* of the differential inclusion (1.1), while the second one concerns *rigidity for approximate solutions*; see [13, Chapter 2] and [19, Chapter 8] for a discussion. Note also that, by a simple laminate construction, if $\text{rk}(A - B) = 1$ then there exists a non-affine $u \in W^{1,\infty}(\Omega; \mathbb{R}^m)$ such that $Du(x) \in \{A, B\}$ for almost all $x \in \Omega$. Moreover, laminates are the only possible solutions of this inclusion.

By recalling that on a simply connected domain $v = Du$ if and only if $\text{curl } v = 0$, the above results can be summarized as follows:

The inclusion

$$\begin{cases} \text{curl } v = 0, \\ v \in \{A, B\} \end{cases}$$

is rigid for both exact and approximate solutions (in $L^\infty(\Omega; \mathbb{R}^{m \times d})$) if and only if $\text{rk}(A - B) \geq 2$.

In view of the above discussion it is natural to ask what happens when we replace the curl with a general differential operator acting on vector-valued functions $v \in C^\infty(\Omega; \mathbb{R}^\ell)$, $\Omega \subset \mathbb{R}^d$, namely

$$\mathcal{A}v := \sum_{|\alpha|=k} A_\alpha \partial^\alpha v, \quad (1.2)$$

where the sum is over all multi-indices $\alpha \in (\mathbb{N} \cup \{0\})^d$, $A_\alpha \in \mathbb{R}^n \otimes \mathbb{R}^\ell$ are (constant) matrices (note that then the equation $\mathcal{A}v = 0$ is actually a *system of equations*).

Since the seminal work of Murat and Tartar [14, 15, 24, 25] it is well understood that the role of rank-one matrices for gradient inclusions should be played by the *wave cone* associated to \mathcal{A} :

$$\Lambda_{\mathcal{A}} := \bigcup_{|\xi|=1} \ker \mathbb{A}(\xi), \quad \mathbb{A}(\xi) := (2\pi i)^k \sum_{|\alpha|=k} A_\alpha \xi^\alpha,$$

where $\xi^\alpha = \xi_1^{\alpha_1} \cdots \xi_d^{\alpha_d}$. Indeed, one may easily check that if $\lambda - \mu \in \ker \mathbb{A}(\xi)$ for some $\xi \neq 0$ and $h : \mathbb{R} \rightarrow \{0, 1\}$ is measurable, then the function

$$v(x) := \lambda h(x \cdot \xi) + \mu(1 - h(x \cdot \xi)) \quad (1.3)$$

is a solution to the differential inclusion

$$\begin{cases} \mathcal{A}v = 0 & \text{in the sense of distributions,} \\ v \in \{\lambda, \mu\} & \text{a.e.} \end{cases}$$

Our main result generalizes the Ball–James Theorem 1.1 to general operators \mathcal{A} as above.

Theorem 1.2. *Let $\Omega \subset \mathbb{R}^d$ be an open, bounded, and connected set, and let \mathcal{A} be as in (1.2). Suppose that $\lambda, \mu \in \mathbb{R}^\ell$ with $\lambda - \mu \notin \Lambda_{\mathcal{A}}$. Then:*

(A) *If $v \in L^\infty(\Omega; \mathbb{R}^\ell)$ is such that*

$$\mathcal{A}v = 0 \quad \text{in the sense of distributions}$$

and

$$v(x) \in \{\lambda, \mu\} \quad \text{for a.e. } x \in \Omega,$$

then either $v \equiv \lambda$ or $v \equiv \mu$.

(B) *Let $(v_j) \subset L^1(\Omega; \mathbb{R}^\ell)$ be a uniformly norm-bounded sequence of maps such that*

$$\mathcal{A}v_j = 0 \quad \text{in the sense of distributions.}$$

Assume that

$$\lim_{j \rightarrow \infty} \int_{\Omega} \text{dist}(v_j(x), \{\lambda, \mu\}) \, dx = 0.$$

Then, up to extracting a subsequence, either

$$\int_{\Omega} |v_j(x) - \lambda| \, dx \rightarrow 0 \quad \text{or} \quad \int_{\Omega} |v_j(x) - \mu| \, dx \rightarrow 0 \quad \text{as } j \rightarrow \infty.$$

Remark 1.3. As mentioned above, if $\lambda - \mu \in \Lambda_{\mathcal{A}}$, then it is always possible to find a non-trivial solution of the exact differential inclusion. However, for operators of order greater or equal than two, not all the solutions are given by (1.3) and more general structures can arise. Classifying all of them would be an interesting problem.

For particular choices of \mathcal{A} , some instances of Theorem 1.2 were already known: as mentioned before, if $\mathcal{A} = \text{curl}$, then it essentially reduces to the Ball–James Theorem (1.1) up to the improvement on the summability assumption on the sequence $(v_j)_j = (Du_j)_j$ (which could also have been obtained in this particular case by the combination of Zhang’s Lemma [26] and the simple Lemma 2.1 below). In the case when $\mathcal{A} = \text{div}$, the above result was obtained by Garroni and Nesi [11]. In the same paper the result is also generalized to some other first-order operators. Other types of first-order operators (essentially combination of the div and curl) have been treated by Barchiesi in his master thesis [3].

We refer to Remark 2.2 for an extension of this theorem to non-homogeneous operators and non-zero (but asymptotically vanishing) right-hand sides in the constraint PDE.

Another interesting choice of operator is the so called “curl curl”-operator, a second-order operator whose kernel identifies those symmetric matrix-valued fields $S: \Omega \subset \mathbb{R}^d \rightarrow (\mathbb{R}^d \otimes \mathbb{R}^d)_{\text{sym}}$, $\Omega \subset \mathbb{R}^d$ a bounded simply-connected Lipschitz domain, for which there exists $u: \Omega \rightarrow \mathbb{R}^d$ such that

$$S = \mathcal{E}u := \frac{Du + (Du)^T}{2}.$$

For this operator the wave cone is given by the elementary symmetrized products,

$$\Lambda_{\text{curl curl}} = \left\{ a \odot \xi := \frac{a \otimes \xi + \xi \otimes a}{2} : a, \xi \in \mathbb{R}^d \right\},$$

see for instance [10, Example 3.10(e)] and [7, Proof of Theorem 1.7]. In this case Theorem 1.2 implies the following:

Corollary 1.4. *Let $(u_j)_j \in W^{1,1}(\Omega; \mathbb{R}^d)$ be a sequence of maps such that*

$$\sup_{j \in \mathbb{N}} \|\mathcal{E}u_j\|_{L^1} < \infty \quad \text{and} \quad \text{dist}(\mathcal{E}u_j, \{A, B\}) \xrightarrow{L^1} 0$$

for some $A, B \in (\mathbb{R}^d \otimes \mathbb{R}^d)_{\text{sym}}$ with $A - B \notin \Lambda_{\text{curl curl}}$. Then, up to extracting a subsequence, either

$$\int_{\Omega} |\mathcal{E}u_j(x) - A| \, dx \rightarrow 0 \quad \text{or} \quad \int_{\Omega} |\mathcal{E}u_j(x) - B| \, dx \rightarrow 0 \quad \text{as } j \rightarrow \infty.$$

Other, more complicated, differential inclusion for (bounded) symmetric gradients have been studied in [4, 5, 8, 18, 20, 21] (but without the linear-growth assumption).

Despite the fact that many interesting cases of Theorem 1.2 were already known, we nevertheless believe that the generality of Theorem 1.2 is of independent interest:

First, it extends all the known results to general operators of any order.

Second, our result for the first time enables one to treat the case of maps that are merely bounded in L^1 . Hence, Theorem 1.2 may be seen as a result of *compensated compactness theory in L^1* .

Third, the previously-known cases of Theorem 1.2 have been proved by *ad hoc* techniques. On the contrary, our proof exploits in quite a clear way the heuristic principle that *when restricted to maps whose image is not in the wave cone, \mathcal{A} is elliptic* (which in this generality is too good to be true); see Section 2 for a more precise statement. Of course, this idea lies at the root of the Murat–Tartar compensated compactness theory, which also provides the conceptual framework of the present work. In the proof of Theorem 1.2 we exploit some ideas which have been

introduced in [7], where the first and third author proved the natural generalization of Alberti's rank-one theorem to the setting of \mathcal{A} -free measures.

We conclude this introduction by mentioning that one can also study more complicated differential inclusions. In the case of $\mathcal{A} = \text{curl}$, this has been a very active and fruitful area of research with several deep results and nice connections with other problems in mathematics. Since here we cannot give a detailed account of the (extensive) literature, we refer the reader to [19, Chapter 8] for a recent survey on the theory.

It is worth noting that in the study of more general inclusions in the general setting of Theorem 1.2, the intrinsic geometry of the operator \mathcal{A} should play an important role. Indeed, for differential inclusions involving more than two states the type of results available seems to be different depending on the particular choice of \mathcal{A} . For instance, when $\mathcal{A} = \text{curl}$ it has been proved by Šverák that the three-state problem is rigid both for approximate and exact solutions [23]. On the contrary, when $\mathcal{A} = \text{div}$, in [11, Section 4] it is shown that the three state-problem is not rigid for approximate solutions, while Palombaro and Ponsiglione [16] showed that it is rigid for exact solutions; see also [17].

Acknowledgements. G. D. P. is supported by the MIUR SIR-grant ‘‘Geometric Variational Problems’’ (RBSI14RVEZ). This project has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement 757254) for the project ‘‘SINGULARITY’’. F. R. also acknowledges the support from an EPSRC Research Fellowship on ‘‘Singularities in Nonlinear PDEs’’ (EP/L018934/1).

2. PROOF OF THEOREM 1.2

In this Section we prove Theorem 1.2. As mentioned in the introduction, the proof is based on the observation (which lies at the very definition of the wave cone) that, if $\lambda \notin \Lambda_{\mathcal{A}}$, then the operator \mathcal{A} restricted to functions whose image lies in $\text{span}\{\lambda\}$ is *elliptic* in the sense that its Fourier symbol does not vanish. Exploiting this elliptic regularization together with some classical estimates in harmonic analysis in the spirit of [7] allows us to deduce the desired rigidity.

We start with an elementary lemma, which relates the L^1 -convergence to zero of $\text{dist}(v_j, \{\lambda, \mu\})$ to the equi-integrability of the sequence $(v_j)_j$. Recall that a sequence of measurable functions $f_j: \Omega \rightarrow \mathbb{R}$ ($j \in \mathbb{N}$) is said to converge to 0 in measure if

$$\int_{\Omega} \min\{|f_j|, 1\} \, dx \rightarrow 0 \quad \text{as } j \rightarrow \infty.$$

Lemma 2.1. *Let $\Omega \subset \mathbb{R}^d$ be an open and bounded set, let $(v_j)_j \subset L^1(\Omega; \mathbb{R}^\ell)$ and let $\lambda, \mu \in \mathbb{R}^\ell$. Then the following are equivalent:*

- (i) $\text{dist}(v_j, \{\lambda, \mu\}) \xrightarrow{L^1} 0$;
- (ii) *the sequence $(v_j)_j$ is equi-integrable, i.e.,*

$$\limsup_{t \rightarrow \infty} \int_{\bigcup_{j \in \mathbb{N}} \Omega \cap \{|v_j| > t\}} |v_j| \, dx = 0,$$

and $\text{dist}(v_j, \{\lambda, \mu\}) \rightarrow 0$ in measure;

- (iii) *the sequence $(v_j)_j$ is equi-integrable and $\text{dist}(v_j, \{\lambda, \mu\}) \xrightarrow{L^1} 0$.*

Proof. For $t \geq 2 \max\{|\lambda|, |\mu|\}$ we have

$$\{\text{dist}(v, \{\lambda, \mu\}) > 2t\} \subset \{|v| > t\} \subset \{\text{dist}(v, \{\lambda, \mu\}) > t/2\}.$$

Hence, $(v_j)_j$ is equi-integrable if and only if $(\text{dist}(v_j, \{\lambda, \mu\}))_j$ is equi-integrable. By Vitali's Theorem (which says that sequence of equi-integrable functions converges to 0 in measure if and only if it converges to 0 in L^1) together with the fact that L^1 -converging sequences are equi-integrable, we immediately get that (i), (ii), and (iii) are equivalent. \square

Proof of Theorem 1.2 (A). Shifting $\tilde{v}(x) := v(x) - \mu$, we can assume that $\mu = 0$ and $v = \lambda \mathbb{1}_E$ with $\lambda \notin \Lambda_{\mathcal{A}}$ and

$$E = \{x \in \Omega : v(x) = \lambda\}.$$

The goal is to show that either $E = \Omega$ or $E = \emptyset$. We will prove that $\mathbb{1}_E \in C^\infty(\Omega)$, which immediately implies the conclusion since Ω is connected. In order to do so, let $\varphi \in C_c^\infty(\Omega)$ and let us consider the function $w := \varphi \mathbb{1}_E \in L^2(\mathbb{R}^d)$. Then,

$$\mathcal{A}(\lambda w) = \varphi \mathcal{A}(\lambda \mathbb{1}_E) + [\mathcal{A}, \varphi](\lambda \mathbb{1}_E) = [\mathcal{A}, \varphi](\psi \lambda \mathbb{1}_E), \quad (2.1)$$

where $\psi \in C_c^\infty(\Omega)$ is identically 1 in $\text{supp } \varphi$ and for $f \in L^2(\mathbb{R}^d)$ we have defined

$$\begin{aligned} [\mathcal{A}, \varphi](f) &:= \mathcal{A}(\varphi f) - \varphi \mathcal{A}(f) \\ &= \sum_{h=1}^{k-1} \sum_{|\beta|=h} B_\beta(x) \partial^\beta f \end{aligned} \quad (2.2)$$

with some coefficient matrices $B_\beta \in C_c^\infty(\text{supp } \varphi; \mathbb{R}^n \otimes \mathbb{R}^\ell)$. The key point is that the commutator $[\mathcal{A}, \varphi]$ is an operator of order $(k-1)$. More precisely, one easily checks that

$$[\mathcal{A}, \varphi] : H^s(\mathbb{R}^d) \rightarrow H^{s-(k-1)}(\mathbb{R}^d)$$

for all $s \in \mathbb{R}$, see for instance [9, Proof of Theorem 9.26] or part (B) of the proof below. Here, $H^s(\mathbb{R}^d)$ is the classical L^2 -based Sobolev space, that is,

$$f \in H^s(\mathbb{R}^d) \iff (\text{id} - \Delta)^{s/2} f \in L^2(\mathbb{R}^d) \iff (1 + 4\pi^2 |\xi|^2)^{s/2} \hat{f}(\xi) \in L^2(\mathbb{R}^d), \quad (2.3)$$

where we denote the Fourier transform of $f \in L^2(\mathbb{R}^d)$ by

$$\hat{f}(\xi) = \mathcal{F}[f](\xi) := \int f(x) e^{-2\pi i x \cdot \xi} dx,$$

which also lies $L^2(\mathbb{R}^d)$ by the Plancherel theorem. For general tempered distributions f we define \hat{f} in the distributional sense, see [12] for details.

Taking the Fourier transform of (2.1), we thus obtain

$$\mathbb{A}(\xi) \lambda \hat{w}(\xi) = \mathcal{F}[[\mathcal{A}, \varphi](\psi \lambda \mathbb{1}_E)](\xi), \quad \xi \in \mathbb{R}^d.$$

Scalar multiplying the above equation with $\overline{\mathbb{A}(\xi) \lambda} \in \mathbb{C}^n$,

$$(1 + |\mathbb{A}(\xi) \lambda|^2) \hat{w}(\xi) = \overline{\mathbb{A}(\xi) \lambda} \cdot \mathcal{F}[[\mathcal{A}, \varphi](\psi \lambda \mathbb{1}_E)](\xi) + \hat{w}(\xi) := \hat{R}(\xi). \quad (2.4)$$

Since by assumption $\lambda \notin \Lambda_{\mathcal{A}}$, there exists $c > 0$ such that $|\mathbb{A}(\xi) \lambda| \geq c |\xi|^k$ for all $\xi \in \mathbb{R}^d$. Hence, by the very definition of Sobolev spaces (2.3), the operator T defined by

$$T(f) := \mathcal{F}^{-1} \left(\frac{\hat{f}(\xi)}{1 + |\mathbb{A}(\xi) \lambda|^2} \right)$$

maps $H^s(\mathbb{R}^d)$ into $H^{s+2k}(\mathbb{R}^d)$ for all $s \in \mathbb{R}$. Since, by our initial assumption, $\mathbb{1}_E \in L^2(\mathbb{R}^d) = H^0(\mathbb{R}^d)$, we have that the right-hand side \hat{R} of (2.4) belongs to $\mathcal{F}(H^{1-2k}(\mathbb{R}^d))$. Thus,

$$w = \varphi \mathbb{1}_E = T(\mathcal{F}^{-1}(\hat{R})) \in H^1(\mathbb{R}^d) \quad \text{for all } \varphi \in C_c^1(\Omega).$$

This in turn yields $\hat{R} \in \mathcal{F}(H^{2-2k}(\mathbb{R}^d))$, which implies that $w = \varphi \mathbb{1}_E \in H^2(\mathbb{R}^d)$ for all $\varphi \in C_c^1(\Omega)$. Iterating, we obtain the desired regularity for $\mathbb{1}_E$ and we conclude the proof (in fact, already the first step implies that $\mathbb{1}_E \in H_{\text{loc}}^1(\Omega)$ and this is enough to conclude since there are no non-constant characteristic functions in $H_{\text{loc}}^1(\Omega)$). \square

Proof of Theorem 1.2 (B). As in the proof for (A) we can assume that $\mu = 0$ and $\lambda \notin \Lambda_{\mathcal{A}}$. By Lemma 2.1 the sequence (v_j) is equi-integrable. Hence, by the Dunford–Pettis compactness theorem (see, for instance, [1, Theorem 1.38]), there exists a (non-relabelled) subsequence and a function $v \in L^1(\Omega)$ such that

$$v_j \xrightarrow{L^1} v.$$

With

$$\varepsilon_j := \|\text{dist}(v_j, \{\lambda, 0\})\|_{L^1}^{1/2} \rightarrow 0$$

let us define

$$E_j := \{x \in \Omega : |v_j(x) - \lambda| \leq \varepsilon_j\}.$$

We first show that

$$\|v_j - \lambda \mathbb{1}_{E_j}\|_{L^1} \rightarrow 0. \quad (2.5)$$

Indeed,

$$A_j := \{x \in \Omega : \text{dist}(v_j(x), \{\lambda, 0\}) \leq \varepsilon_j\} = E_j \cup \{x \in \Omega : |v_j(x)| \leq \varepsilon_j\}$$

and the union is disjoint provided ε_j is small enough. Moreover,

$$|\Omega \setminus A_j| \leq \frac{1}{\varepsilon_j} \int_{\Omega} \text{dist}(v_j, \{\lambda, 0\}) \, dx \rightarrow 0.$$

Hence,

$$\begin{aligned} \int_{\Omega} |v_j - \lambda \mathbb{1}_{E_j}| \, dx &= \int_{E_j} |v_j - \lambda \mathbb{1}_{E_j}| \, dx + \int_{\{|v_j| \leq \varepsilon_j\}} |v_j| \, dx + \int_{\Omega \setminus A_j} |v_j| \, dx \\ &\leq 2\varepsilon_j |\Omega| + \int_{\Omega \setminus A_j} |v_j| \, dx \rightarrow 0 \end{aligned}$$

since the sequence (v_j) is equi-integrable and $|\Omega \setminus A_j| \rightarrow 0$. This shows (2.5) and then also

$$\lambda \mathbb{1}_{E_j} \xrightarrow{L^1} v. \quad (2.6)$$

We now claim:

$$\text{The sequence } \mathbb{1}_{E_j} \text{ is (strongly) pre-compact in } L^1(\Omega). \quad (2.7)$$

Before proving this claim, let us show how this gives the desired conclusion. Indeed, (2.5), (2.6) and (2.7) imply that

$$\|\lambda \mathbb{1}_{E_j} - v\|_{L^1} + \|v_j - v\|_{L^1} \rightarrow 0. \quad (2.8)$$

In particular, $v = \lambda \mathbb{1}_E$ for some set $E \subset \Omega$. Since clearly $\mathcal{A}v = 0$, part (A) implies that either $E = \Omega$ or $E = \emptyset$. Together with (2.8) this concludes the proof.

We are thus left to show the claim (2.7). By the equi-integrability, it is enough to show that $w_j := \varphi \mathbb{1}_{E_j}$ is pre-compact for all $\varphi \in C_c^\infty(\Omega)$. As in part (A) we have

$$\mathcal{A}(\lambda w_j) = \mathcal{A}(\lambda w_j - \varphi v_j) + \mathcal{A}(\varphi v_j) = \mathcal{A}(\lambda w_j - \varphi v_j) + [\mathcal{A}, \varphi](\psi v_j),$$

where $\psi \in C_c^\infty(\Omega)$ is identically 1 on $\text{supp } \varphi$ and $[\mathcal{A}, \varphi]$ is defined in (2.2). Taking the Fourier transform of the above equation and multiplying by $\overline{\mathbb{A}(\xi)\lambda}$, we get

$$(1 + |\mathbb{A}(\xi)\lambda|^2)\hat{w}_j(\xi) = \overline{\mathbb{A}(\xi)\lambda} \cdot \mathbb{A}(\xi)\hat{z}_j(\xi) + \overline{\mathbb{A}(\xi)\lambda} \cdot \mathcal{F}[[\mathcal{A}, \varphi](u_j)](\xi) + \hat{w}_j(\xi),$$

where we have set

$$z_j := \lambda w_j - \varphi v_j, \quad u_j := \psi v_j.$$

Hence,

$$w_j = T_1(z_j) + T_2(u_j) + T_3(w_j), \quad (2.9)$$

where $z_j, u_j, w_j \in L^1(\text{supp } \psi)$ and

$$\begin{aligned} T_1(z) &:= \mathcal{F}^{-1} \left(\frac{\overline{\mathbb{A}(\xi)\lambda} \cdot \mathbb{A}(\xi)\hat{z}(\xi)}{1 + |\mathbb{A}(\xi)\lambda|^2} \right), \\ T_2(u) &:= \mathcal{F}^{-1} \left(\frac{\overline{\mathbb{A}(\xi)\lambda} \cdot \mathcal{F}[[\mathcal{A}, \varphi](u)](\xi)}{1 + |\mathbb{A}(\xi)\lambda|^2} \right), \\ T_3(w) &:= \mathcal{F}^{-1} \left(\frac{\hat{w}(\xi)}{1 + |\mathbb{A}(\xi)\lambda|^2} \right). \end{aligned}$$

We also have, by (2.5),

$$z_j = \varphi(\lambda \mathbb{1}_{E_j} - v_j) \xrightarrow{L^1} 0, \quad \sup_{j \in \mathbb{N}} \int_{\mathbb{R}^d} |w_j| + |u_j| < \infty. \quad (2.10)$$

We now claim that

$$\begin{cases} \|T_1(z_j)\|_{L^{1,\infty}} := \sup_{t>0} t \{ |T_1(z_j)| > t \} \rightarrow 0; \\ T_1(z_j) \rightarrow 0 \quad \text{in the sense of distributions;} \\ (T_2(u_j))_j, (T_3(w_j))_j \text{ are pre-compact in } L^1_{\text{loc}}. \end{cases} \quad (2.11)$$

The above facts *together with the positivity of w_j* imply (2.7), see [7, Proof of Theorem 1.1] and also [6, Lemma 3.3], where the convergence lemma is explicitly stated.

The proof of (2.11) closely follows the proof of [7, Theorem 1.1]. Indeed, by assumption, $|\mathbb{A}(\xi)\lambda| \geq c|\xi|^k$ for some $c > 0$. Thus, the Hörmander-Mihlin multiplier theorem, see [12, Theorem 5.2.7], yields that T_1 is a bounded operator from $L^1(\mathbb{R}^d)$ to $L^{1,\infty}(\mathbb{R}^d)$ and in particular, via (2.10),

$$\|T_1(z_j)\|_{L^{1,\infty}} \leq C \|z_j\|_{L^1} \rightarrow 0.$$

Moreover, for $\zeta \in C_c^\infty(\mathbb{R}^d)$,

$$\langle T_1(z_j), \zeta \rangle = \langle z_j, T_1^*(\zeta) \rangle \rightarrow 0,$$

where $T_1^* : C_c^\infty(\mathbb{R}^d) \rightarrow \mathcal{S}(\mathbb{R}^d)$, the Schwartz space of test functions, is the adjoint of T_1 .

We are thus left with the task to show that $(T_2(u_j))_j$ and $(T_3(w_j))_j$ are pre-compact in L^1_{loc} . To this end note that, by (2.2), $T_2(u_j)$ can be written as a finite sum of terms of the form

$$f_j^\beta = Q \circ (\text{id} - \Delta)^{-\frac{k}{2}} \circ P_\beta \circ (\text{id} - \Delta)^{\frac{|\beta|-k}{2}} [u_j^\beta],$$

where $\beta \in (\mathbb{N} \cup \{0\})^d$, $0 \leq |\beta| \leq (k-1)$, $\sup_{j \in \mathbb{N}} \|u_j^\beta\|_{L^1} \leq C$,

$$Q[u] = \mathcal{F}^{-1} \left[(1 + |\mathbb{A}(\xi)\lambda|^2)^{-1} (1 + 4\pi^2 |\xi|^2)^{k/2} \overline{\mathbb{A}(\xi)\lambda} \hat{u}(\xi) \right],$$

and P_β is the k 'th-order pseudo-differential operator given by

$$P_\beta[u](x) = \int \frac{(2\pi i)^{|\beta|} \xi^\beta}{(1 + 4\pi^2 |\xi|^2)^{\frac{|\beta|-k}{2}}} B_\beta(x) \hat{u}(\xi) e^{2\pi i x \cdot \xi} d\xi.$$

The composition $(\text{id} - \Delta)^{-k/2} \circ P_\beta$ is a pseudo-differential operator of order 0, see [22, Theorem 2, Chapter VI], and thus bounded from $L^p(\mathbb{R}^d)$ to itself, see [22, Proposition 4 in Chapter VI]. By the Hörmander–Mihlin multiplier theorem, also Q is a bounded operator from $L^p(\mathbb{R}^d)$ to itself. Since $|\beta| \leq k-1$, one easily checks that $(\text{id} - \Delta)^{(|\beta|-k)/2}$ is compact from $L^1(\text{supp } \psi)$ to $L^p(\mathbb{R}^d)$ for $1 < p < p(d, k - |\beta|)$, see for instance [7, Lemma 2.1]. We thus infer that $(T_2(u_j))_j$ is pre-compact in $L^1_{\text{loc}}(\mathbb{R}^d)$. Since a similar (and actually easier) argument applies to $(T_3(w_j))_j$, this concludes the proof of (2.11) and thus of the theorem. \square

Remark 2.2. It is clear from the proof that Theorem 1.2 can be extended to include non-homogeneous operators and (asymptotically vanishing) right-hand sides. More precisely, assume that

$$\mathcal{A} = \sum_{|\alpha| \leq k} A_\alpha \partial^\alpha = \sum_{|\alpha|=k} A_\alpha \partial^\alpha + \sum_{|\alpha| \leq (k-1)} A_\alpha \partial^\alpha := \mathcal{A}^k + \mathcal{A}^{<k}$$

and that $(v_j)_j \subset L^1(\Omega, \mathbb{R}^\ell)$ satisfies

$$\mathcal{A}v_j = r_j, \quad \text{dist}(v_j, \{\lambda, \mu\}) \xrightarrow{L^1} 0,$$

where r_j can be written as

$$r_j = \sum_{|\beta|=k} \partial^\beta f_j^\beta \quad \text{with} \quad \|f_j^\beta\|_{L^1} \rightarrow 0. \quad (2.12)$$

Moreover, we assume

$$\lambda - \mu \notin \Lambda_{\mathcal{A}^k} := \bigcup_{|\xi|=1} \ker \mathbb{A}^k(\xi), \quad \mathbb{A}^k(\xi) := (2\pi i)^k \sum_{|\alpha|=k} A_\alpha \xi^\alpha.$$

Then, up to taking a subsequence, either $\|v_j - \lambda\|_{L^1} \rightarrow 0$ or $\|v_j - \mu\|_{L^1} \rightarrow 0$. Note in particular that (2.12) includes the case in which $r_j \rightarrow 0$ strongly in $W^{-k,p'}$ for $p > 1$.

With these assumptions, the proof of Theorem 1.2 can be repeated almost verbatim with \mathcal{A} replaced by \mathcal{A}^k . The presence of the non-homogeneous terms $\mathcal{A}^{<k}v_j$ and r_j can be dealt with by adding them to the right-hand side of (2.9). Indeed, this yields additional terms of the form

$$T_4(v_j) := \mathcal{F}^{-1} \left(\frac{\overline{\mathbb{A}^k(\xi)\lambda} \mathcal{F}[\varphi \mathcal{A}^{<k}v_j](\xi)}{1 + |\mathbb{A}^k(\xi)\lambda|^2} \right),$$

$$T_5(f_j^\beta) := \mathcal{F}^{-1} \left(\frac{(2\pi i)^k \overline{\mathbb{A}^k(\xi)\lambda} \xi^\beta \hat{f}_j^\beta(\xi)}{1 + |\mathbb{A}^k(\xi)\lambda|^2} \right).$$

By the Hörmander–Mihlin theorem, $T_5(f_j^\beta)$ converges to 0 in $L^{1,\infty}$ and in the sense of distributions, while, by the same reasoning used for $(T_2(u_j))_j$, one shows that

$(T_4(u_j))_j$ is pre-compact in L^1_{loc} . Gathering together these facts, we may conclude the proof by the very same arguments as in Theorem 1.2.

REFERENCES

- [1] L. Ambrosio, N. Fusco, and D. Pallara, *Functions of Bounded Variation and Free-Discontinuity Problems*, Oxford Mathematical Monographs, Oxford University Press, 2000.
- [2] J. M. Ball and R. D. James, *Fine phase mixtures as minimizers of energy*, Arch. Ration. Mech. Anal. **100** (1987), 13–52.
- [3] M. Barchiesi, *Inclusioni Differenziali per EDP: Rigidità e Mancanza di Rigidità*, Master’s thesis, Università di Roma “La Sapienza”, 2003.
- [4] A. Capella and F. Otto, *A rigidity result for a perturbation of the geometrically linear three-well problem*, Comm. Pure Appl. Math. **62** (2009), 1632–1669.
- [5] ———, *A quantitative rigidity result for the cubic-to-tetragonal phase transition in the geometrically linear theory with interfacial energy*, Proc. Roy. Soc. Edinburgh Sect. A **142** (2012), 273–327.
- [6] G. De Philippis, A. De Rosa, and F. Ghiraldin, *Rectifiability of varifolds with locally bounded first variation with respect to anisotropic surface energies*, Comm. Pure Appl. Math. (2016), to appear, arxiv: <https://arxiv.org/abs/1609.01908>.
- [7] G. De Philippis and F. Rindler, *On the structure of \mathcal{A} -free measures and applications*, Ann. of Math. **184** (2016), 1017–1039.
- [8] G. Dolzmann and S. Müller, *The influence of surface energy on stress-free microstructures in shape memory alloys*, Meccanica **30** (1995), 527–539, Microstructure and phase transitions in solids (Udine, 1994). MR 1360969
- [9] G. B. Folland, *Real Analysis*, 2nd ed., Pure and Applied Mathematics, Wiley, 1999.
- [10] I. Fonseca and S. Müller, *\mathcal{A} -quasiconvexity, lower semicontinuity, and Young measures*, SIAM J. Math. Anal. **30** (1999), no. 6, 1355–1390.
- [11] A. Garroni and V. Nesi, *Rigidity and lack of rigidity for solenoidal matrix fields*, Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci. **460** (2004), no. 2046, 1789–1806. MR 2067561
- [12] L. Grafakos, *Classical Fourier Analysis*, 3rd ed., Graduate Texts in Mathematics, vol. 249, Springer, 2014.
- [13] S. Müller, *Variational models for microstructure and phase transitions*, Calculus of variations and geometric evolution problems (Cetraro, 1996), Lecture Notes in Mathematics, vol. 1713, Springer, 1999, pp. 85–210.
- [14] F. Murat, *Compacité par compensation*, Ann. Sc. Norm. Sup. Pisa Cl. Sci. **5** (1978), 489–507.
- [15] ———, *Compacité par compensation. II*, Proceedings of the International Meeting on Recent Methods in Nonlinear Analysis (Rome, 1978), Pitagora Editrice Bologna, 1979, pp. 245–256.
- [16] M. Palombaro and M. Ponsiglione, *The three divergence free matrix fields problem*, Asymptot. Anal. **40** (2004), 37–49.
- [17] M. Palombaro and V. P. Smyshlyaev, *Relaxation of three solenoidal wells and characterization of extremal three-phase H -measures*, Arch. Ration. Mech. Anal. **194** (2009), 775–722.
- [18] F. Rindler, *Lower semicontinuity for integral functionals in the space of functions of bounded deformation via rigidity and Young measures*, Arch. Ration. Mech. Anal. **202** (2011), 63–113.
- [19] ———, *Calculus of Variations*, Universitext, Springer, 2018.
- [20] A. Rühländ, *The cubic-to-orthorhombic phase transition: rigidity and non-rigidity properties in the linear theory of elasticity*, Arch. Ration. Mech. Anal. **221** (2016), 23–106.
- [21] ———, *A rigidity result for a reduced model of a cubic-to-orthorhombic phase transition in the geometrically linear theory of elasticity*, J. Elasticity **123** (2016), 137–177.
- [22] E. M. Stein, *Harmonic Analysis*, Princeton University Press, 1993.
- [23] V. Šverák, *On regularity for the Monge-Ampère equation without convexity assumptions*, Tech. report, Heriot-Watt University, 1991.
- [24] L. Tartar, *Compensated compactness and applications to partial differential equations*, Nonlinear analysis and mechanics: Heriot-Watt Symposium, Vol. IV, Res. Notes in Math., vol. 39, Pitman, 1979, pp. 136–212.
- [25] ———, *The compensated compactness method applied to systems of conservation laws*, Systems of nonlinear partial differential equations (Oxford, 1982), NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., vol. 111, Reidel, 1983, pp. 263–285.

- [26] K. Zhang, *A construction of quasiconvex functions with linear growth at infinity*, Ann. Sc. Norm. Sup. Pisa Cl. Sci. **19** (1992), 313–326.

G. De Philippis: SISSA, VIA BONOMEA 265, 34136 TRIESTE, ITALY.
E-mail address: `guido.dephilippis@sissa.it`

L. Palmieri: SISSA, VIA BONOMEA 265, 34136 TRIESTE, ITALY.
E-mail address: `contact@lpalmieri.com`

F. Rindler: MATHEMATICS INSTITUTE, UNIVERSITY OF WARWICK, COVENTRY CV4 7AL, UK,
AND THE ALAN TURING INSTITUTE, BRITISH LIBRARY, 96 EUSTON ROAD, LONDON NW1 2DB
LONDON, UK.

E-mail address: `F.Rindler@warwick.ac.uk`