



SCUOLA INTERNAZIONALE SUPERIORE DI STUDI AVANZATI

SISSA Digital Library

Direct dark matter detection around the corner? Prospects in the constrained MSSM

Original

Direct dark matter detection around the corner? Prospects in the constrained MSSM / Trotta, R.; De Austri, R. R.; Roszkowski, L.. - In: JOURNAL OF PHYSICS. CONFERENCE SERIES. - ISSN 1742-6588. - 60:1(2007), pp. 259-263. (TeV Particle Astrophysics II Workshop 2006 Madison, WI AUG 28-31, 2006) [10.1088/1742-6596/60/1/056].

Availability:

This version is available at: 20.500.11767/116851 since: 2020-12-30T22:15:01Z

Publisher:

Published

DOI:10.1088/1742-6596/60/1/056

Terms of use:

Testo definito dall'ateneo relativo alle clausole di concessione d'uso

Publisher copyright

note finali coverpage

(Article begins on next page)

Direct dark matter detection around the corner? Prospects in the Constrained MSSM

Roberto Trotta

Astrophysics Department, Oxford University
Denys Wilkinson Building, Keble Road, Oxford OX1 3RH, United Kingdom

Roberto Ruiz de Austri

Departamento de Física Teórica C-XI and Instituto de Física Teórica C-XVI,
Universidad Autónoma de Madrid, Cantoblanco, 28049 Madrid, Spain

Leszek Roszkowski

Department of Physics and Astronomy, University of Sheffield,
Sheffield S3 7RH, England

E-mail: rxt@astro.ox.ac.uk

Abstract. We outline the WIMP dark matter parameter space in the Constrained MSSM by performing a comprehensive statistical analysis that compares with experimental data predicted superpartner masses and other collider observables as well as a cold dark matter abundance. We find that $10^{-10} \text{ pb} \lesssim \sigma_p^{SI} \lesssim 10^{-8} \text{ pb}$ for direct WIMP detection (with details slightly dependent on the assumptions made). We conclude that most of the 95% probability region for the cross section will be explored by future one-tonne detectors, that will therefore cover most of the currently favoured region of parameter space.

1. Introduction

Two of the most challenging questions facing particle physics today are the instability of the Higgs mass against radiative corrections (known as the “fine-tuning problem”) and the nature of dark matter. Unlike the Standard Model (SM), weak scale softly broken supersymmetry (SUSY) provides solutions to both of them. Firstly, the fine-tuning problem is addressed via the cancellation of quadratic divergences in the radiative corrections to the Higgs mass. Secondly, assuming R -parity, the lightest supersymmetric particle (LSP) is a leading weakly interactive massive particle (WIMP) candidate for cold dark matter (CDM). Despite these and other attractive features, without a reference to grand unified theories (GUTs), low energy SUSY models suffer from the lack of predictivity due to a large number of free parameters (*e.g.*, over 120 in the Minimal Supersymmetric Standard Model (MSSM)), most of which arise from the SUSY breaking sector.

The MSSM with one particularly popular choice of universal boundary conditions at the grand unification scale is called the Constrained Minimal Supersymmetric Standard Model (CMSSM) [1]. The CMSSM is defined in terms of five free parameters: common scalar (m_0),

gaugino ($m_{1/2}$) and tri-linear (A_0) mass parameters (all specified at the GUT scale) plus the ratio of Higgs vacuum expectation values $\tan\beta$ and $\text{sign}(\mu)$, where μ is the Higgs/higgsino mass parameter whose square is computed from the conditions of radiative electroweak symmetry breaking (EWSB). The economy of parameters in this scheme makes it a useful tool for exploring SUSY phenomenology.

In this work we report results from a Bayesian exploration of the CMSSM parameter space, obtained through the use of Markov Chain Monte Carlo methods. In particular we focus on the prospects for direct neutralino dark matter detection with the next generation of dark matter searches. We refer the reader to [2] for full details. For other works applying a similar approach to the CMSSM, see [3] (with some relevant differences) and more recently [4].

2. Parameter space and data used

We restrict our analysis to the case $\text{sign}(\mu) = +1$, as motivated by the fact that the observed anomalous magnetic moment of the muon is positive, and since the sign of the SUSY contribution to it is the same as the sign of μ . We thus consider the 8 dimensional parameter space (θ, ψ) , where θ is a vector of CMSSM parameters,

$$\theta = (m_0, m_{1/2}, A_0, \tan\beta) \quad (1)$$

while ψ is a vector of relevant SM parameters,

$$\psi = (M_t, m_b(m_b)^{\overline{MS}}, \alpha_{\text{em}}(M_Z)^{\overline{MS}}, \alpha_s(M_Z)^{\overline{MS}}), \quad (2)$$

where M_t is the pole top quark mass, $m_b(m_b)^{\overline{MS}}$ is the bottom quark mass at m_b , while $\alpha_{\text{em}}(M_Z)^{\overline{MS}}$ and $\alpha_s(M_Z)^{\overline{MS}}$ are the electromagnetic and the strong coupling constants at the Z pole mass M_Z , the last three evaluated in the \overline{MS} scheme. We are not interested in constraining the value of the SM parameters, but rather in including the effect of the uncertainty in their experimental determination in our inferences. Therefore we marginalize over them in the end.

We employ a Bayesian approach coupled with a Markov Chain Monte Carlo method for exploring the CMSSM parameter space and derive posterior constraints on the neutralino mass and spin-independent scattering cross section, assuming that all of the cosmological dark matter abundance is made of neutralinos. From the CMSSM and SM parameters η we compute a series of derived observable quantities $f(\eta)$: the W gauge boson mass, the effective leptonic weak mixing angle $\sin^2\theta_{\text{eff}}$, the anomalous magnetic moment of the muon, $a_\mu \equiv (g-2)_\mu$, the branching ratios $BR(\bar{B} \rightarrow X_s \gamma)$ and $BR(B_s \rightarrow \mu^+ \mu^-)$, the cosmological neutralino relic abundance $\Omega_{\text{CDM}} h^2$, the light Higgs mass and the superpartner masses. For all of those quantities, relevant measurements (summarized in the bottom section of Table 1) or experimental limits are included via the likelihood and used to constrain high posterior probability regions of the model. The likelihood also includes estimated theoretical uncertainties in the mapping from CMSSM and SM parameters to derived quantities, another major advantage of employing a Bayesian approach (see [2] for details). We then compute a spin-independent dark matter WIMP elastic scattering cross section on a free proton, σ_p^{SI} , including full supersymmetric contributions which have been derived by several groups (see [5] and references therein), but we do not include current constraints in the likelihood, in view of the uncertainties in the structure of the Galactic halo (*e.g.*, existence of clumps of dark matter and therefore the value of the local halo mass density) as well as in the values of some hadronic matrix elements entering the computation of σ_p^{SI} .

3. Results for neutralino direct detection

Our Bayesian approach allows to easily compute the posterior pdf for the cross section or any other derived variable. In Fig. 1 we present the 2-dimensional posterior pdf for σ_p^{SI} and m_χ ,

Nuisance parameter	Mean value	Uncertainty	
	μ	experimental	theoretical
M_t	172.7 GeV	2.9 GeV	N/A
$m_b(m_b)\overline{MS}$	4.24 GeV	0.11 GeV	N/A
$\alpha_s(M_Z)\overline{MS}$	0.1186	0.002	N/A
$1/\alpha_{em}(M_Z)\overline{MS}$	127.958	0.048	N/A
Derived observable			
M_W	80.425 GeV	34 MeV	13 MeV
$\sin^2 \theta_{eff}$	0.23150	16×10^{-5}	25×10^{-5}
$\delta a_\mu^{SUSY} \times 10^{10}$	25.2	9.2	1
$BR(\bar{B} \rightarrow X_s \gamma) \times 10^4$	3.39	0.30	0.30
$\Omega_\chi h^2$	0.119	0.009	$0.1 \Omega_\chi h^2$

Table 1. Experimental mean μ and standard deviation for the nuisance parameters (top section) and derived observable quantities (bottom section, including a theoretical uncertainty describing the imprecise mapping of CMSSM and SM parameters onto observable quantities) used in the analysis.




Figure 1. The 2-dimensional probability density in the neutralino mass and spin-independent cross section plane in the CMSSM (all other parameters marginalized) with the contours containing 68% and 95% probability also marked. Current 90% experimental upper limits are also shown. A large fraction of the high-probability region lies just below current constraints and it will be probed by the next generation of dark matter searches, starting from the focus point region (horizontal region at $\sigma_p^{SI} \sim 10^{-8}$).

with all other parameters marginalized over. For comparison, we also show current CDMS-II [6], Edelweiss-I [7] and UKDMC ZEPLIN-I [8] 90% CL upper limits, but we stress that this constraint has not been used in our analysis.

The biggest, banana-shaped region of high probability (68% regions delimited by the internal solid, blue curve) shows a well-defined anticorrelation between σ_p^{SI} and m_χ . It results from two allowed regions in the CMSSM parameter space: the bulk and stau coannihilation region and from the A -resonance at large $\tan \beta$. This region covers roughly the range $10^{-10} \lesssim \sigma_p^{SI} \lesssim 10^{-8}$ pb

and $200 \lesssim m_\chi \lesssim 700$ GeV. In both cases the dominant contribution to σ_p^{SI} comes from a heavy Higgs exchange. At small $m_\chi \lesssim 100$ GeV we notice a small vertical band of fairly low probability density ($\lesssim 0.2$) at small σ_p^{SI} , a region where the light Higgs resonance contribute to reducing $\Omega_\chi h^2$ at small $m_{1/2}$ to meet the WMAP measurement. Finally, we can see a well pronounced region of high probability at fairly constant $\sigma_p^{SI} \sim 1.6 \times 10^{-8}$ pb for $m_\chi \lesssim 420$ GeV which at low m_χ partly overlaps with the previous region. At 95% this region extends up to $m_\chi \lesssim 720$ GeV for fairly constant σ_p^{SI} . This “high” σ_p^{SI} band results from the focus point (FP) region, basically independently of $\tan\beta$. This result has to be interpreted carefully, since there are large uncertainties associated with FP region, in particular with its location in the $(m_{1/2}, m_0)$ plane. Despite those outstanding questions, we believe that it is safe to expect that the FP will be the first to be probed by dark matter search experiments.

After marginalizing over all other parameters, we obtain the following 1-dimensional region encompassing 95% of the total probability:

$$0.5 \times 10^{-10} \text{ pb} < \sigma_p^{SI} < 3.2 \times 10^{-8} \text{ pb} \quad (95\% \text{ region}). \quad (3)$$

Currently running experiments (most notably CDMS-II but also Edelweiss-II and ZEPLIN-II) should be able to reach down to a few $\times 10^{-8}$ pb, on the edge of exploring this FP region. A future generation of “one-tonne” detectors is going to reach down to $\sigma_p^{SI} \gtrsim 10^{-10}$ pb, thus exploring much of the 95% interval.

We have checked that the above result is rather robust with respect to a change in the prior range used in the analysis or to the exclusion of the constrain coming from $g - 2$.

4. Conclusions

We have presented a detailed investigation of the prospects for dark matter detection in the framework of the CMSSM parameter using state-of-the-art Bayesian methods. We have shown that the WIMP dark matter direct detection elastic scattering cross section σ_p^{SI} presents a wide spread of values (below today’s limits) at around $10^{-9\pm 1}$ pb and a strong anticorrelation with m_χ . In addition, a region at relatively large $\sigma_p^{SI} \simeq 1.6 \times 10^{-8}$ pb, and fairly independent of m_χ , appears to be a feature of the focus point region (despite large theoretical uncertainties) and will be the first to be tested in direct detection experiments.

We conclude that a large fraction of the high-probability parameter space of neutralino dark matter in the CMSSM will be within reach of currently running upgraded dark matter detector, and most of it will be explored by future one-tonne detectors. Our result is thus highly suggestive of a possible detection by the next generation of direct dark matter searches.

Acknowledgements: R.T. is supported by the Royal Astronomical Society through the Sir Norman Lockyer Fellowship. R.Rda is supported by the program “Juan de la Cierva” of the Ministerio de Educación y Ciencia of Spain. R.T. would like to thank the organizers of the workshop for a very enjoyable and highly interesting meeting.

References

- [1] G. L. Kane, C. F. Kolda, L. Roszkowski and J. D. Wells, *Study of constrained minimal supersymmetry*, *Phys. Rev.* **D49** (1994) 6173.
- [2] R. Ruiz de Austri, R. Trotta and L. Roszkowski, *A Markov Chain Monte Carlo analysis of the CMSSM* (2006) *JHEP* **05** (2006) 002.
- [3] E. A. Baltz and P. Gondolo, *Markov chain monte carlo exploration of minimal supergravity with implications for dark matter*, *JHEP* **0410** (2004) 052.
- [4] B. C. Allanach and C. G. Lester, *Multi-dimensional MSUGRA likelihood maps*, arXiv:hep-ph/0507283. B. C. Allanach, *Naturalness priors and fits to the constrained minimal supersymmetric standard model*, arXiv:hep-ph/0601089.

- [5] Y. G. Kim, T. Nihei, L. Roszkowski and R. Ruiz de Austri, *Upper and lower limits on neutralino WIMP mass and spin-independent scattering cross section, and impact of new $(g - 2)_\mu$ measurement*, *JHEP* **0212** (2002) 034, .
- [6] D. S. Akerib, *et al* , [CDMS Collaboration], *First results from the cryogenic dark matter search in the Soudan underground lab.*, *Phys. Rev. Lett.* 932004211301.
- [7] V. Sanglard *et al* [EDELWEISS Collaboration], *Final results of the EDELWEISS-I dark matter search with cryogenic heat-and-ionization Ge detectors*, *Phys. Rev.* **D71** (2005) 122002.
- [8] G. J. Alner *et al* [UK Dark Matter Collaboration], *First limits on nuclear recoil events from the ZEPLIN-I galactic dark matter detector*, *Ann. Phys., Paris* 232005444.