



SCUOLA INTERNAZIONALE SUPERIORE DI STUDI AVANZATI

SISSA Digital Library

A Bethe Ansatz Type Formula for the Superconformal Index

Original

A Bethe Ansatz Type Formula for the Superconformal Index / Benini, F.; Milan, P.. - In: COMMUNICATIONS IN MATHEMATICAL PHYSICS. - ISSN 0010-3616. - 376:2(2020), pp. 1413-1440. [10.1007/s00220-019-03679-y]

Availability:

This version is available at: 20.500.11767/117582 since: 2021-01-15T20:04:51Z

Publisher:

Published

DOI:10.1007/s00220-019-03679-y

Terms of use:

Testo definito dall'ateneo relativo alle clausole di concessione d'uso

Publisher copyright
Springer

This version is available for education and non-commercial purposes.

note finali coverpage

(Article begins on next page)

A Bethe Ansatz type formula for the superconformal index

Francesco Benini^{1,2,3}, Paolo Milan^{1,2}

¹ *SISSA, Via Bonomea 265, 34136 Trieste, Italy*

² *INFN, Sezione di Trieste, Via Valerio 2, 34127 Trieste, Italy*

³ *ICTP, Strada Costiera 11, 34151 Trieste, Italy*

`fbenini@sissa.it`, `pmilan@sissa.it`

Abstract

Inspired by recent work by Closset, Kim and Willett, we derive a new formula for the superconformal (or supersymmetric) index of 4d $\mathcal{N} = 1$ theories. Such a formula is a finite sum, over the solution set of certain transcendental equations that we dub Bethe Ansatz Equations, of a function evaluated at those solutions.

Contents

1	Introduction and summary	1
2	The 4d superconformal index	4
3	A new Bethe Ansatz type formula	7
3.1	Continuation to generic fugacities	9
3.2	Proof of the formula	10
A	Numerical semigroups and the Fröbenius problem	19
B	A dense set	22
C	Weyl group fixed points	22
C.1	The rank-one case	23
C.2	The higher-rank case	26

1 Introduction and summary

In supersymmetric quantum field theories there are many classes of observables that can be computed exactly and non-perturbatively, making supersymmetry an appealing testing ground for general ideas about quantum field theory and, through holography, also quantum gravity. One of those observables is the superconformal index [1–3] which—in theories with superconformal invariance—counts with signs the number of local operators in short representations of the superconformal algebra. This counting can be done keeping track of the spin and other charges of the operators. Despite its simplicity, the superconformal index is an observable that contains a lot of information about the theory, and indeed it has been studied in all possible dimensions (*i.e.* up to six) and under so many angles (for reviews see [4]). In this note we focus on the four-dimensional superconformal index.

Because the index does not depend on continuous deformations of the theory, and a suitable supersymmetric generalization thereof does not depend on the RG flow, it follows that in theories that are part of a conformal manifold and have a weakly-coupled point on it, and in theories that are asymptotically free, the evaluation of the index can be reduced to a weak coupling computation.¹ This amounts to counting all possible local operators in short representations one can write down, and then restricting to the gauge-invariant ones. In the language of radial quantization, one counts all multi-particle states in short representations

¹There is a small caveat: the IR superconformal R-symmetry must be visible in the UV, *i.e.* it should not be accidental.

on the sphere, and then imposes Gauss law. In the case of the 4d $\mathcal{N} = 1$ superconformal (or supersymmetric) index of a gauge theory with gauge group G and matter chiral multiplets in representation \mathfrak{R} , the counting is captured by the standard formula [1, 2, 5]:

$$\mathcal{I}(p, q; v) = \frac{(p; p)_{\infty}^{\text{rk}(G)} (q; q)_{\infty}^{\text{rk}(G)}}{|\mathcal{W}_G|} \oint_{\mathbb{T}^{\text{rk}(G)}} \frac{\prod_{\rho_a \in \mathfrak{R}} \Gamma((pq)^{r_a/2} z^{\rho_a} v^{\omega_a}; p, q)}{\prod_{\alpha \in \Delta} \Gamma(z^{\alpha}; p, q)} \prod_{i=1}^{\text{rk}(G)} \frac{dz_i}{2\pi i z_i}. \quad (1.1)$$

Here, briefly, p, q are the (complex) fugacities associated to the angular momentum, v collectively indicates the fugacities for flavor symmetries, z indicates the fugacities for the gauge symmetry, r_a are the R-charges, and Γ is the elliptic gamma function. All the details will be reviewed in Section 2.

In this note, inspired by recent work of Closset, Kim and Willett [6, 7], we show that when the fugacities for the angular momentum satisfy

$$q^a = p^b \quad (1.2)$$

for some coprime positive integers a, b , then one can derive an alternative, very different formula for the 4d superconformal index. The condition (1.2) can be rewritten as

$$p = h^a, \quad q = h^b \quad (1.3)$$

for some fugacity h and coprime $a, b \in \mathbb{N}$.

The new formula, that we will explain in great detail in Section 3, is a finite sum over the solution set $\mathfrak{M}_{\text{BAE}}$ to certain transcendental equations—that we dub Bethe Ansatz equations (BAEs)—of a function, closely related to the integrand in (1.1), evaluated at those solutions. Very schematically, we prove that

$$\mathcal{I}(p, q; v) = \frac{(p; p)_{\infty}^{\text{rk}(G)} (q; q)_{\infty}^{\text{rk}(G)}}{|\mathcal{W}_G|} \sum_{z \in \mathfrak{M}_{\text{BAE}}} \sum_{\{m_i\}=1}^{ab} \mathcal{Z}(z/h^m, p, q, v) H(z, p, q, v)^{-1}. \quad (1.4)$$

Here the function \mathcal{Z} is the integrand in the standard formula (1.1); $\mathfrak{M}_{\text{BAE}}$ is the set of solutions, on a torus of exponentiated modular parameter h , to the BAEs which take the schematic form

$$Q_i(z, p, q, v) = 1 \quad \text{for } i = 1, \dots, \text{rk}(G) \quad (1.5)$$

in terms of functions Q_i defined in (3.3); the function H is a “Jacobian”

$$H(z, p, q, v) = \det_{ij} \frac{\partial Q_i(z, p, q, v)}{\partial \log z_j}. \quad (1.6)$$

The precise expressions (in which we use chemical potentials instead of fugacities, in order to deal with single-valued functions) can be found at the beginning of Section 3. A special case of this formula when $p = q$, namely $a = b = 1$, was derived in [6].

The condition (1.2) limits the applicability of the Bethe Ansatz (BA) formula (1.4) in the space of complex fugacities. Yet, as we discuss in Section 3.1, the domain of the formula is rich enough to uniquely fix the index as a continuous function (with poles) of general fugacities. We offer two arguments, one that uses holomorphy of the index and one that just uses continuity. Roughly, the reason is that the set of pairs (p, q) satisfying (1.2) is dense in the space of general complex fugacities (see Appendix B).

In a separate publication [8] we will use the BA formula (1.4) to address the large N limit of the index of a specific theory, namely $\mathcal{N} = 4$ Super-Yang-Mills, finding some differences with previous literature. We will connect the large N limit of the index to the entropy of BPS black holes in AdS₅.² In this way, we will extend the success of the counting of microstates of dyonic black hole in AdS₄ [11–14] to the case of electric rotating black holes in AdS₅ [15–18]. More generally, the new BA formula is much easier to deal with, compared to the standard integral formula, when performing numerical computations. We thus hope that it will be useful in a wider context.

The BA formula (1.4) can be thought of, in some sense, as the “Higgs branch localization” partner of the standard “Coulomb branch localization” integral formula (1.1), using the terminology of [19, 20]. More precisely, the existence of a formula as (1.4) can be justified along the lines of [11, 13, 21, 22, 6, 7]. The superconformal index can be defined as the partition function of the Euclidean theory on $S^3 \times S^1$, with suitable flat connections along S^1 and a suitable complex structure that depends on p, q , and with the Casimir energy [23, 24] stripped off.³ The standard localization computation of the partition function leads to (1.1). However, when p, q satisfy (1.2) the geometry is also a Seifert torus fibration over S^2 . Along the lines of [7], one expects to be able to reduce to the computation of a correlator in an A -twisted theory on S^2 [25], which should give an expression as in (1.4). In any case, we have derived the BA formula (1.4) by standard manipulations of the integral expression and thus we do not rely on any such putative 2d reduction.

The note is organized as follows. In Section 2 we review the standard formula for the 4d superconformal index, carefully stressing its regime of applicability. In Section 3 we present our new BA formula in great detail, and then we derive it in Section 3.2.

²The two recent papers [9, 10] also investigate the entropy of BPS black holes in AdS₅.

³Notice that the superconformal index, up to a change of variables reviewed in Section 3.1, is a single-valued function of the fugacities, while the partition function is not [24].

2 The 4d superconformal index

In order to fix our notation, let us review the standard formulation of the superconformal index [1, 2], which counts local operators in short representations of the 4d $\mathcal{N} = 1$ superconformal algebra (SCA) $\mathfrak{su}(2, 2|1)$. Going to radial quantization, this is the same as counting (with sign) $\frac{1}{4}$ -BPS states of the theory on S^3 .

The bosonic part of the superconformal algebra is $\mathfrak{su}(2, 2) \oplus \mathfrak{u}(1)_R$, where the first factor is the 4d conformal algebra and the second one is the R-symmetry. We pick on S^3 one Poincaré supercharge, specifically $\mathcal{Q} = \overline{Q}_-$, and its conjugate conformal supercharge $\mathcal{Q}^\dagger = S_+$. Together with $\Delta = \frac{1}{2}\{\mathcal{Q}, \mathcal{Q}^\dagger\}$ they form an $\mathfrak{su}(1|1)$ superalgebra. The superconformal index is then equal to the Witten index

$$\mathcal{I}(t) = \text{Tr}_{\mathcal{H}[S^3]} (-1)^F e^{-\beta\Delta} \prod_k t_k^{J_k}, \quad (2.1)$$

where J_k are Cartan generators of the commutant of $\mathfrak{su}(1|1)$ in the full SCA and t_k are the associated complex fugacities. By standard arguments [26], $\mathcal{I}(t)$ counts only states with $\Delta = 0$, *i.e.* annihilated by both \mathcal{Q} and \mathcal{Q}^\dagger , and thus it does not depend on β . On the other hand, it is holomorphic in the fugacities t_k , which serve both as regulators and as refinement parameters.

To be more precise, the states counted by (2.1) have $\Delta = E - 2j_+ - \frac{3}{2}r = 0$, where E is the conformal Hamiltonian or dimension, j_\pm are the Cartan generators of the angular momentum $\mathfrak{su}(2)_+ \oplus \mathfrak{su}(2)_- \subset \mathfrak{su}(2, 2)$, and r is the superconformal $U(1)_R$ charge. Moreover, the subalgebra of $\mathfrak{su}(2, 2|1)$ which commutes with $\mathfrak{su}(1|1)$ has Cartan generators $E + j_+$ and j_- . Therefore we write

$$\mathcal{I}(p, q) = \text{Tr}_{\Delta=0} (-1)^F p^{\frac{1}{3}(E+j_+)+j_-} q^{\frac{1}{3}(E+j_+)-j_-} = \text{Tr}_{\Delta=0} (-1)^F p^{j_1+\frac{r}{2}} q^{j_2+\frac{r}{2}}, \quad (2.2)$$

where $j_{1,2} = j_+ \pm j_-$ parametrize the rotated frame $\mathfrak{u}(1)_1 \oplus \mathfrak{u}(1)_2 \subset \mathfrak{su}(2)_+ \oplus \mathfrak{su}(2)_-$ and p, q are the associated fugacities (up to a shift by $r/2$). Whenever the theory enjoys flavor symmetries, one can introduce fugacities v_α for the Cartan generators of the flavor group. Then, the index will depend holomorphically also on v_α .

The trace formula (2.2) can be exactly evaluated at all regimes in the couplings. Indeed, since \mathcal{I} is invariant under any continuous deformation of the theory, one can explicitly account for the contribution of every gauge-invariant state with $\Delta = 0$ in the free regime [27, 28, 2]. In particular, the contributions of all the multi-particle states are simply encoded in the plethystic exponential [29] of the “single-letter partition functions”, whereas the restriction to the gauge-invariant sector is done by integrating the latter contributions over the gauge group. This procedure yields a finite-dimensional integral formula for the superconformal

index, which can be expressed as an elliptic hypergeometric integral [5].⁴

For concreteness, we consider a generic $\mathcal{N} = 1$ gauge theory with semi-simple gauge group G , flavor symmetry group G_F and non-anomalous $U(1)_R$ R-symmetry. We assume that the theory flows in the IR to a non-trivial fixed point and we parametrize $U(1)_R$ with the superconformal R-charge sitting in the SCA of the IR CFT (assuming this is visible in the UV). Furthermore, the matter content consists of n_χ chiral multiplets Φ_a in representations \mathfrak{R}_a of G , carrying flavor weights ω_a in some representations \mathfrak{R}_F of G_F and with superconformal R-charges r_a . Additionally, we turn on flavor fugacities v_α , with $\alpha = 1, \dots, \text{rk}(G_F)$, parametrizing the maximal torus of G_F . The integral representation of the superconformal index is given by

$$\mathcal{I}(p, q; v) = \frac{(p; p)_\infty^{\text{rk}(G)} (q; q)_\infty^{\text{rk}(G)}}{|\mathcal{W}_G|} \oint_{\mathbb{T}^{\text{rk}(G)}} \frac{\prod_{a=1}^{n_\chi} \prod_{\rho_a \in \mathfrak{R}_a} \Gamma((pq)^{r_a/2} z^{\rho_a} v^{\omega_a}; p, q)}{\prod_{\alpha \in \Delta} \Gamma(z^\alpha; p, q)} \prod_{i=1}^{\text{rk}(G)} \frac{dz_i}{2\pi i z_i}. \quad (2.3)$$

The integration variables z_i parametrize the maximal torus of G , and the integration contour is the product of $\text{rk}(G)$ unit circles. Then ρ_a are the weights of the representation \mathfrak{R}_a , α parametrizes the roots of G and $|\mathcal{W}_G|$ is the order of the Weyl group. Moreover, we have introduced the notation $z^{\rho_a} = \prod_{i=1}^{\text{rk}(G)} z_i^{\rho_a^i}$ and $v^{\omega_a} = \prod_{\alpha=1}^{\text{rk}(G_F)} v_\alpha^{\omega_a^\alpha}$, whereas

$$\Gamma(z; p, q) = \prod_{m,n=0}^{\infty} \frac{1 - p^{m+1} q^{n+1} / z}{1 - p^m q^n z}, \quad |p| < 1, \quad |q| < 1 \quad (2.4)$$

is the elliptic gamma function [33] and

$$(z; q)_\infty = \prod_{n=0}^{\infty} (1 - zq^n), \quad |q| < 1 \quad (2.5)$$

is the q -Pochhammer symbol.

This representation makes manifest the holomorphic dependence of the index on p, q, v_α . It is important to stress that the expression (2.3), which is a contour integral along $\text{rk}(G)$ unit circles, is only valid as long as the fugacities stay within the following

$$\text{Domain:} \quad |p|, |q| < 1, \quad |pq| < |(pq)^{r_a/2} v^{\omega_a}| < 1, \quad \forall a. \quad (2.6)$$

These conditions descend from the requirement of convergence of the plethystic representation of the index, from which (2.3) is derived. The plethystic expansion of the elliptic gamma

⁴An alternative way to obtain the integral formula is to use supersymmetric localization [30]. Indeed, the supersymmetric partition function Z of the theory on a primary Hopf surface $\mathcal{H}_{p,q} \simeq S^1 \times S^3$ can be computed with localization [31, 32] and it is related to the superconformal index through $Z = e^{-E_{\text{SUSY}} \mathcal{I}}$, where E_{SUSY} is the supersymmetric Casimir energy [23, 24].

function,

$$\Gamma(z; p, q) = \exp \left[\sum_{m=1}^{\infty} \frac{1}{m} \frac{z^m - (pq)^m z^{-m}}{(1-p^m)(1-q^m)} \right], \quad (2.7)$$

converges for

$$|pq| < |z| < 1 \quad \text{and} \quad |p|, |q| < 1. \quad (2.8)$$

The domain (2.6) then follows from requiring the integrand of (2.3) to have a convergent expansion. Indeed, within the domain of convergence (2.8), the elliptic gamma function is a single-valued analytic function with no zeros, poles nor branch cuts. Both $\Gamma(z; p, q)$ and $(z; q)_{\infty}$ can be analytically continued to $z \in \mathbb{C}$. However, when we analytically continue the integral (2.3) outside the domain (2.6), the integration contour must be continuously deformed in order to take into account the movement of the various poles of the integrand in the complex plane, in such a way that the poles do not cross the contour. As a result, for generic fugacities the integration contour is not as simple as a product of unit circles. To avoid this complication, throughout this paper we will always work within (2.6)—and perform analytic continuation only at the end, if needed.

It will be useful to set some new notation. We define a set of chemical potentials through

$$p = e^{2\pi i \tau}, \quad q = e^{2\pi i \sigma}, \quad v_{\alpha} = e^{2\pi i \xi_{\alpha}}, \quad z_i = e^{2\pi i u_i}, \quad (2.9)$$

as well as a fictitious chemical potential ν_R for the R-symmetry, whose value is fixed to

$$\nu_R = \frac{1}{2}(\tau + \sigma) \quad (2.10)$$

by supersymmetry. Moreover, we redefine the elliptic gamma function as a (periodic) function of the chemical potentials:

$$\tilde{\Gamma}(u, \tau, \sigma) = \Gamma(e^{2\pi i u}; e^{2\pi i \tau}, e^{2\pi i \sigma}), \quad (2.11)$$

so that the integrand of (2.3) can be expressed as

$$\mathcal{Z}(u; \xi, \nu_R, \tau, \sigma) = \frac{\prod_{a=1}^{n_{\chi}} \prod_{\rho_a \in \mathfrak{R}_a} \tilde{\Gamma}(\rho_a(u) + \omega_a(\xi) + r_a \nu_R; \tau, \sigma)}{\prod_{\alpha \in \Delta} \tilde{\Gamma}(\alpha(u); \tau, \sigma)}. \quad (2.12)$$

At last, we define

$$\kappa_G = \frac{(p; p)_{\infty}^{\text{rk}(G)} (q; q)_{\infty}^{\text{rk}(G)}}{|\mathcal{W}_G|}. \quad (2.13)$$

The integral representation of the index takes then the following compact form:

$$\mathcal{I}(p, q; v) = \kappa_G \int_{\mathbb{T}^{\text{rk}(G)}} \mathcal{Z}(u; \xi, \nu_R, \tau, \sigma) d^{\text{rk}(G)} u. \quad (2.14)$$

The integration contour $\mathbb{T}^{\text{rk}(G)}$ is represented on the u -plane by a product of straight segments of length one on the real axes. In terms of the chemical potentials, the domain (2.6) can be rewritten as:

$$\text{Im } \tau, \text{Im } \sigma > 0, \quad 0 < \text{Im } \omega_a(\xi) < \text{Im}(\tau + \sigma), \quad \forall a. \quad (2.15)$$

The integral formula (2.14) is the starting point of our analysis. In the next Section we will focus our attention to the case where τ/σ is a rational number to derive—from (2.14)—a new formula that expresses the index as a finite sum.

3 A new Bethe Ansatz type formula

The integral representation (2.14) of the superconformal index is valid for generic complex values of the chemical potentials within the domain (2.15). However, if we restrict to a case where

$$\tau/\sigma \in \mathbb{Q}_+, \quad (3.1)$$

we can prove an alternative formula describing the index as a finite sum over the set of solutions to certain transcendental equations, which we call *Bethe Ansatz Equations* (BAEs). We will first present the formula in detail, and then provide a proof. In Section 3.1 we will also discuss the properties of the set of pairs (τ, σ) satisfying (3.1).

Let us take

$$\tau = a\omega, \quad \sigma = b\omega \quad \text{with } a, b \in \mathbb{N} \text{ such that } \gcd(a, b) = 1 \quad (3.2)$$

and $\text{Im } \omega > 0$. This implies (1.2). We can set $p = h^a$ and $q = h^b$ with $h = e^{2\pi i \omega}$, although we will mostly work with chemical potentials. We introduce the BAEs as the set of equations

$$Q_i(u; \xi, \nu_R, \omega) = 1, \quad \forall i = 1, \dots, \text{rk}(G), \quad (3.3)$$

written in terms of “BA operators” defined as

$$Q_i(u; \xi, \nu_R, \omega) = \prod_{a=1}^{n_\chi} \prod_{\rho_a \in \mathfrak{R}_a} P(\rho_a(u) + \omega_a(\xi) + r_a \nu_R; \omega)^{\rho_a^i}. \quad (3.4)$$

The basic BA operator is

$$P(u; \omega) = \frac{e^{-\pi i \frac{u^2}{\omega} + \pi i u}}{\theta_0(u; \omega)}, \quad (3.5)$$

where $\theta_0(u; \omega) = (z; h)_\infty (z^{-1}h; h)_\infty$ with $z = e^{2\pi i u}$ and $h = e^{2\pi i \omega}$.

The BA operators satisfy three important properties. First, they are doubly-periodic in the gauge chemical potentials:

$$Q_i(u + n + m\omega; \xi, \nu_R, \omega) = Q_i(u; \xi, \nu_R, \omega), \quad \forall n_i, m_i \in \mathbb{Z}, \quad i = 1, \dots, \text{rk}(G). \quad (3.6)$$

Second, they are invariant under $SL(2, \mathbb{Z})$ modular transformations of ω :

$$Q_i(u; \xi, \nu_R, \omega) = Q_i(u; \xi, \nu_R, \omega + 1) = Q_i\left(\frac{u}{\omega}; \frac{\xi}{\omega}, \frac{\nu_R}{\omega}, -\frac{1}{\omega}\right) = Q_i(-u; -\xi, -\nu_R, \omega). \quad (3.7)$$

The last equality represents invariance under the center of $SL(2, \mathbb{Z})$. Third, they capture the quasi-periodicity of the index integrand:

$$Q_i(u; \xi, \nu_R, \omega) \mathcal{Z}(u; \xi, \nu_R, a\omega, b\omega) = \mathcal{Z}(u - \delta_i ab\omega; \xi, \nu_R, a\omega, b\omega), \quad (3.8)$$

valid $\forall i$ and where $\delta_i = (\delta_{ij})_{j=1}^{\text{rk}(G)}$ so that $(u - \delta_i ab\omega)_j = u_j - \delta_{ij} ab\omega$.

Because of the double-periodicity of Q_i , the actual number of solutions \hat{u}_i to the system of BAEs (3.3) is infinite. However, the solutions can be grouped into a finite number of equivalence classes $[\hat{u}_i]$ such that $\hat{u}_i \sim \hat{u}_i + 1 \sim \hat{u}_i + \omega$. In other words, the equations and their solutions are well-defined on a torus $\mathbb{T}^{2\text{rk}(G)}$ which is the product of $\text{rk}(G)$ identical complex tori of modular parameter ω , and the number of solutions on the torus is finite. The modular invariance (3.7) confirms that the equations are well-defined on the torus. We define

$$\mathfrak{M}_{\text{BAE}} = \left\{ [\hat{u}_i], i = 1, \dots, \text{rk}(G) \mid Q_i([\hat{u}]; \xi, \nu_R, \omega) = 1, \quad w \cdot [\hat{u}] \neq [\hat{u}] \quad \forall w \in \mathcal{W}_G \right\} \quad (3.9)$$

as the set of solutions (on the torus) that are not fixed by non-trivial elements of the Weyl group. For definiteness we can choose, as representatives, the elements living in a fundamental domain of the torus with modulus ω , *i.e.* with $0 \leq \text{Re } \hat{u}_i < 1$ and $0 \leq \text{Im } \hat{u}_i < \text{Im } \omega$. Notice that, because of (3.7), the solutions must organize into representations of $SL(2, \mathbb{Z})$.

As we prove below, thanks to the properties of the BA operators, we can rewrite the superconformal index as a sum over solutions to the BAEs in the following way:

$$\mathcal{I}(p, q; v) = \kappa_G \sum_{\hat{u} \in \mathfrak{M}_{\text{BAE}}} \mathcal{Z}_{\text{tot}}(\hat{u}; \xi, \nu_R, a\omega, b\omega) H(\hat{u}; \xi, \nu_R, \omega)^{-1}. \quad (3.10)$$

Here

$$\mathcal{Z}_{\text{tot}}(u; \xi, \nu_R, a\omega, b\omega) = \sum_{\{m_i\}=1}^{ab} \mathcal{Z}(u - m\omega; \xi, \nu_R, a\omega, b\omega), \quad (3.11)$$

where \mathcal{Z} is precisely the integrand defined in (2.12) and

$$H(u; \xi, \nu_R, \omega) = \det_{ij} \left[\frac{1}{2\pi i} \frac{\partial Q_i(u; \xi, \nu_R, \omega)}{\partial u_j} \right] \quad (3.12)$$

is the contribution from the Jacobian of the change of variables $u_i \mapsto Q_i(u)$. Notice that both the function H , and the function \mathcal{Z}_{tot} evaluated on the solutions to the BAEs, are doubly-periodic on the product of complex tori of modular parameter ω .

A specialization of this formula to the case $\tau = \sigma$ was derived in [6], while a three-dimensional analog was derived in [7]. In the next Section we will spell out in detail how the BA formula uniquely fixes the index for all values of the complex fugacities, using either holomorphy or continuity. In Section 3.2 we will derive the final formula (3.10), starting from the integral representation (2.14). The proof is rather technical and it does not give new physical insights on the main result. Therefore, uninterested readers may stop here.

3.1 Continuation to generic fugacities

Our BA formula (3.10) can only be applied for special values of the angular fugacities that satisfy (1.2). We will offer two arguments, one based on holomorphy and the other based on just continuity, that this is enough to completely determine the index for all values of the complex fugacities.

Using the standard definition (2.2), the index is not a single-valued function of the angular fugacities p, q —unless the R-charges of chiral multiplets are all even. This is also apparent from the integral formula (2.3). On the other hand, regarded as a function of chemical potentials τ, σ each living on the upper half-plane \mathbb{H} , the index is single-valued and holomorphic. Keeping the flavor fugacities fixed in the argument that follows, the BA formula applies to points $(\tau, \sigma) \in \mathbb{H}^2$ such that $\tau/\sigma \in \mathbb{Q}_+$. Such a set is dense in a hyperplane $\mathcal{J} \cong \mathbb{R}^3$ of real codimension one in \mathbb{H}^2 defined as $\mathcal{J} = \{(\tau, \sigma) \mid \tau/\sigma \in \mathbb{R}_+\}$. Thus, the BA formula determines the index on \mathcal{J} by continuity. On the other hand, we know that the index is a holomorphic function on \mathbb{H}^2 , therefore its restriction to \mathcal{J} completely fixes the function on \mathbb{H}^2 by analytic continuation.

It turns out that we can refine the argument in such a way that we only use continuity, and not holomorphy, of the index. This is because if we think in terms of angular fugacities p, q each living in the open unit disk \mathbb{D} , then the set of points $(p, q) \in \mathbb{D}^2$ such that $q^a = p^b$ for coprime $a, b \in \mathbb{N}$ is dense in \mathbb{D}^2 . This fact is not completely obvious, and we show it in Appendix B.

Unfortunately, the index (2.3) is not a single-valued function of p, q if we keep the flavor fugacities v_α fixed, unless the R-charges are all even. However, it is always possible to find a change of variables which expresses \mathcal{I} as a single-valued function of a set of new fugacities.

The latter is defined by

$$\Delta_a = \omega_a(\xi) + r_a \nu_R \quad \Rightarrow \quad y_a = e^{2\pi i \Delta_a} = v^{\omega_a} (pq)^{\frac{r_a}{2}}, \quad \forall a = 1, \dots, n_\chi. \quad (3.13)$$

This gives us a set of (redundant) chemical potentials Δ_a , one for each chiral multiplet present in the theory, which must satisfy some linear constraint, following the requirement of invariance of the theory under flavor and R-symmetry. Suppose, indeed, the theory has a superpotential given by

$$W(\Phi) = \sum_A W_A(\Phi), \quad (3.14)$$

where each $W_A(\Phi)$ is a gauge-invariant homogeneous polynomial of degree n_A . Then, for each term in (3.14), the following linear constraints must be satisfied:

$$\sum_{a \in A} r_a = 2, \quad \sum_{a \in A} \omega_a^\alpha = 0, \quad \forall \alpha = 1, \dots, \text{rk}(G), \quad (3.15)$$

where we used $a \in A$ to indicate the chiral components Φ_a which are present in W_A . The first equation imposes that the superpotential has R-charge 2. The second equation constrains W to be invariant under G_F . Indeed, $\omega_a = (\omega_a^\alpha)_{\alpha=1}^{\text{rk}(G_F)}$ are the flavor weights carried by Φ_a . A similar role is played by ABJ anomalies.

Translating (3.15) to the definition of Δ_a , we obtain

$$\sum_{a \in A} \Delta_a = 2\nu_R = \tau + \sigma \quad \forall A. \quad (3.16)$$

In such a new set of variables we have

$$\mathcal{Z}(u; \Delta, \tau, \sigma) = \frac{\prod_{a=1}^{n_\chi} \prod_{\rho_a \in \mathfrak{R}_a} \tilde{\Gamma}(\rho_a(u) + \Delta_a; \tau, \sigma)}{\prod_{\alpha \in \Delta} \tilde{\Gamma}(\alpha(u); \tau, \sigma)}, \quad (3.17)$$

showing that the index is now a well-defined, single-valued and continuous function (in fact, also holomorphic) of the fugacities p, q, y_a . Indeed, recall that the elliptic gamma function is a single-valued function of its arguments, and notice that the constraints (3.16) always involve integer combinations of τ, σ , thus never introducing non-trivial monodromies under integer shifts. Once again, the BA formula can be applied whenever $q^a = p^b$ and for generic values of y_a . Since such a set of points is dense in the space of generic fugacities, we conclude that the BA formula fixes the index completely.

3.2 Proof of the formula

We prove the formula (3.10) in three steps. First we verify the properties (3.6) and (3.8) of the BA operators. Then we use them to modify the contour of the integral (2.14) and to reduce it to a sum of simple residues. Finally we prove that the only poles that contribute to the residue formula are determined by the BAEs, thus obtaining (3.10).

3.2.1 Properties of the BA operators

First, we prove the identities (3.6) and (3.8). For later convenience, let us briefly recall the anomaly cancellation conditions that are required to have a well-defined four-dimensional theory. These requirements can be expressed in terms of the anomaly coefficients. In particular, let $\mathbf{i} = (i, \alpha)$ collectively denote the Cartan indices of the gauge \times flavor group, where $i = 1, \dots, \text{rk}(G)$ are the gauge indices and $\alpha = 1, \dots, \text{rk}(G_F)$ are the flavor indices. Moreover, define $\mathbf{a} = (a, \rho_a)$ as running over all chiral multiplets components, where ρ_a are the weights of the gauge representation \mathfrak{R}_a . Then the anomaly coefficients for gauge/flavor symmetries are defined by

$$\mathcal{A}^{ijk} = \sum_{\mathbf{a}} Q_{\mathbf{a}}^i Q_{\mathbf{a}}^j Q_{\mathbf{a}}^k, \quad \mathcal{A}^{ij} = \sum_{\mathbf{a}} Q_{\mathbf{a}}^i Q_{\mathbf{a}}^j, \quad \mathcal{A}^i = \sum_{\mathbf{a}} Q_{\mathbf{a}}^i, \quad (3.18)$$

where $Q_{\mathbf{a}}^i = Q_{(a, \rho_a)}^i = (\rho_a^i, \omega_a^\alpha)$ are the components of the gauge \times flavor weights carried by the chiral multiplets. The first and the last coefficient in (3.18) are associated with the gauge³ and mixed gauge-gravitational² perturbative anomalies. The second term—sometimes called pseudo-anomaly coefficient—describes the non-perturbative or global anomaly [34–36] when the corresponding perturbative anomaly vanishes.

Similarly, the perturbative anomaly coefficients involving the R-symmetry are defined by

$$\begin{aligned} \mathcal{A}^{ijR} &= \sum_{\mathbf{a}} Q_{\mathbf{a}}^i Q_{\mathbf{a}}^j (r_{\mathbf{a}} - 1) + \delta^{ij, ij} \sum_{\alpha \in \Delta} \alpha^i \alpha^j & \mathcal{A}^{iRR} &= \sum_{\mathbf{a}} Q_{\mathbf{a}}^i (r_{\mathbf{a}} - 1)^2 \\ \mathcal{A}^{RRR} &= \sum_{\mathbf{a}} (r_{\mathbf{a}} - 1)^3 + \dim G & \mathcal{A}^R &= \sum_{\mathbf{a}} (r_{\mathbf{a}} - 1) + \dim G, \end{aligned} \quad (3.19)$$

whereas the pseudo R-anomaly coefficients are

$$\mathcal{A}^{iR} = \sum_{\mathbf{a}} Q_{\mathbf{a}}^i (r_{\mathbf{a}} - 1) \quad \mathcal{A}^{RR} = \sum_{\mathbf{a}} (r_{\mathbf{a}} - 1)^2 + \dim G. \quad (3.20)$$

Anomaly cancellation is realized by a set of conditions on the coefficients defined above, that a well-defined quantum gauge theory must satisfy. We will also restrict to the case that the gauge group G is semi-simple. The conditions for the cancellation of the gauge and gravitational anomaly are

$$\mathcal{A}^{ijk} = \mathcal{A}^i = 0 \quad \text{and} \quad \mathcal{A}^{ij} \in 4\mathbb{Z} \quad \text{for } G \text{ semi-simple}. \quad (3.21)$$

The conditions for the cancellation of the ABJ anomalies of G_F and $U(1)_R$, namely that those are global symmetries of the quantum theory, are

$$\mathcal{A}^{ij\alpha} = \mathcal{A}^{ijR} = 0. \quad (3.22)$$

Finally,

$$\mathcal{A}^{i\alpha\beta} = \mathcal{A}^{i\alpha R} = \mathcal{A}^{iRR} = 0 \quad \text{and} \quad \mathcal{A}^{i\alpha} = \mathcal{A}^{iR} = 0 \quad (3.23)$$

simply follow from the restriction to semi-simple gauge group G .

We now focus on describing some properties of the basic BA operator

$$P(u; \omega) = \frac{e^{-\pi i \frac{u^2}{\omega} + \pi i u}}{\theta_0(u; \omega)}. \quad (3.24)$$

First, consider the function

$$\theta_0(u; \omega) = (z; h)_\infty (z^{-1}h; h)_\infty = \prod_{k=0}^{\infty} (1 - zh^k)(1 - z^{-1}h^{k+1}), \quad z = e^{2\pi i u}, \quad h = e^{2\pi i \omega} \quad (3.25)$$

which is holomorphic in z and h , and satisfies the following properties:

$$\begin{aligned} \theta_0(u + n + m\omega; \omega) &= (-1)^m e^{-2\pi i m u - \pi i m(m-1)\omega} \theta_0(u; \omega) \quad \forall n, m \in \mathbb{Z} \\ \theta_0(-u; \omega) &= \theta_0(u + \omega; \omega) = -e^{-2\pi i u} \theta_0(u; \omega). \end{aligned} \quad (3.26)$$

They immediately imply

$$\begin{aligned} P(-u; \omega) &= -P(u; \omega) \\ P(u + n + m\omega; \omega) &= (-1)^{n+m} e^{-\frac{\pi i}{\omega}(2nu+n^2)} P(u; \omega) \quad \forall n, m \in \mathbb{Z}. \end{aligned} \quad (3.27)$$

It turns out that the basic BA operator has also nice modular transformation properties:

$$P(u; \omega + 1) = e^{\pi i \frac{u^2}{\omega(\omega+1)}} P(u; \omega), \quad P\left(\frac{u}{\omega}; -\frac{1}{\omega}\right) = e^{\pi i \left(\frac{u^2}{\omega} - \frac{u}{6} - \frac{1}{6\omega} + \frac{1}{2}\right)} P(u; \omega). \quad (3.28)$$

In order to prove (3.8), we also need to show that

$$\begin{aligned} P(u + r\nu_R; \omega)^m \tilde{\Gamma}(u + r\nu_R; a\omega, b\omega) &= (-1)^{\frac{abm^2}{2} + \frac{m(a+b-1)}{2}} e^{-\frac{\pi i m u^2}{\omega} + \pi i abm^2 u - \pi i m(a+b)(r-1)u} \times \\ &\times h^{-\frac{m^3 ab}{6} + \frac{ab(a+b)m^2(r-1)}{4} - \frac{m(a+b)^2(r-1)^2}{8} + \frac{m(a^2+b^2+2)}{24}} \tilde{\Gamma}(u + r\nu_R - mab\omega; a\omega, b\omega). \end{aligned} \quad (3.29)$$

Here $r \in \mathbb{R}$ mimics the contribution from the R-charge of a generic multiplet in the theory. Notice that all factors in front of $\tilde{\Gamma}$ in the r.h.s. of (3.29) explicitly depend on the fermion R-charge $r - 1$. This will be crucial to ensure anomaly cancellation in the full BA operator.

Proof. The identity (3.29) follows from the properties of the elliptic gamma function. Indeed, for generic τ and σ , we have that

$$\tilde{\Gamma}(u + \tau; \tau, \sigma) = \theta_0(u; \sigma) \tilde{\Gamma}(u; \tau, \sigma), \quad \tilde{\Gamma}(u + \sigma; \tau, \sigma) = \theta_0(u; \tau) \tilde{\Gamma}(u; \tau, \sigma). \quad (3.30)$$

Moreover, there exists a factorization property (see Theorem 5.4 of [33]) which expresses $\tilde{\Gamma}(u; a\omega, b\omega)$ as a product of elliptic gamma functions with equal periods:

$$\tilde{\Gamma}(u; a\omega, b\omega) = \prod_{r=0}^{a-1} \prod_{s=0}^{b-1} \tilde{\Gamma}(u + (as + br)\omega; ab\omega, ab\omega), \quad (3.31)$$

valid for $a, b \in \mathbb{Z}$ (not necessarily coprime). Using both (3.30) and (3.31) we obtain the identity

$$\tilde{\Gamma}(u + ab\omega; a\omega, b\omega) = \left[\prod_{r=0}^{a-1} \prod_{s=0}^{b-1} \theta_0(u + (as + br)\omega; ab\omega) \right] \times \tilde{\Gamma}(u; a\omega, b\omega) \quad (3.32)$$

and its generalizations to $m \in \mathbb{Z}$, given by

$$\begin{aligned} \tilde{\Gamma}(u + mab\omega; a\omega, b\omega) &= (-z)^{-\frac{abm(m-1)}{2}} h^{-\frac{m(m-1)}{2} \frac{ab(2ab-a-b)}{2} - \frac{m(m-1)(m-2)a^2b^2}{6}} \times \\ &\times \left[\prod_{r=0}^{a-1} \prod_{s=0}^{b-1} \theta_0(u + (as + br)\omega; ab\omega)^m \right] \times \tilde{\Gamma}(u; a\omega, b\omega). \end{aligned} \quad (3.33)$$

Now, by enforcing the assumption that $\gcd(a, b) = 1$, we can use the properties of numerical semigroups (see Appendix A for more details) to reduce the periods of the theta functions from $ab\omega$ to ω . In order to do so, let us introduce some notation. We call $\mathcal{R}(a, b)$ the set of non-negative integer linear combinations of a, b :

$$\mathcal{R}(a, b) = \{am + bn \mid m, n \in \mathbb{Z}_{\geq 0}\}. \quad (3.34)$$

Then $\mathcal{R}(a, b)$ forms a numerical semigroup, which can be thought of as a subset of $\mathbb{Z}_{\geq 0}$, closed under addition, with only a finite number of excluded non-vanishing elements. The latter elements form the so-called set of gaps $\overline{\mathcal{R}}(a, b) = \mathbb{N} \setminus \mathcal{R}(a, b)$. The highest element of $\overline{\mathcal{R}}(a, b)$ is the Fröbenius number $F(a, b) = ab - a - b$, whereas the order of $\overline{\mathcal{R}}(a, b)$ is called the genus $\chi(a, b)$ and the sum of all its elements is the weight $w(a, b)$. It is a classic result in mathematics that, in terms of a, b , the latter read

$$\chi(a, b) = \frac{(a-1)(b-1)}{2}, \quad w(a, b) = \frac{(a-1)(b-1)(2ab-a-b-1)}{12}. \quad (3.35)$$

Thanks to the properties of these objects, we can use the following identities (proved in Appendix A):

$$\begin{aligned} \prod_{r=0}^{a-1} \prod_{s=0}^{b-1} (zh^{as+br}; h^{ab})_{\infty} &= \frac{(z; h)_{\infty}}{\prod_{k \in \overline{\mathcal{R}}(a, b)} (1 - zh^k)} \\ \prod_{r=0}^{a-1} \prod_{s=0}^{b-1} (z^{-1}h^{ab-as-br}; h^{ab})_{\infty} &= (z^{-1}h; h)_{\infty} \prod_{k \in \overline{\mathcal{R}}(a, b)} (1 - z^{-1}h^{-k}), \end{aligned} \quad (3.36)$$

which lead to

$$\prod_{r=0}^{a-1} \prod_{s=0}^{b-1} \theta_0(u + (as + br)\omega; \omega) = (-z)^{-\chi(a,b)} h^{-w(a,b)} \theta_0(u; \omega). \quad (3.37)$$

Substituting into (3.33) we obtain

$$\begin{aligned} \tilde{\Gamma}(u + mab\omega; a\omega, b\omega) &= (-z)^{-\frac{abm^2}{2} + \frac{m(a+b-1)}{2}} \times \\ &\times h^{-\frac{abm^3}{6} + \frac{ab(a+b)m^2}{4} - \frac{(a^2+b^2+3ab-1)m}{12}} \theta_0(u; \omega)^m \tilde{\Gamma}(u; a\omega, b\omega). \end{aligned} \quad (3.38)$$

Finally, applying (3.38) to the l.h.s. of (3.29) proves the latter identity. \square

We now turn to analyzing the full BA operators. Notice that, in the definition (3.4), Q_i receive contribution only from the chiral multiplets of the theory. The vector multiplets do not appear in (3.4) because their contribution simply amounts to

$$\prod_{\alpha \in \Delta} P(\alpha(u); \omega)^{-\alpha^i} = \prod_{\alpha > 0} \left[\frac{P(-\alpha(u); \omega)}{P(\alpha(u); \omega)} \right]^{\alpha^i} = (-1)^{\sum_{\alpha > 0} \alpha^i} = 1, \quad (3.39)$$

which holds true if G is semi-simple, as in this case the sum of positive roots is always an even integer. Despite this fact, as far as the proof of (3.10) is concerned, we find it more convenient to write

$$Q_i(u; \xi, \nu_R, \omega) = \prod_{a=1}^{n_\chi} \prod_{\rho_a \in \mathfrak{R}_a} P(\rho_a(u) + \omega_a(\xi) + r_a \nu_R; \omega)^{\rho_a^i} \times \prod_{\alpha \in \Delta} P(\alpha(u); \omega)^{-\alpha^i} \quad (3.40)$$

without simplifying the vector multiplet contribution.

At this point, using (3.27) we can show that Q_i satisfy:

$$\begin{aligned} Q_i(u + n; \xi, \nu_R, \omega) &= (-1)^{\mathcal{A}^{ij} n_j} e^{-\frac{\pi i}{\omega} (\mathcal{A}^{ijk} n_j (2u_k + n_k) + 2\mathcal{A}^{ij\alpha} n_j \xi_\alpha + 2\mathcal{A}^{ijR} n_j \omega)} Q_i(u; \xi, \nu_R, \omega) \\ Q_i(u + m\omega; \xi, \nu_R, \omega) &= (-1)^{\mathcal{A}^{ij} m_j} Q_i(u; \xi, \nu_R, \omega), \end{aligned} \quad (3.41)$$

which, thanks to (3.21)–(3.23), reduce to (3.6) $\forall n_i, m_i \in \mathbb{Z}$ in an anomaly-free theory. Similarly, (3.27) and (3.28) together with the anomaly cancelation conditions (3.21)–(3.23) imply (3.7). Moreover, using (3.40), we can write

$$\begin{aligned} Q_i(u; \xi, \nu_R, \omega) \mathcal{Z}(u; \xi, \nu_R, a\omega, b\omega) &= \\ &= \frac{\prod_{a, \rho_a} P(\rho_a(u) + \omega_a(\xi) + r_a \nu_R; \omega)^{\rho_a^i} \tilde{\Gamma}(\rho_a(u) + \Delta_a; a\omega, b\omega)}{\prod_{\alpha \in \Delta} P(\alpha(u); \omega)^{\alpha^i} \tilde{\Gamma}(\alpha(u); a\omega, b\omega)}. \end{aligned} \quad (3.42)$$

Applying (3.29), the latter equation reduces to

$$\begin{aligned}
Q_i(u; \xi, \nu_R, \omega) \mathcal{Z}(u; \xi, \nu_R, a\omega, b\omega) &= (-1)^{\frac{ab}{2} \mathcal{A}^{ii} + \frac{a+b-1}{2} \mathcal{A}^i} e^{\pi i ab (\mathcal{A}^{ij} u_j + \mathcal{A}^{i\alpha} \xi_\alpha)} \times \\
&\times e^{-\frac{\pi i}{\omega} (\mathcal{A}^{ijk} u_j u_k + \mathcal{A}^{i\alpha\beta} \xi_\alpha \xi_\beta + 2\mathcal{A}^{ij\alpha} u_j \xi_\alpha)} e^{-\pi i (a+b) (\mathcal{A}^{ijR} u_j + \mathcal{A}^{i\alpha R} \xi_\alpha) + \frac{\pi i ab (a+b)}{2} \mathcal{A}^{iiR} \omega} \times \\
&\times e^{-\frac{\pi i (a+b)^2}{4} \mathcal{A}^{iRR} \omega - \frac{\pi i a^2 b^2}{3} \mathcal{A}^{iii} \omega + \frac{\pi i (a^2 + b^2)}{12} \mathcal{A}^i \omega} \mathcal{Z}(u - \delta_i ab\omega; \xi, \nu_R, \omega), \quad (3.43)
\end{aligned}$$

which, by anomaly cancellation, reduces to (3.8).

3.2.2 Residue formula

We now use the BA operators and their properties to modify the contour of integration of the index in (2.14). For our purposes, it is sufficient to implement the following trivial relation:

$$\begin{aligned}
\mathcal{I}(p, q; v) &= \kappa_G \oint \mathcal{Z}(u; \xi, \nu_R, a\omega, b\omega) d^{\text{rk}(G)} u \\
&= \kappa_G \oint \frac{\prod_{i=1}^{\text{rk}(G)} (1 - Q_i(u; \xi, \nu_R, \omega))}{\prod_{i=1}^{\text{rk}(G)} (1 - Q_i(u; \xi, \nu_R, \omega))} \mathcal{Z}(u; \xi, \nu_R, a\omega, b\omega) d^{\text{rk}(G)} u. \quad (3.44)
\end{aligned}$$

The numerator of the integrand can be expanded as

$$\begin{aligned}
\prod_{i=1}^{\text{rk}(G)} (1 - Q_i(u; \xi, \nu_R, \omega)) \times \mathcal{Z}(u; \xi, \nu_R, a\omega, b\omega) &= \\
&= \sum_{n=0}^{\text{rk}(G)} \frac{(-1)^n}{n!} \sum_{i_1 \neq \dots \neq i_n}^{\text{rk}(G)} Q_{i_1}(u; \xi, \nu_R, \omega) \dots Q_{i_n}(u; \xi, \nu_R, \omega) \mathcal{Z}(u; \xi, \nu_R, a\omega, b\omega) \quad (3.45) \\
&= \sum_{n=0}^{\text{rk}(G)} \frac{(-1)^n}{n!} \sum_{i_1 \neq \dots \neq i_n}^{\text{rk}(G)} \mathcal{Z}(u - (\delta_{i_1} + \dots + \delta_{i_n}) ab\omega; \xi, \nu_R, a\omega, b\omega),
\end{aligned}$$

where, in the last line, we have used the shift property (3.8). Plugging the last equation back in (3.44) gives:

$$\mathcal{I}(p, q; v) = \kappa_G \sum_{n=0}^{\text{rk}(G)} \frac{(-1)^n}{n!} \sum_{i_1 \neq \dots \neq i_n}^{\text{rk}(G)} I_{i_1 \dots i_n}(p, q; v), \quad (3.46)$$

with

$$\begin{aligned}
I_{i_1 \dots i_n}(p, q; v) &= \oint_{\mathcal{T}^{\text{rk}(G)}} \frac{\mathcal{Z}(u - (\delta_{i_1} + \dots + \delta_{i_n}) ab\omega; \xi, \nu_R, a\omega, b\omega)}{\prod_{i=1}^{\text{rk}(G)} (1 - Q_i(u; \xi, \nu_R, \omega))} d^{\text{rk}(G)} u \\
&= \oint_{\mathcal{C}_{i_1 \dots i_n}} \frac{\mathcal{Z}(u; \xi, \nu_R, a\omega, b\omega)}{\prod_{i=1}^{\text{rk}(G)} (1 - Q_i(u; \xi, \nu_R, \omega))} d^{\text{rk}(G)} u \quad (3.47)
\end{aligned}$$

and where

$$\mathcal{C}_{i_1 \dots i_n} = \mathbb{T}^{\text{rk}(G)-n} \times \bigcup_{k=1}^n \{|z_{i_k}| = |h|^{-ab}; \circlearrowleft\}. \quad (3.48)$$

This is a contour where z_{i_1}, \dots, z_{i_n} live on circles of radius $|h|^{-ab}$, whereas the other variables z_j parametrize the unit circles in $\mathbb{T}^{\text{rk}(G)-n}$. The second line in (3.47) has been obtained by implementing the change of variables $u_{i_k} \mapsto u_{i_k} + ab\omega$ for $k = 1, \dots, n$ and using the periodicity (3.6).

The series of integrals in (3.46) can be resummed to a unique integral over a composite contour:

$$\mathcal{I}(p, q; v) = \kappa_G \oint_{\mathcal{C}} \frac{\mathcal{Z}(u; \xi, \nu_R, a\omega, b\omega)}{\prod_{i=1}^{\text{rk}(G)} (1 - Q_i(u; \xi, \nu_R, \omega))} d^{\text{rk}(G)} u, \quad (3.49)$$

where

$$\mathcal{C} = \sum_{n=0}^{\text{rk}(G)} \frac{(-1)^n}{n!} \bigcup_{i_1 \neq \dots \neq i_n}^{\text{rk}(G)} \mathcal{C}_{i_1 \dots i_n} \simeq \bigcup_{i=1}^{\text{rk}(G)} \{|z_i| = 1; \circlearrowleft\} \cup \{|z_i| = |h|^{-ab}; \circlearrowleft\} \quad (3.50)$$

is a contour encircling the annulus $\mathcal{A} = \{u_i \mid 1 < |z_i| < |h|^{-ab}, i = 1, \dots, \text{rk}(G)\}$.

We now apply the residue theorem to (3.49). The integrand has simple poles coming from the denominator, whose positions are precisely described by the BAEs (3.3). Obviously, only the poles that lie inside the annulus \mathcal{A} contribute to the contour integral. Moreover, as we do in Appendix C, one can show that whenever a particular solution $[\hat{u}]$ to the BAEs (3.3) is fixed (on the torus) by a non-trivial element of the Weyl group \mathcal{W}_G , namely $w \cdot [\hat{u}] = [\hat{u}]$, then the numerator $\mathcal{Z}(\hat{u}; \xi, \nu_R, a\omega, b\omega)$ is such that cancelations take place and there is no contribution to the integral—more precisely, the function $\mathcal{Z}_{\text{tot}}(\hat{u}; \xi, \nu_R, a\omega, b\omega)$ defined in (3.11) vanishes.⁵ Hence, we define the set of relevant poles by:

$$\mathfrak{M}_{\text{index}} = \left\{ \hat{u}_i \mid [\hat{u}_i] \in \mathfrak{M}_{\text{BAE}} \quad \text{and} \quad 1 < |\hat{z}_i| < |h|^{-ab}, i = 1, \dots, \text{rk}(G) \right\}. \quad (3.51)$$

This includes all points inside the annulus \mathcal{A} such that their class belongs to $\mathfrak{M}_{\text{BAE}}$. In particular, the same equivalence class $[\hat{u}_i] \in \mathfrak{M}_{\text{BAE}}$ appears in $\mathfrak{M}_{\text{index}}$ as many times as the number of its representatives living in \mathcal{A} . For this reason, we employ the following alternative description:

$$\mathfrak{M}_{\text{index}} = \left\{ \hat{u}_i^{(m)} = [\hat{u}_i] - m_i \omega \mid [\hat{u}_i] \in \mathfrak{M}_{\text{BAE}}, m_i = 1, \dots, ab, i = 1, \dots, \text{rk}(G) \right\} \quad (3.52)$$

where, we some abuse of notation, we have denoted as $[\hat{u}_i]$ the representative in the fundamental domain of the torus as after (3.9).

⁵In particular, let us stress that the condition $w \cdot [\hat{u}] \neq [\hat{u}]$ in the definition of $\mathfrak{M}_{\text{BAE}}$ could be relaxed with no harm: in that case, we would simply include more poles in the sum, whose residues however combine to zero.

In addition, the numerator \mathcal{Z} has other poles coming from the elliptic gamma functions. As we show below, as long as the fugacities v_α, p, q are taken within the domain (2.6)—which is necessary in order for the standard contour integral representation (2.3) to be valid—those other poles either lie outside the annulus \mathcal{A} or are not poles of the integrand (because the denominator has a pole of equal or higher degree) and thus do not contribute to the integral.

Therefore, working within the domain (2.6), we can rewrite the index as

$$\mathcal{I}(p, q; v) = (-2\pi i)^{\text{rk}(G)} \kappa_G \sum_{\hat{u}^{(\vec{m})} \in \mathfrak{M}_{\text{index}}} \text{Res}_{u=\hat{u}^{(\vec{m})}} \left[\frac{\mathcal{Z}(u; \xi, \nu_R, a\omega, b\omega)}{\prod_{i=1}^{\text{rk}(G)} (1 - Q_i(u; \xi, \nu_R, \omega))} d^{\text{rk}(G)} u \right]. \quad (3.53)$$

Computing the residues produces the final expression for the supersymmetric index:

$$\mathcal{I}(p, q; v) = \kappa_G \sum_{\hat{u}^{(\vec{m})} \in \mathfrak{M}_{\text{index}}} \mathcal{Z}(\hat{u}^{(\vec{m})}; \xi, \nu_R, a\omega, b\omega) H(\hat{u}^{(\vec{m})}; \xi, \nu_R, \omega)^{-1}, \quad (3.54)$$

where H is defined in (3.12). The residue formula (3.54) can be rewritten, more elegantly, in the final form:

$$\mathcal{I}(p, q; v) = \kappa_G \sum_{\hat{u} \in \mathfrak{M}_{\text{BAE}}} \mathcal{Z}_{\text{tot}}(\hat{u}; \xi, \nu_R, a\omega, b\omega) H(\hat{u}; \xi, \nu_R, \omega)^{-1}, \quad (3.55)$$

where

$$\mathcal{Z}_{\text{tot}}(u; \xi, \nu_R, a\omega, b\omega) = \sum_{\{m_i\}=1}^{ab} \mathcal{Z}(u - m\omega; \xi, \nu_R, a\omega, b\omega). \quad (3.56)$$

To obtain this expression we have split the sum over the poles in $\mathfrak{M}_{\text{index}}$ into a sum over the inequivalent solutions to the BAEs, described by the elements of $\mathfrak{M}_{\text{BAE}}$, and the sum over the “repetitions” of these elements in the annulus \mathcal{A} . Moreover, we have used the double-periodicity of the Jacobian $H(u; \xi, \nu_R, \omega)$ to pull the latter sum inside the definition of \mathcal{Z}_{tot} .

3.2.3 Analysis of the residues

The last step consists in showing that the only residues contributing to (3.49) come from zeros of the denominator. In particular we need to show that, remaining within the domain (2.6), all poles in (3.49) which are not given by the BAEs live outside the annulus \mathcal{A} and thus do not contribute to the integral. We concretely do so by proving that every pole of \mathcal{Z} inside \mathcal{A} is also a pole of the denominator $\prod_i (1 - Q_i)$ with a high enough degree that the integrand of (3.49) is non-singular at those points.

We begin by classifying the poles of \mathcal{Z} . Using (3.26), (3.30) and

$$\tilde{\Gamma}(u; \tau, \sigma) = \frac{1}{\tilde{\Gamma}(\tau + \sigma - u; \tau, \sigma)}, \quad (3.57)$$

we can rewrite \mathcal{Z} as

$$\mathcal{Z}(u; \xi, \nu_R, a\omega, b\omega) = \prod_{\alpha > 0} \theta_0(\alpha(u); a\omega) \theta_0(-\alpha(u); b\omega) \times \prod_{a, \rho_a} \tilde{\Gamma}(\rho_a(u) + \omega_a(\xi) + r_a \nu_R; a\omega, b\omega). \quad (3.58)$$

Since $\theta_0(u; \omega)$ has no poles for finite u , the only singularities of \mathcal{Z} come from the elliptic gamma functions related to the chiral multiplets. These can be read off the product expansion:

$$\tilde{\Gamma}(u; a\omega, b\omega) = \prod_{m=0}^{\infty} \prod_{r=0}^{a-1} \prod_{s=0}^{b-1} \left(\frac{1 - h^{ab(m+2)-as-br} z^{-1}}{1 - h^{abm+as+br} z} \right)^{m+1} \quad (3.59)$$

that follows from (3.31), and so they are given by

$$z^{\rho_a} = v^{-\omega_a} h^{-r_a(a+b)/2-abm-as-br} \quad (3.60)$$

for $0 \leq r \leq a-1$, $0 \leq s \leq b-1$ and $m \geq 0$. The multiplicity of each pole is $\mu_m^a = m+1$.⁶ Notice that one could also write $z^{\rho_a} = v^{-\omega_a} h^{-r_a(a+b)/2-k}$ for $k \in \mathcal{R}(a, b)$.

We now turn to analyzing the denominator. More specifically, we need to find the singularities of $\prod_{i=1}^{\text{rk}(G)} (1 - Q_i(u))$. From (3.4) and (3.5) we see that Q_i has a pole whenever $\theta_0(\rho_a(u) + \omega_a(\xi) + r_a \nu_R; \omega) = 0$ and $\rho_a^i > 0$. Therefore, the singularities of the denominator are given by

$$z^{\rho_a} = v^{-\omega_a} h^{-r_a(a+b)/2+n} \quad \text{for } n \in \mathbb{Z}, \quad (3.61)$$

all with the same multiplicity $\nu^a = \sum_{i \in \mathcal{D}_a^+} \rho_a^i$. Here \mathcal{D}_a^\pm represents the set of indices such that $\rho_a^i > 0$, resp. $\rho_a^i < 0$, thus ν^a is the sum of the positive components of ρ_a . We notice that the denominator poles in (3.61) with $-n \in \mathcal{R}(a, b)$ coincide with the numerator poles. Therefore, the actual singularities of the integrand in (3.49) are only those points in (3.60) such that $\mu_m^a > \nu^a$, or more explicitly

$$m \geq \sum_{i \in \mathcal{D}_a^+} \rho_a^i. \quad (3.62)$$

We now want to show that, when the fugacities satisfy (2.6), the set of actual singularities is always living outside the annulus \mathcal{A} . Therefore, we first study the conditions for which (3.60) belong to the annulus \mathcal{A} . By imposing that $1 < |z_i| < |h|^{-ab}$, we obtain that

$$|h|^{-ab \sum_{i \in \mathcal{D}_a^-} \rho_a^i} < |z^{\rho_a}| < |h|^{-ab \sum_{i \in \mathcal{D}_a^+} \rho_a^i}, \quad \forall a. \quad (3.63)$$

⁶In counting the multiplicity one may worry that there could be different choices of r, s that give the same $abm+as+br$ for fixed m . This is equivalent to finding non-trivial solutions to the equation $as+br = as'+br'$. However, it is easy to see that, as long as $0 \leq r, r' \leq a-1$ and $0 \leq s, s' \leq b-1$, such an equation has no non-trivial solution in \mathbb{Z} .

Then we determine the constraints imposed on (3.60) by requiring (2.6). In the rational case, the latter conditions are expressed by $|h|^{a+b} < |v^{\omega_a} h^{r_a(a+b)/2}| < 1, \forall a$. These inequalities, together with $0 \leq as + br \leq 2ab - a - b$, imply that

$$|h|^{-abm} \leq |h|^{-abm-as-br} < |z^{\rho_a}| < |h|^{-abm-a(s+1)-b(r+1)} \leq |h|^{-ab(m+2)}. \quad (3.64)$$

Furthermore, requiring (3.62) to be satisfied, we obtain that

$$|z^{\rho_a}| > |h|^{-ab \sum_{i \in \mathfrak{D}_a^+} \rho_a^i}, \quad \forall a, \quad (3.65)$$

which is satisfied by all the singularities of (3.49) coming from the numerator \mathcal{Z} .

At this point, we immediately notice that the intersection between (3.63) and (3.65) is empty. This means that, if the flavor fugacities satisfy (2.6), all poles of the integrand (3.49) that come from poles of the numerator \mathcal{Z} live outside the annulus \mathcal{A} , and so the only residues contributing to the integral are those given by the BAEs. This completes the proof of (3.10).

Acknowledgments

F.B. is supported in part by the MIUR-SIR grant RBSI1471GJ “Quantum Field Theories at Strong Coupling: Exact Computations and Applications”.

A Numerical semigroups and the Fröbenius problem

Given a set of non-negative integer numbers $\{a_1, \dots, a_r\}$, the Fröbenius problem consists in classifying which integers can (or cannot) be written as non-negative integer linear combinations of those. This problem has deep roots in the theory of numerical semigroups.

A *semigroup* is an algebraic structure \mathcal{R} endowed with an associative binary operation. Analogously to groups, we denote it as $(\mathcal{R}, *)$. On the other hand, differently from the case of a group, no requirement on the presence of identity and inverse elements is made. A *numerical semigroup* is an additive semigroup $(\mathcal{R}, +)$, where \mathcal{R} consists of all non-negative integers $\mathbb{Z}_{\geq 0}$ except for a finite number of positive elements (thus $0 \in \mathcal{R}$). The set $\{n_1, \dots, n_t\}$ is called a *generating set* for $(\mathcal{R}, +)$ if all elements of \mathcal{R} can be written as non-negative integers linear combinations of n_1, \dots, n_t . We then denote the semigroup with the presentation

$$\mathcal{R} = \langle n_1, \dots, n_t \rangle. \quad (A.1)$$

Among all possible presentations of \mathcal{R} , there exists a *unique minimal presentation*, which contains the minimal number of generators. Such a number is called the *embedding dimension*

$e(\mathcal{R})$ of the semigroup. We now define other important quantities associated with numerical semigroups:

- The *multiplicity* $m(\mathcal{R})$ is the smallest non-zero element of \mathcal{R} .
- The *set of gaps* $\overline{\mathcal{R}} = \mathbb{N} \setminus \mathcal{R}$ is the set of positive integers which are not contained in \mathcal{R} . Equivalently, the gaps are defined as all natural numbers which cannot be written as non-negative integer linear combination of the generators n_1, \dots, n_t of \mathcal{R} .
- The set of gaps $\overline{\mathcal{R}}$ is always a finite set. Its largest element is the *Fröbenius number* $F(\mathcal{R})$. Alternatively, given a presentation $\langle n_1, \dots, n_t \rangle$, the Fröbenius number is defined as the largest integer which cannot be written as a non-negative integer linear combination of the generators.
- The *genus* $\chi(\mathcal{R})$ is the number of gaps, *i.e.* it is the order of the set of gaps: $\chi(\mathcal{R}) = |\overline{\mathcal{R}}|$.
- The *weight* $w(\mathcal{R})$ is the sum of all gaps: $w(\mathcal{R}) = \sum_{k \in \overline{\mathcal{R}}} k$.
- The following inequalities hold:

$$e(\mathcal{R}) \leq m(\mathcal{R}) \quad F(\mathcal{R}) \leq 2\chi(\mathcal{R}) - 1. \quad (\text{A.2})$$

In particular, if $x \in \mathcal{R}$, then $F(\mathcal{R}) - x \notin \mathcal{R}$.

We now study the case where the embedding dimension is $e(\mathcal{R}) = 2$, *i.e.* the minimal presentation is defined by two positive integers a, b with $\gcd(a, b) = 1$. The associated numerical semigroup is denoted by $\mathcal{R}(a, b) = \langle a, b \rangle$ and the set of gaps is $\overline{\mathcal{R}}(a, b) = \mathbb{N} \setminus \mathcal{R}(a, b)$. The multiplicity is simply $m(a, b) = \min\{a, b\}$, whereas the Fröbenius number is given by

$$F(a, b) = ab - a - b. \quad (\text{A.3})$$

The genus and the weight are

$$\chi(a, b) = \frac{(a-1)(b-1)}{2} \quad w(a, b) = \frac{(a-1)(b-1)(2ab - a - b - 1)}{12}. \quad (\text{A.4})$$

Thanks to the properties of $\mathcal{R}(a, b)$, one can prove the following identities:

$$\begin{aligned} \prod_{r=0}^{a-1} \prod_{s=0}^{b-1} (zh^{as+br}; h^{ab})_{\infty} &= \frac{(z; h)_{\infty}}{\prod_{k \in \overline{\mathcal{R}}(a, b)} (1 - zh^k)} \\ \prod_{r=0}^{a-1} \prod_{s=0}^{b-1} (z^{-1}h^{ab-as-br}; h^{ab})_{\infty} &= (z^{-1}h; h)_{\infty} \prod_{k \in \mathcal{R}(a, b)} (1 - z^{-1}h^{-k}). \end{aligned} \quad (\text{A.5})$$

Proof. We begin with the first identity. Using the definition of the q -Pochhammer symbol we can write:

$$\prod_{r=0}^{a-1} \prod_{s=0}^{b-1} (zh^{as+br}; h^{ab})_{\infty} = \prod_{n=0}^{\infty} \prod_{r=0}^{a-1} \prod_{s=0}^{b-1} (1 - zh^{abn+as+br}). \quad (\text{A.6})$$

Using that a, b are coprime, the set of integers $\{as + br \mid r = 0, \dots, a-1, s = 0, \dots, b-1\}$ covers once and only once every class modulo ab . It follows that the set of exponents $\{abn + as + br\}$ is precisely $\mathcal{R}(a, b)$. Then

$$\prod_{r=0}^{a-1} \prod_{s=0}^{b-1} (zh^{as+br}; h^{ab})_{\infty} = \prod_{k \in \mathcal{R}(a, b)} (1 - zh^k) = \frac{\prod_{k=0}^{\infty} (1 - zh^k)}{\prod_{k \in \overline{\mathcal{R}}(a, b)} (1 - zh^k)} = \frac{(z; h)_{\infty}}{\prod_{k \in \overline{\mathcal{R}}(a, b)} (1 - zh^k)}, \quad (\text{A.7})$$

which proves the first equality in (3.36).

The proof of the second identity is a bit trickier. The key point is to notice that the set $\{as + br\}$ does not contain any element of $\overline{\mathcal{R}}(a, b)$ and thus

$$\{as + br\} = \{k + \Delta_k ab \mid k = 0, \dots, ab - 1\} \quad \text{with} \quad \Delta_k = \begin{cases} 0 & \text{if } k \in \mathcal{R}(a, b) \\ 1 & \text{if } k \in \overline{\mathcal{R}}(a, b). \end{cases} \quad (\text{A.8})$$

This implies that $\{ab - as - br\} = \{-k + (1 - \Delta_k)ab \mid k = 0, \dots, ab - 1\}$. Finally, including the freedom of choosing $n \geq 0$, we find that the set of exponents is

$$\{abn + ab - as - ar\} = (-\overline{\mathcal{R}}) \cup \mathbb{Z}_{>0}. \quad (\text{A.9})$$

Then

$$\begin{aligned} \prod_{r=0}^{a-1} \prod_{s=0}^{b-1} (z^{-1}h^{ab-as-br}; h^{ab})_{\infty} &= \prod_{n=0}^{\infty} \prod_{r=0}^{a-1} \prod_{s=0}^{b-1} (1 - z^{-1}h^{ab(n+1)-as-br}) \\ &= \prod_{k \in \overline{\mathcal{R}}(a, b)} (1 - z^{-1}h^{-k}) \times \prod_{k=1}^{\infty} (1 - z^{-1}h^k) = (z^{-1}h; h)_{\infty} \prod_{k \in \overline{\mathcal{R}}(a, b)} (1 - z^{-1}h^{-k}). \end{aligned} \quad (\text{A.10})$$

This completes the proof of (3.36). \square

Thanks to the definition of $\theta_0(u; \omega)$, we can apply (3.36) and we obtain that

$$\prod_{r=0}^{a-1} \prod_{s=0}^{b-1} \theta_0(u + (as + br)\omega; \omega) = \prod_{k \in \overline{\mathcal{R}}(a, b)} \frac{(1 - z^{-1}h^{-k})}{(1 - zh^k)} \theta_0(u; \omega) = \frac{1}{(-z)^{\chi(a, b)} h^{w(a, b)}} \theta_0(u; \omega). \quad (\text{A.11})$$

B A dense set

Here we show that the set of points (p, q) such that

$$q^a = p^b \quad \text{for coprime } a, b \in \mathbb{N} \quad (\text{B.1})$$

is dense in $\{|p| < 1, |q| < 1\}$. We write the fugacities in terms of chemical potentials, $p = e^{2\pi i\sigma}$ and $q = e^{2\pi i\tau}$ with $\text{Im } \sigma, \text{Im } \tau > 0$, and for the sake of this argument we choose the determination on the “strip” $0 \leq \text{Re } \sigma, \text{Re } \tau < 1$. Then the condition (B.1) is equivalent to

$$a(\tau + n) = b(\sigma + m) \quad (\text{B.2})$$

for some $m, n \in \mathbb{Z}$ and $a, b \in \mathbb{N}$ coprime.

We choose an arbitrary point (τ_0, σ_0) in the strip and ask if we can find another point (τ, σ) , arbitrarily close, that satisfies (B.2). Consider a straight line in the complex plane that starts from 0 and goes through $\tau_0 + n$ for some integer n . When winding once around the strip, this line has an imaginary excursion

$$\Delta y = \frac{\text{Im } \tau_0}{\text{Re } \tau_0 + n}. \quad (\text{B.3})$$

We can make this quantity arbitrarily small by choosing n sufficiently large. We define σ' as the closest point to σ_0 that lies on the image of the line on the strip modulo 1, and has $\text{Re } \sigma' = \text{Re } \sigma_0$. It is clear that

$$|\sigma' - \sigma_0| = |\text{Im } \sigma' - \text{Im } \sigma_0| \leq \Delta y/2, \quad (\text{B.4})$$

and, by construction, $(\sigma' + m) = t(\tau_0 + n)$ for some $m, n \in \mathbb{Z}$ and $t \in \mathbb{R}_+$. We see that $|\sigma' - \sigma_0|$ can be made arbitrarily small by increasing n . Next, we approximate t by a fraction $a/b \in \mathbb{Q}_+$. This, for a/b sufficiently close to t , defines a point σ in the strip by

$$(\sigma + m) = \frac{a}{b}(\tau_0 + n). \quad (\text{B.5})$$

It is clear that σ can be made arbitrarily close to σ' by approximating t sufficiently well with a/b . We have thus found a pair $(\sigma, \tau = \tau_0)$, arbitrarily close to (σ_0, τ_0) , that satisfies the constraint (B.2).

C Weyl group fixed points

In this appendix we prove that $\mathcal{Z}_{\text{tot}}(u; \xi, \nu_R, a\omega, b\omega)$ vanishes when evaluated at a point \hat{u} which is fixed, on a torus of modular parameter ω , by a non-trivial element w of the Weyl group \mathcal{W}_G :

$$w \cdot [\hat{u}] = [\hat{u}]. \quad (\text{C.1})$$

This implies that the solutions to the BAEs (3.9) which are fixed points on the torus of an element of the Weyl group, can be excluded from the set $\mathfrak{M}_{\text{BAE}}$ —as is done in (3.9)—because they do not contribute to the BA formula (3.10) for the superconformal index.

C.1 The rank-one case

Let us first consider the case that the gauge group G has rank one, *i.e.*, that $\mathfrak{g} = \mathfrak{su}(2)$. Then there are only two roots, α and $-\alpha$, and the Weyl group is $\mathcal{W}_G = \{1, s_\alpha\} \cong \mathbb{Z}_2$ where s_α is the unique non-trivial Weyl reflection along the root α :

$$s_\alpha(u) = -u \quad \forall u \in \mathfrak{h}. \quad (\text{C.2})$$

We choose a basis element $\{H\}$ for the Cartan subalgebra \mathfrak{h} such that $\rho(H) \equiv \rho \in \mathbb{Z}$ for any weight $\rho \in \Lambda_{\text{weight}}$. In this canonical basis $\alpha = 2$ (while the fundamental weight is $\lambda = 1$). The solutions to $s_\alpha \cdot [\hat{u}] = [\hat{u}]$ are given by⁷

$$\hat{u} = \frac{p + q\omega}{2} \quad \text{with } p, q \in \mathbb{Z}. \quad (\text{C.3})$$

Choosing a representative for $[\hat{u}]$ in the fundamental domain of the torus, the inequivalent solutions are with $p = 0, 1$ and $q = 0, 1$.

The representations of $\mathfrak{su}(2)$ are labelled by a half-integer spin $j \in \mathbb{N}/2$ and their weights are $\rho \in \{\ell\alpha \mid \ell = -j, -j+1, \dots, j-1, j\}$. Therefore, exploiting the expression in (3.58), the function \mathcal{Z} reduces to

$$\begin{aligned} \mathcal{Z}(u; \xi, \nu_R, a\omega, b\omega) &= \\ &= \theta_0(\alpha(u); a\omega) \theta_0(-\alpha(u); b\omega) \prod_a \prod_{\ell_a = -j_a}^{j_a} \tilde{\Gamma}(\ell_a \alpha(u) + \omega_a(\xi) + r_a \nu_R; a\omega, b\omega). \end{aligned} \quad (\text{C.4})$$

Moreover, the function \mathcal{Z}_{tot} defined in (3.11) is a single sum over $m = 1, \dots, ab$.

We want to prove that $\mathcal{Z}_{\text{tot}}(\hat{u}; \xi, \nu_R, a\omega, b\omega) = 0$. To do that, we construct an involutive map $\gamma : m \mapsto m'$ acting on the set of integers $\{1, \dots, ab\}$ according to

$$m' = m \pmod{b}, \quad m' = q - m \pmod{a}, \quad (\text{C.5})$$

which define m' uniquely. It will be convenient to introduce the numbers $r, s \in \mathbb{Z}$ such that $m' = m + sb = q - m + ra$. The map γ has the property that

$$m' - q/2 = \begin{cases} m - q/2 & \pmod{b}, \\ -(m - q/2) & \pmod{a}, \end{cases} = m - q/2 + sb = -(m - q/2) + ra. \quad (\text{C.6})$$

⁷The integers p, q appearing in this appendix should not be confused with the complex angular fugacities appearing in the rest of the paper.

We will prove that

$$\mathcal{Z}(\hat{u} - m'\omega; \xi, \nu_R, a\omega, b\omega) = -\mathcal{Z}(\hat{u} - m\omega; \xi, \nu_R, a\omega, b\omega) . \quad (\text{C.7})$$

In particular, the sum over m inside \mathcal{Z}_{tot} splits into a sum over the fixed points of γ and a sum over the pairs of values related by γ . The property (C.7) guarantees that each term in those sums vanishes, implying that \mathcal{Z}_{tot} vanishes.

Let us adopt the notation

$$\mathcal{Z}_m \equiv \mathcal{Z}(\hat{u} - m\omega; \xi, \nu_R, a\omega, b\omega) = \mathcal{Z}(p/2 - (m - q/2)\omega; \xi, \nu_R, a\omega, b\omega) . \quad (\text{C.8})$$

We define the vector multiplet and the chiral multiplet contribution, respectively, as

$$\begin{aligned} \mathcal{A}_m &= \theta_0(\alpha(p/2) - \alpha(m - q/2)\omega; a\omega) \theta_0(-\alpha(p/2) + \alpha(m - q/2)\omega; b\omega) \\ \mathcal{B}_m &= \prod_a \prod_{\ell_a = -j_a}^{j_a} \tilde{\Gamma}(\ell_a \alpha(p/2) - \ell_a \alpha(m - q/2)\omega + \omega_a(\xi) + r_a \nu_R; a\omega, b\omega) , \end{aligned} \quad (\text{C.9})$$

such that $\mathcal{Z}_m = \mathcal{A}_m \mathcal{B}_m$. Then \mathcal{Z}_{tot} evaluated on \hat{u} can be expressed as

$$\mathcal{Z}_{\text{tot}}\left(\frac{p + q\omega}{2}; \xi, \nu_R, a\omega, b\omega\right) = \sum_{m=1: m'=m}^{ab} \mathcal{Z}_m + \sum_{(m, m'): m' \neq m} (\mathcal{Z}_m + \mathcal{Z}_{m'}) . \quad (\text{C.10})$$

Our goal is to show that $\mathcal{Z}_{m'} = -\mathcal{Z}_m$.

We begin by considering the contribution of \mathcal{A}_m . Using (C.6) we can write

$$\begin{aligned} \mathcal{A}_{m'} &= \theta_0(p + (2m - q)\omega - 2ra\omega; a\omega) \theta_0(-p + (2m - q)\omega + 2sb\omega; b\omega) \\ &= \theta_0(-p - (2m - q)\omega + (2r + 1)a\omega; a\omega) \theta_0(-p + (2m - q)\omega + 2sb\omega; b\omega) . \end{aligned} \quad (\text{C.11})$$

In the second equality we used the second relation in (3.26). Using the first relation in (3.26), the identity $2m - q - ra + sb = 0$ and reinstating α , with some algebra we obtain

$$\mathcal{A}_{m'} = -e^{-2\pi i \alpha(r) \alpha(s) \nu_R} \mathcal{A}_m . \quad (\text{C.12})$$

Then we turn to \mathcal{B}_m and, using (C.6), write

$$\begin{aligned} \mathcal{B}_{m'} &= \prod_a \prod_{\ell_a = -j_a}^{j_a} \tilde{\Gamma}(\ell_a p + \ell_a (2m - q)\omega + \omega_a(\xi) + r_a \nu_R - 2\ell_a r a\omega; a\omega, b\omega) \\ &= \prod_a \prod_{\ell_a = -j_a}^{j_a} \tilde{\Gamma}(\ell_a p - \ell_a (2m - q)\omega + \omega_a(\xi) + r_a \nu_R + 2\ell_a r a\omega; a\omega, b\omega) . \end{aligned} \quad (\text{C.13})$$

We recall that j_a can be integer or half-integer. In the second equality we simply redefined $\ell_a \rightarrow -\ell_a$ and shifted the argument by the integer $2\ell_a p$. Using the identity (3.30) repeatedly and distinguishing the cases $\ell_a \leq 0$, we obtain

$$\mathcal{B}_{m'} = \Theta \times \mathcal{B}_m \quad (\text{C.14})$$

where the factor Θ equals

$$\Theta = \prod_a \prod_{\ell_a > 0}^{j_a} \prod_{k=0}^{2\ell_a r - 1} \frac{\theta_0(\ell_a p - \ell_a(2m - q)\omega + \omega_a(\xi) + r_a \nu_R + k a \omega; b \omega)}{\theta_0(-\ell_a p + \ell_a(2m - q)\omega + \omega_a(\xi) + r_a \nu_R + (k - 2\ell_a r) a \omega; b \omega)}. \quad (\text{C.15})$$

The second product starts from 1 or $\frac{1}{2}$ depending on j_a being integer or half-integer. Using $2m - q - r a + s b = 0$ at denominator and shifting the arguments by integers, we rewrite

$$\Theta = \prod_a \prod_{\ell_a > 0}^{j_a} \prod_{k=0}^{2\ell_a r - 1} \frac{\theta_0(\ell_a p - \ell_a(2m - q)\omega + \omega_a(\xi) + r_a \nu_R + k a \omega; b \omega)}{\theta_0(\ell_a p - \ell_a(2m - q)\omega + \omega_a(\xi) + r_a \nu_R + k a \omega - 2\ell_a s b \omega; b \omega)}. \quad (\text{C.16})$$

Finally we use the first relation in (3.26) at denominator, to obtain

$$\Theta = \prod_a \prod_{\ell_a > 0}^{j_a} (-1)^{4\ell_a^2 r s + 8\ell_a^3 r s p} e^{-8\pi i \ell_a^2 r s \omega_a(\xi)} e^{-8\pi i \ell_a^2 r s (r_a - 1) \nu_R}. \quad (\text{C.17})$$

Reinstating the root α , this factor can be written as

$$\Theta = \prod_a \prod_{\ell_a > 0}^{j_a} (-1)^{\ell_a^3 \alpha(r) \alpha(s) \alpha(p)} \times \prod_{a, \rho_a \in \mathfrak{R}_a} e^{\pi i \rho_a(r) \rho_a(s) (\frac{1}{2} - \omega_a(\xi) - (r_a - 1) \nu_R)}. \quad (\text{C.18})$$

Combining with (C.12), the factor picked up by \mathcal{Z} can be expressed in terms of the anomaly coefficients (3.18) and (3.19):

$$\mathcal{Z}_{m'} = -e^{2\pi i \phi} e^{\pi i r s (\frac{1}{2} \mathcal{A}^{ii} - \mathcal{A}^{ii \alpha} \xi_\alpha - \mathcal{A}^{ii R} \nu_R)} \mathcal{Z}_m. \quad (\text{C.19})$$

Here i is the gauge index taking a single value. We recall the anomaly cancelation conditions $\mathcal{A}^{ii \alpha} = \mathcal{A}^{ii R} = 0$ and $\mathcal{A}^{ii} \in 4\mathbb{Z}$, implying that the second exponential equals 1. In the first exponential we defined

$$\phi = \frac{1}{2} \alpha(r) \alpha(s) \alpha(p) \sum_a \sum_{\ell_a > 0}^{j_a} \ell_a^3 = 4r s p \sum_a \sum_{\ell_a > 0}^{j_a} \ell_a^3. \quad (\text{C.20})$$

It remains to show that $\phi \in \mathbb{Z}$, so that also the first exponential equals 1.

For each chiral multiplet in the theory, indicized by a , in order to evaluate the second sum in (C.20) we should distinguish different cases:

$$\psi_j \equiv 4 \sum_{\ell > 0}^j \ell^3 = \begin{cases} j^2(j+1)^2 & \in 4\mathbb{Z} & \text{if } j \in \mathbb{Z} \\ 2(k+1)^2(8k^2+16k+7) & \in 2\mathbb{Z} & \text{if } j = 2k + \frac{3}{2} \in 2\mathbb{Z} + \frac{3}{2} \\ \frac{1}{2}(2k+1)^2(8k^2+8k+1) & \in 4\mathbb{Z} + \frac{1}{2} & \text{if } j = 2k + \frac{1}{2} \in 2\mathbb{Z} + \frac{1}{2}. \end{cases} \quad (\text{C.21})$$

Therefore, chiral multiplets whose gauge representation has spin $j \in \mathbb{Z}$ or $j \in 2\mathbb{Z} + \frac{3}{2}$ give integer contribution to ϕ . On the other hand, chiral multiplets with $j \in 2\mathbb{Z} + \frac{1}{2}$ can give half-integer contribution. However, because of the Witten anomaly [34], the total number of such multiplets must be even. This is reproduced by the condition (3.21) on the pseudo-anomaly coefficient \mathcal{A}^{ii} . Indeed, the contribution of a chiral multiplet to the pseudo-anomaly is

$$\mathcal{A}_{(j)}^{ii} = \sum_{\ell=-j}^j (2\ell)^2 = \frac{4}{3} j(j+1)(2j+1) \in \begin{cases} 4\mathbb{Z} & \text{if } j \in \mathbb{Z} \text{ or } j \in 2\mathbb{Z} + \frac{3}{2} \\ 4\mathbb{Z} + 2 & \text{if } j \in 2\mathbb{Z} + \frac{1}{2}, \end{cases} \quad (\text{C.22})$$

and the condition $\mathcal{A}^{ii} \in 4\mathbb{Z}$ requires that the total number of chiral multiplets with $j \in 2\mathbb{Z} + \frac{1}{2}$ be even. This implies that $\phi \in \mathbb{Z}$, and thus that $\mathcal{Z}_{m'} = -\mathcal{Z}_m$. In turn, using (C.10), this implies that

$$\mathcal{Z}_{\text{tot}}(\hat{u}; \xi, \nu_R, a\omega, b\omega) = 0 \quad (\text{C.23})$$

whenever \hat{u} is fixed on the torus by the non-trivial element s_α of the Weyl group of $\mathfrak{su}(2)$.

C.2 The higher-rank case

Let us now move to the case of a generic semi-simple gauge algebra \mathfrak{g} of rank $\text{rk}(G)$. The Weyl group \mathcal{W}_G is a finite group generated by the Weyl reflections

$$s_\alpha(u) = u - 2 \frac{\alpha(u)}{(\alpha, \alpha)} \tilde{\alpha} \quad \forall u \in \mathfrak{h}, \quad (\text{C.24})$$

where $\tilde{\alpha}$ is the image of the root α under the isomorphism $\mathfrak{h}^* \rightarrow \mathfrak{h}$ induced by the non-degenerate scalar product (\cdot, \cdot) on \mathfrak{h}^* . Suppose that there exists a non-trivial element w of \mathcal{W}_G such that $w \cdot \hat{u} = \hat{u}$. It is a standard theorem that the Weyl group acts freely and transitively on the set of Weyl chambers. Therefore, \hat{u} cannot belong to a Weyl chamber but must instead lie on a boundary between two or more chambers. Such boundaries are the hyperplanes fixed by the Weyl reflections, $\{u | s_\alpha(u) = u\}$, and their intersections. We conclude that there must exist at least one root $\hat{\alpha}$ such that $s_{\hat{\alpha}}(\hat{u}) = \hat{u}$.

On the other hand, we are interested in points \hat{u} such that their equivalence class on the torus is fixed by a non-trivial element of the Weyl group, $w \cdot [\hat{u}] = [\hat{u}]$. In this case, for each w we can always identify (at least) one root $\hat{\alpha}$ such that $s_{\hat{\alpha}}[\hat{u}] = [\hat{u}]$, and moreover we can choose a set of simple roots that contains $\hat{\alpha}$. Let us fix a basis of simple roots $\{\alpha_l\}_{l=1, \dots, \text{rk}(G)}$ for \mathfrak{g} that contains $\hat{\alpha}$. The fundamental weights λ_l are defined by

$$2 \frac{(\lambda_k, \alpha_l)}{(\alpha_l, \alpha_l)} = \delta_{kl}. \quad (\text{C.25})$$

We choose a basis $\{H^i\}$ for the Cartan subalgebra \mathfrak{h} such that the fundamental weights have components $\lambda_l^i = \lambda_l(H^i) = \delta_l^i$. In this basis $\rho(H^i) \equiv \rho^i \in \mathbb{Z}$ for any weight $\rho \in \Lambda_{\text{weight}}$.

Moreover, the double periodicity of the gauge variables $u = u_i H^i$ is $u_i \sim u_i + 1 \sim u_i + \omega$. From (C.24), the fixed points should satisfy

$$2 \frac{\hat{\alpha}(\hat{u})}{(\hat{\alpha}, \hat{\alpha})} \tilde{\alpha} = p + q\omega \quad \text{for} \quad p = p_i H^i, \quad q = q_i H^i \quad \text{and} \quad p_i, q_i \in \mathbb{Z}. \quad (\text{C.26})$$

Here $\tilde{\alpha}$ is dual to $\hat{\alpha}$. It is clear that p, q should be aligned with $\tilde{\alpha}$, therefore we set

$$p = \frac{2\hat{p}}{(\hat{\alpha}, \hat{\alpha})} \tilde{\alpha}, \quad q = \frac{2\hat{q}}{(\hat{\alpha}, \hat{\alpha})} \tilde{\alpha}, \quad \text{with} \quad \hat{p}, \hat{q} \in \mathbb{Z}. \quad (\text{C.27})$$

In the basis $\{H^i\}$ we have chosen, the components of $\tilde{\alpha}$ are $(\lambda_i, \hat{\alpha}) = \delta_{il} (\hat{\alpha}, \hat{\alpha})/2$, where l is such that $\hat{\alpha} = \alpha_l$ and we have used (C.25). Only one component of $\tilde{\alpha}$ is non-zero, which implies that the integer components of p, q are $p_i = \hat{p} \delta_{il}$ and $q_i = \hat{q} \delta_{il}$. This proves integrality of \hat{p}, \hat{q} . The general solution to (C.26) can then be written as

$$\hat{u} = \hat{u}_0 + \frac{p + q\omega}{2}, \quad (\text{C.28})$$

where \hat{u}_0 is such that $\hat{\alpha}(\hat{u}_0) = 0$.

Now, consider the explicit expression (3.11) for \mathcal{Z}_{tot} , in terms of \mathcal{Z} given in (2.12). Given any representation \mathfrak{R} of \mathfrak{g} , we can always decompose it into irreducible representations of the $\mathfrak{su}(2)_{\hat{\alpha}}$ subalgebra associated with $\hat{\alpha}$. The set of weights (with multiplicities) $\Lambda_{\mathfrak{R}}$ corresponding to \mathfrak{R} can be organized as a union $\Lambda_{\mathfrak{R}} = \cup_I \Lambda_{\mathfrak{R}, I}$ of subsets $\Lambda_{\mathfrak{R}, I}$, each corresponding to a representation of $\mathfrak{su}(2)_{\hat{\alpha}}$. Concretely, each $\Lambda_{\mathfrak{R}, I}$ is associated to a representation of $\mathfrak{su}(2)_{\hat{\alpha}}$ of spin j_I , so that its elements can be expressed as an $\hat{\alpha}$ -chain:

$$\Lambda_{\mathfrak{R}, I} = \{ \hat{\rho}_I + \ell_I \hat{\alpha} \mid \ell_I = -j_I, -j_I + 1, \dots, j_I - 1, j_I \}. \quad (\text{C.29})$$

Here $\hat{\rho}_I$ is the central point, which is orthogonal to $\hat{\alpha}$, *i.e.* such that $(\hat{\rho}_I, \hat{\alpha}) = 0$. Notice that, in general, $\hat{\rho}_I$ is not a weight.⁸ The product over all weights ρ of the representation \mathfrak{R} can then be expressed as a product over the representations of $\mathfrak{su}(2)_{\hat{\alpha}}$ contained in \mathfrak{R} . In particular we can write

$$\begin{aligned} \prod_a \prod_{\rho_a \in \mathfrak{R}_a} \tilde{\Gamma}(\rho_a(u) + \omega_a(\xi) + r_a \nu_R; a\omega, b\omega) &= \\ &= \prod_{a, I} \prod_{\ell_{aI} = -j_{aI}}^{j_{aI}} \tilde{\Gamma}(\hat{\rho}_{aI}(u) + \ell_{aI} \hat{\alpha}(u) + \omega_a(\xi) + r_a \nu_R; a\omega, b\omega). \end{aligned} \quad (\text{C.30})$$

When specifying \mathfrak{R} to the adjoint representation, we obtain a similar decomposition for the roots of \mathfrak{g} . Besides the roots $\hat{\alpha}$ and $-\hat{\alpha}$ of $\mathfrak{su}(2)_{\hat{\alpha}}$, the other roots organize into $\hat{\alpha}$ -chains that we indicate as

$$\Lambda_{\text{roots}, J} = \{ \hat{\beta}_J + \ell_J \hat{\alpha} \mid \ell_J = -j_J, -j_J + 1, \dots, j_J - 1, j_J \}, \quad (\text{C.31})$$

⁸Indeed, $\hat{\rho}_I$ is guaranteed to be a weight (and in particular a root) only if the spin j_I is integer.

where $\hat{\beta}_J$ is the non-vanishing central point orthogonal to $\hat{\alpha}$ (once again, $\hat{\beta}_J$ is in general not a weight). Notice that, for each subset $\Lambda_{\text{roots},J}$ of the set of roots, there is a disjoint conjugate subset $\overline{\Lambda}_{\text{roots},J}$ with the same spin j_J but opposite central point $-\hat{\beta}_J$.⁹ For this reason, we have that

$$\begin{aligned} \mathcal{Z}(u; \xi, \nu_R, a\omega, b\omega) &= \theta_0(\hat{\alpha}(u); a\omega) \theta_0(-\hat{\alpha}(u); b\omega) \times \\ &\times \frac{\prod_{a,I} \prod_{\ell_{aI}=-j_{aI}}^{j_{aI}} \tilde{\Gamma}(\hat{\rho}_{aI}(u) + \ell_{aI}\hat{\alpha}(u) + \omega_a(\xi) + r_a\nu_R; a\omega, b\omega)}{\prod_J \prod_{\ell_J=-j_J}^{j_J} \tilde{\Gamma}(\hat{\beta}_J(u) + \ell_J\hat{\alpha}; a\omega, b\omega) \tilde{\Gamma}(-\hat{\beta}_J(u) + \ell_J\hat{\alpha}; a\omega, b\omega)}. \end{aligned} \quad (\text{C.32})$$

Similarly to the rank one case, we want to prove that $\mathcal{Z}_{\text{tot}}(\hat{u}; \xi, \nu_R, a\omega, b\omega) = 0$ for \hat{u} in (C.28). Thus, we construct an involutive map $\gamma : m \mapsto m'$, acting on the set \mathcal{M} of vectors $m = m_i H^i$ with integer components $1 \leq m_i \leq ab$. The map is constructed in such a way that it leaves m invariant along the directions orthogonal to $\tilde{\alpha}$, whereas it shifts the component parallel to $\tilde{\alpha}$ by an integer amount. To be precise, take two vectors $r, s \in \mathfrak{h}$ such that

$$r = \frac{2\hat{r}}{(\hat{\alpha}, \hat{\alpha})} \tilde{\alpha}, \quad s = \frac{2\hat{s}}{(\hat{\alpha}, \hat{\alpha})} \tilde{\alpha}, \quad \text{with } \hat{r}, \hat{s} \in \mathbb{Z}, \quad (\text{C.33})$$

meaning that r, s are parallel to $\tilde{\alpha}$ and have integer components $r_i = \hat{r} \delta_{il}$, $s_i = \hat{s} \delta_{il}$. Then, we construct m' as

$$m' = m + s b, \quad (\text{C.34})$$

which implies that m' differs from m only by integer shifts along the direction of $\tilde{\alpha}$. For \hat{s} we take the unique integer such that $m' \in \mathcal{M}$ and

$$\hat{\alpha}(m') = \hat{\alpha}(m) + \hat{\alpha}(s) b = \hat{\alpha}(q - m) + \hat{\alpha}(r) a. \quad (\text{C.35})$$

Indeed, consider the following equation in r and s : $2\hat{\alpha}(m) - \hat{\alpha}(q) = \hat{\alpha}(r) a - \hat{\alpha}(s) b$. Using (C.27) and (C.33), it reduces to $\hat{\alpha}(m) - \hat{q} = \hat{r} a - \hat{s} b$. Since a, b are coprime, this equation always admits an infinite number of solutions in the pair (\hat{r}, \hat{s}) , which can be parametrized as $(\hat{r}_0 + kb, \hat{s}_0 + ka)$ with $k \in \mathbb{Z}$. There is however one and only one solution such that m' has components $1 \leq m'_i \leq ab$. We define $\gamma(m) = m'$ in such a way. One can easily check that it is an involution.

As in the rank-one case, we adopt the notation

$$\mathcal{Z}_m \equiv \mathcal{Z}(\hat{u} - m\omega; \xi, \nu_R, a\omega, b\omega) = \mathcal{Z}(\hat{u}_0 + p/2 - (m - q/2)\omega; \xi, \nu_R, a\omega, b\omega), \quad (\text{C.36})$$

⁹It is easy to prove that $\Lambda_{\text{roots},J}$ and $\overline{\Lambda}_{\text{roots},J}$ are disjoint. Suppose, on the contrary, that there exists some common element $\hat{\beta}_J + \ell_J \hat{\alpha} = -\hat{\beta}_J + k_J \hat{\alpha}$ for some ℓ_J, k_J . This would imply that $\hat{\beta}_J = (k_J - \ell_J) \hat{\alpha} / 2$, but since $(\hat{\beta}_J, \hat{\alpha}) = 0$, then $\hat{\beta}_J = 0$. Since the only roots proportional to $\hat{\alpha}$ are $-\hat{\alpha}$ and $\hat{\alpha}$ itself, we have reached a contradiction.

and, for later convenience, split \mathcal{Z} into the vector multiplet and chiral multiplet contributions:

$$\mathcal{A}_m = \theta_0(\hat{\alpha}(p/2) - \hat{\alpha}(m - q/2)\omega; a\omega) \theta_0(-\hat{\alpha}(p/2) + \hat{\alpha}(m - q/2)\omega; b\omega) \quad (\text{C.37})$$

$$\mathcal{C}_m^\pm = \prod_J \prod_{\ell_J = -j_J}^{j_J} \tilde{\Gamma}(\pm \hat{\beta}_J(\hat{u}_0 - m\omega) + \ell_J \hat{\alpha}(p/2) - \ell_J \hat{\alpha}(m - q/2)\omega; a\omega, b\omega)$$

$$\mathcal{B}_m = \prod_{a,I} \prod_{\ell_{aI} = -j_{aI}}^{j_{aI}} \tilde{\Gamma}(\hat{\rho}_{aI}(\hat{u}_0 - m\omega) + \ell_{aI} \hat{\alpha}(p/2) - \ell_{aI} \hat{\alpha}(m - q/2)\omega + \omega_a(\xi) + r_a \nu_R; a\omega, b\omega)$$

such that $\mathcal{Z}_m = \mathcal{A}_m \mathcal{B}_m / \mathcal{C}_m^+ \mathcal{C}_m^-$. We will prove that $\mathcal{Z}_{m'} = -\mathcal{Z}_m$, which implies that $\mathcal{Z}_{\text{tot}}(\hat{u})$ vanishes because γ is an involution.

We begin by considering the contribution of $\mathcal{A}_{m'}$. Following the same steps as in (C.11) and using (C.35) and (3.26), we can show

$$\mathcal{A}_{m'} = -e^{-2\pi i \hat{\alpha}(r) \hat{\alpha}(s) \nu_R} \mathcal{A}_m . \quad (\text{C.38})$$

We also used that $\hat{\alpha}(r), \hat{\alpha}(s), \hat{\alpha}(q) \in 2\mathbb{Z}$, which is guaranteed by (C.27) and (C.33). We now turn to $\mathcal{B}_{m'}$. Eqn. (C.34) implies that $\hat{\rho}_{aI}(m') = \hat{\rho}_{aI}(m)$ for any $\hat{\rho}_{aI}$ orthogonal to $\hat{\alpha}$. Using the identity (3.30) repeatedly and distinguishing the cases $\ell_{aI} \leq 0$, we obtain

$$\mathcal{B}_{m'} = \prod_{a,I} \prod_{\ell_{aI} > 0}^{j_{aI}} (-1)^{\ell_{aI}^3 \hat{\alpha}(r) \hat{\alpha}(s) \hat{\alpha}(p)} \prod_{\alpha, \rho_\alpha} (-1)^{\frac{1}{2} \rho_\alpha(r) \rho_\alpha(s)} e^{-\pi i \rho_\alpha(r) \rho_\alpha(s) (\rho_\alpha(\hat{u}_0 - m\omega) + \omega_\alpha(\xi) + (r_\alpha - 1) \nu_R)} \mathcal{B}_m . \quad (\text{C.39})$$

The analysis of \mathcal{C}_m^\pm is analogous to the one for \mathcal{B}_m and it gives the following:

$$\mathcal{C}_{m'}^\pm = \prod_J \prod_{\ell_J > 0}^{j_J} (-1)^{\ell_J^3 \hat{\alpha}(r) \hat{\alpha}(s) \hat{\alpha}(p)} \prod_{\alpha \neq \pm \hat{\alpha}} (-1)^{\frac{1}{2} \alpha(r) \alpha(s)} e^{\pi i \alpha(r) \alpha(s) \nu_R} \times \mathcal{C}_m^\pm . \quad (\text{C.40})$$

Combining (C.38) with the latter, we obtain that the vector-multiplet contribution is

$$\mathcal{A}_{m'} / \mathcal{C}_{m'}^+ \mathcal{C}_{m'}^- = -e^{-2\pi i \sum_{\alpha \in \Delta} \alpha(r) \alpha(s) \nu_R} \mathcal{A}_m / \mathcal{C}_m^+ \mathcal{C}_m^- . \quad (\text{C.41})$$

We used $\hat{\alpha}(r) \hat{\alpha}(s) \in 4\mathbb{Z}$, as well as $\sum_{\alpha \in \Delta} \alpha(r) \alpha(s) \in 4\mathbb{Z}$ for any semi-simple Lie algebra \mathfrak{g} , and that $2\ell_J^3 \hat{\alpha}(r) \hat{\alpha}(s) \hat{\alpha}(p) \in 2\mathbb{Z}$ for any integer or half-integer spin. Including now also the contribution from \mathcal{B}_m , the factor picked up by \mathcal{Z} can be expressed in terms of the anomaly coefficients (3.18) and (3.19):

$$\mathcal{Z}_{m'} = -e^{2\pi i \phi} e^{\pi i r_i s_j (\frac{1}{2} \mathcal{A}^{ij} - \mathcal{A}^{ijk} (\hat{u}_0 - m\omega)_k - \mathcal{A}^{ij\alpha} \xi_\alpha - \mathcal{A}^{ijR} \nu_R)} \mathcal{Z}_m . \quad (\text{C.42})$$

The anomaly cancellation conditions $\mathcal{A}^{ijk} = \mathcal{A}^{ij\alpha} = \mathcal{A}^{ijR} = 0$ and $\mathcal{A}^{ij} \in 4\mathbb{Z}$ imply that the second exponential equals 1. In the first exponential we defined

$$\phi = \frac{1}{2} \hat{\alpha}(r) \hat{\alpha}(s) \hat{\alpha}(p) \sum_{a,I} \sum_{\ell_{aI} > 0}^{j_{aI}} \ell_{aI}^3 = 4\hat{r} \hat{s} \hat{p} \sum_{a,I} \sum_{\ell_{aI} > 0}^{j_{aI}} \ell_{aI}^3 . \quad (\text{C.43})$$

Once again, in an anomaly-free theory $\phi \in \mathbb{Z}$. Indeed, labelling the chiral multiplets by a , their $\mathfrak{su}(2)_{\hat{\alpha}}$ representations by I and dubbing their spin j_{aI} , the only non-integer contributions to ϕ come from representations with $j_{aI} \in 2\mathbb{Z} + \frac{1}{2}$. On the other hand, the contribution of an $\mathfrak{su}(2)_{\hat{\alpha}}$ representation to the pseudo-anomaly coefficient is

$$\mathcal{A}_{aI}^{ij} = \sum_{\ell=-j_{aI}}^{j_{aI}} (\hat{\rho}_{aI} + \ell\hat{\alpha})^i (\hat{\rho}_{aI} + \ell\hat{\alpha})^j . \quad (\text{C.44})$$

Since generic vectors r, s (C.33) have integer components, the condition $\mathcal{A}^{ij} \in 4\mathbb{Z}$ implies that also $\mathcal{A}^{ij} r_i s_j \in 4\mathbb{Z}$ for any choice of r, s . Contracting with the vectors, we obtain

$$\mathcal{A}_{aI}^{ij} r_i s_j = \frac{4}{3} \hat{r} \hat{s} j_{aI} (j_{aI} + 1) (2j_{aI} + 1) \in \begin{cases} 4\mathbb{Z} & \text{if } j_{aI} \in \mathbb{Z} \text{ or } j_{aI} \in 2\mathbb{Z} + \frac{3}{2} \\ 4\mathbb{Z} + 2 & \text{if } j_{aI} \in 2\mathbb{Z} + \frac{1}{2} . \end{cases} \quad (\text{C.45})$$

Therefore, the condition $\mathcal{A}^{ij} \in 4\mathbb{Z}$ requires that the number of $\mathfrak{su}(2)_{\hat{\alpha}}$ representations with $j_{aI} \in 2\mathbb{Z} + \frac{1}{2}$ be even, and this guarantees that $\phi = 0$.

References

- [1] C. Romelsberger, “Counting chiral primaries in $\mathcal{N}=1$, $d=4$ superconformal field theories,” *Nucl. Phys.* **B747** (2006) 329–353, [arXiv:hep-th/0510060](#) [[hep-th](#)].
- [2] J. Kinney, J. M. Maldacena, S. Minwalla, and S. Raju, “An index for 4 dimensional super conformal theories,” *Commun. Math. Phys.* **275** (2007) 209–254, [arXiv:hep-th/0510251](#) [[hep-th](#)].
- [3] J. Bhattacharya, S. Bhattacharyya, S. Minwalla, and S. Raju, “Indices for Superconformal Field Theories in 3,5 and 6 Dimensions,” *JHEP* **02** (2008) 064, [arXiv:0801.1435](#) [[hep-th](#)].
- [4] V. Pestun *et al.*, “Localization techniques in quantum field theories,” *J. Phys.* **A50** (2017) 440301, [arXiv:1608.02952](#) [[hep-th](#)].
- [5] F. A. Dolan and H. Osborn, “Applications of the Superconformal Index for Protected Operators and q -Hypergeometric Identities to $\mathcal{N}=1$ Dual Theories,” *Nucl. Phys.* **B818** (2009) 137–178, [arXiv:0801.4947](#) [[hep-th](#)].
- [6] C. Closset, H. Kim, and B. Willett, “ $\mathcal{N}=1$ supersymmetric indices and the four-dimensional A-model,” *JHEP* **08** (2017) 090, [arXiv:1707.05774](#) [[hep-th](#)].

- [7] C. Closset, H. Kim, and B. Willett, “Seifert fibering operators in 3d $\mathcal{N}=2$ theories,” *JHEP* **11** (2018) 004, [arXiv:1807.02328 \[hep-th\]](#).
- [8] F. Benini and P. Milan, “Black holes in 4d $\mathcal{N}=4$ Super-Yang-Mills,” [arXiv:1812.09613 \[hep-th\]](#).
- [9] A. Cabo-Bizet, D. Cassani, D. Martelli, and S. Murthy, “Microscopic origin of the Bekenstein-Hawking entropy of supersymmetric AdS₅ black holes,” [arXiv:1810.11442 \[hep-th\]](#).
- [10] S. Choi, J. Kim, S. Kim, and J. Nahmgoong, “Large AdS black holes from QFT,” [arXiv:1810.12067 \[hep-th\]](#).
- [11] F. Benini and A. Zaffaroni, “A topologically twisted index for three-dimensional supersymmetric theories,” *JHEP* **07** (2015) 127, [arXiv:1504.03698 \[hep-th\]](#).
- [12] F. Benini, K. Hristov, and A. Zaffaroni, “Black hole microstates in AdS₄ from supersymmetric localization,” *JHEP* **05** (2016) 054, [arXiv:1511.04085 \[hep-th\]](#).
- [13] F. Benini and A. Zaffaroni, “Supersymmetric partition functions on Riemann surfaces,” *Proc. Symp. Pure Math.* **96** (2017) 13–46, [arXiv:1605.06120 \[hep-th\]](#).
- [14] F. Benini, K. Hristov, and A. Zaffaroni, “Exact microstate counting for dyonic black holes in AdS₄,” *Phys. Lett.* **B771** (2017) 462–466, [arXiv:1608.07294 \[hep-th\]](#).
- [15] J. B. Gutowski and H. S. Reall, “Supersymmetric AdS₅ black holes,” *JHEP* **02** (2004) 006, [arXiv:hep-th/0401042 \[hep-th\]](#).
- [16] J. B. Gutowski and H. S. Reall, “General supersymmetric AdS₅ black holes,” *JHEP* **04** (2004) 048, [arXiv:hep-th/0401129 \[hep-th\]](#).
- [17] Z. W. Chong, M. Cvetič, H. Lu, and C. N. Pope, “General non-extremal rotating black holes in minimal five-dimensional gauged supergravity,” *Phys. Rev. Lett.* **95** (2005) 161301, [arXiv:hep-th/0506029 \[hep-th\]](#).
- [18] H. K. Kunduri, J. Lucietti, and H. S. Reall, “Supersymmetric multi-charge AdS₅ black holes,” *JHEP* **04** (2006) 036, [arXiv:hep-th/0601156 \[hep-th\]](#).
- [19] F. Benini and S. Cremonesi, “Partition Functions of $\mathcal{N}=(2, 2)$ Gauge Theories on S² and Vortices,” *Commun. Math. Phys.* **334** (2015) 1483–1527, [arXiv:1206.2356 \[hep-th\]](#).
- [20] F. Benini and W. Peelaers, “Higgs branch localization in three dimensions,” *JHEP* **05** (2014) 030, [arXiv:1312.6078 \[hep-th\]](#).

- [21] C. Closset and H. Kim, “Comments on twisted indices in 3d supersymmetric gauge theories,” *JHEP* **08** (2016) 059, [arXiv:1605.06531 \[hep-th\]](#).
- [22] C. Closset, H. Kim, and B. Willett, “Supersymmetric partition functions and the three-dimensional A-twist,” *JHEP* **03** (2017) 074, [arXiv:1701.03171 \[hep-th\]](#).
- [23] B. Assel, D. Cassani, L. Di Pietro, Z. Komargodski, J. Lorenzen, and D. Martelli, “The Casimir Energy in Curved Space and its Supersymmetric Counterpart,” *JHEP* **07** (2015) 043, [arXiv:1503.05537 \[hep-th\]](#).
- [24] N. Bobev, M. Bullimore, and H.-C. Kim, “Supersymmetric Casimir Energy and the Anomaly Polynomial,” *JHEP* **09** (2015) 142, [arXiv:1507.08553 \[hep-th\]](#).
- [25] N. A. Nekrasov and S. L. Shatashvili, “Bethe/Gauge correspondence on curved spaces,” *JHEP* **01** (2015) 100, [arXiv:1405.6046 \[hep-th\]](#).
- [26] E. Witten, “Constraints on Supersymmetry Breaking,” *Nucl. Phys.* **B202** (1982) 253.
- [27] B. Sundborg, “The Hagedorn transition, deconfinement and $\mathcal{N}=4$ SYM theory,” *Nucl. Phys.* **B573** (2000) 349–363, [arXiv:hep-th/9908001 \[hep-th\]](#).
- [28] O. Aharony, J. Marsano, S. Minwalla, K. Papadodimas, and M. Van Raamsdonk, “The Hagedorn/deconfinement phase transition in weakly coupled large N gauge theories,” *Adv. Theor. Math. Phys.* **8** (2004) 603–696, [arXiv:hep-th/0310285 \[hep-th\]](#).
- [29] S. Benvenuti, B. Feng, A. Hanany, and Y.-H. He, “Counting BPS Operators in Gauge Theories: Quivers, Syzygies and Plethystics,” *JHEP* **11** (2007) 050, [arXiv:hep-th/0608050 \[hep-th\]](#).
- [30] V. Pestun, “Localization of gauge theory on a four-sphere and supersymmetric Wilson loops,” *Commun. Math. Phys.* **313** (2012) 71–129, [arXiv:0712.2824 \[hep-th\]](#).
- [31] C. Closset and I. Shamir, “The $\mathcal{N}=1$ Chiral Multiplet on $T^2 \times S^2$ and Supersymmetric Localization,” *JHEP* **03** (2014) 040, [arXiv:1311.2430 \[hep-th\]](#).
- [32] B. Assel, D. Cassani, and D. Martelli, “Localization on Hopf surfaces,” *JHEP* **08** (2014) 123, [arXiv:1405.5144 \[hep-th\]](#).
- [33] G. Felder and A. Varchenko, “The elliptic gamma function and $sl(3, \mathbb{Z}) \ltimes \mathbb{Z}^3$,” *Adv. Math.* **156** (2000) 44, [arXiv:math/9907061](#).
- [34] E. Witten, “An $SU(2)$ Anomaly,” *Phys. Lett.* **B117** (1982) 324–328.

- [35] S. Elitzur and V. P. Nair, “Nonperturbative Anomalies in Higher Dimensions,” *Nucl. Phys.* **B243** (1984) 205.
- [36] H. Zhang, S. Okubo, and Y. Tosa, “Global Gauge Anomalies for Simple Lie Algebras,” *Phys. Rev.* **D37** (1988) 2946.