

RECEIVED: September 6, 2011

REVISED: October 9, 2011

ACCEPTED: October 9, 2011

PUBLISHED: November 9, 2011

Aspects of AdS/BCFT

Mitsutoshi Fujita,^a Tadashi Takayanagi^a and Erik Tonni^b

^a*Institute for the Physics and Mathematics of the Universe (IPMU), University of Tokyo, Kashiwa, Chiba 277-8582, Japan*

^b*Center for Theoretical Physics, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139, U.S.A.*

E-mail: mitsutoshi.fujita@ipmu.jp, tadashi.takayanagi@ipmu.jp, tonni@mit.edu

ABSTRACT: We expand the results of arXiv:1105.5165, where a holographic description of a conformal field theory defined on a manifold with boundaries (so called BCFT) was proposed, based on AdS/CFT. We construct gravity duals of conformal field theories on strips, balls and also time-dependent boundaries. We show a holographic g-theorem in any dimension. As a special example, we can define a ‘boundary central charge’ in three dimensional conformal field theories and our holographic g-theorem argues that it decreases under RG flows. We also computed holographic one-point functions and confirmed that their scaling property agrees with field theory calculations. Finally, we give an example of string theory embedding of this holography by inserting orientifold 8-planes in $\text{AdS}_4 \times \text{CP}^3$.

KEYWORDS: AdS-CFT Correspondence, Boundary Quantum Field Theory, Conformal Field Models in String Theory

ARXIV EPRINT: [1108.5152](https://arxiv.org/abs/1108.5152)

Contents

1	Introduction	2
2	Holographic dual of BCFT	3
2.1	Neumann boundary condition	3
2.2	Construction of AdS/BCFT	4
3	AdS₃/CFT₂ and boundary entropy	6
3.1	Boundary entropy from disk partition function	6
3.2	Boundary entropy from holographic entanglement entropy	8
4	Holographic dual of intervals and Hawking-page transition	9
4.1	Low temperature phase	9
4.2	High temperature phase	10
4.3	Phase transition	12
5	Holographic g-theorem	13
5.1	Holographic g-theorem in 2D CFT	13
5.2	Holographic g-theorem in higher dimensions	14
6	AdS/BCFT in higher dimension	15
6.1	Holographic dual of balls	15
6.1.1	AdS ₅ case	16
6.1.2	AdS ₄ case	18
6.2	Holographic dual of half space	19
6.3	Holographic dual of strips	19
7	Time-dependent configurations	21
7.1	Description in terms of entangled pair	21
7.2	Comments on g-theorem and topological censorship	22
8	Correlation functions	23
8.1	One point function	23
9	An example of string theory embedding	26
10	Conclusions and discussions	29
A	Calculations of extrinsic curvatures	30
B	Possibility of connected boundary for the annulus	31
C	Boundaries in rotating BTZ black holes	32
D	Details about the one point function	33
E	Discrete symmetries of ABJM theory	35

1 Introduction

The holographic principle [1, 2] has been a very powerful idea which relates the unknown physics of quantum gravity to that of a more familiar non-gravitational theory. It also benefits us in an opposite way. We can conveniently study a strongly coupled quantum field theory by replacing it with a classical gravity theory. One of the most well studied holographic setups is the AdS/CFT correspondence [3–5]. In most of the examples in AdS/CFT studied so far, the holography works as an equivalence between a gravity or string theory on a Anti de-Sitter space (AdS) and a conformal field theory (CFT) on a compact manifold. Therefore it is intriguing to ask what will happen if there are boundaries on the manifold on which the CFT is defined. Such a CFT is called a boundary conformal field theory (BCFT) if a part of conformal symmetry (called boundary conformal symmetry) is preserved at the boundaries.

In the recent paper [6], an effective description of holographic dual of BCFT (AdS/BCFT) has been generally considered and several physical quantities including the partition functions have been computed. We would like to note that in specific setups, a holography construction of BCFT, which is very similar to ours, has already been mentioned in the earlier papers [7, 8]. In this construction, we introduce an extra boundary Q in addition to the asymptotic AdS boundary M such that the boundary of Q coincides with that of M (refer to figure 1). One of the crucial points is that we impose the Neumann boundary condition instead of the standard Dirichlet one in the gravity sector. A new ingredient in this holography is that we can add degrees of freedom localized at the boundary. In the two dimensional CFT, this is measured by so called the boundary entropy or g-function [9]. We would also like to mention that different constructions of holographic dual of field theories with boundaries can be found in [10–12].

The purpose of this paper is to further study the properties of AdS/BCFT. At the same time, we will also expose detailed calculations on the results reported briefly in the letter [6]. Some of new results in this paper are described as follows. We construct gravity duals of conformal field theories on strips, balls and also time-dependent boundaries in any dimension. We prove a holographic g-theorem in any dimension. As a particular example, we introduce a ‘boundary central charge’ in three dimensional conformal field theories. Our holographic g-theorem argues that it decreases under RG flows. We also computed holographic one-point functions. Finally, we give an example of string theory embedding of this holography.

This paper is organized as follows. In section two, we review the general prescription of AdS/BCFT. In section three, we explain the holographic calculation of boundary entropy in two dimensional conformal field theories. In section four, we study the gravity dual of a CFT on an interval and analyze the Hawking-Page phase transition. In section five, we show the holographic g-theorem in any dimension. In section six, we present holographic duals of CFTs defined on balls or strips in higher dimensions. In section seven, we consider examples of AdS/BCFT with time-dependent boundaries. In section eight, we calculate the holographic one point functions. In section nine, we present an example of string theory embedding of AdS/BCFT. In section ten, we summarize conclusions and discuss future problems.




Figure 1. A schematic setup of AdS/BCFT. The CFT lives on M , which has the boundary ∂M . Its gravity dual is denoted by N and its asymptotically AdS is M . The boundary ∂M is extended into the bulk AdS, which constitutes the boundary Q .

2 Holographic dual of BCFT

We would like to formulate a holographic dual of CFT defined on a manifold M with a boundary ∂M . We argue that this is given by generalizing the AdS/CFT correspondence [3] in the following way. To have a gravity dual, we extend d dimensional manifold M to a $d + 1$ dimensional manifold N so that $\partial N = M \cup Q$, where Q is a manifold homologous to M . The conformal invariance in the bulk of M requires that N is a part of AdS space.

In the standard AdS/CFT, we impose the Dirichlet boundary condition at the boundary of AdS and following this we assume the Dirichlet boundary condition on M . On the other hand, we impose a Neumann boundary condition on Q [6] as we will explain later. The reason for this is that this boundary should be dynamical from the viewpoint of holography and there is no natural definite metric on Q specified from the data in the CFT side. Notice that here the boundary Q is no longer asymptotically AdS. On the other hand, if we impose the Neumann boundary condition on a boundary which is parallel with the AdS boundary M , the setup coincides with that of the Randall-Sundrum models [13]. Also, we would like to note that in principle, it is also possible to adopt the Neumann boundary condition at the AdS boundary M as discussed in [14].

2.1 Neumann boundary condition

To make the variational problem sensible, it is conventional to add the Gibbons-Hawking boundary term [15] to the Einstein-Hilbert action:

$$I = \frac{1}{16\pi G_N} \int_N \sqrt{-g}(R - 2\Lambda) + \frac{1}{8\pi G_N} \int_Q \sqrt{-h}K. \quad (2.1)$$

The metric of N and Q is denoted by g and h . $K = h^{ab}K_{ab}$ is the trace of extrinsic curvature. The extrinsic curvature K_{ab} is defined by

$$K_{ab} = \nabla_a n_b, \quad (2.2)$$

where n is the unit vector normal to Q and here we implicitly assume a projection onto Q from N . For example, in the coordinate system (so called Gaussian normal coordinate)

$$ds^2 = dr^2 + h_{ab}dx^a dx^b, \quad (2.3)$$

we can explicitly show that

$$K_{ab} = \frac{1}{2} \frac{\partial h_{ab}}{\partial r}. \tag{2.4}$$

Now let us consider the variation of metric in the above action. After a partial integration, we find

$$\delta I = \frac{1}{16\pi G_N} \int_Q \sqrt{-h} \left(K_{ab} \delta h^{ab} - K h_{ab} \delta h^{ab} \right). \tag{2.5}$$

Notice that the terms which involve the derivative of δh_{ab} cancel out thanks to the boundary term. In this way, by imposing the Neumann boundary condition instead of the Dirichlet one, we obtain the boundary condition

$$K_{ab} - h_{ab} K = 0. \tag{2.6}$$

It is also possible to add some matter fields localized on Q and consider a generalized action by adding

$$I_Q = \int \sqrt{-h} L_Q. \tag{2.7}$$

This modifies (2.6) into

$$K_{ab} - h_{ab} K = 8\pi G_N T_{ab}^Q, \tag{2.8}$$

where we defined

$$T^{Qab} = \frac{2}{\sqrt{-h}} \frac{\delta I_Q}{\delta h_{ab}}. \tag{2.9}$$

Before we go on, we would like to briefly remind us of the standard treatment of the asymptotically AdS boundary M . We introduce the same Gibbons-Hawking boundary term on another boundary M as usual. In this case we do not need to require (2.9) and instead we impose the Dirichlet boundary condition $\delta h_{ab} = 0$, regarding the Brown-York tensor [16, 17]

$$\tau_{ab} = \frac{1}{8\pi G_N} (K_{ab} - h_{ab} K) - T_{ab}^M \tag{2.10}$$

as the holographic energy stress tensor [18–20].

2.2 Construction of AdS/BCFT

As a simple class of examples, we would like to assume that the boundary matter lagrangian L_Q is simply a constant. This leads us to consider the following action (we omit the boundary terms for the AdS boundary M)

$$I = \frac{1}{16\pi G_N} \int_N \sqrt{-g} (R - 2\Lambda) + \frac{1}{8\pi G_N} \int_Q \sqrt{-h} (K - T). \tag{2.11}$$

The constant T is interpreted as the tension of the boundary Q . In AdS/CFT, a $d + 1$ dimensional AdS space (AdS_{d+1}) is dual to a d dimensional CFT. The geometrical $\text{SO}(2, d)$ symmetry of AdS is equivalent to the conformal symmetry of the CFT. When we put a $d - 1$ dimensional boundary to a d dimensional CFT such that the presence of the boundary breaks $\text{SO}(2, d)$ into $\text{SO}(2, d - 1)$, this is called a boundary conformal field theory

(BCFT) [21, 22]. Note that this symmetry structure looks the same as that in the holography for defect or Janus CFTs [7, 23–26].

The boundary condition (2.8) for the system (2.11) leads to

$$K_{ab} = (K - T)h_{ab}. \quad (2.12)$$

By taking its trace, we obtain

$$K = \frac{d}{d-1}T. \quad (2.13)$$

To realize expected symmetries, we employ the following ansatz of the metric

$$ds^2 = d\rho^2 + \cosh^2 \frac{\rho}{R} \cdot ds_{AdS_d}^2. \quad (2.14)$$

If we assume that ρ takes all values from $-\infty$ to ∞ , then (2.14) is equivalent to the AdS_{d+1} . To see this, let us assume the Poincare metric of AdS_d by setting

$$ds_{AdS_d}^2 = R^2 \frac{-dt^2 + dy^2 + d\vec{w}^2}{y^2}, \quad (2.15)$$

where $\vec{w} \in \mathbb{R}^{d-2}$. Remember that the cosmological constant Λ is related to the AdS radius R by

$$\Lambda = -\frac{d(d-1)}{2R^2}. \quad (2.16)$$

If we define new coordinates z and x by

$$z = \frac{y}{\cosh \frac{\rho}{R}}, \quad x = y \tanh \frac{\rho}{R}, \quad (2.17)$$

then we obtain the familiar form of the Poincare metric of AdS_{d+1}

$$ds^2 = R^2 \frac{dz^2 - dt^2 + dx^2 + d\vec{w}^2}{z^2}. \quad (2.18)$$

Now to realize a gravity dual of BCFT, we will put a boundary Q at $\rho = \rho_*$ and this means that we restrict the spacetime to the region $-\infty < \rho < \rho_*$ as depicted in figure 2. The extrinsic curvature on Q can be found from (2.4)

$$K_{ab} = \frac{1}{R} \tanh \left(\frac{\rho}{R} \right) h_{ab}. \quad (2.19)$$

By using (2.12), ρ_* is determined by the tension T as follows

$$T = \frac{d-1}{R} \tanh \frac{\rho_*}{R}. \quad (2.20)$$

This leads to the constraint $-(d-1) \leq TR \leq d-1$.




Figure 2. The holographic dual of a half line. The spacetime dual to BCFT is restricted to the region $-\infty < \rho < \rho_*$ and is surrounded by the shaded region.

3 AdS₃/CFT₂ and boundary entropy

Here let us concentrate on the case of $d = 2$, which describes the two dimensional BCFT. The reason that this setup is special is that it has been well studied in the subject of two dimensional CFT [27]. Moreover, the BCFT has an interesting quantity called the boundary entropy (or g -function) introduced in [9].

The boundary state of a BCFT with a boundary condition α is denoted by $|B_\alpha\rangle$ below. The function called g is defined by the disk amplitude [9]

$$g_\alpha = \langle 0|B_\alpha\rangle, \quad (3.1)$$

where $|0\rangle$ is the vacuum state. The boundary entropy $S_{bdy}^{(\alpha)}$ is simply defined by

$$S_{bdy}^{(\alpha)} = \log g_\alpha. \quad (3.2)$$

Below we will present two different calculations of the boundary entropy, which turns out to coincide as expected. Later, in section 4.2 we will provide a third independent holographic calculation of the boundary entropy.

3.1 Boundary entropy from disk partition function

Consider a holographic dual of a CFT on the round disk defined by $t_E^2 + x^2 \leq r_D^2$ in the Euclidean AdS₃ spacetime

$$ds^2 = R^2 \frac{dz^2 + d\tau^2 + dx^2}{z^2}, \quad (3.3)$$

where τ is the Euclidean time. In the Euclidean formulation, the action (2.11) is now replaced by

$$I_E = -\frac{1}{16\pi G_N} \int_N \sqrt{g}(R - 2\Lambda) - \frac{1}{8\pi G_N} \int_Q \sqrt{h}(K - T). \quad (3.4)$$

Note that ρ_* is related to the tension T of the boundary via (2.20). When the BCFT is defined on the half space $x < 0$, its gravity dual has been found in previous section. Therefore we can find the gravity dual of the BCFT on the round disk by first applying

the conformal map

$$\begin{aligned} x'_\mu &= \frac{x_\mu + c_\mu x^2}{1 + 2(c \cdot x) + c^2 \cdot x^2}, \\ z' &= \frac{z}{1 + 2(c \cdot x) + c^2 \cdot x^2}, \end{aligned} \tag{3.5}$$

where $c^\mu = (c^\tau, c^x)$ are arbitrary constants, and then performing a proper translation [28]. Finally, we obtain the following domain in AdS₃

$$\tau^2 + x^2 + (z - \sinh(\rho_*/R)r_D)^2 - r_D^2 \cosh^2(\rho_*/R) \leq 0. \tag{3.6}$$

In this way we found that the holographic dual of BCFT on a round disk is given by a part of the two dimensional round sphere as described in figure 3. A larger value of tension corresponds to the larger radius.

Now we would like to calculate the disk partition function in order to obtain the boundary entropy. For this we just need to evaluate (3.4) in the domain (3.6). In the end we find

$$I_E = \frac{R}{4G_N} \left(\frac{r_D^2}{2\epsilon^2} + \frac{r_D \sinh(\rho_*/R)}{\epsilon} + \log(\epsilon/r_D) - \frac{1}{2} - \frac{\rho_*}{R} \right), \tag{3.7}$$

where we introduced the cutoff such that $z > \epsilon$. By adding the counter term on the AdS boundary, we can subtract the divergent terms in (3.7). The difference of the partition function between $\rho = 0$ and $\rho = \rho_*$ is given by

$$I_E(\rho_*) - I_E(0) = -\frac{\rho_*}{4G_N}. \tag{3.8}$$

Since the partition function is given by $Z = e^{-S_E}$, we obtain the boundary entropy

$$S_{bdy} = -I_E = \frac{\rho_*}{4G_N} = \frac{c}{6} \operatorname{arctanh}(RT), \tag{3.9}$$

where implicitly we assumed $S_{bdy} = 0$ at $T = 0$ because the boundary contributions vanish in this case.

By employing the integrals computed to get (3.7), we can evaluate the action in the case of a domain M in the boundary given by an annulus. We consider as $Q = Q_1 \cup Q_2$ the disconnected configuration shown in figure 4 and we discuss a connected configuration in the appendix B.

Since the orientation of the two surfaces Q_1 and Q_2 of the disconnected boundaries are opposite, the definition of tension is opposite. This means that $\rho_{*,1} > 0$ at Q_1 corresponds to $T > 0$ (or equally large boundary entropy), while $\rho_{*,2} > 0$ at Q_2 corresponds to $T < 0$ (or smaller boundary entropy). Thus, the result is given by the difference of two contributions like (3.7)

$$I_E = \frac{R}{4G_N} \left[\frac{r_{D,2}^2 - r_{D,1}^2}{2\epsilon^2} + \frac{r_{D,2} \sinh(\rho_{*,2}/R) - r_{D,1} \sinh(\rho_{*,1}/R)}{\epsilon} + \log \left(\frac{r_{D,1}}{r_{D,2}} \right) - \frac{\rho_{*,2} - \rho_{*,1}}{R} \right] \tag{3.10}$$

where the logarithmic divergence is canceled in the difference.




Figure 3. The holographic dual of a disk.

3.2 Boundary entropy from holographic entanglement entropy

Another way to extract the boundary entropy is to calculate the entanglement entropy. The entanglement entropy S_A with respect to the subsystem A is defined by the von Neumann entropy $S_A = -\text{Tr} \rho_A \log \rho_A$ for the reduced density matrix ρ_A . The reduced density matrix ρ_A is defined by tracing out the subsystem B , which is the complement of A . In quantum field theories, we specify the subsystem A by dividing a time slice into two regions. For a two dimensional CFT with a boundary (i.e. BCFT) we can generally obtain the following result [29]

$$S_A = \frac{c}{6} \log \frac{l}{\epsilon} + \log g_\alpha, \quad (3.11)$$

where c is the central charge and ϵ is the UV cut off (or lattice spacing); the subsystem A is chosen to be an interval with length l such that it includes the boundary.

In AdS/CFT, the holographic entanglement entropy [30–32] is given in terms of the area of the codimension two minimal surface (called γ_A) which ends at ∂A

$$S_A = \frac{\text{Area}(\gamma_A)}{4G_N}. \quad (3.12)$$

This calculation of boundary entropy from the holographic entanglement entropy has been first applied to Janus CFTs in [33]. Moreover, for supersymmetric Janus CFTs, the excellent agreement between the holographic result and the CFT result has been confirmed in [34, 35].

Consider the gravity dual of two dimensional BCFT on a half space $x < 0$ in the coordinate (3.3). By taking the time slice $\tau = 0$, we define the subsystem A by the interval $-l \leq x \leq 0$. In this case, the minimal surface (or geodesic line) γ_A is given by $x^2 + z^2 = l^2$. If we go back to the coordinate system (2.14) and (2.15), then γ_A is simply given by $\tau = 0, y = l$ and $-\infty < \rho \leq \rho_*$. Then

$$S_A = \frac{1}{4G_N} \int_{-\rho_\infty}^{\rho_*} d\rho = \frac{\rho_* + \rho_\infty}{4G_N}. \quad (3.13)$$

Here ρ_∞ is related to the UV cut off using (2.17) via $\rho_\infty = R \log \frac{2l}{\epsilon}$. By subtracting the bulk contribution which is divergent as in (3.11), we finally find

$$S_{bdy} = S_A(\rho_*) - S_A(0) = \frac{\rho_*}{4G_N}. \quad (3.14)$$

This indeed agrees with (3.9).




Figure 4. The holographic dual of an annulus made by two disconnected surfaces in the bulk.

Actually, in the formula (3.12), we need to choose the end point of the geodesic γ_A on Q such that the total length takes the minimum value. Indeed, we can check explicitly that this minimum is realized when γ_A is a circle with radius l as is assumed in (3.13).

4 Holographic dual of intervals and Hawking-page transition

So far we studied the holographic dual of BCFT in the presence of a single boundary. As a next step, we would like to analyze the holographic dual of two dimensional CFT on an interval in the setup of AdS₃/CFT₂ as one of the simplest examples with multiple boundaries. We assume such a system at finite temperature and there are two candidates for the bulk geometry, one of them is the thermal AdS₃ and the other is the BTZ black hole (AdS₃ black hole) [36]. In the absence of the boundaries, the former is favored at low temperature, while the latter at high temperature. It is natural to expect a similar phase structure and indeed we will confirm this below.

4.1 Low temperature phase

At low temperature, the bulk geometry is expected to be given by the thermal AdS₃ defined by the metric

$$ds^2 = R^2 \left(\frac{d\tau^2}{z^2} + \frac{dz^2}{h(z)z^2} + \frac{h(z)}{z^2} dx^2 \right), \tag{4.1}$$

where $h(z) = 1 - (z/z_0)^2$. The periodicity of the Euclidean time τ , denoted by the inverse temperature $1/T_{BCFT} (\equiv 2\pi z_H)$, can be chosen arbitrary, while that of the space direction x is determined to be $2\pi z_0$ by requiring the smoothness.

We can describe the boundary Q by the curve $x = x(z)$. The space-like unit vector n^μ normal to the surface Q is given by

$$(n^\tau, n^z, n^x) = (0, -x'(z)h(z)^2, 1) \cdot \frac{z}{R\sqrt{h(z)(1 + h(z)^2 x'(z)^2)}}. \tag{4.2}$$

The extrinsic curvature K_{ab} can be computed by following the procedure explained in the appendix A. In the end, the boundary condition (2.12) leads to the following constraint on

the profile of Q

$$\frac{dx}{dz} = \frac{RT}{h(z)\sqrt{h(z) - R^2T^2}}, \quad (4.3)$$

which is solved (fixing the constant shift by setting $x(0) = 0$)

$$x(z) = z_0 \cdot \arctan\left(\frac{RTz}{z_0\sqrt{h(z) - R^2T^2}}\right). \quad (4.4)$$

Notice that $x'(z)$ gets divergent at $z_* = z_0\sqrt{1 - R^2T^2}$ and thus this should be the turning point. Thus the boundary Q extends from $x = 0$ to $x = \pi z_0$. Assuming $T > 0$, the bulk spacetime N is defined by the sum of $(-\pi z_0 \leq x \leq 0, 0 < z \leq z_0)$ and $(0 < x \leq \pi z_0, z(x) < z < z_0)$, where $z(x)$ is the inverse function of (4.4) and its extension to $\frac{\pi}{2}z_0 < x < \pi z_0$. This is described in figure 5(a).

Now the Euclidean action (3.4) reads

$$I_E = \frac{Rz_H}{G_N} \int_{\epsilon}^{z_*} \frac{dz}{z^3} \left(x(z) + \frac{\pi z_0}{2}\right) + \int_{z_*}^{z_0} \frac{dz}{z^3} (\pi z_0) - \frac{z_H T R^2}{2G_N} \int_{\epsilon}^{z_*} \frac{dz}{z^2 \sqrt{h(z) - R^2T^2}}, \quad (4.5)$$

where ϵ is the UV cut off as before. To evaluate (4.5) by eliminating the divergence, we need to be careful in that we have to regard $2\pi\tilde{z}_0$ as the physical radius, defined by

$$\tilde{z}_0 = \sqrt{f(\epsilon)} z_0, \quad (4.6)$$

by matching the asymptotic geometry at $z = \epsilon$. Also the contribution Gibbons-Hawking term at the AdS boundary M is vanishing as usual, by using the boundary integral of $K - K^{(0)}$ instead of that of K , where $K^{(0)}$ is the trace of extrinsic curvature for the pure AdS₃ (3.3). In the end, we obtain the result

$$I_E = -\frac{\pi R z_H}{8G_N z_0} = -\frac{\pi}{24} \cdot \frac{c}{\Delta x \cdot T_{BCFT}}, \quad (4.7)$$

where we employed the well-known relation [37] between the AdS₃ radius R and the central charge c of CFT_2 , given by

$$c = \frac{3R}{2G_N}. \quad (4.8)$$

Note that the final result (4.7) does not depend on the tension T and we can confirm that (4.7) is also correct when $T < 0$.

4.2 High temperature phase

We expect that in the higher temperature phase the bulk is described by a part of the BTZ black hole

$$ds^2 = R^2 \left(\frac{f(z)}{z^2} d\tau^2 + \frac{dz^2}{f(z)z^2} + \frac{dx^2}{z^2} \right), \quad (4.9)$$

where $f(z) = 1 - (z/z_H)^2$. The Euclidean time τ is compactified on a circle such that $\tau \sim \tau + 2\pi z_H$ and thus the temperature in the dual BCFT is $T_{BCFT} = \frac{1}{2\pi z_H}$. The length of the interval is again denoted by $\Delta x = \pi z_0$.




Figure 5. The holographic dual of an interval at low temperature (a) and high temperature (b).

We specify the boundary Q by the profile $x = x(z)$.

The space-like unit vector n^μ normal to the surface Q is given by

$$(n^\tau, n^z, n^x) = (0, -x'(z)f(z), 1) \cdot \frac{z}{R\sqrt{h(z)(1+f(z)x'(z)^2)}}. \quad (4.10)$$

The extrinsic curvature K_{ab} can be again computed by following the procedure explained in the appendix A. In the end, we find that the boundary condition (2.12) requires

$$\frac{dx}{dz} = \frac{RT}{\sqrt{1-R^2T^2f(z)}}. \quad (4.11)$$

This is solved as

$$x(z) = z_H \cdot \operatorname{arcsinh} \left(\frac{RTz}{z_H \sqrt{1-R^2T^2}} \right). \quad (4.12)$$

To realize the holographic dual of the interval we need two boundaries as described in figure 5(b). In the appendix C we study the rotating BTZ black hole generalizing (4.12) for that case (see (C.6)).

Now we are able to evaluate the Euclidean action (3.4) in the form

$$I_E = 2I_{bdy} + I_{bulk}, \quad (4.13)$$

where $2I_{bdy}$ is the contribution from the presence of two boundaries which is vanishing if we set $T = 0$. I_{bulk} is the bulk contribution which does not depend on T . To calculate them we subtract the divergence in a standard holographic renormalization. For the bulk part, we obtain

$$I_{bulk} = -\frac{\pi c}{6} \Delta x \cdot T_{BCFT}, \quad (4.14)$$

where we have to be careful about the issue like (4.6). This result (4.14) clearly agrees with what we expect from the standard CFT results. On the other hand, each of two boundary contributions is found to be

$$\begin{aligned} \frac{Rz_H}{2G_N} \int_\epsilon^{z_H} dz \frac{x(z)}{z^3} - \frac{TR^2z_H}{4G_N} \int_\epsilon^z dz \frac{1}{z^2 \sqrt{1-R^2T^2f(z)}} &= -\frac{Rz_H}{4G_N} \left[\frac{x(z)}{z^2} \right]_\epsilon^{z_H} \\ &= -\frac{\rho_*}{4G_N} + (\text{divergent terms}), \end{aligned} \quad (4.15)$$

where we relate the tension T to ρ_* via the previous relation (2.20) and the index α describes each of the two boundary. Therefore we obtain the contribution to the Euclidean action from each of the boundaries

$$I_{bdy} = -\frac{\rho_*}{4G_N} = -\frac{c}{6}\operatorname{arctanh}(RT). \quad (4.16)$$

Thus the total action is found to be

$$I_E = -\frac{\pi c}{6}\Delta x \cdot T_{BCFT} - \frac{c}{3}\operatorname{arctanh}(RT). \quad (4.17)$$

The total entropy of this thermal system is found from (4.14) and (4.16)

$$S_{thermal} = \frac{\pi}{3}c\Delta x \cdot T_{BCFT} + \frac{c}{3}\operatorname{arctanh}(RT) \quad (4.18)$$

We would also like to point out that this calculation offers one more different calculation of boundary entropy S_{bdy} in AdS/CFT. Consider a BCFT at a finite temperature T_{BCFT} , in other words, a CFT defined on a cylinder, the two boundary conditions imposed on its two boundaries are denoted by α and β . They are described by the boundary states $|B_\alpha\rangle$ and $|B_\beta\rangle$. The partition function $Z_{\alpha\beta}$ on a cylinder, whose length is denoted by Δx , gets factorized in the high temperature limit $T_{BCFT}\Delta x \gg 1$

$$\langle B_\alpha | e^{-H\Delta x} | B_\beta \rangle \simeq g_\alpha g_\beta e^{-E_0\Delta x}, \quad (4.19)$$

where H is the Hamiltonian (in closed string channel) and E_0 is the ground state energy. The final factor $e^{-E_0\Delta x}$ is interpreted as the thermal energy for the CFT as is clear in opens string channel. Therefore the contribution from the presence of boundary is the product of g-function $g_\alpha g_\beta$ [9]. In our holographic calculation, this means $g = e^{S_{bdy}} = e^{-I_{bdy}}$ and this is indeed true by comparing (4.16) and (3.9).

4.3 Phase transition

Let us examine when either of the two phases is favored. To see this we compare (4.7) and (4.17) and pick up the smaller one. In this way we find that the black hole phase is realized if and only if

$$\Delta x \cdot T_{BCFT} > -\frac{1}{\pi}\operatorname{arctanh}(RT) + \sqrt{\frac{1}{4} + \frac{1}{\pi^2}\operatorname{arctanh}^2(RT)}, \quad (4.20)$$

as plotted in figure 6. At lower temperature, the thermal AdS phase is favored. At vanishing tension $T = 0$, the phase boundary $z_0 = z_H$ coincides with that of the Hawking-Page transition [38]. As the tension gets larger, the critical temperature gets lower. This is consistent with the fact that the entropy S_{bdy} carried by the boundary increases as the tension does. Note also that the critical temperature gets vanishing at the maximum tension $TR = 1$ and that it gets divergent at the smallest tension $TR = -1$. This phase transition is first order and can be regarded as a generalization of the Hawking-Page phase transition dual to the confinement/deconfinement transition in gauge theories [39].




Figure 6. The plot of the critical temperature of the phase transition between black hole phase (the region above the curve) and thermal AdS phase (the region below the curve).

5 Holographic g-theorem

In two dimension, the central charge c is a very useful quantity which characterizes the degrees of freedom of a given CFT. Moreover, there is a well-known fact, so called c-theorem, that the central charge decreases under the RG flow. We can construct a c-function which interpolates the two central charges in two CFTs which are connected by a RG flow such that it is a monotonically decreasing function [40]. Holographic proofs of c-theorems have been obtained e.g. in [41, 42].

In the case of BCFT, an analogous quantity is actually known and is called g-function or boundary entropy. At fixed points of boundary RG flow, they are reduced to that of BCFT as already mentioned in (3.2). The monotonicity similar to the c function has been conjectured in [9] and been shown in [43]. This is called the g-theorem. Here we would like to study the holographic proof of this g-theorem. Refer to [44] for an earlier work where the holographic c-theorem has been studied in a probe approximation.

5.1 Holographic g-theorem in 2D CFT

Consider the $AdS_3/BCFT_2$ setup and we just consider a pure gravity in the bulk as we would like to keep the bulk conformal invariance. Since all solutions to the Einstein equation with a negative cosmological constant are locally AdS_3 , we can assume that the gravity dual is given just by cutting a AdS_3 along a boundary Q . We describe the boundary Q by the curve $x = x(z)$ in the metric (3.3) as before. We assume generic matter fields on Q and this leads to the energy stress tensor T_{ab}^Q term in the boundary condition as explained in (2.8). It is easy to check the energy conservation $\nabla^a T_{ab}^Q = 0$ because $\nabla^a (K_{ab} - K h_{ab}) = R_{nb} \propto g_{nb} = 0$ in Einstein manifolds, where n is the Gaussian normal coordinate which is normal to Q .

Now we would like to require that the matter fields on the boundary is physically sensible. In particular, we impose the null energy condition (or equally weaker energy condition) as is done usually for the holographic proof of c-theorem. It is given by the

following inequality for any null vector N^a

$$T_{ab}^Q N^a N^b \geq 0. \tag{5.1}$$

In our case, we can choose

$$N^t = \pm 1, \quad N^z = \frac{1}{\sqrt{1 + (x'(z))^2}}, \quad N^x = \frac{x'(z)}{\sqrt{1 + (x'(z))^2}}. \tag{5.2}$$

Then the condition (5.1) leads to

$$(K_{ab} - Kh_{ab})N^a N^b = -\frac{R \cdot x''(z)}{z(1 + (x'(z))^2)^{3/2}} \geq 0. \tag{5.3}$$

Thus we obtained the condition

$$x''(z) \leq 0. \tag{5.4}$$

Since at a fixed point the boundary entropy is given by $S_{bdy} = \frac{\rho_*}{4G_N}$ and we have the relation $\frac{x}{z} = \sinh(\rho_*/R)$ on the boundary Q , we would like to propose the following g -function

$$\log g(z) = \frac{R}{4G_N} \cdot \operatorname{arcsinh}\left(\frac{x(z)}{z}\right). \tag{5.5}$$

By taking derivative, we get

$$\frac{\partial \log g(z)}{\partial z} = \frac{x'(z)z - x(z)}{\sqrt{z^2 + x^2}}. \tag{5.6}$$

Indeed we can see that $x'z - x$ is negative because this is vanishing at $z = 0$ and (5.4) leads to $(x'z - x)' = x''z \leq 0$. In this way, we manage to derive the g -theorem in our setup. Notice also that we can choose $x(z)$ such that $g(z)$ flows from g_{UV} to g_{IR} . In this case we always have $g_{UV} > g_{IR}$.

5.2 Holographic g -theorem in higher dimensions

Consider a d dimensional CFT on $\mathbb{R}^{1,d-1}$, whose coordinate is denoted by (t, x_1, \dots, x_{d-1}) . We put a boundary at $x_1 = 0$ and consider the theory on the half space defined by $x_{d-1} < 0$. We assume a boundary relevant perturbation and would like to find its holographic dual. This gravity dual should have the translation invariance and the Lorentz invariance $SO(1, d-2)$ in the directions $(x_0, x_1, \dots, x_{d-2})$. Therefore we can assume the following form of the metric for the holographic dual

$$ds^2 = A(x_{d-1}, z)dx_{d-1}^2 + B(x_{d-1}, z)dx_{d-1}dz + C(x_{d-1}, z)dz^2 + D(x_{d-1}, z)(-dt^2 + dx_1^2 + \dots + dx_{d-2}^2). \tag{5.7}$$

By coordinate transformations of x_{d-1} and z , we can set $B(x_{d-1}, z) = 0$, $D(x_{d-1}, z) = \frac{1}{z^2}$. Then by requiring the vacuum Einstein equation (6.2), we find that allowed solutions are either pure AdS space or AdS Schwarzschild black holes. Since we are interested in zero

temperature setup, the metric should be simply given by that of the pure AdS_{d+1} . We specify the boundary Q by $x_{d-1} = x(z)$ again. We can choose the null vector as follows:

$$N^t = \pm 1, \quad N^z = \frac{1}{\sqrt{1+(x'(z))^2}}, \quad N^{x_{d-1}} = \frac{x'(z)}{\sqrt{1+(x'(z))^2}}, \quad N^{x_i} = 0, \quad (i = 1, 2, \dots, d-2). \quad (5.8)$$

Then the condition (5.1) leads to

$$(K_{ab} - Kh_{ab})N^a N^b = -\frac{R \cdot x''(z)}{z(1+(x'(z))^2)^{3/2}} \geq 0. \quad (5.9)$$

As in the AdS_3 case, this means that ρ_* is a monotonically decreasing function under the RG flow. Therefore we can, for example, choose (5.5) to be an analogue of holographic g-function in higher dimension. The precise relation to the partition function on a ball will be analyzed in the next section.

6 AdS/BCFT in higher dimension

6.1 Holographic dual of balls

Consider a $d + 1$ dimensional (Euclidean) AdS space with the Poincare metric

$$ds^2 = R^2 \frac{dz^2 + dx_\mu dx^\mu}{z^2}. \quad (6.1)$$

This satisfies the Einstein equation $R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} + \Lambda g_{\mu\nu} = 0$. We find

$$R_{\mu\nu} + \frac{d}{R^2}g_{\mu\nu} = 0, \quad R = -\frac{d(d+1)}{R^2}, \quad \Lambda = -\frac{d(d-1)}{2R^2}. \quad (6.2)$$

To find a holographic dual of d dimensional ball B_d with radius r_B , we can act the conformal transformation [28]

$$\begin{aligned} x'_\mu &= \frac{x_\mu + cx^2}{1 + 2(c \cdot x) + c^2 \cdot x^2}, \\ z' &= \frac{z}{1 + 2(c \cdot x) + c^2 \cdot x^2}, \end{aligned} \quad (6.3)$$

on the d dimensional half space. In this way, we find that the following surface satisfies the constraint (2.12)

$$x_0^2 + x_1^2 + \dots + x_{d-1}^2 + (z - r_B \sinh(\rho_*/R))^2 - r_B^2 \cosh^2(\rho_*/R) = 0, \quad (6.4)$$

where ρ_* is related to the tension T as in (2.20). Then, we define $r(z)$ as the radius of the sphere at a slice of z as follows:

$$r(z) = \sqrt{r_B^2 \cosh^2\left(\frac{\rho_*}{R}\right) - \left(z - r_B \sinh\left(\frac{\rho_*}{R}\right)\right)^2}. \quad (6.5)$$

The Euclidean action (3.4) is evaluated as follows

$$\begin{aligned}
 I_E &= \frac{dR^{d-1}B_d}{8\pi G_N} \int_\epsilon^{r_B e^{\rho_*/R}} dz \frac{r(z)^d}{z^{d+1}} - \frac{TR^d S_d}{8\pi(d-1)G_N} \int_\epsilon^{r_B e^{\rho_*/R}} dz \frac{r(z)^{d-1} \sqrt{r(z)'^2 + 1}}{z^d} \quad (6.6) \\
 &= \frac{R^{d-1}}{8\pi G_N} \frac{2\pi^{d/2}}{\Gamma(d/2)} \int_\epsilon^{r_B e^{\rho_*/R}} dz \left(\frac{r(z)^d}{z^{d+1}} - r_B \sinh \frac{\rho_*}{R} \frac{r(z)^{d-2}}{z^d} \right), \quad S_d = \frac{2\pi^{(d+1)/2}}{\Gamma((d+1)/2)},
 \end{aligned}$$

where B_d and S_d are the volume of d dimensional unit ball and sphere, respectively. In the last line of (6.6), the formula $dB_d = S_{d-1}$ and (2.20) are used. Notice that for $d = 2$, the analysis has been already done in 3.1.

6.1.1 AdS₅ case

In particular we consider the AdS₅ case $d = 4$. By subtracting the divergences simply,¹ the result is expressed as

$$I_E = \frac{\pi R^3}{16G_N} \left(5 - 2e^{-\frac{2\rho_*}{R}} + 4\frac{\rho_*}{R} + 4\log r_B \right). \quad (6.7)$$

For a 4D CFT with the central charge a , this can be rewritten as follows

$$I_E = \frac{a}{2} \left(5 - 2e^{-\frac{2\rho_*}{R}} + 4\frac{\rho_*}{R} + 4\log r_B \right). \quad (6.8)$$

Notice that the holographic g-theorem (5.9) argues that this increases monotonically under the RG flow as opposed to the AdS₃ case.

Using the Euclidean action (6.8) and holography, we can reproduce the central charge of the even dimensional theory via the trace anomaly. Since we consider the theory on the $d = 4$ ball (a disk for $d = 2$) for the field theory side, we have the scale r_B , namely the radius of the ball. The conformal invariance is quantum mechanically broken because of the trace anomaly. Under the Weyl transformation $\delta g_{\mu\nu} = 2\delta r_B g_{\mu\nu}/r_B$, the trace anomaly for CFT's is then described by [30–32, 45]

$$\begin{aligned}
 r_B \frac{\delta \log Z(B_d)}{\delta r_B} &= \int_{B_d} d^d x \sqrt{g} \langle T^\mu{}_\mu \rangle = -2(-1)^{d/2} a \int_{B_d} \sqrt{g} E_d \\
 &= -2(-1)^{d/2} A,
 \end{aligned} \quad (6.9)$$

where $Z(B_d) = e^{-I_E(B_d)}$ is the partition function for the CFT, A is the central charge in particular $A = \frac{c}{12}$ for $d = 2$ and $A = a$ for $d = 4$. The energy stress tensor is normalized such that $T_{\mu\nu} \equiv -(2/\sqrt{g})\delta I/\delta g^{\mu\nu}$ for the CFT's action I . Here, we used the fact that for theories on the ball, the following normalized Euler density E_d in d dimension only contributes to the trace anomaly:

$$\int_{S_d} d^d x \sqrt{g} E_d = 2, \quad \int_{B_d} d^d x \sqrt{g} E_d = 1. \quad (6.10)$$

¹We just subtract terms of the form $\sum_{i=1}^4 a_i \epsilon^{-i} + a_0 \log \epsilon$ to make the action finite.

Substituting the Euclidean action I_E (6.8) or (3.7) into $-\log Z(B_d)$ in (6.9), we indeed obtain via holography

$$A = \begin{cases} a & \text{for } d = 4, \\ \frac{R}{8G_N} = \frac{c}{12} & \text{for } d = 2. \end{cases} \quad (6.11)$$

Hence, (6.11) reproduces the correct central charges in two and four dimensions.

For even $d = 2k$ ($k \geq 1$), we can write the action (6.6) more explicitly. As for the contribution from the bulk term, the indefinite integral we need to compute reads

$$\int \frac{r(z)^{2k}}{z^{2k+1}} dz = (-1)^k \log z + \sum_{r=0}^{k-1} \sum_{s=0}^{k-r} a_{r,s}^{(k)}(\rho_*/R) \left(\frac{r_B}{z}\right)^{2k-2r-s} \quad (6.12)$$

where we have introduced

$$a_{r,s}^{(k)}(\alpha) \equiv (-1)^r \binom{k}{r} \binom{k-r}{s} \frac{(2 \sinh \alpha)^s}{2r+s-2k}. \quad (6.13)$$

The result (6.12) has been obtained by expanding the polynomial $r(z)^2$ (see (6.5)). Since $0 \leq 2r+s \leq 2k-1$, all the terms in the double sum are power like divergent when $z \rightarrow 0$. As for the boundary term we have to consider (now $k \geq 0$)

$$\int \frac{r(z)^{2k}}{z^{2k+2}} dz = \frac{(-1)^{k+1}}{z} + \frac{1}{r_B} \sum_{r=0}^{k-1} \sum_{s=0}^{k-r} \tilde{a}_{r,s}^{(k)}(\rho_*/R) \left(\frac{r_B}{z}\right)^{2k+1-2r-s} \quad (6.14)$$

where the expansion coefficients read

$$\tilde{a}_{r,s}^{(k)}(\alpha) \equiv (-1)^r \binom{k}{r} \binom{k-r}{s} \frac{(2 \sinh \alpha)^s}{2r+s-2k-1}. \quad (6.15)$$

In the double sum occurring in (6.14) we have all the powers from z^{-2k-1} to z^{-2} .

By employing (6.12) and (6.14) into (6.6), we find that all the terms computed at $z = \epsilon$ are divergent and therefore must be subtracted. Thus $O(1)$ term in the ϵ expansion of I_E is given only by the contribution at the upper extremum and it reads

$$I_E = \frac{R^{2k-1}}{8\pi G_N} \frac{2\pi^k}{\Gamma(k)} \left[(-1)^k \log(r_B e^{\rho_*/R}) + \sum_{r=0}^{k-1} \sum_{s=0}^{k-r} a_{r,s}^{(k)}(\rho_*/R) e^{-(2k-2r-s)\rho_*/R} \right. \\ \left. - \sinh(\rho_*/R) \left((-1)^k e^{-\rho_*/R} + \sum_{r=0}^{k-2} \sum_{s=0}^{k-1-r} \tilde{a}_{r,s}^{(k-1)}(\rho_*/R) e^{-(2k-1-2r-s)\rho_*/R} \right) \right]. \quad (6.16)$$

Notice also that, since the upper extremum in (6.6) is proportional to r_B , the dependence on r_B cancels in all the power-like terms and enters only through the logarithmic term occurring in (6.12) computed at the upper extremum. This is consistent with the CFT dual as the violation of conformal symmetry only comes from the conformal anomaly.

6.1.2 AdS₄ case

In this case, by evaluating I_E and subtracting any divergences which are polynomials of ϵ , we finally obtain

$$I_E = \frac{R^2}{2G_N} \left[\frac{\pi}{2} + \arctan \left(\sinh \frac{\rho_*}{R} \right) - \frac{1}{24} \sinh \frac{3\rho_*}{R} + \left(\log \frac{\epsilon}{r_B} + \log \cosh \frac{\rho_*}{R} - \frac{33}{24} - \log 2 \right) \sinh \frac{\rho_*}{R} \right]. \quad (6.17)$$

Notice that there is a logarithmic term which should be interpreted as some conformal anomaly. Even though this theory is dual to three dimensional CFT, this appearance of the anomaly is expected because there is the two dimensional boundary which can lead to the conformal anomaly. Indeed the coefficient of the log term is proportional to $\sinh \frac{\rho_*}{R}$ and therefore it vanishes for the trivial boundary $\rho_* = 0$. We can define the effective boundary central charge c_{bdy} as

$$r_B \frac{\partial \log Z}{\partial r_B} = -\frac{1}{2\pi} \left\langle \int dx^2 \sqrt{g} T_\mu^\mu \right\rangle = \frac{c_{bdy}}{6} \chi(\Sigma), \quad (6.18)$$

where $\chi(\Sigma)$ is the Euler number of Σ , which is the boundary of M i.e. given by S^2 . In this way, we obtain

$$c_{bdy} = \frac{3R^2}{2G_N} \sinh \frac{\rho_*}{R}. \quad (6.19)$$

According to the discussion in the analysis in section 5.2, ρ_* decreases under RG flows. Therefore our boundary central charge c_{bdy} also decreases under RG flows.

As done in the section 6.1.1, we can consider the generic case with even number of dimensions, i.e. odd $d = 2k - 1$ ($k \geq 2$). In this case the analysis is more complicated because a square root remains in the integrands of the terms involved in the computation. In order to shorten the expressions, from (6.5) we find it convenient to introduce

$$r(z) \equiv r_B \tilde{r}(\tilde{z}) \quad \tilde{z} \equiv \frac{z}{r_B} \quad (6.20)$$

Then, the action (6.6) in our case becomes

$$I_E = \frac{R^{2(k-1)}}{8\pi G_N} \frac{2\pi^{(2k-1)/2}}{\Gamma((2d-1)/2)} \left[\int_{\tilde{\epsilon}}^{e^{\rho_*/R}} \frac{\tilde{r}(\tilde{z})^{2k}}{\tilde{z}^{2k} \tilde{r}(\tilde{z})} d\tilde{z} - \sinh \beta \int_{\tilde{\epsilon}}^{e^{\rho_*/R}} \frac{\tilde{r}(\tilde{z})^{2(k-1)}}{\tilde{z}^{2(k-1)+1} \tilde{r}(\tilde{z})} d\tilde{z} \right]. \quad (6.21)$$

We proceed as in the section 6.1.1 and for (6.20) we get

$$\tilde{r}(\tilde{z})^{2h} = \sum_{r=0}^h \sum_{s=0}^h b_{r,s}^{(h)}(\rho_*/R) \tilde{z}^{r+s} \quad b_{r,s}^{(h)}(\beta) \equiv (-1)^r \binom{h}{r} \binom{h}{s} e^{\beta(r-s)}. \quad (6.22)$$

This allows us to write the term contained between the square brackets in (6.21) as follows

$$\sum_{r=0}^k \sum_{s=0}^k b_{r,s}^{(k)}(\rho_*/R) \int_{\tilde{\epsilon}}^{e^{\rho_*/R}} \frac{d\tilde{z}}{\tilde{z}^{2k-r-s} \tilde{r}(\tilde{z})} - \sinh(\rho_*/R) \sum_{r=0}^{k-1} \sum_{s=0}^{k-1} b_{r,s}^{(k-1)}(\rho_*/R) \int_{\tilde{\epsilon}}^{e^{\rho_*/R}} \frac{d\tilde{z}}{\tilde{z}^{2k-1-r-s} \tilde{r}(\tilde{z})} \quad (6.23)$$

where we remark that, in the first double sum, for the power of z^{-1} we have $0 \leq 2k - r - s \leq 2k$, while in the second double sum, coming from the boundary term, we have $1 \leq 2k - 1 - r - s \leq 2k - 1$. As for the integrals occurring in (6.23) we have

$$\int_{\tilde{\epsilon}}^{e^{\rho_*/R}} \frac{d\tilde{z}}{\tilde{r}(\tilde{z})} = \frac{\pi}{2} + \arctan(\sinh(\rho_*/R)) + \dots \quad \int_{\tilde{\epsilon}}^{e^{\rho_*/R}} \frac{d\tilde{z}}{\tilde{z}\tilde{r}(\tilde{z})} = \log r_B - \log\left(\frac{\cosh(\rho_*/R)}{2}\right) + \dots \quad (6.24)$$

By computing other examples, we recognize the following structure for $q \geq 2$

$$\begin{aligned} \int_{\tilde{\epsilon}}^{e^{\rho_*/R}} \frac{d\tilde{z}}{\tilde{z}^q \tilde{r}(\tilde{z})} &= \left[C_q(\rho_*/R) \log\left(\frac{\tilde{z}}{1 + \tilde{z} \sinh(\rho_*/R) + \tilde{r}(\tilde{z})}\right) - P_{q-2}(\tilde{z}, \rho_*/R) \frac{\tilde{r}(\tilde{z})}{\tilde{z}^{q-1}} \right] \Big|_{\tilde{\epsilon}}^{e^{\rho_*/R}} \\ &= C_q(\rho_*/R) \left[\log r_B - \log\left(\frac{\cosh(\rho_*/R)}{2}\right) \right] + \gamma_q(\rho_*/R) + \dots \end{aligned} \quad (6.25)$$

where the dots denote vanishing or diverging terms when $\epsilon \rightarrow 0$. The polynomial $P_{q-2}(\tilde{z}, \beta)$ has degree $q-2$ in terms of \tilde{z} and its coefficients depend on ρ_*/R . We recall that $\tilde{r}(e^{\rho_*/R}) = 0$ and $\tilde{\epsilon} = \epsilon/r_B$. Unfortunately we are not able to get closed forms for the function $C_q(\rho_*/R)$ and $P_{q-2}(\tilde{z}, \rho_*/R)$ and, in order to obtain at least the dependence on r_B of the finite part of the action for any k we need to know $C_q(\rho_*/R)$ for any positive integer $q \geq 2$.

6.2 Holographic dual of half space

Consider a holographic dual of a half space in the pure AdS_{d+1}

$$ds^2 = R^2 \frac{dz^2 + \sum_{i=0}^{d-2} dx_i^2 + dx_{d-1}^2}{z^2}. \quad (6.26)$$

According to (2.17), the boundary Q is defined by

$$x = \sinh \frac{\rho_*}{R} \cdot z, \quad (6.27)$$

where ρ_* is related to the tension via $TR = (d-1) \tanh \frac{\rho_*}{R}$. The length in the Euclidean time and the one in x_{d-1} direction are defined to be β and L . The Euclidean action I_E is given by

$$\begin{aligned} I_E &= \frac{dR^{d-1}V_{d-2}\beta L}{8\pi G_N} \int_{\epsilon}^{\infty} \frac{dz}{z^{d+1}} + \frac{(d-1)R^{d-1}V_{d-2}\beta}{8\pi G_N} \sinh \frac{\rho_*}{R} \int_{\epsilon}^{\infty} \frac{dz}{z^d} \\ &= \frac{R^{d-1}V_{d-2}\beta L}{8\pi G_N \epsilon^d} + \frac{R^{d-1}V_{d-2}\beta}{8\pi G_N \epsilon^{d-1}} \sinh \frac{\rho_*}{R}. \end{aligned} \quad (6.28)$$

Both divergent terms are completely canceled by the counter terms. Therefore $I_E = 0$.

6.3 Holographic dual of strips

Consider a holographic dual of a strip in $\mathbb{R}^{1,d-1}$ which is defined by restricting to the range $0 < x_{d-1} < \Delta x$ with other coordinates x_0, x_1, \dots, x_{d-2} taking any values. We are interested in the zero temperature case and thus we consider the AdS soliton geometry

$$ds^2 = R^2 \frac{h(z)^{-1} dz^2 + \sum_{i=0}^{d-2} dx_i^2 + h(z) dx_{d-1}^2}{z^2}, \quad h(z) = 1 - \left(\frac{z}{z_0}\right)^d. \quad (6.29)$$

We compactify x_{d-1} as $x_{d-1} \sim x_{d-1} + \frac{4\pi}{d} z_0$. We describe the boundary Q by $x_{d-1} = x(z)$. The constraint (2.12) leads to the differential equation

$$x'(z) = \pm \frac{RT}{h(z)\sqrt{(d-1)^2 h(z) - R^2 T^2}}, \quad (6.30)$$

where + (or -) sign corresponds to the case where the bulk spacetime N is situated in the side of smaller (or larger) x . We fix the integration constant by requiring $x(z_*) = 0$, where z_* is defined by $(d-1)^2 h(z_*) = R^2 T^2$. Notice that the bulk spacetime N is given by

$$-x(z) \leq x_{d-1} \leq x(z), \quad (6.31)$$

for $T \leq 0$. For $T > 0$, N is given by the complement of (6.31) at $T = -|T|$ with respect to the total manifold of the AdS soliton i.e. $|x_{d-1}| \leq \frac{2\pi}{d} z_0$.

If we define $x_0 \equiv x(0)$, the length Δx of the AdS boundary M is given by

$$\begin{aligned} \Delta x &= 2x_0 && (\text{when } T \leq 0), \\ \Delta x &= \frac{4\pi}{d} z_0 - 2x_0 && (\text{when } T \geq 0). \end{aligned} \quad (6.32)$$

We can calculate x_0 as follows

$$\begin{aligned} x_0 &= \int_0^{z_*} \left(\frac{dx}{dz} \right) dz = \frac{RTz_0}{d(d-1)} \int_0^{w_*} dw \frac{w^{\frac{1-d}{d}}}{(1-w)\sqrt{1 - \frac{R^2 T^2}{(d-1)^2} - w}}, \\ &= \frac{\Gamma(1/d)\Gamma(1/2)}{\Gamma(1/2 + 1/d)} \cdot \frac{RTz_0}{d(d-1)} \cdot w_*^{1/d-1/2} \cdot F(1, 1/d, 1/2 + 1/d; w_*), \end{aligned} \quad (6.33)$$

where $w_* \equiv 1 - \frac{R^2 T^2}{(d-1)^2}$ and $F(a, b, c; z)$ is the standard hypergeometric function.

The entanglement entropy can be calculated as follows (we first assume $T < 0$):

$$\begin{aligned} I_E &= -\frac{1}{16\pi G_N} \int_N \sqrt{g}(R - 2\Lambda) - \frac{1}{8\pi G_N(d-1)} \int_Q \sqrt{h}T, \\ &= \frac{dR^{d-1}L^{d-2}\beta}{4\pi G_N} \int_\epsilon^{z_*} dz \frac{x(z)}{z^{d+1}} - \frac{TR^d L^{d-2}\beta}{4\pi G_N} \int_\epsilon^{z_*} \frac{dz}{z^d \sqrt{(d-1)^2 h(z) - R^2 T^2}}, \\ &= -\frac{R^{d-1}L^{d-2}\beta}{4\pi G_N} \left[\frac{h(z)x(z)}{z^d} \right]_\epsilon^{z_*}, \\ &= \frac{R^{d-1}L^{d-2}\beta}{4\pi G_N} x_0 \cdot \left(\frac{1}{\epsilon^d} - \frac{1}{z_0^d} \right) - \frac{R^{d-1}L^{d-2}\beta}{4\pi G_N \epsilon^{d-1}} \cdot \frac{RT}{\sqrt{(d-1)^2 - R^2 T^2}}. \end{aligned} \quad (6.34)$$

Here β and L is the length of x_0 and x_i directions.

We would like to cancel the divergences by adding local counter terms. The bulk counter term which is proportional to ϵ^{-d} leads to the subtraction (remember the factor $\sqrt{h(\epsilon)}$ in the x_{d-1} direction):

$$-\frac{R^{d-1}L^{d-2}\beta}{4\pi G_N} x_0 \cdot \frac{\sqrt{h(\epsilon)}}{\epsilon^d}. \quad (6.35)$$

Also the final term in (6.34) is simply canceled by the boundary counter term. In this way, by taking into account the finite contribution in (6.35) we obtain the final result

$$I_E = -\frac{R^{d-1}L^{d-2}\beta}{16\pi G_N} \cdot \frac{\Delta x}{z_0^d}. \tag{6.36}$$

It is possible to check that this result (6.36) remains the same even if $T > 0$.

7 Time-dependent configurations

So far we have focused on the holography for static systems where the time evolution of boundaries in a CFT is trivial. Thus here we would like to study time dependent configurations especially in the AdS₃/BCFT₂ setup. In most of the arguments below, higher dimensional generalizations are straightforward.

7.1 Description in terms of entangled pair

Consider a pure AdS₃

$$ds^2 = R^2 \frac{-dt^2 + dz^2 + dx^2}{z^2}, \tag{7.1}$$

and a boundary Q specified by $z = z(x, t)$. The solutions to the boundary equation of motion (2.12) are obtained from (3.6) by a double Wick rotation. The gravity dual is given by the region

$$(z - A)^2 + (x - \alpha)^2 - (t - \beta)^2 \geq \gamma^2, \tag{7.2}$$

where α, β, γ and A are arbitrary constants. This is depicted in the right figure in figure 7. The tension T is related to these parameters via $T = -\frac{A}{R|\gamma|}$. We also need to assume $|A| < |\gamma|$ so that the boundary of (7.2) is time-like. Using the translational invariance we simply set $\alpha = \beta = 0$ below. The AdS boundary of this spacetime (7.2) consists of two disconnected regions $x \geq \sqrt{\gamma^2 - A^2 + t^2}$ and $x \leq -\sqrt{\gamma^2 - A^2 + t^2}$, which are connected in the bulk of AdS (see the left figure in figure 7).

It is also possible to flip the sign of the inequality in (7.2). This leads to the region $(z - A)^2 + (x - \alpha)^2 - (t - \beta)^2 \leq \gamma^2$. In this case, the region looks like a half of three dimensional cylinder and this is dual to a CFT lives on an interval defined by $|x| \leq \sqrt{\gamma^2 - A^2 + t^2}$.

Let us consider the holographic interpretation of the solution (7.2). Since end point of each of boundaries is accelerated much like a Rindler observer, the light cone defined by $t = \pm x$ plays the role of Rindler horizon. Therefore the two regions at the AdS boundary are causally disconnected. However, they are connected in the bulk AdS. This means that they are entangled with each other as in the AdS Schwarzschild black holes [46]. This interpretation gets clearer if we perform the following familiar coordinate transformation:

$$\begin{aligned} t - x &= -e^{\theta-u} \sqrt{1 - \frac{r_+^2}{r^2}}, \\ t + x &= e^{\theta+u} \sqrt{1 - \frac{r_+^2}{r^2}}, \\ z &= \frac{r_+ e^\theta}{r}. \end{aligned} \tag{7.3}$$

This leads to the BTZ black hole metric

$$ds^2 = R^2 \left[- \left(\frac{r^2}{r_+^2} - 1 \right) du^2 + \frac{dr^2}{r^2 - r_+^2} + \frac{r^2}{r_+^2} d\theta^2 \right], \quad (7.4)$$

where the horizon is situated at $r = r_+$. In this coordinate system, (7.2) can be rewritten as

$$e^{2\theta} - 2 \frac{Ar_+ e^\theta}{r} \geq \gamma^2 - A^2, \quad (7.5)$$

which coincides with (4.12).² The region $|x| \geq \sqrt{\gamma^2 - A^2 + t^2}$, where the CFT is defined, is now mapped to the half line $\theta \geq \frac{1}{2} \log(\gamma^2 - A^2)$. The horizon $r = r_+$ is mapped to the Rindler horizon in (7.1).

The holographic entanglement entropy between the CFT on the two half lines (we choose the subsystem A as one of the half lines) are computed as follows³

$$S_A = \frac{R}{4G_N} \int_{\gamma+A}^{z_{IR}} \frac{dz}{z}, \quad (7.6)$$

where z_{IR} is the IR cut off, which is very large. We can easily see that this is equivalent to the entropy of the BTZ black hole (7.4) restricted inside of the boundary Q . Since the condition (7.5) is estimated as $e^\theta \geq A + \gamma$, we find the black hole entropy is given by

$$S_{BH} = \frac{R}{4G_N} \int_{\log(\gamma+A)}^{\log z_{IR}} d\theta. \quad (7.7)$$

Therefore S_A and S_{BH} are identical as we wanted to show.

These correspondences are naturally understood as an example of Unruh effect because a CFT on two half lines which are entangled due to a uniform acceleration, is equivalent to a CFT on a half line at finite temperature.

It is straightforward to generalize our setups so that it is dual to a CFT on the two disconnected intervals, instead of half lines. This is obtained by considering the gravity on the region

$$\gamma_1^2 \leq (z - A)^2 + x^2 - t^2 \leq \gamma_2^2, \quad (7.8)$$

assuming $|\gamma_1| < |\gamma_2|$. After the coordinate transformation (7.3), the region (7.8) is mapped to an interval in the BTZ black hole.

7.2 Comments on g-theorem and topological censorship

In the standard AdS/CFT, a Lorentzian asymptotic AdS spacetime with multiple AdS boundaries which are connected without encountering any horizons, is prohibited once we assume an appropriate energy condition [48]. This is called the topological censorship. It is

²See also appendix in which we also get a surface in the rotating BTZ BH using the coordinate transformation.

³Strictly speaking, we need to apply the covariant prescription of holographic entanglement entropy [47] in such a Lorentzian time-dependent spacetime. However, since here we restrict to the time symmetric surface at $t = 0$, we just need to calculate the geodesic length on that surface.




Figure 7. The left figure denotes a two dimensional CFT defined on a spacetime with time-dependent boundaries. The right figure describes its holographic dual.

interesting to study a similar property in our AdS/BCFT setups. We only consider static setups and assume that the AdS boundary consists of two disconnected intervals.

In the presence of two intervals, we can find a wormhole like geometry if we can connect them in the bulk AdS. However, this is not allowed once the null energy condition is imposed. As discussed in section 5.2, the g-theorem is holographically equivalent to $x''(z) \leq 0$. This requires the boundary Q eventually evolves toward the inside direction. Therefore it prevents the boundaries from connecting smoothly with each other. In this way, we find that the g-theorem is holographically interpreted as the topological censorship. Notice that in our previous example of section 7.1, the two disconnected regions at the AdS boundary can connect inside of the bulk AdS because the two regions are connected through a horizon.

8 Correlation functions

8.1 One point function

Considering the setup given by the metric (2.14), we employ the metric (2.18). The boundary Q is specified by

$$x = \tan \theta z, \quad (8.1)$$

where $\tan \theta = \sinh \frac{\rho^*}{R}$ (see (2.17)).

We are interested in a one point function $\langle O \rangle$ of a scalar operator O and the scalar field dual to O is denoted by ϕ . The bulk action of ϕ is given by

$$\begin{aligned} S &= \frac{1}{2} \int (dx)^{d+1} \sqrt{g} (\partial^\mu \phi \partial_\mu \phi + m^2 \phi^2) \\ &= \frac{1}{2} \int dz dx (dw)^{d-1} \left[\frac{R^{d-1}}{z^{d-1}} \left((\partial_z \phi)^2 + (\partial_x \phi)^2 + (\partial_\tau \phi)^2 + (\partial_{\vec{w}} \phi)^2 \right) + \frac{R^{d+1}}{z^{d+1}} m^2 \phi^2 \right]. \end{aligned} \quad (8.2)$$

where $(dw)^{d-1} = d\tau d\vec{w}$. On the boundary Q , we assume the following coupling of ϕ

$$S_Q = \int_Q (d\xi)^d \sqrt{h} (a\phi + \dots), \quad (8.3)$$

where the coordinates ξ parameterize the boundary Q and \dots means higher terms in ϕ which are not considered here; a is just a constant.

It is useful to define a rotated coordinate \tilde{x} and \tilde{z}

$$\tilde{x} = \cos \theta x - \sin \theta z, \quad \tilde{z} = \sin \theta x + \cos \theta z. \quad (8.4)$$

The boundary Q is described by $\tilde{x} = 0$. By considering the variation $\delta\phi$ of ϕ with the equation of motion imposed, we find

$$\delta S = R^{d-1} \int d\tilde{z} (dw)^{d-1} \delta\phi \left(\frac{\partial_{\tilde{x}}\phi}{z^{d-1}} + \frac{aR}{z^d} \right). \quad (8.5)$$

In this way, we find that the boundary condition on Q for the scalar field ϕ reads

$$(\cos \theta \partial_x - \sin \theta \partial_z)\phi + \frac{aR}{z} = 0, \quad (8.6)$$

where the operator acting on ϕ is $\partial_{\tilde{x}}$. Notice that to calculate the one point function, we can neglect the dependence on w .

Requiring the regularity at $z = \infty$, we can expand the solution to the equation of motion of (8.2) as follows

$$\phi(z, x) = z^{d/2} \int_{-\infty}^{\infty} dk A(k) K_{\nu}(|k|z) e^{ikx}, \quad (8.7)$$

where $\nu = \sqrt{d^2/4 + m^2 R^2}$. The function $A(k)$ can be found by plugging (8.7) into the boundary condition (8.6) which becomes

$$\begin{aligned} & \int_{-\infty}^{+\infty} A(k) \left[ik K_{\nu}(|k|z) \cos \theta - \left(\frac{d}{2z} K_{\nu}(|k|z) + \partial_z K_{\nu}(|k|z) \right) \sin \theta \right] e^{ikz \tan \theta} dk \\ &= \int_{-\infty}^{\infty} dk \left[ik A(k) \cos \theta K_{\nu}(|k|z) e^{ikz \tan \theta} \right. \\ & \quad \left. - \frac{\sin \theta}{z} \left(\frac{dA(k) e^{ikz \tan \theta}}{2} - \left(k A(k) e^{ikz \tan \theta} \right)' \right) K_{\nu}(|k|z) \right] \\ &= -\frac{aR}{z^{d/2+1}}, \end{aligned} \quad (8.8)$$

where in the last line, we performed a partial integration in terms of k using $\partial_z K_{\nu}(|k|z) = k \partial_k K_{\nu}(|k|z)/z$. In the appendix D we give a detailed treatment of this equation.

By taking the AdS boundary limit $z \rightarrow 0$ of ϕ , we get the well known behavior

$$\phi \rightarrow z^{d-\Delta} \alpha(x) + z^{\Delta} \beta(x), \quad (8.9)$$

where $\Delta = d/2 + \nu$ is the conformal dimension of the dual operator. According to the standard dictionary in AdS/CFT [49], we find the one point function as

$$\langle O \rangle = (2\Delta - d)\beta(x). \quad (8.10)$$

Note that we have imposed two boundary conditions to determine the behavior of the solution (8.7). The boundary condition of the regularity at $z = \infty$ is included in the

boundary condition (8.6) and the boundary condition at the AdS boundary limit $z \rightarrow \infty$ is also imposed on ϕ as seen in (8.9). Thus, the boundary condition above is enough to determine the solution of ϕ . The dependence on k of $A(k)$ turns out to be⁴

$$A(k) \equiv c_\theta \frac{|k|^{d/2}}{k} \quad (8.11)$$

and in the following we will find c_θ . In the simplest case of $\theta = 0$, from (8.8) we have

$$c_0 = \frac{iaR}{2^{d/2} \Gamma\left(\frac{d-\Delta+1}{2}\right) \Gamma\left(\frac{\Delta+1}{2}\right)} \quad \text{for } d/2 + 1 > \nu \quad (8.12)$$

(see the appendix D for details). Thus we find

$$\begin{aligned} \beta(x) &= \frac{\langle O \rangle}{2\Delta - d} = c_0 \int_{-\infty}^{\infty} dk \frac{-\pi k^{-1} |k|^\Delta}{2 \sin \pi \left(\Delta - \frac{d}{2}\right) \Gamma\left(\Delta - \frac{d}{2} + 1\right) 2^{\Delta-d/2}} e^{ikx} \\ &= \frac{\pi aR \sin\left(\frac{\pi\Delta}{2}\right) \Gamma(\Delta) x^{-\Delta}}{2^\Delta \sin \pi \left(\Delta - \frac{d}{2}\right) \Gamma\left(\Delta - \frac{d}{2} + 1\right) \Gamma\left(\frac{d-\Delta+1}{2}\right) \Gamma\left(\frac{\Delta+1}{2}\right)}. \end{aligned} \quad (8.13)$$

In this way we get $\langle O \rangle \sim 1/x^\Delta$ as expected. Our result can be extended for $\theta \neq 0$. Assuming (8.11), the boundary condition (8.8) is solved by

$$c_\theta = \frac{iaR}{2^{d/2} \Gamma\left(\frac{\Delta+1}{2}\right) \Gamma\left(\frac{d-\Delta+1}{2}\right) \sqrt{1+v^2} F\left(\frac{\Delta+1}{2}, \frac{d-\Delta+1}{2}, \frac{1}{2}; -v^2\right)} \quad (8.14)$$

(again, see the appendix D for details) where $v = \sinh(\rho_*/R)$ and therefore we get

$$\begin{aligned} \langle O \rangle &\equiv \frac{aR\gamma}{x^\Delta} = \frac{aR(2\Delta - d)\pi \sin\left(\frac{\pi\Delta}{2}\right) \Gamma(\Delta)}{2^\Delta \sin \pi \left(\Delta - \frac{d}{2}\right) \Gamma\left(\Delta - \frac{d}{2} + 1\right) \Gamma\left(\frac{d-\Delta+1}{2}\right) \Gamma\left(\frac{\Delta+1}{2}\right)} \\ &\quad \cdot \frac{x^{-\Delta}}{\sqrt{1+v^2} F\left(\frac{\Delta+1}{2}, \frac{d-\Delta+1}{2}, \frac{1}{2}; -v^2\right)}. \end{aligned} \quad (8.15)$$

Thus, we find that the scaling of the one-point function is independent of θ . We can also see from above that the coefficient of the one-point function should be real by considering the odd/even property of the Fourier integral (8.8). We plotted the coefficient of the one-point function γ as the function of v in figure 8.

For $d = 4$, it is interesting to compare the one-point function for $\theta = 0$ (8.10) with the one-point function obtained via the AdS/dCFT correspondence [23]. Substituting $aR = \lambda^{1/2}$ into (8.13) and using some formulas for the Gamma function, we can confirm that our one-point function agrees completely with the one-point function in [23] of the following form:

$$\langle O \rangle = \lambda^{1/2} \frac{\Gamma\left(\frac{\Delta-3}{2}\right) \Gamma\left(\frac{\Delta}{2}\right) \Gamma\left(\frac{3}{2}\right)}{\pi \Gamma(\Delta - 2)} x^{-\Delta}. \quad (8.16)$$

⁴If we choose the ansatz for the function $A(k) \sim |k|^{d/2-1}$, the left hand side of the boundary condition (8.8) vanishes.




Figure 8. Setting $R = 1$, $v(= \sinh \rho_*/R)$ dependence of γ for $d = 4$ is plotted, where γ is the coefficient of the one-point function. $|\gamma|$ for the irrelevant operator ($\Delta > 4$) seems to be small for any v .

As pointed out in [23], the power of λ in (8.16) at strong coupling does not need to agree with field theory results at weak coupling. For general BCFT with the gravity dual, this power at strong coupling seems to be related with the parameter aR .

9 An example of string theory embedding

As a final study in this paper, we would like to present a concrete setup in string theory which can be regarded as an example of AdS/BCFT. In our model described below, orientifolds with D-branes play the role of the boundary Q . In a geometry which looks like $AdS \times M$ (M is a compact manifold), we would like to wrap the orientifolds and branes completely on M so that the backreaction inside M gets trivial and that only the AdS part becomes relevant as assumed in our effective descriptions of AdS/BCFT.

We would like to consider $AdS_4 \times CP^3$ solution in type IIA string theory dual to $N = 6$ supersymmetric Chern-Simons theory (ABJM theory) [50]. We write the metric of AdS_4 as follows

$$ds^2 = R^2 \frac{dz^2 - dt^2 + dx^2 + dy^2}{z^2}. \tag{9.1}$$

The ABJM theory lives on (t, x, y) and consists of the $U(N)_k \times U(N)_{-k}$ gauge field, denoted by $(A_\mu^{(1)}, A_\mu^{(2)})$ and four chiral multiplets (its bi-fundamental scalars and fermions are denoted by X^I and Ψ^I ($I = 1, 2, 3, 4$)). We introduce the two orientifold 8-planes ($O8$ -planes) so that they are situated at $y = 0$ and $y = L$, extending in the nine directions other than y . Notice that all of such configurations preserve the $SU(4)$ R-symmetry. There are four different types of $O8$ -planes denoted by $O8^-$, $\overline{O8}^-$, $O8^+$ and $\overline{O8}^+$ (for such a classification

of orientifolds see e.g. [51–53]). Their differences are clear in the boundary state formalism. If we write the cross-cap state in the NSNS and RR sector as $|\Omega_{NSNS}\rangle$ and $|\Omega_{RR}\rangle$, then we can represent

$$\begin{aligned} |O8^-\rangle &= -|\Omega_{NSNS}\rangle - |\Omega_{RR}\rangle, & |\overline{O8^-}\rangle &= -|\Omega_{NSNS}\rangle + |\Omega_{RR}\rangle, \\ |O8^+\rangle &= |\Omega_{NSNS}\rangle + |\Omega_{RR}\rangle, & |\overline{O8^+}\rangle &= |\Omega_{NSNS}\rangle - |\Omega_{RR}\rangle. \end{aligned} \tag{9.2}$$

In string theory setup with orientifolds, we need to impose the tadpole cancellation, which requires that the total RR charge vanishes. For example, the RR-charge of an $O8^\mp$ -plane is given by ∓ 8 times that of a D8-brane.

The basic setup is the $O8^- - O8^-$ system, i.e. a $O8^-$ is at $y = 0$ and another $O8^-$ is at $y = L$ (refer to figure 9). We need to add totally 16 D8-branes. If we place a half of them at $y = 0$ and the other half at $y = L$, then the RR charge source completely cancels out and the solution will coincide with the one in AdS/BCFT with the vanishing tension $T = 0$ as the NS tadpole is also completely canceled out. In this case of two $O8^-$ planes, the Z_2 orientifold projection acts as a parity transformation which exchanges the two gauge groups

$$\begin{aligned} A_{t,x}^{(1,2)}(t, x, y) &\rightarrow -\left(A_{t,x}^{(2,1)}(t, x, -y)\right)^T, \\ A_y^{(1,2)}(t, x, y) &\rightarrow \left(A_y^{(2,1)}(t, x, -y)\right)^T, \\ X^I(t, x, y) &\rightarrow X^I(t, x, -y)^T, \\ \Psi^I(t, x, y) &\rightarrow P \cdot \Psi^I(t, x, -y)^T, \end{aligned} \tag{9.3}$$

where M^T denotes the matrix transposition of M ; the matrix P is the standard one which describes the parity transformation of three dimensional fermions. For more details refer to the appendix E. Notice that the $SU(4)_R$ symmetry which rotates the index I is preserved after this projection.

The Z_2 projection (9.3) is also clear from the T-dualized setup of N D3-branes with a NS5-brane and $(1, k)$ 5-brane considered in [50]. The D3-branes are wrapped on a circle and they are divided into two parts (called fractional D3-branes), cut by the two 5-branes. We insert two $O7$ -planes so that they coincide with two 5-branes. The orientifold action now exchanges two different fractional D3-branes and therefore it also exchanges the two $U(N)$ gauge group. This setup preserves a half of the original supersymmetries.

The presence of 8 D8-branes at each of the boundaries $y = 0$ and $y = L$ introduces 8 massless chiral (complex) fermions at each of them. This comes from the D2-D8 open strings with the orientifold projection. In the ABJM setup without the orientifold, an appearance of Weyl fermions have been studied in [54, 55] as an example of holographic edge states of quantum Hall effect. In summary, this three dimensional gauge theory dual to the AdS_4 with two $O8^-$ planes is defined by the ABJM theory projected by the action (9.3) at two boundaries $y = 0$ and $y = L$ with 8 chiral fermions at each of the boundaries. This supersymmetric Chern-Simons theory lives on $R^{1,2} \times S^1/Z_2$ and note that S^1/Z_2 describes an interval. The holography tells us that this theory preserves the boundary conformal

symmetry at the two boundaries and therefore it offers us an example of AdS/BCFT. The tension of the boundaries is vanishing as the local tadpole cancelation between O8-planes and D8-branes.

Before we go on, it is instructive to why these edge modes need to appear in this model. To answer this, we would like to look at the gauge anomaly when we reduce the theory on the interval and consider its two dimensional field theory in the low energy limit. It is useful to note that the matrix P can be regarded as the chirality matrix when it is reduced to two dimension. Thus the condition (9.3) tells us that the right-moving mode of the fermion should be symmetric, while the left-moving one should be antisymmetric. In this way, the low energy two dimensional theory is not a non-chiral theory and we have to worry about the gauge anomaly. Now we want to estimate the anomaly of $U(N)$ gauge symmetry which is obtained from the projection of $U(N) \times U(N)$ gauge symmetry by (9.3). First consider the $SU(N)$ part. The gauge anomaly induced matter in the representation R in two dimension is proportional to $\text{Tr}_R[T^a T^b]_{\text{right}} - \text{Tr}_R[T^a T^b]_{\text{left}}$, where the difference means that between the left-moving sector contribution and right-moving one. The relation of such traces between antisymmetric (A), symmetric (S) and fundamental representation (F) is given by

$$\text{Tr}_A[T^a T^b] = (N - 2)\text{Tr}_F[T^a T^b], \quad \text{Tr}_S[T^a T^b] = (N + 2)\text{Tr}_F[T^a T^b]. \quad (9.4)$$

Since there are four Dirac fermions Ψ^I which satisfy the projection (9.3), the $SU(N)$ gauge anomaly is given by

$$4\text{Tr}_S[T^a T^b] - 4\text{Tr}_A[T^a T^b] = 16\text{Tr}_F[T^a T^b]. \quad (9.5)$$

This anomaly is completely canceled if we add extra 16 chiral left-moving fermions with the fundamental representation, which indeed coincide with the edge modes from the D8-branes. For the anomaly for the $U(1)$ part of $U(N)$, we can similarly confirm the cancelation.

Moreover, we can move some of the D8-branes from one of $O8^-$ planes to another with the supersymmetries kept. This offers us an example where the boundary Q has non-vanishing tension. However, we need to be careful to identify its bulk geometry because the boundary also gives rise to the dilaton gradient. It will be an intriguing future problem to work out its back-reacted solution in type IIA supergravity.

So far we studied the case where both of the O8-planes are the $O8^-$ type. We would also like to briefly discuss some other cases (refer to figure 9). If one of them is $O8^-$ and the other is its anti-plane i.e. $\overline{O8^-}$, then the supersymmetries are completely broken and we do not need any D8-branes for the RR tadpole cancelation. In this case, no chiral fermion appears and there is no anomaly. This is also consistent with the fact that the fermion boundary condition is twisted and no zero mode remains. Notice that the Z_2 projection for $\overline{O8^-}$ is given by (9.3) with the sign in front of fermion transformation flipped. Since the tension of these orientifolds is negative, it is bent toward internal direction and eventually should smoothly connect with each other. This can be regarded as a boundary version of the AdS soliton.




Figure 9. Schematic Pictures of $AdS_4/BCFT_3$ setups with various $O8$ -planes. The vertical line in the left side in each of figures represents the AdS boundary.

On the other hand, if we consider a system with a $O8^-$ and a $O8^+$, then a half of the original supersymmetries are preserved. In this case, since both NSNS and RR tadpole are canceled, we do not need any D8-branes. The Z_2 projection for $O8^+$ is given by replacing the SO projection in (9.3) with the Sp projection. In this case, we can indeed confirm that the remained fermion zero modes are non-chiral after we impose the two boundary conditions. Finally if we consider two $O8^+$ -planes, then we need to insert 8 anti D-branes at each boundaries. This again leads to eight chiral (complex) fermions at each of them.

Finally we would like to notice that if we take the strong coupling limit, we will find the Horava-Witten model [56, 57], where the boundary Q is now introduced in the $AdS_4 \times S^7/Z_k$ background of M-theory.

10 Conclusions and discussions

In this paper we studied various aspects of the holographic dual of BCFT ($AdS/BCFT$). We explicitly constructed holographic duals of conformal field theories on balls.

In $AdS_5/BCFT_4$, we confirmed that the coefficient of the logarithmic term in the partition function agrees with that computed from the conformal anomaly in the field theory. In $AdS_4/BCFT_3$, we noticed there is again a logarithmic term and we found that this is associated with the boundary degrees of freedom as in the central charge in CFT_2 . Thus we called this a boundary central charge. It will be an intriguing future direction to study this quantity from a field theory side. In $AdS_3/BCFT_2$, the partition function is interpreted as the g -function.

Next we showed a holographic g -theorem in any dimension, which argued that the holographically defined g -function decreases under RG flows if we assume the null energy condition. As a special example of this, we found that the boundary central charge in $BCFT_3$ decreases under RG flows. We also pointed out this g -theorem is closely related to the topological censorship which prohibits the existence of AdS wormholes.

If we heat up such systems, it is natural to expect a Hawking-Page type phase transition. We explicitly show this transition in the $AdS_3/BCFT_2$ case. We find that as the boundary entropy gets larger, the transition temperature gets lower. In higher dimen-

sions, though we found a solution in the low temperature phase, we did not in the high temperature phase. This leaves an interesting future problem.

Moreover, we presented a time-dependent example of AdS/BCFT. In this example, the two disconnected regions at the AdS boundary are accelerated so that they cannot communicate with each other. These two regions are connected in the bulk AdS. We confirmed that the entanglement entropy between them is equal to the black hole entropy after a coordinate transformation which makes the background static.

We also computed holographic one-point functions and confirmed that their scaling property is the same as what we expect from field theory calculations. Finally, we gave an example of string theory embedding of this holography, based on the type IIA AdS₄×CP³ geometry with O8-planes. This is dual to a ABJM theory on an interval. The gauge anomaly cancelation requires massless chiral fermions localized at the two dimensional boundary, being consistent with the D-brane analysis. In this example, even though we can change the value of the tension T , it is possible that the back reactions of the D8-branes and O8-planes break the conformal invariance. This issue deserves future study. At the same time, it is an important future problem to find some other examples in string theory and realize AdS/BCFT with a non-vanishing tension T .

Acknowledgments

The authors are grateful to L. Gomi, M. Headrick, Y. Hikida, A. Karch, J. Maldacena, J. McGreevy, R. Myers, T. Nishioka, N. Ogawa, S. Ryu, T. Ugajin and E. Witten for valuable comments. MF and TT are supported by World Premier International Research Center Initiative (WPI Initiative) from the Japan Ministry of Education, Culture, Sports, Science and Technology (MEXT). MF is supported by the postdoctoral fellowship program of the Japan Society for the Promotion of Science (JSPS), and partly by JSPS Grant-in-Aid for JSPS Fellows No.22·1028. TT thanks very much the Strings 2011 conference in Uppsala, where a part of this work has been announced and conducted. TT and ET are grateful to the Aspen center for physics and the Aspen workshop “Quantum Information in Quantum Gravity and Condensed Matter Physics,” where a part of this work was conducted. TT is partly supported by JSPS Grant-in-Aid for Scientific Research No.20740132 and by JSPS Grant-in-Aid for Creative Scientific Research No.19GS0219. ET is supported by Istituto Nazionale di Fisica Nucleare (INFN) through the “Bruno Rossi” fellowship and by funds of the U.S. Department of Energy under the cooperative research agreement DE-FG02-05ER41360.

A Calculations of extrinsic curvatures

Here we would like to explain the calculations of extrinsic curvatures in $d + 1$ dimensional spacetime. If we define $h_{\mu\nu}$ ($\mu, \nu = 0, 1, 2, \dots, d$) to be the induced metric of d -dimensional submanifold Q and n^μ to be a space-like unit normal vector on Q (toward the outside direction), then we have the relation

$$h_{\mu\nu} = g_{\mu\nu} - n_\mu n_\nu. \tag{A.1}$$




Figure 10. A connected surface in the bulk associated to the annulus.

The extrinsic curvature $K_{\mu\nu}$ is defined by

$$K_{\mu\nu} = h_{\mu}^{\rho} h_{\nu}^{\lambda} \nabla_{\rho} n_{\lambda}, \quad (\text{A.2})$$

where $\nabla_{\rho} n_{\lambda}$ is the standard covariant derivative in the $d + 1$ dimensional spacetime. Its trace is given by

$$K = g^{\mu\nu} K_{\mu\nu} = h^{\mu\nu} K_{\mu\nu}. \quad (\text{A.3})$$

These expressions using the $d + 1$ dimensional coordinate are degenerate because $K_{\mu\nu} n^{\nu} = h_{\mu\nu} n^{\nu} = 0$. In this convention, the boundary equation of motion (2.12) is equivalent to

$$K_{\mu\nu} = (K - T) h_{\mu\nu}. \quad (\text{A.4})$$

After projecting to the d dimensional coordinate, we obtain K_{ab} and h_{ab} ($a, b = 0, 1, 2, \dots, d - 1$).

B Possibility of connected boundary for the annulus

In this appendix we consider the boundary theory defined on an annulus and the connected surface Q extending in the bulk shown in figure 10 as a possible solution of (2.12) in Euclidean AdS_3 (3.3).

We find it convenient to parameterize this surface by two angular coordinates (θ, ϕ) as follows

$$z = r_1(\sin \phi + \sin \delta) \quad \tau = (r_0 + r_1 \cos \phi) \cos \theta \quad x = (r_0 + r_1 \cos \phi) \sin \theta \quad (\text{B.1})$$

where $0 \leq \theta \leq 2\pi$ and $-\delta \leq \phi \leq \pi + \delta$ while $\delta \in (-\pi/2, \pi/2)$ is a parameter. The circumference in the middle of the boundary annulus has radius r_0 while $r_1 < r_0$ is the radius of the circumference arch shown in figure 10. The parameter δ tells us whether the center of this circumference arch has $z > 0$ ($\delta > 0$) or $z < 0$ ($\delta < 0$). The surface Q is generated by rotating this arch around a vertical axis defined by $\tau = 0$ and $x = 0$. The thickness of the boundary annulus is $2r_1 \cos \delta$.

The surface Q is equivalently described by the following constraint

$$\left(\sqrt{\tau^2 + x^2} - r_0 \right)^2 + (z - r_1 \sin \delta)^2 = r_1^2. \quad (\text{B.2})$$

The induced metric on Q reads

$$h_{ab} = \frac{R^2}{z^2} \begin{pmatrix} (r_0 + r_1 \cos \phi)^2 & 0 \\ 0 & r_1^2 \end{pmatrix} \quad (\text{B.3})$$

while from the expression of K_{ab} for Q we find that

$$K_{ab} - K h_{ab} = \frac{R}{z^2} \begin{pmatrix} -(r_0 + r_1 \cos \phi)^2 \sin \delta & 0 \\ 0 & \frac{r_0 \sin \phi - r_1 \sin \delta \cos \phi}{r_0 + r_1 \cos \phi} r_1^2 \end{pmatrix}. \quad (\text{B.4})$$

It is clear that we cannot solve both the two equations given by $K_{ab} - K h_{ab} = -T h_{ab}$ through the parameter δ only. Indeed, the first equation provides the tension $T = \sin \delta / R$ but then the other one leads to $r_0 = 0$, which makes the annulus collapse to a circle and Q becomes the surface shown in figure 3.

C Boundaries in rotating BTZ black holes

In this appendix we find a surface in the rotating BTZ satisfying the boundary equation (A.4) by using the fact that it is locally equivalent to AdS_3 . The metric of the rotating BTZ is given by

$$ds^2 = \frac{r^2}{(r^2 - r_+^2)(r^2 - r_-^2)} dr^2 + r^2 dx^2 + \left(-\frac{(r^2 - r_+^2)(r^2 - r_-^2)}{r^2} + \frac{r_+^2 r_-^2}{r^2} \right) dt^2 + 2r_+ r_- dt dx, \quad (\text{C.1})$$

where we used the convention of the unit AdS radius $R = 1$.

This metric is obtained by a coordinate transformation of the pure AdS_3 metric $ds^2 = (dw_+ dw_- + dz^2) / z^2$ as follows:

$$w_+ = \sqrt{\frac{r^2 - r_+^2}{r^2 - r_-^2}} \exp((x+t)(r_+ + r_-)) \equiv T + V, \quad (\text{C.2})$$

$$w_- = \sqrt{\frac{r^2 - r_+^2}{r^2 - r_-^2}} \exp((x-t)(r_+ - r_-)) \equiv -T + V, \quad (\text{C.3})$$

$$z = \sqrt{\frac{r_+^2 - r_-^2}{r^2 - r_-^2}} \exp(xr_+ + tr_-). \quad (\text{C.4})$$

This transformation changes the following sphere to a surface:

$$w_+ w_- + (z - r_B \sinh(\rho_*))^2 - r_B^2 \cosh(\rho_*)^2 = 0. \quad (\text{C.5})$$

Remind that R is set to be 1 and this surface is a solution of the gravity dual of BCFT. Employing the coordinate transformation above, the surface in the rotating BTZ black hole is given by

$$0 = F(t, x, r) \equiv x + \frac{1}{r_+} \left[\rho_* + t r_- - \log\left(\frac{r_B}{2}\right) - \log(\tilde{f}(r)) \right] \quad (\text{C.6})$$

where

$$\tilde{f}(r) \equiv (e^{2\rho_*} - 1) \sqrt{\frac{r_+^2 - r_-^2}{r^2 - r_-^2}} + \sqrt{\frac{e^{4\rho_*}(r_+^2 - r_-^2) + 2e^{2\rho_*}(2r^2 - r_+^2 - r_-^2) + r_+^2 - r_-^2}{r^2 - r_-^2}}. \quad (\text{C.7})$$

Remind that for the case $r_- = 0$, the rotating BTZ black hole becomes the BTZ black hole. Setting $r_- = 0$, indeed, the surface (C.6) becomes the surface of the BTZ BH found in (4.12):

$$x + \frac{1}{r_+} \left[\rho_* + -\log\left(\frac{r_B}{2}\right) - \log\left((e^{2\rho_*} - 1) \frac{r_+}{r} + \frac{\sqrt{4e^{2\rho_*}r^2 + (e^{2\rho_*} - 1)^2 r_+^2}}{r} \right) \right] = 0. \quad (\text{C.8})$$

Taking the derivative and using the coordinate transformation $r \rightarrow 1/z$,

$$\frac{dx}{dz} = \frac{T}{\sqrt{1 - T^2(1 - r_+^2 z^2)}}. \quad (\text{C.9})$$

In addition, the surface (C.6) for the rotating BTZ (C.1) is also a solution of the boundary equation.

To check this, first, the surface normal reads

$$\hat{n}_\mu = \partial_\mu F(t, x, r), \quad n_\mu = \frac{\hat{n}_\mu}{\sqrt{\hat{n}_\nu \hat{n}^\nu}} \quad (\text{C.10})$$

while the induced metric is defined in (A.1). The second Christoffel symbol for the rotating BTZ black hole is obtained as

$$\begin{aligned} \Gamma_{13}^3 &= \frac{r^3}{(r^2 - r_+^2)(r^2 - r_-^2)}, & \Gamma_{12}^3 &= \frac{r_+ r_- r}{(r^2 - r_+^2)(r^2 - r_-^2)}, \\ \Gamma_{13}^2 &= -\frac{r_+ r_- r}{(r^2 - r_+^2)(r^2 - r_-^2)}, & & \end{aligned} \quad (\text{C.11})$$

$$\Gamma_{12}^2 = \frac{(r^2 - r_-^2 - r_+^2) r}{(r^2 - r_+^2)(r^2 - r_-^2)}, \quad \Gamma_{33}^1 = \frac{r^4 - r^2 r_-^2 - r_+^2 r^2 + r_+^2 r_-^2}{r}, \quad (\text{C.12})$$

$$\Gamma_{22}^1 = \frac{r^4 - r^2 r_-^2 - r_+^2 r^2 + r_+^2 r_-^2}{r}, \quad \Gamma_{11}^1 = -\frac{r^4 - r_+^2 r_-^2}{r(r^2 - r_+^2)(r^2 - r_-^2)}. \quad (\text{C.13})$$

The extrinsic curvature are obtained by using (A.2) and (A.3). After a short computation, the surface (C.6) is found to be the solution as follows:

$$K = 2T, \quad K_{\mu\nu} - \frac{K}{2} h_{\mu\nu} = 0. \quad (\text{C.14})$$

D Details about the one point function

In this appendix we give some technical details on the derivation of the results of the section 8.

Given (8.7), it is useful to recall that the modified Bessel function of the second kind $K_\nu(z)$ is related to the modified Bessel function of the first kind $I_\nu(z)$ as follows

$$K_\nu(z) = \frac{\pi I_{-\nu}(z) - I_\nu(z)}{2 \sin \nu \pi} \quad I_\nu(z) = \left(\frac{z}{2}\right)^\nu \sum_{n=0}^{\infty} \frac{(z/2)^{2n}}{n! \Gamma(\nu + n + 1)}. \quad (\text{D.1})$$

where we have also provided the expansion of $I_\nu(z)$. Let us consider the boundary condition (8.8)

$$\int_{-\infty}^{+\infty} A(k) \left[ik K_\nu(|k|z) \cos \theta - \left(\frac{d}{2z} K_\nu(|k|z) + \partial_z K_\nu(|k|z) \right) \sin \theta \right] e^{ikz \tan \theta} dk = -\frac{aR}{z^{d/2+1}}. \quad (\text{D.2})$$

By using that

$$\partial_z K_\nu(z) = -\frac{K_{\nu-1}(z) + K_{\nu+1}(z)}{2} \quad (\text{D.3})$$

and (8.11), the boundary condition (D.2) reads (we change the integration variable $\tilde{k} \equiv kz$ here)

$$\begin{aligned} \frac{c_\theta}{2} \int_{-\infty}^{+\infty} |\tilde{k}|^{d/2} \left\{ 2K_\nu(|\tilde{k}|) \cos \theta \right. \\ \left. + \left[\frac{d}{\tilde{k}} K_\nu(|\tilde{k}|) - \frac{|\tilde{k}|}{\tilde{k}} \left(K_{\nu-1}(|\tilde{k}|) + K_{\nu+1}(|\tilde{k}|) \right) \right] i \sin \theta \right\} e^{i\tilde{k} \tan \theta} d\tilde{k} = iaR. \end{aligned} \quad (\text{D.4})$$

By exploiting the parity of the integrand, we can write (D.4) as follows

$$\begin{aligned} c_\theta \int_0^{+\infty} \tilde{k}^{d/2} \left\{ 2K_\nu(\tilde{k}) \cos \theta \left(e^{i\tilde{k} \tan \theta} + e^{-i\tilde{k} \tan \theta} \right) \right. \\ \left. + i \sin \theta \left[\frac{d}{\tilde{k}} K_\nu(\tilde{k}) - K_{\nu-1}(\tilde{k}) - K_{\nu+1}(\tilde{k}) \right] \left(e^{i\tilde{k} \tan \theta} - e^{-i\tilde{k} \tan \theta} \right) \right\} d\tilde{k} = 2i aR \end{aligned} \quad (\text{D.5})$$

where we can recognize $\cos(\tilde{k} \tan \theta)$ and $\sin(\tilde{k} \tan \theta)$ in the expressions between the round brackets. The integral in the l.h.s. of (D.5) clearly provides a real number, thus c_θ is purely imaginary.

Let us consider first the simplest case of $\theta = 0$ i.e. $x = 0$. The condition (D.5) then becomes

$$2c_0 \int_0^{+\infty} k^{d/2} K_\nu(k) dk = iaR. \quad (\text{D.6})$$

By using the following integral in (D.6)

$$\int_0^\infty k^{\mu-1} K_\nu(k) dk = 2^{\mu-2} \Gamma\left(\frac{\mu+\nu}{2}\right) \Gamma\left(\frac{\mu-\nu}{2}\right) \quad \begin{cases} \text{Re}(\mu) > \text{Re}(\nu) \\ \text{Re}(\mu+\nu) > 0 \end{cases} \quad (\text{D.7})$$

and also (8.11), we get 8.12. In the general case of $\theta \geq 0$, by employing the following integrals

$$\int_0^\infty k^{\mu-1} K_\nu(k) \cos(bk) dk = 2^{\mu-2} \Gamma\left(\frac{\mu+\nu}{2}\right) \Gamma\left(\frac{\mu-\nu}{2}\right) F\left(\frac{\mu+\nu}{2}, \frac{\mu-\nu}{2}; \frac{1}{2}; -b^2\right) \quad (\text{D.8})$$

$$\begin{aligned} \int_0^\infty k^{\mu-1} K_\nu(k) \sin(bk) dk = \\ = 2^{\mu-1} b \Gamma\left(\frac{\mu+\nu+1}{2}\right) \Gamma\left(\frac{\mu-\nu+1}{2}\right) F\left(\frac{\mu+\nu+1}{2}, \frac{\mu-\nu+1}{2}; \frac{3}{2}; -b^2\right) \end{aligned} \quad (\text{D.9})$$

the boundary condition (D.5) becomes

$$\frac{2^{d/2}}{\cos \theta} \Gamma(\alpha_+) \Gamma(\alpha_-) \left\{ \cos^2 \theta F(\alpha_+, \alpha_-; 1/2; -v^2) - \sin^2 \theta \left[(2\alpha_- - 1) F(\alpha_+, \alpha_-; 3/2; -v^2) - 2\alpha_- F(\alpha_+, \alpha_- + 1; 3/2; -v^2) \right] \right\} = \frac{i a R}{c_\theta} \quad (\text{D.10})$$

where

$$\alpha_+ \equiv \frac{d/2 + \nu + 1}{2} \quad \alpha_- \equiv \frac{d/2 - \nu + 1}{2} \quad v \equiv \tan \theta. \quad (\text{D.11})$$

By employing the following identity of the hypergeometric function

$$(c-1)F(a, b; c-1; z) - bF(a, b+1; c; z) + (b-c+1)F(a, b; c; z) = 0 \quad (\text{D.12})$$

the boundary condition (D.10) simplifies to

$$2^{d/2} \Gamma(\alpha_+) \Gamma(\alpha_-) \sqrt{1+v^2} F(\alpha_+, \alpha_-; 1/2; -v^2) = \frac{i a R}{c_\theta}. \quad (\text{D.13})$$

From this equation we obtain c_θ , given in (8.14). For $\theta = 0$ we recover (8.12) as special case.

E Discrete symmetries of ABJM theory

Here we summarize the discrete symmetries of the ABJM theory. The two $U(N)$ gauge fields are denoted by $A_\mu^{(1)}$ and $A_\mu^{(2)}$. The matter fields consists of bi-fundamental four complex scalars X^I and bi-fundamental four Dirac fermions Ψ^I ($I = 1, 2, 3, 4$). See also [50, 58] for earlier discussions on the discrete symmetries. The charge conjugation symmetry C is given by

$$\begin{aligned} A_\mu^{(1,2)}(t, x, y) &\rightarrow -A_\mu^{(1,2)}(t, x, y)^*, \\ X^I(t, x, y) &\rightarrow X^I(t, x, y)^*, \\ \Psi^I(t, x, y) &\rightarrow C \cdot \Psi^I(t, x, y)^*, \end{aligned} \quad (\text{E.1})$$

where T denotes the transposition of the matrix and C denotes the matrix which describes the parity transformation of three dimensional fermions. Notice that the two gauge groups are not interchanged in this C transformation. The parity symmetry P is given by

$$\begin{aligned} A_{t,x}^{(1,2)}(t, x, y) &\rightarrow -\left(A_{t,x}^{(2,1)}(t, x, -y)\right)^T, \\ A_y^{(1,2)}(t, x, y) &\rightarrow \left(A_y^{(2,1)}(t, x, -y)\right)^T, \\ X^I(t, x, y) &\rightarrow X^I(t, x, -y)^T, \\ \Psi^I(t, x, y) &\rightarrow P \cdot \Psi^I(t, x, -y)^T, \end{aligned} \quad (\text{E.2})$$

where the two gauge groups are exchanged. Finally, the time reversal symmetry T is found to be

$$\begin{aligned}
 A_t^{(1,2)}(t, x, y) &\rightarrow \left(A_t^{(2,1)}(-t, x, y) \right)^T, \\
 A_{x,y}^{(1,2)}(t, x, y) &\rightarrow - \left(A_{x,y}^{(2,1)}(-t, x, y) \right)^T, \\
 X^I(t, x, y) &\rightarrow X^I(t, x, -y)^T, \\
 \Psi^I(t, x, y) &\rightarrow T \cdot \Psi^I(t, x, -y)^T,
 \end{aligned}
 \tag{E.3}$$

where the two gauge groups are again exchanged. Remember that a standard Chern-Simons gauge theory with a single gauge group does not have either parity or time reversal symmetry [59].

For a Dirac fermion description with the signature $\eta_{\mu\nu} = (1, -1, -1)$, the gamma matrices can be defined by $\gamma_0 = \sigma_1, \gamma_1 = -i\sigma_2, \gamma_2 = i\sigma_3$ ($\sigma_{1,2,3}$ are the Pauli matrices). In this convention, the matrix C, P, T is given by $C = i\gamma_1, P = i\gamma_2$ and $T = i\gamma_1$.

References

- [1] G. 't Hooft, *Dimensional reduction in quantum gravity*, [gr-qc/9310026](#) [INSPIRE].
- [2] L. Susskind, *The World as a hologram*, *J. Math. Phys.* **36** (1995) 6377 [[hep-th/9409089](#)] [INSPIRE].
- [3] J.M. Maldacena, *The Large- N limit of superconformal field theories and supergravity*, *Adv. Theor. Math. Phys.* **2** (1998) 231 [*Int. J. Theor. Phys.* **38** (1999) 1133] [[hep-th/9711200](#)] [INSPIRE].
- [4] S. Gubser, I.R. Klebanov and A.M. Polyakov, *Gauge theory correlators from non-critical string theory*, *Phys. Lett. B* **428** (1998) 105 [[hep-th/9802109](#)] [INSPIRE].
- [5] E. Witten, *Anti-de Sitter space and holography*, *Adv. Theor. Math. Phys.* **2** (1998) 253 [[hep-th/9802150](#)] [INSPIRE].
- [6] T. Takayanagi, *Holographic Dual of BCFT*, *Phys. Rev. Lett.* **107** (2011) 101602 [[arXiv:1105.5165](#)] [INSPIRE].
- [7] A. Karch and L. Randall, *Locally localized gravity*, *JHEP* **05** (2001) 008 [[hep-th/0011156](#)] [INSPIRE].
- [8] A. Karch and L. Randall, *Open and closed string interpretation of SUSY CFT's on branes with boundaries*, *JHEP* **06** (2001) 063 [[hep-th/0105132](#)] [INSPIRE].
- [9] I. Affleck and A.W. Ludwig, *Universal noninteger 'ground state degeneracy' in critical quantum systems*, *Phys. Rev. Lett.* **67** (1991) 161 [INSPIRE].
- [10] O. Aharony, D. Marolf and M. Rangamani, *Conformal field theories in anti-de Sitter space*, *JHEP* **02** (2011) 041 [[arXiv:1011.6144](#)] [INSPIRE].
- [11] O. Aharony, L. Berdichevsky, M. Berkooz and I. Shamir, *Near-horizon solutions for $D3$ -branes ending on 5 -branes*, [arXiv:1106.1870](#) [INSPIRE].
- [12] M. Chiodaroli, E. D'Hoker, Y. Guo and M. Gutperle, *Exact half-BPS string-junction solutions in six-dimensional supergravity*, [arXiv:1107.1722](#) [INSPIRE].

- [13] L. Randall and R. Sundrum, *A Large mass hierarchy from a small extra dimension*, *Phys. Rev. Lett.* **83** (1999) 3370 [[hep-ph/9905221](#)] [[INSPIRE](#)].
- [14] G. Compere and D. Marolf, *Setting the boundary free in AdS/CFT*, *Class. Quant. Grav.* **25** (2008) 195014 [[arXiv:0805.1902](#)] [[INSPIRE](#)].
- [15] G. Gibbons and S. Hawking, *Action Integrals and Partition Functions in Quantum Gravity*, *Phys. Rev. D* **15** (1977) 2752 [[INSPIRE](#)].
- [16] J. Brown and J.W. York, *Quasilocal energy and conserved charges derived from the gravitational action*, *Phys. Rev. D* **47** (1993) 1407 [[gr-qc/9209012](#)] [[INSPIRE](#)].
- [17] J. Brown, S. Lau and J.W. York, *Action and energy of the gravitational field*, [gr-qc/0010024](#) [[INSPIRE](#)].
- [18] V. Balasubramanian and P. Kraus, *A Stress tensor for Anti-de Sitter gravity*, *Commun. Math. Phys.* **208** (1999) 413 [[hep-th/9902121](#)] [[INSPIRE](#)].
- [19] R.C. Myers, *Stress tensors and Casimir energies in the AdS/CFT correspondence*, *Phys. Rev. D* **60** (1999) 046002 [[hep-th/9903203](#)] [[INSPIRE](#)].
- [20] S. de Haro, S.N. Solodukhin and K. Skenderis, *Holographic reconstruction of space-time and renormalization in the AdS/CFT correspondence*, *Commun. Math. Phys.* **217** (2001) 595 [[hep-th/0002230](#)] [[INSPIRE](#)].
- [21] J.L. Cardy, *Conformal Invariance and Surface Critical Behavior*, *Nucl. Phys. B* **240** (1984) 514 [[INSPIRE](#)].
- [22] D. McAvity and H. Osborn, *Conformal field theories near a boundary in general dimensions*, *Nucl. Phys. B* **455** (1995) 522 [[cond-mat/9505127](#)] [[INSPIRE](#)].
- [23] O. DeWolfe, D.Z. Freedman and H. Ooguri, *Holography and defect conformal field theories*, *Phys. Rev. D* **66** (2002) 025009 [[hep-th/0111135](#)] [[INSPIRE](#)].
- [24] C. Bachas, *Asymptotic symmetries of AdS₂-branes*, [hep-th/0205115](#) [[INSPIRE](#)].
- [25] D. Bak, M. Gutperle and S. Hirano, *A Dilatonic deformation of AdS₅ and its field theory dual*, *JHEP* **05** (2003) 072 [[hep-th/0304129](#)] [[INSPIRE](#)].
- [26] A. Clark, D. Freedman, A. Karch and M. Schnabl, *The Dual of Janus (($\langle : \rangle$) \leftrightarrow ($: \rangle$)) an interface CFT*, *Phys. Rev. D* **71** (2005) 066003 [[hep-th/0407073](#)] [[INSPIRE](#)].
- [27] J.L. Cardy, *Boundary conformal field theory*, [hep-th/0411189](#) [[INSPIRE](#)].
- [28] D.E. Berenstein, R. Corrado, W. Fischler and J.M. Maldacena, *The Operator product expansion for Wilson loops and surfaces in the large N limit*, *Phys. Rev. D* **59** (1999) 105023 [[hep-th/9809188](#)] [[INSPIRE](#)].
- [29] P. Calabrese and J.L. Cardy, *Entanglement entropy and quantum field theory*, *J. Stat. Mech.* **0406** (2004) P06002 [[hep-th/0405152](#)] [[INSPIRE](#)].
- [30] S. Ryu and T. Takayanagi, *Holographic derivation of entanglement entropy from AdS/CFT*, *Phys. Rev. Lett.* **96** (2006) 181602 [[hep-th/0603001](#)] [[INSPIRE](#)].
- [31] S. Ryu and T. Takayanagi, *Aspects of Holographic Entanglement Entropy*, *JHEP* **08** (2006) 045 [[hep-th/0605073](#)] [[INSPIRE](#)].
- [32] T. Nishioka, S. Ryu and T. Takayanagi, *Holographic Entanglement Entropy: An Overview*, *J. Phys. A* **42** (2009) 504008 [[arXiv:0905.0932](#)] [[INSPIRE](#)].

- [33] T. Azeyanagi, A. Karch, T. Takayanagi and E.G. Thompson, *Holographic calculation of boundary entropy*, *JHEP* **03** (2008) 054 [[arXiv:0712.1850](#)] [[INSPIRE](#)].
- [34] M. Chiodaroli, M. Gutperle and L.-Y. Hung, *Boundary entropy of supersymmetric Janus solutions*, *JHEP* **09** (2010) 082 [[arXiv:1005.4433](#)] [[INSPIRE](#)].
- [35] M. Chiodaroli, M. Gutperle, L.-Y. Hung and D. Krym, *String Junctions and Holographic Interfaces*, *Phys. Rev. D* **83** (2011) 026003 [[arXiv:1010.2758](#)] [[INSPIRE](#)].
- [36] M. Bañados, C. Teitelboim and J. Zanelli, *The Black hole in three-dimensional space-time*, *Phys. Rev. Lett.* **69** (1992) 1849 [[hep-th/9204099](#)] [[INSPIRE](#)].
- [37] J. Brown and M. Henneaux, *Central Charges in the Canonical Realization of Asymptotic Symmetries: An Example from Three-Dimensional Gravity*, *Commun. Math. Phys.* **104** (1986) 207 [[INSPIRE](#)].
- [38] S.W. Hawking and D.N. Page, *Thermodynamics of Black Holes in anti-de Sitter Space*, *Commun. Math. Phys.* **87** (1983) 577 [[INSPIRE](#)].
- [39] E. Witten, *Anti-de Sitter space, thermal phase transition and confinement in gauge theories*, *Adv. Theor. Math. Phys.* **2** (1998) 505 [[hep-th/9803131](#)] [[INSPIRE](#)].
- [40] A. Zamolodchikov, *Irreversibility of the Flux of the Renormalization Group in a 2D Field Theory*, *JETP Lett.* **43** (1986) 730 [*Pisma Zh. Eksp. Teor. Fiz.* **43** (1986) 565] [[INSPIRE](#)].
- [41] D. Freedman, S. Gubser, K. Pilch and N. Warner, *Renormalization group flows from holography supersymmetry and a c theorem*, *Adv. Theor. Math. Phys.* **3** (1999) 363 [[hep-th/9904017](#)] [[INSPIRE](#)].
- [42] R.C. Myers and A. Sinha, *Holographic c-theorems in arbitrary dimensions*, *JHEP* **01** (2011) 125 [[arXiv:1011.5819](#)] [[INSPIRE](#)].
- [43] D. Friedan and A. Konechny, *On the boundary entropy of one-dimensional quantum systems at low temperature*, *Phys. Rev. Lett.* **93** (2004) 030402 [[hep-th/0312197](#)] [[INSPIRE](#)].
- [44] S. Yamaguchi, *Holographic RG flow on the defect and g theorem*, *JHEP* **10** (2002) 002 [[hep-th/0207171](#)] [[INSPIRE](#)].
- [45] R.C. Myers and A. Sinha, *Holographic c-theorems in arbitrary dimensions*, *JHEP* **01** (2011) 125 [[arXiv:1011.5819](#)] [[INSPIRE](#)].
- [46] J.M. Maldacena, *Eternal black holes in anti-de Sitter*, *JHEP* **04** (2003) 021 [[hep-th/0106112](#)] [[INSPIRE](#)].
- [47] V.E. Hubeny, M. Rangamani and T. Takayanagi, *A Covariant holographic entanglement entropy proposal*, *JHEP* **07** (2007) 062 [[arXiv:0705.0016](#)] [[INSPIRE](#)].
- [48] G. Galloway, K. Schleich, D. Witt and E. Woolgar, *The AdS/CFT correspondence conjecture and topological censorship*, *Phys. Lett. B* **505** (2001) 255 [[hep-th/9912119](#)] [[INSPIRE](#)].
- [49] I.R. Klebanov and E. Witten, *AdS/CFT correspondence and symmetry breaking*, *Nucl. Phys. B* **556** (1999) 89 [[hep-th/9905104](#)] [[INSPIRE](#)].
- [50] O. Aharony, O. Bergman, D.L. Jafferis and J. Maldacena, *N=6 superconformal Chern-Simons-matter theories, M2-branes and their gravity duals*, *JHEP* **10** (2008) 091 [[arXiv:0806.1218](#)] [[INSPIRE](#)].
- [51] E. Witten, *Baryons and branes in anti-de Sitter space*, *JHEP* **07** (1998) 006 [[hep-th/9805112](#)] [[INSPIRE](#)].

- [52] O. Aharony, O. Bergman and D.L. Jafferis, *Fractional M2-branes*, *JHEP* **11** (2008) 043 [[arXiv:0807.4924](#)] [[INSPIRE](#)].
- [53] D. Gao and K. Hori, *On The Structure Of The Chan-Paton Factors For D-branes In Type II Orientifolds*, [arXiv:1004.3972](#) [[INSPIRE](#)].
- [54] M. Fujita, W. Li, S. Ryu and T. Takayanagi, *Fractional Quantum Hall Effect via Holography: Chern-Simons, Edge States and Hierarchy*, *JHEP* **06** (2009) 066 [[arXiv:0901.0924](#)] [[INSPIRE](#)].
- [55] Y. Hikida, W. Li and T. Takayanagi, *ABJM with Flavors and FQHE*, *JHEP* **07** (2009) 065 [[arXiv:0903.2194](#)] [[INSPIRE](#)].
- [56] P. Hořava and E. Witten, *Heterotic and type-I string dynamics from eleven-dimensions*, *Nucl. Phys. B* **460** (1996) 506 [[hep-th/9510209](#)] [[INSPIRE](#)].
- [57] P. Hořava and E. Witten, *Eleven-dimensional supergravity on a manifold with boundary*, *Nucl. Phys. B* **475** (1996) 94 [[hep-th/9603142](#)] [[INSPIRE](#)].
- [58] S.-J. Rey, T. Suyama and S. Yamaguchi, *Wilson Loops in Superconformal Chern-Simons Theory and Fundamental Strings in Anti-de Sitter Supergravity Dual*, *JHEP* **03** (2009) 127 [[arXiv:0809.3786](#)] [[INSPIRE](#)].
- [59] G.V. Dunne, *Aspects of Chern-Simons theory*, [hep-th/9902115](#) [[INSPIRE](#)].