

SCUOLA INTERNAZIONALE SUPERIORE DI STUDI AVANZATI

SISSA Digital Library

Multilevel Analysis of Locomotion in Immature Preparations Suggests Innovative Strategies to Reactivate Stepping after Spinal Cord Injury

Original

Multilevel Analysis of Locomotion in Immature Preparations Suggests Innovative Strategies to Reactivate Stepping after Spinal Cord Injury / Brumley, Michele R.; Guertin, Pierre A.; Taccola, Giuliano. - In: CURRENT PHARMACEUTICAL DESIGN. - ISSN 1381-6128. - 23:12(2017), pp. 1764-1777. [10.2174/1381612822666161214151051]

Availability:

This version is available at: 20.500.11767/53641 since: 2017-07-06T09:47:53Z

Publisher:

Published DOI:10.2174/1381612822666161214151051

Terms of use:

Testo definito dall'ateneo relativo alle clausole di concessione d'uso

Publisher copyright

note finali coverpage

(Article begins on next page)



SCUOLA INTERNAZIONALE SUPERIORE DI STUDI AVANZATI



SISSA Digital Library

Publication title: Current Pharmaceutical Design

Citation	Brumley, M.R., Guertin, P.A., Taccola, G. Multilevel analysis of locomotion in immature preparations suggests innovative strategies to reactivate stepping after spinal cord injury (2017) Current Pharmaceutical Design, 23 (12), pp. 1764-1777.
	DOI: 10.2174/1381612822666161214151051
Link to the Publisher (es. DOI)	
Publisher	Bentham Science Publishers
Version (Postprint, Preprint)	Postprint
Terms of Use (Licence)	Embargo 12 mesi

Multilevel Analysis of Locomotion in Immature Preparations Suggests Innovative Strategies to Reactivate Stepping after Spinal Cord Injury

Michele R. Brumley¹, Pierre A. Guertin², & Giuliano Taccola³

¹ Department of Psychology, Idaho State University, Pocatello, ID, USA; ² Department of Psychiatry and Neurosciences, Faculty of Medicine, Laval University, Quebec City, QC, Canada; ³ Neuroscience Department, International School for Advanced Studies (SISSA), Trieste,

Italy

Corresponding author: Michele R. Brumley Department of Psychology Idaho State University 921 S 8th Ave, Stop 8112 Pocatello, ID 83209-8112 Phone: (208) 282-4751 Fax: (208) 282-4832 Email: brummich@isu.edu

Abstract

Locomotion is one of the most complex motor behaviors. Locomotor patterns change during early life, reflecting development of numerous peripheral and hierarchically organized central structures. Among them, the spinal cord is of particular interest since it houses the central pattern generator (CPG) for locomotion. This main command center is capable of eliciting and coordinating complex series of rhythmic neural signals sent to motoneurons and to corresponding target-muscles for basic locomotor activity. For a long-time, the CPG has been considered a black box. In recent years, complementary insights from in vitro and in vivo animal models have contributed significantly to a better understanding of its constituents, properties and ways to recover locomotion after a spinal cord injury (SCI). This review discusses key findings made by comparing the results of in vitro isolated spinal cord preparations and spinal-transected in vivo models from neonatal animals. Pharmacological, electrical, and sensory stimulation approaches largely used to further understand CPG function may also soon become therapeutic tools for potent CPG reactivation and locomotor movement induction in persons with SCI or developmental neuromuscular disorder.

Basic structure of locomotion: biomechanical basis and neural organization

Walking is a relatively stereotyped motor behavior that allows terrestrial movement of limbed animals. In these organisms, locomotion consists of the rhythmic reiteration of a basic motor scheme, the gait, which is characterized by the alternated activation of pairs of appendages and, within each limb, by the transition between the phase in which the base of support remains in contact with the ground (stance) and when it swings forwards (swing) [1]. The different number of limbs between bipeds and quadrupeds influences complexity of gait patterns. There is a large repertoire of locomotor gaits expressed by quadrupeds, mainly in relation to the speed of locomotion [2]. However, the most common type of locomotion is characterized by the double alternation between hind- and fore-limbs, and between ipsilateral limbs [3], particularly during to low-to-moderate speeds of locomotion.

Although different tetrapod species may exhibit different gaits as adults (i.e., walking, trotting, bounding, etc.), a pattern of interlimb coordination characteristic of walking (alternating homologous limbs) is shown by newborns of many species, including kittens, rabbits, rats, jerboas, gerbils, jirds, kangaroo rats, dormice, and voles [4-11]. Similarities in locomotor coordination may be due to similarities in body size and morphologies (short limbs, wide stance), and relative immaturity of the CNS (central nervous system), PNS (peripheral nervous system), and skeletal system. Specialization in locomotor patterns subsequently emerges in animals experiencing geometric and allometric growth and continued development of neural and motor systems. Even human infants, which can show a variety of crawling patterns (i.e., hands-and-knees, hands-and-feet, creeping, scooting, and mixes of these patterns), predominately exhibit an alternating interlimb pattern during crawling [12] that is kinematically similar to non-human primates and other mammals [13]. Furthermore, the development of bipedal walking locomotion in humans shows many parallels—e.g., gradual reduction of step cycle duration and variability, hyperflexion of leg joints, training effects—as with other animals [14].

In bipedal organisms, locomotion faces a further challenge. Notably, each step continuously pushes the center of mass forward and this requires a series of sophisticated systems of postural control to recover balance in response to this continuous instability [15]. Nevertheless, the imbalance following each step seems to obey to a functional significance. Indeed, the forward propulsion of the body to recover the center of mass becomes the necessary consequence to maintain the equilibrium, which is then compromised once again at the end of each step and at the beginning of the following one [16].

This rhythmic nature of locomotion requires a phasic activation of osteo-articular and muscular actuators in the periphery. As a matter of fact, muscles are recruited only in distinct phases to alternate the two limbs. Furthermore, flexor and extensor muscles around the joints of each limb are sequentially activated to allow swing and stance. Phasic muscular activation generates metabolic advantages with respect to a postural massive tonic contraction. Indeed, albeit the effort of moving the body in space, the energetic consumption during gait is quite similar to static posture [17]. The energetic cost of locomotion is concomitantly reduced also by passive elastic structures (e.g. tendons, ligaments and muscular components) that temporarily store the propulsive energy lost at one stage of the stride and return it in the following phase of gait [18]. From a kinetic point of view, limbed animals typically use gaits that are energetically favorable for body propulsion [19]. Note that further details about energetic considerations and locomotion in spinal cord-injured persons may be found in this issue in the paper by Nash and colleagues.

The most economical locomotor pattern is selected by continuously processing sensory input, including proprioceptive afferents that provide information about body mechanics [20; 21]. Nevertheless, continuous fluctuations from the nominal preferred gait naturally occur during normal walking, regardless of the increase in energy expenditure [22]. Step-by-step variability also plays an important role in optimizing locomotor performance, as it represents a robust control system that promptly adjusts the pattern in response to environmental perturbations [23]. This same logic of efficiency of movement and dynamic sensorimotor integration governs the systems responsible for generating and organizing rhythmic interlimb coordination during locomotion.

A remarkable network of spinal interneurons, mainly localized in the upper lumbar segments of the spinal cord, is responsible for producing the fundamental neural commands underlying basic locomotion [24-26]. Once activated, this network, named central pattern generator (CPG), sustains itself rather automatically [27]. This hierarchical arrangement thus reduces the need for constant supraspinal modulation of the locomotor rhythm. Supraspinal mechanisms may then be mostly limited to planning, triggering, and terminating locomotion [28; 29], modulating movement and posture in response to visual and auditory stimuli [30], and allocating neural resources toward the control of vital and cognitive functions. Although the locomotor CPG represents only a small portion of the spinal cord, a wider network of propriospinal neurons reverberates its rhythmic pattern along the whole axis [31], to integrate other rhythmic tasks, such as respiration or movement of upper limbs [32; 33].

4

During the embryonic stage of prenatal development, intrinsic rhythmogenicity of spinal networks and basic elements of the locomotor pattern (i.e. the double alternation among the two sides of the cord) are already expressed [34]. Shortly before and after birth the spinal CPG is tuned by descending fibers [35]. Hence, activity-dependent mechanisms of plasticity mediate the processing of afferent inputs and their regulation of the locomotor pattern [36]. This is especially apparent in immature animals that are undergoing continual development of muscle-skeletal actuators, overall body growth, and physiological systems. Even the earliest attempts to perform locomotion reveal a dynamic interplay between form and function. For example, rabbits [11] and some rodents [8] show quadrupedal walking during the early postnatal period, before the development of elongated hindlimbs and other forms of locomotion such as bounding and ricochetal locomotion. This suggests that although locomotor mechanisms are structurally in place and capable of functioning, that locomotor behavior is assembled in a dynamic fashion, and is dependent upon multiple factors that are necessary to physically support and move the animal's body.

Maximal efficiency in the hierarchical organization and sensorimotor integration of the neuromotor system is, in turn, supported to some extent by the redundancy of structures, which renders locomotion more resistant to occasional failures [37] and reduces vulnerability in response to peripheral or central lesions. Indeed, synergies in the activation of multiple muscles [38; 39] allow the alternated activation of limbs, even in the presence of localized muscular deficits [40; 41]. At the same time, the intrinsic variability in recruiting CPG interneurons [42] physiologically drives plastic rearrangements and neuronal compensation mechanisms, enabling gait to be expressed even after discrete neurological lesions [43]. Together, task efficiency and redundancy of structure sustain the function of the neuromotor system, even if the two principles are in contrast with one another because the maintenance of supernumerary replacing elements in case of damage requires a more consuming structure. Despite the redundancy, the compromise between these two elements reached by the neuromotor system still exposes locomotion to serious functional deficits in cases of severe impairments.

Although problems with peripheral actuators (such as osteoarticular, muscle or peripheral nerve lesions) may be allayed to some degree by using prostheses or orthoses, to permit locomotion, on the other hand, central damage is currently incurable, and may lead to paralysis. In this situation, the study of the neuronal bases of locomotion, including cellular substitution or reconfiguration of residual spinal circuits, can support targeted interventions to exploit spinal mechanisms of self-repair and plasticity. Such a multi-level understanding of CPG functions and supraspinal-peripheral system contributions to its

modulation is likely to yield the development of a multidisciplinary approach for functional recovery in persons with a spinal cord injury (SCI). Although locomotion in neurologically intact individuals involves continuous integration of neural networks throughout the CNS, including cortical, sub-cortical, cerebellar, and spinal areas, we chose to focus primarily in this review on spinal mechanisms, CPG properties, and clinically relevant research tools.

Initiation and modulation of locomotion: considerations for recovery of locomotor activity following SCI

In individuals with a complete SCI, several sensory inputs can still access and modulate the output of the spinal locomotor CPG, including afferent feedback from muscle proprioceptors, joint receptors, and cutaneous afferents. Sensory stimulation and activity-dependent feedback has been shown to facilitate locomotion for those with an incomplete SCI (e.g., [44-47]). See also the paper from Pearcey and colleagues in this issue, for further details on cutaneous contribution to locomotion. However afferent input alone is not likely to lead to a functional recovery of locomotor ability in humans with complete SCI [48]. Therefore, it is critical to consider modulatory effects on spinal networks in combination (sensory, electrical, and pharmacological approaches) for the development of therapeutic techniques, as combined methodologies may ultimately achieve greater functional outcomes through synergistic actions. Here we review the sensory afferent control, electrical initiation, and pharmacological neuromodulation of locomotor activity in spinal preparations, which together represents a promising avenue for examining the function and plasticity of spinal locomotor networks.

Sensory afferent control of locomotion

Early experiments by Graham Brown [49; 50] were critical in determining that the spinal cord contains the necessary elements to produce basic, phasic locomotor activity produced by the limbs, devoid of sensory inputs. However, since the time of these crucial studies that discovered the independence of central mechanisms from external stimuli, it has become widely recognized that sensory afferent stimulation plays an important role in modulating spinal locomotor networks and plays a key role in the recovery of locomotion for individuals with SCI [51; 52]. Sensory stimulation not only facilitates the expression of locomotion, but it permits adaptation of locomotion to the environment, regulates reflex activity, promotes transitions between different phases of the locomotor cycle, and helps to induce plasticity in the injured spinal cord.

Proprioceptive stimulation from muscle and Golgi tendon organ receptors play an important role in regulating reflex activity in the spinal cord and during locomotion. Because central excitability is typically decreased or impaired following neural damage, understanding how reflexes may alter activity and plasticity in spinal locomotor networks is essential. It is well established that hip joint afferents activate appropriate patterns of leg muscle activity during walking and are important for initiating the transition from stance to swing. This has been shown to be the case for spinal cats [53-55], human infants [56], and adult humans with SCI [57; 58]. Additionally, in spinal cats, activation of group la and group Ib afferents from ankle extensors entrains the locomotor rhythm, prolongs extensors bursts, and inhibits flexor activity [59; 60], such that a reduction in extensor muscle lb activity promotes the transition from stance to swing during locomotion [61]. Stimulation of ankle group la afferents and cutaneous nerve stimulation (delivered to the nerve innervating the plantar foot) also prolongs extensor activity. This suggests that both lb and la afferent activity continually shape amplitude and timing of the locomotor step cycle [60]. Proprioceptive feedback from the hindlimbs is also thought to be important for regulating interlimb coordination and locomotor speed adaptations, as on a treadmill, for intact as well as spinal animals [62-65].

Strong cutaneous stimulation, delivered to the perineum [65-67] or to the tail (tailpinch; [68]), can induce some locomotor stepping in spinal animals. In fact, recently it was shown that perineal stimulation alone was sufficient to induce stepping movements on a treadmill in spinal rats, thus permitting treadmill training to occur [67]. Modulation of locomotor behavior, such as altered foot contact and limb activity, occurs following mechanical stimulation of the skin on the back [69], section of cutaneous nerves innervating the foot [70], and electrical stimulation delivered to the foot dorsum [71]. Such studies indicate that cutaneous stimulation likely alters excitability of spinal circuits for locomotion and weight-supported posture, and are important for inducing plasticity following SCI (see [72] for review). In adult humans with chronic incomplete SCI, excitation of plantar cutaneous afferents modulated walking in a phase-dependent manner, suggesting interactions among locomotor mechanisms, peripheral afferents, and segmental reflex circuits [73].

Given that the spinal cord is capable of sensory-induced functional plasticity, activity-dependent mechanisms in the spinal cord are often exploited to try and rehabilitate locomotor function. For example, operant conditioning of the H-reflex modifies spinal reflex pathways in various animals [74-76], as well as improves locomotion in incomplete spinal rats [77] and chronic incomplete spinal humans [46]. Cycle training has been shown to

7

normalize spinal reflex excitability in spinal adult rats [78], as well as determine gait (an alternating or synchronous pattern) following several days of anti- or in-phase cycle training in young spinal rabbits [11]. More commonly, daily treadmill training is used to help restore locomotion in animals with SCI [62; 63; 65; 79].

It is likely that both proprioceptive and cutaneous afferents are involved in cycle and treadmill training effects. However, the necessary and sufficient mechanisms promoting activity-dependent functional plasticity in the spinal cord remain largely elusive. Possible neural mechanisms involve plastic changes (i.e., neural reconfigurations, receptor and transporter up- and down-regulation, axonal sprouting, long-term potentiation or depression, presynaptic modulation) occurring at the level of locomotor CPG, interneurons downstream from the locomotor CPG, or motoneurons. Regardless of the exact mechanisms [80], it is clear that the isolated or damaged spinal cord is capable of dynamic, sensorimotor integration that is dependent upon both endogenous and exogenous factors [81], and that understanding these mechanisms provides important opportunities for facilitating recovery and limiting further damage.

In fact, sensory afferent stimulation and use-dependent plasticity is a hallmark of physical therapy treatments. For decades now, stepping on a treadmill or use of gait orthoses has helped to restore gait in individuals with SCI (e.g., [82; 83]). For those with incomplete SCI, locomotor training improves many aspects of locomotion, including: interlimb coordination, endurance, walking speed, and limb kinematics (for review see [48]). However outcomes are typically better for individuals with an incomplete rather than complete SCI, indicating that supraspinal structures likely play a role in recovery of function for incomplete lesions. Although locomotor training typically has not resulted in recovery of walking locomotion in complete SCI individuals [57; 84], a case report of a 33-year-old man with complete SCI showed some over-ground walking function following task-specific practice coupled with robotic locomotor training as part of an intensive physical therapy program [85]. The authors asserted that intensive physical therapy and locomotor training together was likely more effective than locomotor training alone, and that training intensity, frequency, and task-specificity are likely important factors for improving motor outcomes.

Another promising application of afferent stimulation and use-dependent plasticity in promoting locomotor function can be seen with partial body-weight supported treadmill training (BWSTT) in infants that have developmental neuromuscular disorders. Parents of infants with Down Syndrome were provided small treadmills for the home and engaged their babies in treadmill-induced stepping practice 5 days a week, between the ages of 8-

8

10 months, in addition to traditional physical therapy. Although infants with Down Syndrome often start walking at 2 years of age (which is about one year later compared to typically developing infants), infants that received treadmill training learned to walk independently significantly earlier compared to infants that received physical therapy alone [86; 87]. Infants also showed improvement in other motor milestones, such as pulling to stand [86]. Similar early intervention strategies using BWSTT are currently being examined in infants that have myelomeningocele (MMC) [88]. MMC is the most severe form of spinal bifida in which the developing spine and neural tube do not close properly during prenatal development. This typically results in a small part of the lower spinal cord and meninges (forming a sac) protruding from the back of the individual, accompanied by severe motor and sensory deficits including bladder dysfunction and paralysis below the level of spinal damage, which is usually at the lumbar or sacral level. Infants with MMC start walking around 2.5-5 years of age [89], if they are able to walk at all. After 6 months of BWSTT as described above for infants with Down Syndrome, MMC infants showed earlier mean onset ages for motor items on the Bayley Scales of Infant Development, and higher bone mineral content in the legs compared to MMC infants who did not receive treadmill training [88]. Furthermore, enhancing sensory feedback via increasing overall friction on the treadmill belt increased the step rate on the treadmill for infants with MMC [90], suggesting that synergistic approaches may be more effective at triggering locomotor plasticity in the injured, developing spinal cord.

Electrostimulation facilitates locomotion

In humans, spinal locomotor circuits can be directly activated, even in the absence of any voluntary control, by relatively nonspecific stimuli such as direct electrical nonpatterned stimulation of the lumbar cord [91], continuous vibration of the quadriceps and hamstring muscle groups [92], tonic electrical stimulation of the peroneal or sural nerves [93], transdermal spinal cord stimulation (see companion paper from Minassian and colleague) or electromagnetic stimulation at the level of the lumbosacral spinal cord [94]. The automatic stepping movements generated by these approaches suggests that the CPG can function independently from brain control and thus opens the door to new paradigms for the recovery of posture and locomotion in individuals with severe SCI.

Among these methods, epidural stimulation dorsally applied over lumbosacral segments promotes reproducible locomotor patterns that can be recorded from adult spinal rats in vivo [95-97]. In humans, epidural stimulation is a minimally invasive technique that has been used for several years to alleviate spasticity and pain. Clinical use

confirmed that epidural stimulation of the thoraco-lumbar spinal cord enables bursts of electromyographic activity in lower limb muscles and few step-like alternating flexion and extension movements after complete SCI [91]. More recently, epidural stimulation associated with intense training reactivated motor functions in persons with a chronic spinal lesion [98; 99]. In these cases, electrical stimulation was not able to automatically trigger locomotion *per se*, but facilitated locomotor-like patterns evoked by afferent stimuli and reactivated voluntary commands, but only during protocol delivery [98; 99]. Likewise, transcutaneous electrical stimulation generated similar results in five SCI subjects [100]. Overall, electrical stimulation of the spinal cord enabled all of the nine subjects tested with complete paralysis to voluntarily move their lower limbs. Therefore, it now represents one of the most promising strategies to restore locomotor function following SCI.

However, the potential of electrical stimulation has not been fully disclosed yet. In fact, while research on both animals and humans has assessed the best parameters of intensity, frequency and location for stimulation, there has not been a full exploration of stimulating patterns and their motor consequences. For example, only trains of square pulses have been used [101], without varying the wave shape of single pulses. Another issue that requires further study is the combined use of electrical stimulation with pharmacology, in order to find agents more specifically targeted to enhancing locomotor CPG function. Indeed, some experimental pharmacological interventions have already been associated with potential recovery of locomotion in individuals with SCI [102]. More recently, results of a phase I/IIa trial with a first oral CPG activator called Spinalon[™] has provided evidence of safety as well as promising preliminary efficacy data (induced rhythmic EMG activity in both legs) in 45 complete SCI persons (paper from Radhakrishna and colleagues, this issue; for corresponding preclinical results in mice, see [103]). It is thus straightforward to consider the adoption of complementary and synergistic strategies as a logical direction for translating some basic biological concepts into clinical settings.

The possibility of identifying a methodology for reactivating human spinal locomotor mechanisms after SCI does not imply that spinal injured persons could easily and safely just get up and walk voluntarily. An essential component of successful over ground locomotion are the neural mechanisms for maintaining posture and recovering stability after an occasional imbalance, which are severely compromised following a spinal lesion, both in animals [104] and in humans [105]. Nevertheless, epidural electrical stimulation significantly improved posture and recovery after a loss of balance, when applied to the lumbar segments of spinal animals [106; 107] as well as to spinal cord injured persons [98;

99]. These findings suggest that electrical stimulation may therefore be a promising component to a rehabilitation strategy for recovering walking.

Pharmacological modulation of locomotion

In this section we briefly highlight findings on in vitro and in vivo animal models regarding some of the main neurotransmitters and neuromodulators that are known to stimulate and modulate synaptic spinal locomotor function. These chemical signals principally influence locomotor CPG functioning by altering motoneuron and CPG interneuron electrical properties, altering synaptic responses between motoneurons and CPG interneurons, or both. It is important to note that the effect of neuromodulators on locomotor network activity occasionally differs among species. For a more comprehensive review of pharmacological neuromodulation of locomotor networks, see Miles and Sillar [108] or Guertin [109].

The rhythmic activity produced within the spinal locomotor CPG is mainly produced by glutamate-mediated excitation and GABA- and glycine-mediated inhibition between spinal interneurons. Both ionotropic [110; 111] and metabotropic glutamate receptors [112] modulate aspects of CPG activity such as excitability, speed, and rhythmicity. Inhibitory neurotransmission regulates the left-right alternating pattern, and the speed and stability of the locomotor rhythm [113]. Renshaw cells, Ia inhibitory neurons, inhibitory commissural neurons, and several other classes of inhibitory neurons are involved [113-115].

Monoaminergic systems also play a key role in activating and modulating spinal locomotor networks. Activation of 5-HT receptors induces locomotor activity in the isolated rodent spinal cord in vitro (e.g., [116; 117]) and in spinal rodents in vivo (e.g., [118-121]). Depending on the receptor class that is activated, 5-HT receptor activation in some cases also increases the frequency and amplitude of locomotor bursts, increases the regularity of stepping, and can decrease stepping (reviewed in [108]). Activation of dopamine receptors stimulates locomotor activity in spinal rodents in vivo [122], but in the isolated spinal cord in vitro the rhythm is slower than that which is induced by 5-HT [123]. Stimulation of noradrenergic receptors induces locomotor activity in spinal cats (e.g., [124; 125]) and modulates network activity, such as increasing tonic spinal activity and locomotor bursts, in the isolated spinal cord of the neonatal rat in vitro [126; 127]. Additional modulators of spinal locomotor circuits are acetylcholine [128; 129] various neuropeptides [130; 131], and trace amines [132].

In animals with SCI, the availability of some neuromodulators to influence spinal circuits changes drastically. Acutely after SCI, glutamate and aspartate levels increase to

>400% in the spinal cord and this contributes to tissue injury [133]; their levels, and levels of GABA and glycine decrease thereafter [134]. Thus following SCI, the balance between excitation and inhibition in the spinal cord is disrupted [81]. Furthermore, the brain is the main source of monoaminergic-containing cells in the CNS, including 5-HT [135]. Following SCI, these substances no longer can be released from supraspinal projections caudal to the site of injury. Part of the consequence then is the up-regulation of 5-HT and noradrenergic receptors caudal to the lesion [136-139]. Hence, levels of endogenous neuromodulators, and likewise receptor levels, after a SCI change in relation to the time of injury, and may thereby influence responses to both drugs and sensory or electrical stimulation.

Pharmacological modulation is one way to help induce plasticity in the injured spinal cord, though combination efforts may be more fruitful than drug treatments alone. For instance, spinal adult rats treated with subthreshold doses of serotonergic agonists, provided electric epidural stimulation, and step-trained improved their hindlimb stepping coordination and muscle activation patterns within one week following SCI [97]. Comparable results without electrical stimulation may be obtained using higher doses of synergistic combinations with 5-HT agonists and NA/DA agonists or precursors in spinaltransected mice and turtles [103; 140]. As with other combinatorial approaches, the potent CPG-activating effects of suprathreshold doses of proper drug cocktails (e.g., SpinalonTM) can further improve overtime with repeated training (drug administration 3-5 times/week) [141; 142]. These findings suggest that sensory afferent feedback from step training interacts with electrical and/or pharmacological activation of spinal networks to induce neuronal plasticity changes following SCI. CPG activation through locomotor training increases the percentage of active motoneurons in the spinal cord [143]. In turn, these results suggest that afferent feedback may act on enhanced motoneuron excitability, induced by serotonergic receptors and electrical stimulation. Serotonergic stimulation also has been shown to influence spinal reflex pathways and to presynaptically influence segmental afferent projections (reviewed in [144]). Although using multiple, concurrent treatments in humans with SCI may not be the most desired approach to reinstating locomotor behavior, experimental paradigms such as those using rodent or cat models are explicating many important principles of reawakening spinal locomotor networks that are important to understand in approaching this challenge in humans. In fact, recent work with humans indicates that coupling stimulation with training promotes more adaptive plasticity and improves motor performance, and suggests that augmenting training with stimulation helps to better activate spinal circuitry [145]. Understanding and treating the

pharmacological bases of this plasticity should help to further facilitate improvements in function.

Examining spinal mechanisms of locomotion from different levels of analysis in immature preparations: novel strategies for activating locomotor stepping following SCI

Numerous experimental paradigms have been developed to study locomotion. Here we focus on the isolated spinal cord in vitro, and behavioral analysis in vivo, discussing recent insights provided by our laboratories using electrostimulation, pharmacological, and sensory feedback manipulations. Our research illustrates how an integrative approach to the study of locomotor mechanisms in immature animals reveals important dynamic interactions among levels of analysis, and strengths and limitations of specific experimental approaches. Together this work has important implications for neurorehabilitation strategies for SCI, including opening new avenues for combinatorial approaches.

Selective electrostimulation of dorsal roots triggers locomotor patterns in the isolated spinal cord

Electrostimulation through a bipolar hook electrode selectively applied to dorsal roots (DRs) cut distally from the spinal cord has been shown to evoke bouts of locomotion in spinal cats ([146]; see also Lev Tov and colleagues, this issue). A similar outcome was observed more consistently on in vitro spinal cords isolated from neonatal rats. In these preparations, electrical stimulation with stereotyped trains of square impulses triggered brief episodes of electrical oscillations, alternating between flexor and extensor motor pools on both sides of the cord (fictive locomotion rhythm, FL; [25]), when selectively delivered through tight fitting electrodes to either DRs [147] or sacrocaudal afferents [148]. In addition, activation of multiple DRs with staggered pulses [149; 150] effectively generated FL, indicating a multi-segmental convergence of afferent inputs on neuronal circuits during electrical spinal cord stimulation, as also reported in both in vivo animals [151] and in humans [100; 152].

However, still to be defined are both the neurophysiological mechanisms of electrical stimulation for triggering locomotor activity and the involved spinal wiring. Supposedly, the origin of FL episodes in response to DR stimulation may relate to the cumulative depolarization of distinct post-synaptic sites able to vary extracellular ionic concentrations [153] and facilitate the release of neurotransmitters that selectively activate network

elements crucial for generating the locomotor pattern. Indeed, selective stimulation of a subpopulation of spinal neurons is sufficient to trigger the locomotor pattern [154]. Modeling studies also have demonstrated that it is possible to effectively activate the CPG through even a few afferent projections [155]. Functional projections from the periphery to the CPG have been identified in both la afferents from muscle spindles and, mostly, in lb afferents from Golgi tendon organs [156].

A peculiarity of locomotor episodes evoked by electrical stimulation in the spinal cord in vitro is that they spontaneously decay, regardless of continuous delivery of trains, and only can be transiently rescued by varying either intensity or stimulation site. The cause of this failure is not related to impairment of action potential invasion toward afferent terminals, nor to changes in the passive properties of the motoneuron membrane [157]. On the other hand, at the presynaptic level, stimulation with trains of impulses decreases glutamate release [157], even though this effect does not seem to be linked to the disappearance of locomotor cycles [147]. Rather, progressive deterioration of FL episodes and pattern ceasing during continuous DR electrostimulation can be caused, at the postsynaptic level, by the membrane shunt determined by the depolarization that derives from increased potassium concentrations [153] and by the release of inhibitory neurotransmitters eventually reducing FL oscillations [158; 159]. High frequency stimulation may also involve receptor desensitization, since recovery (e.g. for glutamate receptors) can require up to hundreds of milliseconds [160], and depend on the quantity of the receptor agonist and the composition of the receptor subunit [161].

Overall, the spontaneous cessation of the pattern induced by afferent stimulation may be a property of the functional organization of the spinal locomotor circuit, which attributes a triggering role to afferents, with intrinsic self-limiting properties. Indeed, volleys in afferent fibers induce presynaptic inhibition on their own terminals, thus stopping excitation [162], in a manner dependent upon the frequency of incoming input [163]. Moreover, spinal interneurons that are rhythmically active during locomotor activity are modulated by the ongoing phasic rhythm and might filter sensory input out of phase with their oscillation frequency, thus stopping the pattern shortly after its onset [164]. Cellular properties, and peculiar channel expression patterns shown by crucial classes of dorsal interneurons, also may be involved in sensory motor integration and gating [165].

However, in vitro studies allow induction of FL through different experimental modalities to optimally trace the dynamics of CPG recruitment [116; 166]. A comparison between electrically- and pharmacologically-induced FL patterns indicates that neurochemicals added to the perfusion bath generate FL with a much slower onset, but

that once established remains stable for many hours. Moreover, unlike electrical stimulation, it is possible to finely modulate frequency of pharmacologically-induced FL by titrating concentrations of pharmacological agents [147]. This might imply that modulation of the locomotor pattern requires involvement of a more widespread region of the spinal cord rather than the few segments activated by electrical stimulation of a single DR [167].

Albeit variations in frequency of stereotyped trains of pulses within a relatively broad range (1-25 Hz) does not affect number (nor periodicity) of locomotor cycles [147], stimulation with trains of distinct pairs of frequencies, even simultaneously delivered to different DRs, activates longer episodes of FL [150]. This suggests that, rather than the selection of a specific frequency, optimal DR electrostimulation to evoke FL must provide a minimum level of input range variability. Several studies suggest that critical levels of variability in CPG input are required to engage neural control mechanisms, even in a highly repetitive motor task. For example, lack of variation in step trajectories interferes with the normal cycle progression that the networks execute, which can result in an inability to learn or improve the performance of motor tasks [42; 168; 169].

Innovative protocols of electrostimulation exploit the intrinsic rhythmogenic potential of spinal circuits

Locomotor-like activity in the in vitro spinal cord (Fig. 1 A) has been optimally evoked by stimulating one DR or the cauda equina with intrinsically variable asynchronous (i.e. noisy) patterns, obtained by sampling biosignals corresponding to rhythmic motor patterns in vitro or in vivo, from either a ventral root (VR, Fig. 1 B), a muscle, or a single motoneuron [131; 150; 170; 173; 174]. The clear advantage of this approach relies on stimulation strength, which, unlike canonical electrostimulation, is much lower than the minimum one required to induce a reflex response (i.e., sub-threshold). Moreover, when compared to the classic protocols of electrical stimulation [147], noisy biosignals induce locomotor-like oscillations of longer duration and with a greater number of cycles [170; 173], although the pattern still does not last throughout the protocol. The reason behind the improved efficiency of the protocols that use noisy biosignals is still under investigation. A possible explanation could rely on the presence of an intrinsic variability in amplitude and frequency of noisy protocols that accommodates the variability required by the locomotor network, mimicking the volley of physiological input that reach the spinal cord during locomotion [175]. Noise-derived high variability of the stimulus per se is not sufficient to elicit FL, as a phasic component in the lower frequencies seems also to be required, as demonstrated by the inefficacy of stimulation using either the sole Gaussian

noise (Fig. 1 C) or biosignals sampled during tonic muscle activation [174]. At the same time, FL could not even be induced by noise-free phasic input such as pure sinusoids (Fig. 1 D), or artificial noisy waveforms, software-designed by adding to a pure sinusoid either the spontaneous baseline activity at rest [170] or Gaussian noise (Fig. 1 E). These results indicate that input able to optimally trigger the CPG must contain both the low frequency component of rhythmic motor tasks and the high frequency spectral density of motor-related biosignals. As a result, effectiveness of noisy waveforms might be linked to the relative contribution of such distinct stimulus frequencies particularly efficient in activating frequency-dependent CPG elements [150]. Moreover, variability in the amplitude of the stimulating patterns might play a crucial role reminiscent of the control over sacral network output, using amplitude-modulated signals delivered to the peripheral nerve [171; 172]. The possibility to deliver these protocols at subthreshold intensity makes them an elective tool to exploit the intrinsic rhythmogenic potential of spinal circuits.

Pharmacological synergism of electrically-induced locomotor patterns

In spinal animals, superior locomotor performances so far have been found with suprathreshold doses of specific drug cocktails or with subthreshold doses of 5-HT agonists combined with electrical stimulation of the spinal cord [103; 141; 142; 176-179]. This suggests that innovative neurorehabilitation strategies to improve sensorimotor functions following neuromotor disorders could combine pharmacotherapy, training and electrical stimulation. In neonatal rat isolated spinal cords, FL was activated by the association of neurochemicals at low doses and noisy protocols at weak intensity (but not conventional trains of rectangular pulses), both unable to generate a locomotor pattern on their own. Moreover, this combination modulated cycle frequency and increased duration of FL episodes beyond the limits of electrical stimulation alone, even if delivered at optimal intensity [173]. However, these effects were not seen in the presence of a generic increase in the overall neuronal excitability of the spinal cord mediated by a shift in extracellular ionic concentrations [173], indicating that locomotor circuits, once optimally triggered by low intensity noisy patterns, can be modulated by a likewise selective (low concentration) pharmacological stimulation.

In this regard, it has been recently demonstrated that even nanomolar concentrations of the neuropeptide oxytocin, which alone is unable to elicit FL, can synergize with weak noisy stimulating protocols to elicit locomotor network activation [131]. These findings suggest that combining low doses of oxytocin with direct sub-threshold electrical stimulation helps to exploit the automatic locomotor capacities of isolated spinal circuits.

This perspective is even more interesting, in light of the ongoing clinical trials targeting safety of oxytocin for spinal cord dysfunction (http://clinicaltrials.gov).

Strengths and limitations of the in vitro approach

Newborn rat spinal cord networks are organized in a very similar way as adult networks [180], but the former ones allow advantages in terms of easier surgical isolation of the spinal cord, technical access to multiple electrophysiological recordings and electrical stimulations, as well as a longer in vitro availability compared to older tissue [181]. In addition, spinal cord isolation reduces the basic modulatory tone [182], in turn increasing consistency of motor output. As a result, it is possible to unveil even the slightest modulatory effects that could only be barely identified in vivo even using a very high number of repetitions. In general, however, the in vitro approach also allows recordings of the motor output with a pure neuronal origin, thus excluding any influence from the activation of either compensatory muscle contractions or modulators of peripheral circulation. Moreover, the clear distinction between input from DRs and motor output from VRs makes the isolated spinal cord an elective model for assessing the recruitment of locomotor networks by afferent electrical pulses. As a result, we can carefully determine the efficacy of the different protocols of stimulation, by quantifying the number of FL oscillations or by assessing the minimum duration of stimulation required to induce an episode of FL. For example, the most selective protocols available in vitro are efficient even when delivered for periods as short as 500 ms [150].

Nevertheless, the in vitro model does have a few limitations. For example, it does not allow a full analysis of motor control in terms of fine-tuning abilities, such as kinematic analysis, which is available with in vivo animal preparations. Furthermore, using in vitro preparations, we cannot identify the neuronal output that corresponds with maintenance of standing posture nor to the different coordination among muscle groups, considering the complexity of the motor behavior displayed by the behaving animal [3]. Thus for example, this does not permit confirmation of whether distinct protocols of electrical stimulation can generate different motor behaviors in vivo. For all these reasons, in order to propose innovative strategies to reactivate stepping after spinal cord damage, and to consider pediatric incidence of SCI [183], it would be profitable to adopt a multilevel analysis of locomotion in immature preparations. An extremely useful research approach could thus consider the serial application of the same experimental treatments to the same animals in each setup, to integrate initial kinematic assessments of real behavior and electrophysiological recordings of spinal network activity, after spinal cord isolation.

17

Stimulation of stepping behavior in vivo

To confirm the function of spinal circuits in vivo, behavioral paradigms in animals have been developed. The in vivo complement to the isolated spinal cord in vitro comes in the form of air-stepping. During air-stepping, animals typically are provided body-weight support by being held in a sling, with limbs unobstructed so they can move in the air (Fig. 2). Using the air-stepping paradigm, the function of locomotor circuits may be examined in immature and SCI animals that may not have the postural control or muscle strength for independent walking. To evoke air-stepping, pharmacological, sensory, or electrical stimulation is often used.

For example, when newborn rats are suspended in a sling, air-stepping may be evoked by treatment with the dopamine precursor L-DOPA [184] or the 5-HT_{2A} receptor agonist quipazine [185]. Both L-DOPA-induced and quipazine-induced air-stepping produce alternating limb kinematic patterns consistent with walking locomotion [186; 187]. A mid- or low-thoracic spinal transection eliminates L-DOPA-induced hindlimb stepping [188], however it does not eliminate quipazine-induced stepping [120; 121; 188; 189], suggesting that 5-HT receptors in the spinal cord engage spinal locomotor networks. Pharmacological stimulation of air-stepping has led to better understanding of the development [120; 190; 191], mechanisms [188; 192], function [187], and sensory modulation [121; 185] of locomotor circuits in vivo, including for animals with SCI [119; 189; 193; 194].

Sensory stimulation such as tail-pinch [68] and olfactory stimulation (bedding material; [195]), and electrical stimulation delivered by epidural [196] or intraspinal methods [197], also stimulates air-stepping. Air-stepping is not a phenomenon limited to rodents, as it has been reported in cats [198], dogs [199], monkeys [197], and human infants [201] and adults [202].

There are several advantages for using the air-stepping paradigm to examine locomotion. First, air-stepping occurs in a living, animal body that is equipped with a complex anatomy and physiology for supporting behavior. Thus compared to in vitro models, it is more behaviorally relevant. Second, because of this complex physiology, it allows examination of interactions among factors that may influence ongoing locomotor behavior, such as neurotransmitter receptor stimulation and movement-produced sensory feedback. Third, it permits investigation of locomotor activity without the need for balance control and body-weight support via reduction of external resistance. This is useful for studying developing animals that have immature postural systems and weak muscles, and humans and animals that have weakened or damaged sensorimotor systems such as with SCI. Fourth, and related to the reduction of external resistance, it allows for study of the integrity of locomotor mechanisms separate from postural mechanisms. This separation may be useful to understand in some situations where balance and posture problems may interfere with phasic limb patterning.

However, the air-stepping paradigm alone will not reveal all mechanisms involved with locomotion. Techniques at additional levels of analysis, and use of other paradigms such as the isolated spinal cord in vitro, are necessary to more precisely identify cellular properties, molecular signaling cascades, and genetic regulation of spinal locomotor networks. Further, while air-stepping resembles locomotor behavior in terms of alternating limb activity, it is still not actual locomotion. True locomotion involves integration among sensory, motor and cognitive systems and movement of the body center of mass through space. Thus air-stepping is a rather contrived experimental situation that is quite removed from the complex, dynamic interactions experienced by walking individuals. Therefore it is necessary to combine findings from behavioral experiments using the air-stepping paradigm with more reductionist, as well as more sophisticated, paradigms and preparations to more accurately depict the control and regulation of locomotion. This kind of multilevel analysis of locomotion is necessary for approaching the myriad factors that are necessary for addressing SCI.

Synergistic effects of pharmacological and sensory stimulation on locomotor behavior in developing rats in vivo

Recent research has focused on the development and regulation of locomotor behavior in the developing nervous system, using the in vivo perinatal rat as a model system. Understanding how such factors promote development and shaping of locomotor mechanisms during ontogeny has implications for facilitating recovery of function following SCI or developmental neuromuscular disorders, particularly as we now recognize that these mechanisms are activity-dependent [88].

In rats, the neural mechanisms controlling locomotion begin developing during the prenatal period [120; 203; 204], with much continued development occurring during the early postnatal period [4; 205]. During this early time in development, the spinal cord exhibits remarkable plasticity. For instance, following a spinal cord transection, immature rats recover significantly more motor function compared to older animals, mainly due to increased synaptogenesis and decreased denervation and spinal shock [206-209]. Thus

by studying locomotor function in spinal cord transected immature rats, the function of the isolated spinal cord in vivo may be evaluated at the height of spinal plasticity.

For example, in a series of studies, how newborn rats adapt their stepping behavior to a range of motion (ROM) restriction manipulation was examined. In these studies alternating air-stepping behavior was induced with the 5-HT_{2A} receptor agonist quipazine (3.0 mg/kg), and ROM restriction was imposed by placing a Plexiglas plate beneath the limbs of the rats at a distance of 50% of limb length when the limbs were fully extended. Intact postnatal day 1 (P1; 24 hr after birth) and P10 rats adapted their stepping behavior to the ROM restriction, such that they accommodated the ROM restriction task by altering intralimb coordination to apparently preserve the alternating pattern of interlimb coordination [191]. Specifically subjects made larger hindlimb step cycle excursions moving their limbs more towards the front and back of the body, rather than directly underneath the body. When subjects were administered a low-thoracic spinal cord transection on P1, such that hindlimb locomotor networks were now isolated from the rest of the CNS, hindlimb stepping behavior on P10 was abundant and intralimb adaptations to the ROM restriction also were made in these spinal subjects [121]. In fact, hindlimb stepping in spinal subjects (~450 bilateral hindlimb steps per 5 min bin) occurred approximately three times as much compared to intact subjects. This may be due in part to an up-regulation of 5-HT receptors in the caudal spinal cord following a spinal cord transection [137-139]. But in spinal subjects that received ROM restriction, frequencies of hindlimb stepping decreased to intact levels of stepping (~150 bilateral hindlimb steps per 5 min bin) during, but not after, ROM restriction (Fig. 3 A). Hence the cutaneous and proprioceptive stimulation provided by ROM restriction may have acted to specifically reduce stepping behavior or, alterations in intralimb coordination may have compromised the ability to maintain such high levels of alternating interlimb coordination in the isolated spinal cord. Intralimb adaptations to ROM restriction were much more drastic in spinal compared to sham subjects [121]. Together, these studies are suggestive of strong synergistic actions between pharmacological stimulation and sensory afferent feedback in permitting locomotor adaptations to environmental perturbations in the isolated spinal cord in vivo. To establish if 5-HT_{2A} receptor up-regulation is a mechanism of hindlimb behavioral supersensitivity producing these effects, specifically in the area of the hindlimb locomotor CPG, an investigation is underway which is examining hindlimb stepping parameters and 5-HT_{2A} receptor density in the lumbar cord, in relation to age at spinal cord transection.

Additionally, because it is becoming clear that sensory and pharmacological stimulation may often have synergistic effects on spinal function, recently the effect of

quipazine on sensory responsiveness in acute spinal transected rats was examined (unpublished data by Swann, Kauer, Allmond & Brumley). Response to tail pinch was recorded in newborn rats that were prepared by mid-thoracic spinal transection and pretreated with quipazine, and compared to controls. All subjects showed an immediate and robust motor response to tail pinch that consisted mainly of hindlimb steps (Fig. 3 B). In shams, both quipazine-treated and saline-treated subjects showed persistent effects of the tail pinch. However in spinal animals it was only quipazine-treated subjects that showed persistent effects, while saline-treated subjects did not. This study suggests that serotonergic stimulation in spinal subjects helps to recover sensory responsiveness to sham levels. However, it is important to note that quality of movement was different in spinal and sham subjects: spinal subjects including those treated with quipazine showed a higher percentage of low amplitude and smaller excursion hindlimb steps in response to tail pinch, whereas sham subjects showed a high percentage of high amplitude and large excursion steps. Thus serotonergic stimulation may help to restore excitation in the spinal cord, but not necessarily the amplitude and kinematics of leg movements.

Examination of non-neural factors in the regulation of locomotor function in spinal injured rats also has been investigated in the immature rat model in vivo. In this study, rats were treated with a thoracic hemisection on P3 and injected into the lesion site with human placental pericytes (unpublished data by Mayo, Kauer, Brumley and Bearden). Pericytes are cells of the microvascular wall that have been shown to stimulate angiogenesis in vitro [210; 211], promote functional recovery in ischemic heart repair [212], muscle regeneration following injury [213], and regulate blood-brain barrier permeability [214]. On P10, spinal injured subjects were examined for locomotor function. Pericyte treatment significantly improved hindlimb locomotor function and increased neurofilament density in both male and female rats. Additionally, placental pericytes were found in the tissue of all subjects, and migrated both rostral and caudal from the site of injury. Vessel density increased only in males. These results indicate that vascular changes within the spinal cord play a role in locomotor recovery from SCI in rats, and suggest some possible sex differences in vascular organization, function, or timing of repair in spinal tissue (unpublished data by Mayo, Kauer, Brumley and Bearden). Thus pericytes may be useful as a therapeutic cell treatment following SCI, perhaps limiting vascular dysfunction and/or playing a role in supporting neuronal reconfigurations. Intriguingly, assays with endothelial cells or spinal cord tissue culture showed faster wound healing and greater vascular density when pericytes were stimulated with CoCl₂ (to activate hypoxia-inducible pathways known for stimulating capillary growth) in vitro [211]. However when examined in spinal tissue in vivo (described above), naïve pericytes, but not pericytes stimulated with $CoCl_2$, promoted better recovery of locomotor function in SCI subjects (unpublished data by Mayo, Kauer, Brumley and Bearden). Thus results at one level of analysis may not necessarily be predicative of results at another level of analysis (i.e., cellular/in vitro \neq behavioral/in vivo; under exact conditions), though each approach can reveal important insights to inform a different level of analysis (i.e., increasing angiogenic activity in vitro and improving locomotor function in vivo; under modified conditions).

While the spinal in vivo neonatal rat preparation is more directly relevant to SCI, the intact neonatal rat offers important insights into general issues of neurobehavioral development and plasticity as well. For example, it has been shown that locomotor behavior in intact newborn rats is modulated by the substrate that the animal is stepping on [188], ROM restriction [191], treadmill speed [215], posture [216], and testing environment [187]. Thus even before the onset of independent walking and maturation of neural pathways (e.g., corticospinal tract development, myelination), it is clear that locomotor mechanisms demonstrate plasticity and are responsive to the environment. This principle is evident in developing humans as well [88]. Understanding how the development of locomotion typically occurs at multiple levels of analysis and factors that go into the shaping of locomotor circuits is crucial for developing therapies of locomotor recovery for infants and children that experience motor dysfunction due to pediatric SCI, stroke, or congenital disorders (e.g., neural tube defects such as spina bifida). For example, basic research has yielded insights of clinical significance, such as early identification and empirically-based treatments of motor dysfunction to optimize neurobehavioral outcomes in children [217]. Implementation of activity-based treatments for infants with Down syndrome and MMC were discussed earlier in this review. To further our understanding of these disorders, mechanisms affected, and treatment options, experimental paradigms with animals such as the in vivo perinatal rat is crucial as it permits testing at earlier ages, cellular and systems manipulations, and evaluation of possible treatments.

Conclusion/Perspective

The spinal locomotor system is complex and, undoubtedly, still incompletely understood. From the seminal work of Sherrington and Graham Brown a century ago, which suggested the existence of this 'black box' for locomotion in the spinal cord, up to the pivotal insights since the 1980s about cellular and pharmacological properties of the CPG gained with the development of different in vitro isolated spinal cord preparations (e.g, isolated spinal cords from lampreys, tadpoles, turtles, rats and mice both wild-type and genetically-engineered), significant advances have been made. As challenging as it is, carefully comparing data from in vitro and in vivo approaches has already begun to yield the development of promising combinatorial approaches that remain to be clinically tested. If one day, some of these CPG-activating approaches get approval by regulatory authorities, they may not cure SCI, but, combined with proper training, they may lead to significant benefits on health, as holistic approaches designed to prevent or reverse metabolic diseases, cardiovascular problems and other chronic illnesses associated generally with SCI-related physical inactivity.

Acknowledgments

GT is supported by funding from the European Union's Horizon 2020 research and innovation program under the Marie Sklodowska-Curie (grant agreement No. 661452). PAG is supported by Natural Sciences and Engineering Research Council.

Author Disclosure Statement

MRB declares having no conflict of interest. PAG declares having no conflict of interest. GT declares having no conflict of interest.



Figure 1. Innovative protocols of electrostimulation applied to a DR optimally trigger fictive locomotion patterns in the isolated spinal cord. A: The isolated spinal cord from a neonatal rat (one day post-natal) continuously perfused with physiological solution remains longlastingly viable, allowing multiple recordings and stimulations through suction glass electrodes connected to ventral and dorsal roots, respectively. **B**: A 60 s trace sampled from VRrL5 during a stable FL induced by NMDA (5 µM) + 5HT (10 µM) is exported through off line analysis to a programmable electrical stimulator, to design the protocol named FListim (Fictive Locomotion induced stimulation). FListim is delivered (6 µA, 0.3 threshold, Th, defined as the minimum intensity required to induce a reflex response using a single square pulse) to the DRIL6 of the same isolated spinal cord, now perfused in physiological solution after extensive wash out from neurochemicals. In response to stimulation, a cumulative depolarization appears superimposed by an episode of fictive locomotion (FL) pattern, consisting of 15 oscillations fully alternated among the bilateral L2 VRs (see magnification on B_1). After 30 seconds, traces repolarize to baseline, while FL cycles fade away despite continuous stimulation. C: Delivery of a trace of Gaussian noise artificially created through software failed to elicit FL, which is replaced by multiple synchronous bursts. D: A pure sinusoid of the same main frequency and amplitude of FListim induces a first cumulative depolarization that eventually ceases, while FL cycles are replaced by synchronous discharges time-locked with peaks of the stimulating waves. E: An artificial noisy waveform, constructed by adding the Gaussian noise to a pure sinusoid, does not induce any alternating cycles but only a cumulative depolarization with few synchronous cycles.



Figure 2. Air-stepping in the neonatal rat. Photograph of a 1-day-old rat showing alternating stepping behavior, following treatment with the 5-HT2A receptor agonist quipazine. The subject was secured to a horizontal bar, injected with quipazine, and recorded from a camera at a lateral angle. Behavioral testing occurred inside of an infant incubator that is temperature- and humidity-controlled.



15 30 45 Time (15-sec bins)

-15

0

60

Figure 3. Alternating hindlimb stepping in neonatal rats following pharmacological and sensory stimulation. **A**: Rats were given a low thoracic spinal transection or sham surgery on P1, and tested for quipazine-induced hindlimb stepping on P10. Following a 5-min baseline, half of the subjects experienced ROM restriction (shaded region), whereby a Plexiglas plate was placed beneath their limbs. They were also injected with 3.0 mg/kg quipazine (arrow) to induce stepping behavior. Note that spinal subjects showed significantly more hindlimb stepping across the test session, except for ROM-restricted subjects during the period of restriction (they fell to sham levels). **B**: Rats were prepared by acute mid-thoracic spinal transection and tested for sensory responsiveness to a tail pinch on P1. Ten minutes before tail pinch, subjects were pretreated with 3.0 mg/kg quipazine. Tail pinch (dashed line) was administered by gently squeezing forceps around the base of the tail. Response to tail pinch occurred immediately and persisted for about 1-min in sham subjects and spinal subjects pretreated with quipazine. Points show means; bars depict SEM.

References

- [1] Afelt Z, Kasicki S. Limb coordinations during locomotion in cats and dogs. Acta Neurobiol Exp (Wars) 1975; 35: 369-78.
- [2] Muybridge E. The science of the horse's motions. Sci Am 1878; 39: 241.
- [3] Lemieux M, Josset N, Roussel M, Couraud S, Bretzner F. Speed-dependent modulation of the locomotor behavior in adult mice reveals attractor and transitional gaits. Front Neurosci 2016; 10: 42.
- [4] Altman J, Sudarshan K. Postnatal development of locomotion in the laboratory rat. Anim Behav 1975; 23: 896-920.
- [5] Blumberg-Feldman H, Eilam D. Postnatal development of synchronous stepping in the gerbil (Gerbillus dasyurus). J Exp Biol 1995; 198: 363-372.
- [6] Butterworth BB. A comparative study of growth and development of the kangaroo rats, Dipodomys deserti Stephens and Dipodomys merriami Mearns. Growth 1961; 25: 127-38.
- [7] Chew RM, Butterworth BB. Growth and development of Merriam's kangaroo rat, Dipodomys merriami. Growth 1959; 23:75-95.
- [8] Eilam D. Postnatal development of body architecture and gait in several rodent species. J Exp Biol 1997; 200: 1339-1350.
- [9] Eilam D, Shefer G. The developmental order of bipedal locomotion in the jerboa (Jaculus orientalis): pivoting, creeping, quadrupedalism, and bipedalism. Dev Psychobiol 1997; 31: 137-142.
- [10] Peters SE. Postnatal development of gait behaviour and functional allometry in the domestic cat (Felis catus). J Zool 1983; 199: 461-486.
- [11] Viala D, Viala G, Fayein N. Plasticity of locomotor organization in infant rabbits spinalized shortly after birth. In: ME Goldberger, A Gorio A, M Murray (Eds). Development and plasticity of the mammalian spinal cord (pp 301-10). 1986. Padova, Italy: Liviana Press.
- [12] Patrick SK, Noah JA, Yang JF. Developmental constraints of quadrupedal coordination across crawling styles in human infants. J Neurophysiol 2012; 107: 3050-3061.
- [13] Righetti L, Nylen A, Rosander K, Ijspeert AJ. Kinematic and gait similarities between crawling human infants and other quadruped mammals. Front Neurol 2015; 6: 17.

- [14] Yang JF, Mitton M, Musselman KE, Patrick SK, Tajino J. Characteristics of the developing human locomotor system: similarities to other mammals. Dev Psychobiol 2015; 57: 397-408.
- [15] Musienko P, Courtine G, Tibbs JE, Kilimnik V, Savochin A, Garfinkel A, Roy RR, Edgerton VR, Gerasimenko Y. Somatosensory control of balance during locomotion in decerebrated cat. J Neurophysiol 2012; 107: 2072-2082.
- [16] Saibene F, Minetti AE. Biomechanical and physiological aspects of legged locomotion in humans. Eur J Appl Physiol 2003; 88: 297-316.
- [17] DeJaeger D, Willems PA, Heglund NC. The energy cost of walking in children. Pflugers Arch 2001; 441: 538-543.
- [18] Alexander RM. Elastic energy stores in running vertebrates. American Zoologist 1984; 24: 85–94.
- [19] Nishii J. An analytical estimation of the energy cost for legged locomotion. J Theor Biol 2006; 238: 636-645.
- [20] Dean JC. Proprioceptive feedback and preferred patterns of human movement. Exerc Sport Sci Rev 2013; 41: 36-43.
- [21] Hubbuch JE, Bennett BW, Dean JC. Proprioceptive feedback contributes to the adaptation toward an economical gait pattern. J Biomech 2015; 48:2925-2931.
- [22] O'Connor SM, Xu HZ, Kuo AD. Energetic cost of walking with increased step variability. Gait Posture 2012; 36: 102-107.
- [23] Dingwell JB, John J, Cusumano JP. Do humans optimally exploit redundancy to control step variability in walking? PLoS Comput Biol 2010; 6:e1000856.
- [24] Grillner S. Control of locomotion in bipeds, tetrapods, and fish. In: Brookhart, Mountcastle (Eds.), Handbook of Physiology. Besthesda. 1981; 1179-1236.
- [25] Guertin PA. The mammalian central pattern generator for locomotion. Brain Res Review 2009; 62: 45-56.
- [26] Kiehn O. Locomotor circuits in the mammalian spinal cord. Annu Rev Neurosci 2006; 29: 279–306.
- [27] Edgerton VR, Tillakaratne NJ, Bigbee AJ, de Leon RD, Roy RR. Plasticity of the spinal neural circuitry after injury. Annu Rev Neurosci 2004; 27: 145-167.
- [28] Takakusaki K. Neurophysiology of gait: from the spinal cord to the frontal lobe. Mov Disord 2013; 28: 1483-1491.
- [29] Grillner S, Robertson B. The basal ganglia downstream control of brainstem motor centres--an evolutionarily conserved strategy. Curr Opin Neurobiol 2015; 33: 47-52.

- [30] Grillner S, Ekeberg, El Manira A, Lansner A, Parker D, Tegnér J, Wallén P. Intrinsic function of a neuronal network - a vertebrate central pattern generator. Brain Res Brain Res Rev 1998; 26: 184-197.
- [31] Falgairolle M, Ceccato JC, Seze Md, Herbin M, Cazalets JR. Metachronal propagation of motor activity. Front Biosci (Landmark Ed) 2013; 18: 820-837.
- [32] Johnson SM, Mitchell GS. Activity-dependent plasticity of descending synaptic inputs to spinal motoneurons in an in vitro turtle brainstem-spinal cord preparation. J Neurosci 2000; 20: 3487-3495.
- [33] de Kam D, Rijken H, Manintveld T, Nienhuis B, Dietz V, Duysens J. Arm movements can increase leg muscle activity during submaximal recumbent stepping in neurologically intact individuals. J Appl Physiol (1985) 2013; 115: 34-42.
- [34] O'Donovan MJ, Wenner P, Chub N, Tabak J, Rinzel J. Mechanisms of spontaneous activity in the developing spinal cord and their relevance to locomotion. Ann N Y Acad Sci 1998; 860: 130-141.
- [35] Brocard F, Vinay L, Clarac F. Gradual development of the ventral funiculus input to lumbar motoneurons in the neonatal rat. Neuroscience 1999; 90: 1543-1554.
- [36] Chen Y, Chen L, Wang Y, Wolpaw JR, Chen XY. Operant conditioning of rat soleus H-reflex oppositely affects another H-reflex and changes locomotor kinematics. J Neurosci 2011; 31: 11370-11375.
- [37] Lafreniere-Roula M, McCrea DA. Deletions of rhythmic motoneuron activity during fictive locomotion and scratch provide clues to the organization of the mammalian central pattern generator. J Neurophysiol 2005; 94: 1120-1132.
- [38] Mussa-Ivaldi FA, Giszter SF, Bizzi E. Linear combinations of primitives in vertebrate motor control. Proc Natl Acad Sci U S A 1994; 91: 7534-7538.
- [39] Dominici N, Ivanenko YP, Cappellini G, d'Avella A, Mondì V, Cicchese M, Fabiano A, Silei T, Di Paolo A, Giannini C, Poppele RE, Lacquaniti F. Locomotor primitives in newborn babies and their development. Science 2011; 334: 997-999.
- [40] Maas H, Gregor RJ, Hodson-Tole EF, Farrell BJ, English AW, Prilutsky BI. Locomotor changes in length and EMG activity of feline medial gastrocnemius muscle following paralysis of two synergists. Exp Brain Res 2010; 203: 681-692.
- [41] Wang R, Gutierrez-Farewik EM. Compensatory strategies during walking in response to excessive muscle co-contraction at the ankle joint. Gait Posture 2014; 39: 926-932.
- [42] Ziegler MD, Zhong H, Roy RR, Edgerton VR. Why variability facilitates spinal learning. J Neurosci 2010; 30: 10720-10726.

- [43] Hubli M, Dietz V. The physiological basis of neurorehabilitation--locomotor training after spinal cord injury. J Neuroeng Rehabil 2013; 10: 5.
- [44] Dietz V, Colombo G, Jensen L. Locomotor activity in spinal man. Lancet 1994; 1260-3.
- [45] Field-Fote EC, Lindley SD, Sherman AL. Locomotor training approaches for individuals with spinal cord injury: a preliminary report of walking-related outcomes. J Neurol Phys Ther 2005; 29: 127-137.
- [46] Thompson AK, Pomerantz FR, Wolpaw JR. Operant conditioning of a spinal reflex can improve locomotion after spinal cord injury in humans. J Neurosci 2013; 33: 2365-2375.
- [47] Wernig A, Muller S, Nanassy A, Cagol E. Laufband therapy based on 'rules of spinal locomotion' is effective in spinal cord injured persons. Eur J Neurosci 1995; 7: 823-829.
- [48] Knikou M. Neural control of locomotion and training-induced plasticity after spinal and cerebral lesions. Clin Neurophysiol 2010a; 121: 1655-1668.
- [49] Brown TG. The intrinsic factors in the act of progression in the mammal. Proc R Soc Lond B Biol 1911; 84: 308-319.
- [50] Brown TG. Ton the nature of the fundamental activity of the nervous centres; together with an analysis of the conditioning rhythmic activity in progression, and a theory of evolution of function in the nervous system. J Physiol Lond 1914; 48: 18-46.
- [51] Frigon A, Rossignol S. Functional plasticity following spinal cord lesions. Prog Brain Res 2006; 231-260.
- [52] Rossignol S. Plasticity of connections underlying locomotor recovery after central and/or peripheral lesions in the adult mammals. Phil Trans R Soc B 2006; 361: 1647-1671.
- [53] Grillner S, Rossignol S. On the initiation of the swing phase of locomotion in chronic spinal cats. Brain Res 1978; 146: 269-277.
- [54] Andersson O, Grillner S. Peripheral control of the cat's step cycle. I. Phase dependent effects of ramp-movements of the hip during 'fictive locomotion'. Acta Physiol Scand 1981; 113: 89-101.
- [55] Andersson O, Grillner S. Peripheral control of the cat's step cycle. II. Entrainment of the central pattern generators for locomotion by sinusoidal hip movements during 'fictive locomotion. Acta Physiol Scand 1983; 118: 229-239.

- [56] Pang MY, Yang JF. The initiation of the swing phase in human infant stepping: importance of hip position and leg loading. J Physiol 2000; 528: 389-404.
- [57] Dobkin BH, Harkema S, Requejo R, Edgerton VR. Modulation of locomotor-like EMG activity in subjects with complete and incomplete spinal cord injury. J Neurol Rehabil 1995; 9: 183-190.
- [58] Dietz V, Müller R, Colombo G. Locomotor activity in spinal man: significance of afferent input from joint and load receptors. Brain 2002; 125: 2626-2634.
- [59] Pearson KG, Ramirez JM, Jiang W. Entrainment of the locomotor rhythm by group Ib afferents from ankle extensor muscles in spinal cats. Exp Brain Res 1992; 90: 557-566.
- [60] Guertin PA, Angel MJ, Perrault MC, McCrea DA. Ankle extensor group I afferents excite extensors throughout the hindlimb during fictive locomotion in the cat. J Physiol 1995; 487: 197-209.
- [61] Conway BA, Hultborn H, Kiehn O. Proprioceptive input resets central locomotor rhythm in the spinal cat. Exp Brain Res 1987; 68: 643-656.
- [62] Forssberg H, Grillner S, Halbertsma J, Rossignol S. The locomotion of the low spinal cat. II. Interlimb coordination. Acta Physiol Scand 1980; 108: 283-295.
- [63] Forssberg H, Grillner S, Halbertsma J. The locomotion of the low spinal cat. II.Coordination within a hindlimb. Acta Physiol Scand 1980; 108: 269-281.
- [64] Halbertsma JM. The stride cycle of the cat: the modeling of locomotion by computerized analysis of automatic recordings. Acta Physiol Scand Suppl. 1983; 521: 1-75.
- [65] Barbeau H, Rossignol S. Recovery of locomotion after chronic spinalization in the adult cat. Brain Res 1987; 412: 84-95.
- [66] Afelt Z. Reflex activity in chronic spinal cats. Acta Neurobiol Exp (Wars) 1970; 30: 129-144.
- [67] Alluin O, Delivet-Mongrain H, Rossignol S. Inducing hindlimb locomotor recovery in adult rat after complete thoracic spinal cord section using repeated treadmill training with perineal stimulation only. J Neurophysiol 2015; 114: 1931-1946.
- [68] Norreel JC, Pflieger JF, Pearlstein E, Simeoni-Alias J, Clarac F, Vinay L. Reversible disorganization of the locomotor pattern after neonatal spinal cord transection in the rat. J Neurosci 2003; 23: 1924-1932.
- [69] Hurteau MF, Thibaudier Y, Dambreville C, Desaulniers C, Frigon A. Effect of stimulating the lumbar skin caudal to a complete spinal cord injury on hindlimb locomotion. J Neurophysiol 2015; 113: 669-676.

- [70] Bouyer L, Rossignol S. Contributions of cutaneous inputs from the hindpaw to the control of locomotion. II. Spinal cats. J Neurophysiol 2003; 90: 3640-3653.
- [71] Forssberg H, Grillner S, Rossignol S. Phasic gain control of reflexes from the dorsum of the paw during spinal locomotion. Brain Res 1977; 132: 121-139.
- [72] Panek I, Bui T, Wright AT, Brownstone RM. Cutaneous afferent regulation of motor function. Acat Neurobiol Exp (Wars) 2014; 74: 158-171.
- [73] Knikou M. Plantar cutaneous afferents normalize the reflex modulation patterns during stepping in chronic human spinal cord injury. J Neurophysiol 2010b; 103: 1304-1314.
- [74] Wolpaw JR. Operant conditioning of primate spinal reflexes: the H-reflex. J Neurophysiol 1987; 57: 443-459.
- [75] Chen XY, Wolpaw. Operant conditioning of H-reflex in freely moving rats. J Neurophysiol 1995; 73: 411-415.
- [76] Carp JS, Tennissen AM, Chen XY, Wolpaw JR. H-reflex operant conditioning in mice. J Neurophysiol 2006; 96: 1718-1727.
- [77] Chen Y, Chen XY, Jakeman LB, Chen L, Stokes BT, Wolpaw JR. Operant conditioning of H-reflex can correct a locomotor abnormality after spinal cord injury in rats. J Neurosci 2006; 26: 12537-12543.
- [78] Skinner RD, Houle JD, Reese NB, Berry CL, Garcia-Rill E. Effects of exercise and fetal spinal cord implants on the H-reflex in chronically spinalized adult rats. Brain Res 1996; 729: 127-131.
- [79] Cha J, Heng C, Reinkensmeyer DJ, Roy RR, Edgerton VR, de Leon RR. Locomotor ability in spinal rats is dependent on the amount of activity imposed on the hindlimbs during treadmill training. J Neurotrauma 2007; 24: 1000-1012.
- [80] Lapointe NP, Ung RV, Guertin PA. Plasticity in sublesionally located neurons following SCI. J Neurophysiol 2007; 98: 2497-2500.
- [81] Rossignol S, Frigon A. Recovery of locomotion after spinal cord injury: some facts and mechanisms. Ann Rev Neurosci 2011; 34: 413-440.
- [82] Barbeau H, Wainberg M, Finch L. Description and application of a system for locomotor rehabilitation. Med Biol Eng Comput 1987; 25: 341-344.
- [83] Colombo G, Wirz M, Dietz V. Driven gait orthosis for improvement of locomotor training in paraplegic patients. Spinal Cord 2001; 39: 252-255.
- [84] Dietz V, Colombo G, Jensen L, Baumgartner L. Locomotor capacity of spinal cord in paraplegic patients. Ann Neurol 1995; 37: 574-582.

- [85] Manella KJ, Torres J, Field-Fote EC. Restoration of walking function in an individual with chronic complete (AIS A) spinal cord injury. J Rehab Med 2010; 42: 795-798.
- [86] Ulrich DA, Ulrich BD, Angulo-Kinzler RM, Yun J. Treadmill training of infants with Down syndrome: evidence-based developmental outcomes. Pediatrics 2001; 108: E84.
- [87] Ulrich DA, Lloyd MC, Tiernan CW, Looper JE, Angulo-Barroso RM. Effects of intensity of treadmill training on developmental outcomes and stepping in infants with Down syndrome: a randomized trial. Phys Ther 2008; 88: 114-122.
- [88] Teulier C, Lee DK, Ulrich BD. Early gait development in human infants: plasticity and clinical applications. Dev Psychobiol 2015; 57: 447-458.
- [89] Williams EN, Broughton NS, Menelaus MB. Age-related walking in children with spina bifida. Dev Med Child Neurol 1999; 41: 446-449.
- [90] Pantall A, Teulier C, Ulrich, BD. Changes in muscle activation patterns in response to enhanced sensory input during treadmill stepping in infants born with myelomeningocele. Hum Mov Sci 2012; 31: 1670-1687.
- [91] Dimitrijevic MR, Gerasimenko Y, Pinter MM. Evidence for a spinal central pattern generator in humans. Ann N Y Acad Sci 1998; 860: 360-376.
- [92] Gurfinkel VS, Levik YS, Kazennikov OV, Selionov VA. Locomotor-like movements evoked by leg muscle vibration in humans. Eur J Neurosci 1998; 10:1608-1612.
- [93] Selionov VA, Ivanenko YP, Solopova IA, Gurfinkel VS. Tonic central and sensory stimuli facilitate involuntary air-stepping in humans. J Neurophysiol 2009; 101: 2847-2858.
- [94] Gerasimenko Y, Gorodnichev R, Machueva E, Pivovarova E, Semyenov D, Savochin A, Roy RR, Edgerton VR. Novel and direct access to the human locomotor spinal circuitry. J Neurosci 2010; 30: 3700-3708.
- [95] Ichiyama RM, Gerasimenko YP, Zhong H, Roy RR, Edgerton VR. Hindlimb stepping movements in complete spinal rats induced by epidural spinal cord stimulation. Neurosci Lett 2005; 383: 339-344.
- [96] Lavrov I, Dy CJ, Fong AJ, Gerasimenko Y, Courtine G, Zhong H, Roy RR, Edgerton VR. Epidural stimulation induced modulation of spinal locomotor networks in adult spinal rats. J Neurosci 2008; 28: 6022-6029.
- [97] Courtine G, Gerasimenko Y, van den Brand R, Yew A, Musienko P, Zhong H, Song B, Ao Y, Ichiyama RM, Lavrov I, Roy RR, Sofroniew MV, Edgerton VR. Transformation of nonfunctional spinal circuits into functional states after the loss of brain input. Nat Neurosci 2009; 12: 1333-1342.

- [98] Harkema S, Gerasimenko Y, Hodes J, Burdick J, Angeli C, Chen Y, Ferreira C, Willhite A, Rejc E, Grossman RG, Edgerton VR. Effect of epidural stimulation of the lumbosacral spinal cord on voluntary movement, standing, and assisted stepping after motor complete paraplegia: a case study. Lancet 2011; 377: 1938-1947.
- [99] Angeli CA, Edgerton VR, Gerasimenko YP, Harkema SJ. Altering spinal cord excitability enables voluntary movements after chronic complete paralysis in humans. Brain 2014; 137: 1394–1409.
- [100] Gerasimenko Y, Gorodnichev R, Puhov A, Moshonkina T, Savochin A, Selionov V, Roy RR, Lu DC, Edgerton VR. Initiation and modulation of locomotor circuitry output with multisite transcutaneous electrical stimulation of the spinal cord in noninjured humans. J Neurophysiol 2015; 113: 834-42.
- [101] Gerasimenko Y, Roy RR, Edgerton VR. Epidural stimulation: comparison of the spinal circuits that generate and control locomotion in rats, cats and humans. Exp Neurol 2008; 209: 417-425.
- [102] Norman KE, Pépin A, Barbeau H. Effects of drugs on walking after spinal cord injury. Spinal Cord 1998; 36: 699-715.
- [103] Guertin PA, Ung RV, Rouleau P. Oral administration of a tri-therapy for central pattern generator activation in paraplegic mice: proof-of-concept of efficacy. Biotechnol J 2010; 5: 421-426.
- [104] Lyalka VF, Zelenin PV, Karayannidou A, Orlovsky GN, Grillner S, Deliagina TG. Impairment and recovery of postural control in rabbits with spinal cord lesions. J Neurophysiol 2005; 94: 3677-3690.
- [105] Thigpen MT, Cauraugh J, Creel G, Day K, Flynn S, Fritz S, Frost S, Respess R, Gardner-Smith P, Brack M, Behrman A. Adaptation of postural responses during different standing perturbation conditions in individuals with incomplete spinal cord injury. Gait Posture 2009; 29: 113-118.
- [106] Musienko PE, Zelenin PV, Orlovsky GN, Deliagina TG. Facilitation of postural limb reflexes with epidural stimulation in spinal rabbits. J Neurophysiol 2010; 103: 1080-1092.
- [107] Lavrov I, Gerasimenko Y, Burdick J, Zhong H, Roy RR, Edgerton VR. Integrating multiple sensory systems to modulate neural networks controlling posture. J Neurophysiol 2015; 114: 3306-1334.
- [108] Miles GB, Sillar KT. Neuromodulation of vertebrate locomotor control networks. Physiol 2011; 26: 393-411.

- [109] Guertin PA. Preclinical evidence supporting the clinical development of central pattern generator-modulating therapies for chronic spinal cord-injured patients. Front Hum Neurosci 2014; 8: 272.
- [110] Beato M, Bracci E, Nistri A. Contribution of NMDA and non-NMDA glutamate receptors to locomotor pattern generation in the neonatal rat spinal cord. Proc Biol Sci 1997; 264: 877-884.
- [111] Tapalar AE, Kiehn O. Glutamergic mechanisms for speed control and network operation in the rodent locomotor CPG. Front Neural Circuits 2010; 4: 1-14.
- [112] Nistri A, Ostroumov K, Sharifullina E, Taccola G. Tuning and playing a motor rhythm: how metabotropic glutamate receptors orchestrate generation of motor patterns in the mammalian central nervous system. J Physiol 2006; 57: 323-334.
- [113] Nishimaru H, Kakizaki M. The role of inhibitory neurotransmission in locomotor circuits of the developing mammalian spinal cord. Acta Physiol 2009; 197: 83-97.
- [114] Jankowska E. Spinal interneuronal systems: identification, multifunctional character and reconfigurations in mammals. J Physiol 2001; 533: 31-40.
- [115] Jankowska E. Spinal interneuronal networks in the cat: elementary components. Brain Res Rev 2008; 57: 46-55.
- [116] Cazalets JR, Sqalli-Houssaini Y, Clarac F. Activation of the central pattern generators for locomotion by serotonin and excitatory amino acids in neonatal rat. J Physiol 1992; 455: 187-204.
- [117] Madriaga MA, McPhee LC, Chersa T, Kristie KJ, Whelan PJ. Modulation of locomotor activity by multiple 5-HT and dopaminergic receptor subtypes in the neonatal mouse spinal cord. J Neurophysiol 2004; 92: 1566-1576.
- [118] Feraboli-Lohnherr D, Barthe JY, Orsal D. Serotonin-induced activation of the network for locomotion in adult spinal rats. J Neurosci Res 1999; 55: 87-98.
- [119] Landry ES, Guertin PA. Differential effects of 5-HT1 and 5-HT2 receptor agonists on hindlimb movements in paraplegic mice. Prog Neuropsychopharamacol Biol Psychiatry 2004; 28: 1053-1060.
- [120] Brumley MR, Robinson SR. The serotonergic agonists quipazine, CGS-12066A, and alpha-methylserotonin alter motor activity and induce hindlimb stepping in the intact and spinal rat fetus. Behav Neurosci 2005; 199: 821-833.
- [121] Strain MM, Kauer SD, Kao T, Brumley MR. Inter- and intralimb adaptations to a sensory perturbation during activation of the serotonin system after a low spinal cord transection in neonatal rats. Front Neural Circuits 2014; 80: 1-13.

- [122] Lapointe NP, Rouleau P, Ung RV, Geurtin PA. Specific role of dopamine D1 receptors in spinal network activation and rhythmic movement induction in vertebrates. J Physiol 2009; 587: 1499-1511.
- [123] Kiehn O, Kjaerulff O. Spatiotemporal characteristics of 5-HT and dopamine-induced rhythmic activity in the in vitro neonatal rat. J Neurophysiol 1996; 75: 1472-1482.
- [124] Barbeau H, Rossignol S. Initiation and modulation of the locomotor pattern in the adult chronic spinal cat by noradrenergic, serotonergic, and dopaminergic drugs. Brain Res 1991; 546: 250-260.
- [125] Kiehn O, Hultborn H, Conway BA. Spinal locomotor activity in acutely spinalized cats induced by intrathecal application of noradrenaline. Neurosci Lett 1992; 143: 243-246.
- [126] Kiehn O, Sillar KT, Kjaerulff O, McDearmid JR. Effects of noradrenaline on locomotor rhythm-generating networks in the isolated neonatal rat spinal cord. J Neurophysiol 1999; 82: 741-746.
- [127] Sqalli-Houssaini Y, Cazalets JR. Noradrenergic control of locomotor networks in the in vitro spinal cord of the neonatal rat. Brain Res 2000; 852: 100-109.
- [128] Huang A, Noga BR, Carr PA, Fedirchuk B, Jordan LM. Spinal cholinergic neurons activated during locomotion: localization and electrophysiological characterization. J Neurophysiol 2000; 83: 3537-3547.
- [129] Jordan LM, McVagh JR, Noga BR, Cabaj AM, Majczynksi H, Slawinska U, Provencher J, Leblond H, Rossignol S. Cholinergic mechanisms in spinal locomotion—potential target for rehabilitation approaches. Front Neural Circuits 2014; 8: 1-26.
- [130] Barriere G, Bertrand S, Cazalets JR. Peptidergic neuromodulatoin of the lumbar locomotor network in the neonatal rat spinal cord. Peptides 2005; 26: 277-286.
- [131] Dose F, Zanon P, Coslovich T, Taccola G. Nanomolar oxytocin synergizes with weak electrical afferent stimulation to activate the locomotor CpG of the rat spinal cord in vitro. PLoS One 2014; 9: e92967.
- [132] Gozal EA, O'Neill BE, Sawchuk MA, Zhu H, Halder M, Chou CC, Hochman S. Anatomical and functional evidence for trace amines as unique modulators of locomotor function in the mammalian spinal cord. Front Neural Circuits 2014; 134: 1-12.
- [133] Panter SS, Yum SW, Faden AI. Alteration in extracellular amino acids after traumatic spinal cord injury. Ann Neurol 1990; 27: 96-99.

- [134] Demediuk P, Daly MP, Faden AI. Effect of impact trauma on neurotransmitter and nonneurotransmitter amino acids in rat spinal cord. J Neurochem 1989; 52: 1529-1536.
- [135] Lakke EA. The projections to the spinal cord of the rat during development: a timetable of descent. Adv Anat Embryol Cell Biol 1997; 135: 1-143.
- [136] Giroux N, Rossignol S, Reader TA. Autoradiographic study of alpha1- and alpha2noradrenergic and serotonin 1A receptors in the spinal cord of normal and chronically transected cats. J Comp Neuro 1999; 406: 402-414.
- [137] Kong XY, Wienecke J, Hultborn H, Zhang M. Robust upregulation of serotonin 2A receptors after chronic spinal transection of rats: an immunohistochemical study. Brain Res 2010; 1320: 60-68.
- [138] Kong XY, Wienecke J, Chen M, Hultborn H, Zhang M. The time course of serotonin 2A receptor expression after spinal transection of rats: an immunohistochemical study. Neurosci 2011; 177: 114-126.
- [139] Ren LQ, Wienecke J, Chen M, Møller M, Hultborn H, Zhang M. The time course of serotonin 2C receptor expression after spinal transection of rats: an immunohistochemical study. Neurosci 2013; 236: 31-46.
- [140] Lapointe NP, Guertin PA. Synergistic effects of D1/5 and 5-HT1A/7 receptor agonists on locomotor movement induction in complete spinal cord-transected mice. J Neurophysiol 2008; 100: 160-168.
- [141] Guertin PA, Ung RV, Rouleau P, Steuer I. Effects on locomotion, muscle, bone, and blood induced by a combination therapy eliciting weight-bearing stepping in nonassisted spinal cord-transected mice. Neurorehabil Neural Repair 2011; 25: 234-242
- [142] Ung RV, Rouleau P, Guertin PA. Functional and physiological effects of treadmill training induced by buspirone, carbidopa, and L-DOPA in clenbuterol-treated paraplegic mice. Neurorehabil Neural Repair 2012; 26: 385-394.
- [143] Duru PO, Tillakaratne NJK, Kim, JA, Zhong H, Stauber SM, Pham TT, Xiao MS, Edgerton VR, Roy RR. Spinal neuronal activation during locomotor-like activity enabled by epidural stimulation and 5-hydroxytryptamine agonists in spinal rats. J Neurosci Res 2015; 93: 1229-1239.
- [144] Schmidt BJ, Jordan LM. The role of serotonin in reflex modulation and locomotor rhythm production in the mammalian spinal cord. Brain Res 2000; 53:689-710.
- [145] Field-Fote EC. Exciting recovery: augmenting practice with stimulation to optimize outcomes after spinal cord injury. Prog Brain Res 2015; 218: 103-126.

- [146] Barthélemy D, Leblond H, Rossignol S. Characteristics and mechanisms of locomotion induced by intraspinal microstimulation and dorsal root stimulation in spinal cats. J Neurophysiol 2007; 97: 1986-2000.
- [147] Marchetti C, Beato M, Nistri A. Alternating rhythmic activity induced by dorsal root stimulation in the neonatal rat spinal cord in vitro. J Physiol 2001a; 530: 105-112.
- [148] Etlin A, Blivis D, Ben-Zwi M, Lev-Tov A. Long and short multifunicular projections of sacral neurons are activated by sensory input to produce locomotor activity in the absence of supraspinal control. J Neurosci 2010; 30:10324-10336.
- [149] Dose F, Deumens R, Forget P, Taccola G. Staggered multi-site low-frequency electrostimulation effectively induces locomotor patterns in the isolated rat spinal cord. Spinal Cord 2016; 54: 93-101.
- [150] Dose F, Taccola G. Two Distinct Stimulus Frequencies Delivered Simultaneously at Low Intensity Generate Robust Locomotor Patterns. Neuromodulation 2016; 19: 563-575.
- [151] Shah PK, Sureddi S, Alam M, Zhong H, Roy RR, Edgerton VR, Gerasimenko Y. Unique Spatiotemporal Neuromodulation of the Lumbosacral Circuitry Shapes Locomotor Success after Spinal Cord Injury. J Neurotrauma. In press.
- [152] Sayenko DG, Atkinson DA, Floyd TC, Gorodnichev RM, Moshonkina TR, Harkema SJ, Edgerton VR, Gerasimenko YP. Effects of paired transcutaneous electrical stimulation delivered at single and dual sites over lumbosacral spinal cord. Neurosci Lett 2015; 609: 229-234.
- [153] Marchetti C, Beato M, Nistri A. Evidence for increased extracellular K(+) as an important mechanism for dorsal root induced alternating rhythmic activity in the neonatal rat spinal cord in vitro. Neurosci Lett 2001b; 304:77-80.
- [154] Hagglund M, Dougherty KJ, Borgius L, Itohara S, Iwasato T, Kiehn O. Optogenetic dissection reveals multiple rhythmogenic modules underlying locomotion. Proc Natl Acad Sci U S A 2013; 110: 11589–11594.
- [155] Bui TV, Brownstone RM. Sensory-evoked perturbations of locomotor activity by sparse sensory input: a computational study. J Neurophysiol 2015; 113: 2824-39.
- [156] Hultborn H, Conway BA, Gossard JP, Brownstone R, Fedirchuk B, Schomburg ED, Enríquez-Denton M, Perreault MC. How do we approach the locomotor network in the mammalian spinal cord? Ann N Y Acad Sci 1998; 860: 70-82.
- [157] Lev-Tov A, Pinco M. In vitro studies of prolonged synaptic depression in the neonatal rat spinal cord. J Physiol 1992; 447: 149-169.

- [158] Dale N, Gilday D. Regulation of rhythmic movements by purinergic neurotransmitters in frog embryos. Nature 1996; 383: 259-263.
- [159] Taccola G, Olivieri D, D'Angelo G, Blackburn P, Secchia L, Ballanyi K. A(1) adenosine receptor modulation of chemically and electrically evoked lumbar locomotor network activity in isolated newborn rat spinal cords. Neuroscience 2012; 222: 191-204.
- [160] Heckmann M, Dudel J. Desensitization and resensitization kinetics of glutamate receptor channels from Drosophila larval muscle. Biophys J 1997; 72: 2160-9.
- [161] Schlesinger F, Tammena D, Krampfl K, Bufler J. Desensitization and resensitization are independently regulated in human recombinant GluR subunit coassemblies. Synapse 2005; 55: 176-182.
- [162] Eccles JC, Schmidt R, Willis WD. Pharmacological studies on presynaptic inhibition. J Physiol 1963; 168: 500–530.
- [163] Clair JM, Anderson-Reid JM, Graham CM, Collins DF. Postactivation depression and recovery of reflex transmission during repetitive electrical stimulation of the human tibial nerve. J Neurophysiol 2011; 106: 184-192.
- [164] Sillar KT. Spinal pattern generation and sensory gating mechanisms. Curr Opin Neurobiol 1991; 1: 583-589.
- [165] Nanou E, Alpert MH, Alford S, El Manira A. Differential regulation of synaptic transmission by pre- and postsynaptic SK channels in the spinal locomotor network. J Neurophysiol 2013; 109: 3051-3059.
- [166] Klein DA, Patino A, Tresch MC. Flexibility of motor pattern generation across stimulation conditions by the neonatal rat spinal cord. J Neurophysiol 2010; 103: 1580-90.
- [167] Grillner S. Biological pattern generation: the cellular and computational logic of networks in motion. Neuron 2006; 52:751-766.
- [168] Shah PK, Gerasimenko Y, Shyu A, Lavrov I, Zhong H, Roy RR, Edgerton VR. Variability in step training enhances locomotor recovery after a spinal cord injury. Eur J Neurosci 2012; 36: 2054-2062.
- [169] van den Brand R, Heutschi J, Barraud Q, DiGiovanna J, Bartholdi K, Huerlimann M, Friedli L, Vollenweider I, Moraud EM, Duis S, Dominici N, Micera S, Musienko P, Courtine G. Restoring voluntary control of locomotion after paralyzing spinal cord injury. Science 2012; 336:1182-1185.

- [170] Taccola G. The locomotor central pattern generator of the rat spinal cord in vitro is optimally activated by noisy dorsal root waveforms. J Neurophysiol 2011; 106: 872-884.
- [171] Shen B, Roppolo JR, Subbaroyan J, Diubaldi A, Wahlgren S, de Groat WC, Tai C. Neuromodulation of bladder activity by stimulation of feline pudendal nerve using a transdermal amplitude modulated signal (TAMS). Neurourol Urodyn 2011; 30: 1686-1694.
- [172] Lyon TD, Ferroni MC, Kadow BT, Slater RC, Zhang Z, Chang V, Lamm V, Shen B, Wang J, Roppolo JR, de Groat WC, Tai C. Pudendal but not tibial nerve stimulation inhibits bladder contractions induced by stimulation of pontine micturition center in cats. Am J Physiol Regul Integr Comp Physiol 2016; 310: R366-74.
- [173] Dose F, Taccola G. Coapplication of noisy patterned electrical stimuli and NMDA plus serotonin facilitates fictive locomotion in the rat spinal cord. J Neurophysiol 2012; 108: 2977-2990.
- [174] Dose F, Menosso R, Taccola G. Rat locomotor spinal circuits in vitro are activated by electrical stimulation with noisy waveforms sampled from human gait. Physiol Rep 2013; 1: e00025.
- [175] Loeb GE, Bak MJ, Duysens J. Long-term unit recording from somatosensory neurons in the spinal ganglia of the freely walking cat. Science 1977; 197: 1192-1194.
- [176] Gerasimenko YP, Ichiyama RM, Lavrov IA, Courtine G, Cai L, Zhong H, Roy RR, Edgerton VR. Epidural spinal cord stimulation plus quipazine administration enable stepping in complete spinal adult rats. J Neurophysiol 2007; 98: 2525-2536.
- [177] Ichiyama RM, Gerasimenko Y, Jindrich DL, Zhong H, Roy RR, Edgerton VR. Dose dependence of the 5-HT agonist quipazine in facilitating spinal stepping in the rat with epidural stimulation. Neurosci Lett 2008; 438: 281-285.
- [178] Fong AJ, Roy RR, Ichiyama RM, Lavrov I, Courtine G, Gerasimenko Y, Tai YC, Burdick J, Edgerton VR. Recovery of control of posture and locomotion after a spinal cord injury: solutions staring us in the face. Prog Brain Res 2009; 175: 393-418.
- [179] Musienko P, van den Brand R, Märzendorfer O, Roy RR, Gerasimenko Y, Edgerton VR, Courtine G. Controlling specific locomotor behaviors through multidimensional monoaminergic modulation of spinal circuitries. J Neurosci 2011; 31: 9264-9278.
- [180] Vinay L, Brocard F, Fellippa-Marques S, Clarac F. Antidromic discharges of dorsal root afferents in the neonatal rat. J Physiol 1999; 93: 359–367.

- [181] Juvin L, Simmers J, Morin D. Locomotor rhythmogenesis in the isolated rat spinal cord: a phase-coupled set of symmetrical flexion extension oscillators. J Physiol 2007; 583: 115-128.
- [182] Schiemann J, Puggioni P, Dacre J, Pelko M, Domanski A, van Rossum MC, Duguid
 I. Cellular mechanisms underlying behavioral state-dependent bidirectional modulation of motor cortex output. Cell Rep 2015; 11: 1319-1330.
- [183] Saunders LL, Selassie A, Cao Y, Zebracki K, Vogel LC. Epidemiology of Pediatric Traumatic Spinal Cord Injury in a Population-Based Cohort, 1998-2012. Top Spinal Cord Inj Rehabil 2015; 21: 325-332.
- [184] Van Hartesveldt C, Sickles AE, Porter JD, Stehouwer DJ. L-dopa induced airstepping in developing rats. Brain Res Dev Brain Res 1991: 58; 251-255.
- [185] Brumley MR, Roberto ME, Strain, MM. Sensory feedback modulates quipazineinduced stepping behavior in the newborn rat. Behav Brain Res 2012; 229: 257-264.
- [186] Stehouwer DJ, McCrea AE, Van Hartesveldt C. L-dopa-induced air-stepping in preweanling rats. II. Kinematic analyses. Brain Res Dev Brain Res 1994; 82: 143-151.
- [187] Swann HE, Kempe RB, Van Orden AM, Brumley MR. Serotonergic activation of locomotor behavior and posture in one-day-old rats. Behav Brain Res 2016; 302: 104-114.
- [188] McEwen ML, Van Hartesveldt C, Stehouwer DJ. L-DOPA and quipazine elicit airstepping in neonatal rats with spinal cord transections. Behav Neurosci 1997; 11: 825-833.
- [189] Guertin PA. Role of NMDA receptor activation in serotonin agonist-induced airstepping in paraplegic mice. Spinal Cord 2004; 42: 185-190.
- [190] McCrea AE, Stehouwer DJ, Van Hartesveldt C. L-dopa-induced air-stepping in preweanling rats. I. Effects of dose and age. Brain Res Dev Brain Res 1994; 82: 136-142.
- [191] Strain MM, Brumley MR. Range of motion (ROM) restriction influences quipazineinduced stepping behavior in postnatal day one and ten rats. Behav Brain Res 2014; 274: 365-381.
- [192] Staup MA, Stehouwer DJ. Ontogeny of L-DOPA-induced locomotion: Expression of c-fos in the brainstem and basal ganglia of rats. Brain Res 2006; 1068: 56-64.
- [193] Guertin PA, Steuer I. Ionotropic 5-HT3 receptor agonist-induced motor responses in the hindlimbs of paraplegic mice. J Neurophysiol 2005; 94: 3397-3405.

- [194] Slawinska U, Majczynski H, Dai Y, Jordan LM. The upright posture improves plantar stepping and alters responses to serotonergic drugs in spinal rats. J Physiol 2012; 590: 1721-1736.
- [195] Fady JC, Jamon M, Clarac F. Early olfactory-induced rhythmic limb activity in the newborn rat. Brain Res Dev Brain Res 1998; 108: 111-123.
- [196] Iwahara T, Atsuta Y, Garcia-Rill E, Skinner RD. Spinal cord stimulation-induced locomotion in the adult cat. Brain Res Bull 1992; 28: 99-105.
- [197] Barthélemy D, Leblond H, Provencher J, Rossignol S. Nonlocomotor and locomotor hindlimb responses evoked by electrical microstimulation of the lumbar cord in spinalized cats. J Neurophysiol 2006; 96: 3273-3292.
- [198] AuYong N, Ollivier-Lanvin K, Lemay MA. Preferred locomotor phase of activity of lumbar interneurons during air-stepping in subchronic spinal cats. J Neurophysiol 2011; 105: 1011-1022.
- [199] Goltz F, Freusberg A. Uber die Funktionen des Lendenmarkes des Hundes. Pflugers Physiol 1874; 8: 460-498.
- [200] Vilensky JA, Wilson P, Gankiewicz E, Townsend DW. An analysis of air-stepping in normal infant vervet monkeys. J Mot Behav 1989; 21: 429-456.
- [201] Barbu-Roth M, Anderson DI, Desprès A, Streeter RJ, Cabrol D, Trujillo M, Campos JJ, Provasi J. Air stepping in response to optic flows that move toward and away from the neonate. Dev Psychobiol 2014; 56: 1142-1149.
- [202] Solopova IA, Selionov VA, Sylos-Labini F, Gurfinkel VS, Lacquaniti F, Ivanenko YP. Tapping into rhythm generation circuitry in humans during simulated weightlessness conditions. Front Syst Neurosci 2015; 9: 1-9.
- [203] Bekoff A, Lau B. Interlimb coordination in 20-day old rat fetuses. J Exp Zool 1980; 214: 173-175.
- [204] Nakayama K, Nishimaru H, Kudo N. Developmental changes in 5hydroxytryptamine-induced rhythmic activity in the spinal cord of rat fetuses in vitro. Neurosci Lett 2001; 307: 1-4.
- [205] Clarac F, Brocard F, Vinay L. The maturation of locomotor networks. Prog Brain Res 2004; 143: 57-66.
- [206] Stelzner DJ, Weber ED, BryzGornia WF. Sparing of function in developing spinal cord: Anatomical substrate. In ME Goldstein, A Gorio, M Murray (Eds). Development and plasticity of the mammalian spinal cord (vol 3, pp81-99). 1986. Padova: Liviana Press.

- [207] Tillakaratne NJ, Guu K, De Leon JJ, Bigbee RD, London AJ, Zhong NJ. Functional recovery of stepping in rats after a complete neonatal spinal transection is not due to regrowth across the lesion site. Neurosci 2010; 166: 23-33.
- [208] Weber ED, Stelzner DJ. Behavioral effects of spinal cord transection in the developing rat. Brain Res 1977; 125: 241-255.
- [209] Weber ED, Stelzner DJ. Synaptogenesis in the intermediate gray region of the lumbar spinal cord in the postnatal rat. Brain Res 1980; 185: 17-37.
- [210] Chang WG, Andrejecsk JW, Kluger MS, Saltzman WM, Pober JS. Pericytes modulate endothelial sprouting. Cardiovasc Rese 2013; 100: 492-500.
- [211] Mayo JN, Bearden SE. Driving the hypoxia-inducible pathway in human pericytes promotes vascular density in an exosome-dependent manner. Microcirculation 2015; 22: 711-723.
- [212] Chen CW, Okada M, Proto JD, Gao X, Sekiya N, Beckman SA, Corselli M, Crisan M, Saparov A, Tobita K. Péault B, Huard J. Human pericytes for ischemic heart repair. Stem Cells 2013; 31: 305-316.
- [213] Birbrair A, Zhang T, Wang ZM, Messi ML, Enikolopov GN, Mintz A, Delbono O. Role of pericytes in skeletal muscle regeneration and fat accumulation. Stem Cells Dev 2013; 22: 2298-2314.
- [214] Lai CH, Kuo KH. The critical component to establish in vitro BBB model: Pericyte. Brain Res Brain Res Rev 2005; 50: 258-265.
- [215] Brumley MR, Kauer SD, Swann HE. Developmental plasticity of coordinated action patterns in the perinatal rat. Dev Psychobiol 2015; 57: 409-420.
- [216] Mendez-Gallardo V, Roberto ME, Kauer SD, Brumley MR. Posture effects on spontaneous limb movements, alternated stepping, and the leg extension response in neonatal rats. Physiol Behav 2016; 155: 122-130.
- [217] Ulrich BD. Opportunities for early intervention based on theory, basic neuroscience, and clinical science. Phys Ther 2010; 90: 1868-1880.